

Electroweak phase transition in the reduced minimal 3-3-1 model

Vo Quoc Phong* and Vo Thanh Van†

Department of Theoretical Physics, Ho Chi Minh City University of Science, Vietnam

Hoang Ngoc Long‡

Institute of Physics, Vietnamese Academy of Science and Technology,

10 Dao Tan, Ba Dinh, Hanoi, Vietnam

'Dat pt b r
trow a p as trans t on s on s r n a wor of r u n
a o ' - tru tur oop as trans t on n t s o s v
nto two pro s +rst pro s t p as trans t on $SU' \rightarrow SU'$ at
s a an t s on on s $SU' \rightarrow U'$ w s t tan ar o
trow a p as trans t on n ass of n utra H..s boson 'h₁ sta n to
qua to t LHC va u m_{h₁} G t n t s p as trans t on s ar t +rst
or r p as trans t on s t ass of Z₂ s about . an w n t r on o
para t r spa w t t +rst or r p as trans t on at v_{ρ₀} G s a a
n to an t v pot n t a w r ass of ar H..s boson s n ran o
. < m_{h₊₊} < . r or w t t s appoa n w bosons ar t
tr.. rs of +rst or r trow a p as trans t on w t s n + an p at on s
or t y ab ty of trow a baryo n s s nar os
AC nu b rs Ex. Fr. Cq

Keywords: Spontaneous breaking of gauge symmetries, Extensions of electroweak Higgs sector, Particle-theory models (Early Universe)

* vqp on us u vn
† vtvan us u vn
‡ n on op vast a vn

I. INTRODUCTION

Electroweak phase transition (EWPT) is a type of symmetry-breaking phase transition which plays an important role at early stage of expanding universe. Particularly, the EWPT is important to explaining baryon asymmetry of our Universe. As proposed by Sakharov [1], three necessary conditions that a baryon-generating interaction in a theoretical model must satisfy to produce an excess of baryons over antibaryons are: baryon number violation, C and CP violations, and deviation from thermal equilibrium [1].

If baryon number (B) is conserved and is equal to zero, it will equal to zero forever. In contrast, B will be vanished in the state of thermal equilibrium. Therefore we need the third condition about the deviation from thermal equilibrium. The second condition is appropriate for ensuring a different decay rate for particles and antiparticles [1].

The baryon number, C and CP violations can be showed throughout the sphaleron rate and the CKM-matrix in models [2]. The sphaleron rate tells us about the baryon number violation and the non-zero phases of CKM-matrix tell us about CP violation.

It is well-known that in order to ensure sufficient the third condition, deviations from thermal equilibrium should be large enough, and therefore, the EWPT should be the first order phase transition. The EWPT is the transition between symmetric phase to asymmetric one in order to generate masses for particles. Therefore, the EWPT is related to the mass of the Higgs boson [1].

In the basic model of particles, the first and second conditions can be satisfied, but conditions on thermal imbalance is difficult to satisfy. So the analysis of the third condition is the only approach at present in order to explain the baryon asymmetry.

Why is the first order phase transition? The effective potential is a function of temperature and vacuum expectation values (VEVs). For very large temperature, it has only one minimum at the zero, and the symmetry is restored. As temperature goes to T_0 , the non-zero second minimum appears, this is the sign of symmetry breaking. When temperature at the critical temperature ($T_c < T_0$), the values of the effective potential at two minimums are equal together, the symmetry breaking is turned on. And at the critical temperature, if the two minimums are separated by a potential barrier, the phase transition will occur with bubble nucleations. Inside the bubbles, the scalar field stores a non-zero expectation value. If the bubble nucleation rate exceeds the universe's expansion rate, the bubbles collide and

eventually fill all space [1]. Such a transition is called the first order phase transition. It is very violent and one can expect large deviations from thermal equilibrium [1]. The other possible scenario takes place if the two minimums are never separated by a potential barrier. In this case, the phase transition is a smooth transition, not violent or the second order phase transition.

To study the EWPT, one considers the high-temperature effective potential as follows

$$V_{eff} = D.(T^2 - T_0^2)v^2 - E.Tv^3 + \frac{\lambda_T}{4}v^4,$$

where v is the VEV of Higgs. In order to have the first order phase transition, the strength of phase transition should be larger than the unit, i.e., $\frac{v_c}{T_c} \geq 1$.

The EWPT has been investigated in the Standard Model (SM) [1, 2] and in various extended models [3–10]. Also, a very interesting research has showed that dark matter as triggers of the electroweak phase transition [11]. For the SM, the strength of the EWPT is larger than the unit at the electroweak scale, appears too weak for the experimentally allowed mass of the SM scalar Higgs boson [1, 2]; therefore, it seems that electroweak baryogenesis requires a new physics beyond the SM at weak scale [3].

Before the neutral Higgs was found, the study of phase transitions in most of models has been focused on two basic issues: determining the order of phase transition and the neutral Higgs mass. For the SM, the first order phase transition problem has one variable which is the mass of the neutral Higgs boson. However, for the extended models, this problem has at least two variables, the first one is the Higgs mass and the rest includes the masses of heavy particles. Recently, the neutral Higgs has been discovered by the LHC [12], so the electroweak phase transition problem is reduced by one variable. This gives a lot of hope for the extended models in examining the EWPT. The remarkable successes of the last survey are:

- The SM cannot explain this phenomenon: sources of CP violation is smaller than Baryon Asymmetry of Universe (BAU) and no the first order phase transition with the large mass of the neutral Higgs or other word speaking the SM is not enough triggers for the first order phase transition to turn on [1].
- The extended models such as the two Higgs doublet model (TDHM) or MSSM, can explain the baryon asymmetry, because sources of CP violation in these models are

stronger than in the SM and they have the first order phase transition with the mass of neutral Higgs about 120 GeV. Triggers for the first order phase transition in these models are heavy bosons, or dark matter candidates [7–9, 11].

Among the extended models, the models based on $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$ gauge group (called 3-3-1 for short) [13, 15] have some intriguing features such as they can answer on the generation problem [13, 15], provide the electric charge quantization [16], etc. According to the above summarizing, we hope that the 3-3-1 models can also answer the problem of baryon asymmetry in our Universe.

The current 3-3-1 models have many different forms, but the core is based on the above mentioned gauge group. The most disadvantage of the 3-3-1 models is the complication in the Higgs sector, namely these models need at least three Higgs triplets to generate masses of fermions. Recently there are attempts to solve this problem, and some models with the simplest Higgs sector (with only two $SU(3)_L$ Higgs triplets) have been constructed.

With such a group structure, the 3-3-1 models must have at least two Higgs triplets [17, 18]. Therefore, the number of bosons in the 3-3-1 models will many more than in the SM and symmetry breaking structure is different to the SM.

In the present work, we consider the EWPT in the reduced minimal 3-3-1 (RM331) model [18] because of its simplicity. This model has the minimal leptonic content (with only the SM leptons) and the presence of bileptons: a singly and a doubly charged gauge bosons, V^\pm and $U^{\pm\pm}$, the heavy neutral boson Z_2 and the exotic quarks. This model also has two Higgs triplets. Therefore, the physical scalar spectrum of the RM331 model is composed by a doubly charged scalar h^{++} and two neutral scalars h_1 and h_2 [18]. These new particles and exotic quarks can be triggers for the first order phase transition.

The plan of the paper is as follows. In Sec.II we give a review of the RM331 about the boson, lepton and Higgs sectors. In Sec. III, we find the effective potential in the RM331 that has the contribution from heavy bosons and a contribution part like in the SM. In Sec. IV, we calculate in details structure of phase transition in the RM331, find the first order phase transition and show constraints on mass of charged Higgs boson. Finally, we summarize and make outlooks in Sec. V.

II. A REVIEW OF THE RM331 MODEL

The fermion content of the RM331 model is the same as of the minimal 3-3-1 model [13]. The difference is only in the Higgs sector.

A. Higgs potential

The Higgs potential in the RM331 [18] is given by

$$V(\chi, \rho) = \mu_1^2 \rho^\dagger \rho + \mu_2^2 \chi^\dagger \chi + \lambda_1 (\rho^\dagger \rho)^2 + \lambda_2 (\chi^\dagger \chi)^2 + \lambda_3 (\rho^\dagger \rho)(\chi^\dagger \chi) + \lambda_4 (\rho^\dagger \chi)(\chi^\dagger \rho). \quad (1)$$

The scalar sector contains only two Higgs scalar triplets [18]

$$\rho = \begin{pmatrix} \rho^+ \\ \rho^0 \\ \rho^{++} \end{pmatrix} \sim (\mathbf{3}, 1), \quad \chi = \begin{pmatrix} \chi^- \\ \chi^{--} \\ \chi^0 \end{pmatrix} \sim (\mathbf{3}, -1). \quad (2)$$

Expansion of ρ^0 and χ^0 around their VEVs is usually

$$\rho^0, \chi^0 \rightarrow \frac{1}{\sqrt{2}}(v_{\rho, \chi} + R_{\rho, \chi} + iI_{\rho, \chi}). \quad (3)$$

This potential immediately gives us two charged Goldstone bosons ρ^\pm and χ^\pm which are eaten by the gauge bosons W^\pm and V^\pm .

Let us resume content of the Higgs sector: the physical scalar spectrum of the RM331 model is composed by a doubly charged scalar h^{++} and two neutral scalars h_1 and h_2 . Since the lightest neutral field, h_1 , is basically a $SU(2)_L$ component, we identify it as the SM Higgs boson. In the effective limit $v_\chi \gg v_\rho$, the Higgs content can be summarized as follows

$$\rho = \begin{pmatrix} G_{W^+} \\ \frac{v_\rho}{\sqrt{2}} + \frac{1}{\sqrt{2}}(h_1 + iG_Z) \\ h^{++} \end{pmatrix}, \quad \chi = \begin{pmatrix} G_{V^-} \\ G_{U^{--}} \\ \frac{v_\chi}{\sqrt{2}} + \frac{1}{\sqrt{2}}(h_2 + iG_{Z'}) \end{pmatrix}, \quad (4)$$

where the Higgs masses are given by:

$$M_{h_1}^2 = \left(\lambda_1 - \frac{\lambda_3^2}{4\lambda_2} \right) v_\rho^2, \quad M_{h_2}^2 = \lambda_2 v_\chi^2 + \frac{\lambda_3^2}{4\lambda_2} v_\rho^2, \quad (5)$$

$$M_{h^{--}}^2 = \frac{\lambda_4}{2}(v_\chi^2 + v_\rho^2). \quad (6)$$

B. Gauge boson sector

The masses of gauge bosons appear in the Lagrangian part

$$\mathcal{L} = (\mathcal{D}_\mu \chi)^\dagger (\mathcal{D}^\mu \chi) + (\mathcal{D}_\mu \rho)^\dagger (\mathcal{D}^\mu \rho), \quad (7)$$

where

$$\mathcal{D}_\mu = \partial_\mu - igA_\mu^a \frac{\lambda^a}{2} - ig_X X \frac{\lambda_9}{2} B_\mu, \quad (8)$$

with $\lambda_9 = \sqrt{\frac{2}{3}} \text{diag}(1, 1, 1)$ so $\text{Tr}(\lambda_9 \lambda_9) = 2$. The couplings of $SU(3)_L$ and $U(1)_X$ satisfy the relation:

$$\frac{g_X^2}{g^2} = \frac{6s_W^2}{1 - 4s_W^2},$$

where $c_W = \cos \theta_W$, $s_W = \sin \theta_W$, $t_W = \tan \theta_W$ with θ_W is the Weinberg angle.

Substitution of the expansion in Eq. (3) into (7) leads to the following result

$$\begin{aligned} W^\pm &= \frac{A^1 \mp iA^2}{\sqrt{2}} \quad \rightarrow \quad m_{W^\pm}^2 = \frac{g^2 v_\rho^2}{4}, \\ V^\pm &= \frac{A^4 \pm iA^5}{\sqrt{2}} \quad \rightarrow \quad m_{V^\pm}^2 = \frac{g^2 v_\chi^2}{4}, \\ U^{\pm\pm} &= \frac{A^6 \pm iA^7}{\sqrt{2}} \quad \rightarrow \quad m_{U^{\pm\pm}}^2 = \frac{g^2 (v_\rho^2 + v_\chi^2)}{4} \end{aligned} \quad (9)$$

From (9), it follows that $v_\rho = 246$ GeV and the relation

$$m_U^2 - m_V^2 = m_W^2.$$

In the neutral gauge boson sector, with the basis $(A_\mu^3, A_\mu^8, B_\mu)$, mass matrix is given by

$$M^2 = \frac{g^2}{4} \begin{pmatrix} v_\rho^2 & -\frac{v_\rho^2}{\sqrt{3}} & -2\kappa v_\rho^2 \\ -\frac{v_\rho^2}{\sqrt{3}} & \frac{1}{3}(v_\rho^2 + 4v_\chi^2) & \frac{2}{\sqrt{3}}(v_\rho^2 + 2v_\chi^2) \\ -2\kappa v_\rho^2 & \frac{2}{\sqrt{3}}(v_\rho^2 + 2v_\chi^2) & 4\kappa^2(v_\rho^2 + v_\chi^2) \end{pmatrix},$$

where $\kappa = \frac{g_X}{g}$. We can easily identify the photon field A_μ as well as the massive neutral Z and Z' bosons [19]

$$\begin{aligned} A_\mu &= s_W A_\mu^3 + c_W (\sqrt{3} t_W A_\mu^8 + \sqrt{1 - 3t_W^2} B_\mu), \\ Z_\mu &= c_W A_\mu^3 - s_W (\sqrt{3} t_W A_\mu^8 + \sqrt{1 - 3t_W^2} B_\mu), \end{aligned}$$

and

$$Z'_\mu = -\sqrt{1 - 3t_W^2} A_\mu^8 + \sqrt{3} t_W B_\mu$$

where the mass-square matrix for $\{Z, Z'\}$ is given by

$$\begin{pmatrix} m_Z^2 & m_{ZZ'}^2 \\ m_{ZZ'}^2 & m_{Z'}^2 \end{pmatrix}$$

with

$$\begin{aligned} m_Z^2 &= \frac{1}{4} \frac{g^2}{\cos^2 \theta_W} v_\rho^2, \\ m_{Z'}^2 &= \frac{1}{3} g^2 \left[\frac{\cos^2 \theta_W}{1 - 4 \sin^2 \theta_W} v_\chi^2 + \frac{1 - 4 \sin^2 \theta_W}{4 \cos^2 \theta_W} v_\rho^2 \right], \\ m_{ZZ'}^2 &= \frac{1}{4\sqrt{3}} g^2 \frac{\sqrt{1 - 4 \sin^2 \theta_W}}{\cos^2 \theta_W} v_\rho^2. \end{aligned}$$

Diagonalizing the mass matrix gives the mass eigenstates Z_1 and Z_2 which can be taken as mixtures

$$\begin{aligned} Z_1 &= Z \cos \phi - Z' \sin \phi, \\ Z_2 &= Z \sin \phi + Z' \cos \phi. \end{aligned}$$

The mixing angle ϕ is given by

$$\tan 2\phi = \frac{m_Z^2 - m_{Z_1}^2}{m_{Z_2}^2 - m_Z^2},$$

where m_{Z_1} and m_{Z_2} are the *physical* mass eigenvalues

$$\begin{aligned} m_{Z_1}^2 &= \frac{1}{2} \left\{ m_{Z'}^2 + m_Z^2 - [(m_{Z'}^2 - m_Z^2)^2 - 4(m_{ZZ'}^2)^2]^{1/2} \right\}, \\ m_{Z_2}^2 &= \frac{1}{2} \left\{ m_{Z'}^2 + m_Z^2 + [(m_{Z'}^2 - m_Z^2)^2 - 4(m_{ZZ'}^2)^2]^{1/2} \right\}. \end{aligned}$$

In diagonalization, the mass of Z_1 is approximately proportional to v_ρ (because $v_\chi \gg v_\rho$), so Z_1 likes the neutral gauge boson Z in the SM. The mass of the new heavy boson Z_2 depends on v_ρ and v_χ ; in addition, $m_W = 80.39$ GeV and $v_{\rho_0} = 246$ GeV [20]. Choosing $v_{\chi_0} = 4$ TeV [14, 18], we obtain $m_V = 1307.15$ GeV and $m_U = 1309.62$ GeV. If we choose $s_W^2 = 0.23116$ [20], we derive $m_{Z_1} \approx m_Z = 91.68$ GeV and $m_{Z_2} = 4.821$ TeV. Hence we can approximate $m_{Z_2} \approx 1.2v_\chi$.

C. Fermion sector

The fermion sector in the model under consideration is the same as in the minimal 3-3-1 model [13]. The Yukawa couplings give the exotic quark masses [18]

$$L_{Yuk}^{exot} = \lambda_{11}^T \bar{Q}_{1L} \chi T_R + \lambda_{ij}^D \bar{Q}_{iL} \chi^* D_{jR} + H.c.$$

$$\begin{aligned}
&= \lambda_{11}^T (\bar{u}_{1L}\chi^- + \bar{d}_{1L}\chi^{--} + \bar{T}_L\chi^0) T_R \\
&\quad + \lambda_{ij}^D (\bar{d}_{iL}\chi^+ - \bar{u}_{iL}\chi^{++} + \bar{D}_{iL}\chi^{0*}) D_{jR} + H.c.
\end{aligned} \tag{10}$$

When the χ field develops its VEV, these couplings lead to the mass matrix in the basis (T, D_2, D_3)

$$M_J = \frac{v_\chi}{\sqrt{2}} \begin{pmatrix} \lambda_{11}^T & 0 & 0 \\ 0 & \lambda_{22}^D & \lambda_{23}^D \\ 0 & \lambda_{32}^D & \lambda_{33}^D \end{pmatrix}$$

So the exotic quarks have masses around few TeVs because their masses are proportional to v_χ . Therefore we see that masses of exotic quarks approximately equal to m_{Z_2} , but they only involve in the transition phase $SU(3) \rightarrow SU(2)$.

For usual quarks, the Yukawa couplings give the masses of them through the triplet ρ . Therefore, also like the SM, the usual quarks only involve in the transition phase $SU(2) \rightarrow U(1)$.

However, the charged lepton masses arise from the effective dimension five operator through the couplings of both χ and ρ with the following Lagrangian [18]

$$L_{Yuk}^l = \frac{\kappa_l}{\Lambda} (\bar{f}_L^c \rho^*) (\chi^\dagger f_L) + H.c. \tag{11}$$

From Lagrangian (11), we obtain $m_l = \frac{v_\chi}{\Lambda} \kappa_l v_\rho$, the coupling constant $v_\chi/\Lambda \approx 1$, so $m_l \approx \kappa_l v_\rho$. Finally the masses of charged leptons depend only on v_ρ . Therefore, they only involve in the transition phase $SU(2) \rightarrow U(1)$. Taking into account $m_e = 0.5$ MeV, $m_\mu = 105$ MeV, $m_\tau = 1.77$ GeV, ones get $k_e = 2 \times 10^{-5}$, $k_\mu = 4.3 \times 10^{-3}$ and $k_\tau = 7.2 \times 10^{-2}$.

III. EFFECTIVE POTENTIAL IN RM331

From the Higgs potential we obtain V_0 depending on VEVs as follows

$$V_0(v_\chi, v_\rho) = \mu_1^2 v_\chi^2 + \mu_2^2 v_\rho^2 + \lambda_1 v_\chi^4 + \lambda_2 v_\rho^4 + \lambda_3 v_\chi^2 v_\rho^2.$$

Here V_0 has quartic form like in the SM, but it depends on two variables v_χ and v_ρ and has the mixing between v_χ and v_ρ . However, developing the potential (1), we obtain two minimum equations. Therefore, we can transform the mixing between v_χ and v_ρ to the form that depends only on v_χ or v_ρ . Hence, we can write $V_0(v_\chi, v_\rho) = V_0(v_\chi) + V_0(v_\rho)$.

In order to derive effective potential, starting from the Higgs Lagrangian and using the principle of least action, we arrive at the equation of motion for fields. Expanding Higgs fields around VEVs and averaging over space for all fields, we obtain the one-loop effective potential (for details, see Ref.[1]).

The full Higgs Lagrangian in the RM331 model is given by

$$\mathcal{L} = (\mathcal{D}_\mu \chi)^\dagger (\mathcal{D}^\mu \chi) + (\mathcal{D}_\mu \rho)^\dagger (\mathcal{D}^\mu \rho) + V(\chi, \rho),$$

where

$$\begin{aligned} V(\chi, \rho) = & \mu_1^2 \rho^\dagger \rho + \mu_2^2 \chi^\dagger \chi + \lambda_1 (\rho^\dagger \rho)^2 + \lambda_2 (\chi^\dagger \chi)^2 \\ & + \lambda_3 (\rho^\dagger \rho)(\chi^\dagger \chi) + \lambda_4 (\rho^\dagger \chi)(\chi^\dagger \rho). \end{aligned}$$

Expanding ρ and χ around v_ρ and v_χ which are considered as variables [24], so we obtain

$$\mathcal{L} = \frac{1}{2} \partial^\mu v_\chi \partial_\mu v_\chi + \frac{1}{2} \partial^\mu v_\rho \partial_\mu v_\rho + V_0(v_\chi, v_\rho) + \sum_{boson} m_{boson}^2(v_\chi, v_\rho) W^\mu W_\mu,$$

where W runs over all gauge fields and Higgs bosons. Through the boson mass formulations as in the above sections, we can split masses of particles into two parts as follows

$$m_{boson}^2(v_\chi, v_\rho) = m_{boson}^2(v_\chi) + m_{boson}^2(v_\rho).$$

The effective potential is the function that depends on VEVs and temperature. The masses of particles depend on VEV of Higgs bosons. Therefore, when we consider the effective potential, we must consider contributions from fermions and bosons. However, for fermions, we have retained here only the top quark and exotic quarks, which dominate over the contributions from the other fermions [1]. And in the RM331 model, there are two VEVs so we have two motion equations according to v_χ and v_ρ

$$\partial^\mu v_\chi \partial_\mu v_\chi + \frac{\partial V_0(v_\chi)}{\partial v_\chi} + \sum \frac{\partial m_{bosons}^2(v_\chi)}{\partial v_\chi} W^\mu W_\mu + \sum \frac{\partial m_{exotic-quarks}(v_\chi)}{\partial v_\chi} Q \bar{Q} = 0, \quad (12)$$

$$\partial^\mu v_\rho \partial_\mu v_\rho + \frac{\partial V_0(v_\rho)}{\partial v_\rho} + \sum \frac{\partial m_{bosons}^2(v_\rho)}{\partial v_\rho} W^\mu W_\mu + \frac{\partial m_{top-quark}(v_\rho)}{\partial v_\rho} t \bar{t} = 0. \quad (13)$$

The RM331 has the gauge bosons: two massive like the SM bosons Z_1 and W^\pm and the new heavy neutral boson Z_2 , the singly and doubly charged gauge bosons $U^{\pm\pm}$ and V^\pm ; and two doubly charged Higgses h^{++} and h^{--} , one heavy neutral Higgs h_2 and one like-SM Higgs h_1 . Masses of gauge bosons and Higgses in the RM 331 model are presented in Table

TABLE I Masses of various bosons in the model

Bosons	m^2, v_χ, v_ρ	m^2, v_χ	m^2, v_ρ
$m_{W^\pm}^2$	$\frac{g^2 v_\rho^2}{4}$		$\frac{g^2 v_\rho^2}{4}$
$m_{V^\pm}^2$	$\frac{g^2 v_\chi^2}{4}$	$\frac{g^2 v_\chi^2}{4}$	
$m_{U^{\pm\pm}}^2$	$\frac{g^2 (v_\rho^2 + v_\chi^2)}{4}$	$\frac{g^2 v_\chi^2}{4}$	$\frac{g^2 v_\rho^2}{4}$
$m_{Z_1}^2 \sim m_Z^2$	$\frac{1}{4} \frac{g^2}{\cos^2 \theta_W} v_\rho^2$		$\frac{1}{4} \frac{g^2}{\cos^2 \theta_W} v_\rho^2$
$m_{Z_2}^2 \sim m_{Z'}^2$	$\frac{1}{3} g^2 \left[\frac{\cos^2 \theta_W}{1-4 \sin^2 \theta_W} v_\chi^2 + \frac{1-4 \sin^2 \theta_W}{4 \cos^2 \theta_W} v_\rho^2 \right]$	$\frac{1}{3} g^2 v_\chi^2$	$\frac{1}{3} g^2 v_\rho^2$
$m_{h_1}^2$	$\left(\lambda_1 - \frac{\lambda_3}{4\lambda_2} \right) v_\rho^2$		$\frac{\lambda_3}{4\lambda_2} v_\rho^2$
$m_{h^{--}}^2$	$\frac{\lambda_4}{2} v_\chi^2 + v_\rho^2$	$\frac{\lambda_4}{2} v_\chi^2$	$\frac{\lambda_4}{2} v_\rho^2$
$m_{h_2}^2$	$\lambda_2 v_\chi^2 + \frac{\lambda_3}{4\lambda_2} v_\rho^2$	$\lambda_2 v_\chi^2$	$\frac{\lambda_3}{4\lambda_2} v_\rho^2$

From two equations (12) and (13) and averaging over space by using Bose-Einstein and Fermi-Dirac distributions for bosons and fermions, we obtain the following effective potentials

$$\begin{aligned}
V_{eff}(v_\chi) = & V_0(v_\chi) + \frac{3}{64\pi^2} \left(m_{Z_2}^4(v_\chi) \ln \frac{m_{Z_2}^2(v_\chi)}{Q^2} - 12m_Q^4(v_\chi) \ln \frac{m_Q^2(v_\chi)}{Q^2} \right) \\
& + \frac{1}{64\pi^2} \left(m_{h_2}^4(v_\chi) \ln \frac{m_{h_2}^2(v_\chi)}{Q^2} + 2m_{h^{++}}^4(v_\chi) \ln \frac{m_{h^{++}}^2(v_\chi)}{Q^2} \right) \\
& + \frac{3}{64\pi^2} \left(2m_U^4(v_\chi) \ln \frac{m_U^2(v_\chi)}{Q^2} + 2m_V^4(v_\chi) \ln \frac{m_V^2(v_\chi)}{Q^2} \right) \\
& + \frac{T^4}{4\pi^2} \left[F_- \left(\frac{m_{h_2}(v_\chi)}{T} \right) + 2F_- \left(\frac{m_{h^{++}}(v_\chi)}{T} \right) + 12F_+ \left(\frac{m_Q(v_\chi)}{T} \right) \right] \\
& + \frac{3T^4}{4\pi^2} \left[F_- \left(\frac{m_{Z_2}(v_\chi)}{T} \right) + 2F_- \left(\frac{m_U(v_\chi)}{T} \right) + 2F_- \left(\frac{m_V(v_\chi)}{T} \right) \right],
\end{aligned}$$

and

$$\begin{aligned}
V_{eff}(v_\rho) = & V_0(v_\rho) + \frac{3}{64\pi^2} \left(m_{Z_1}^4(v_\rho) \ln \frac{m_{Z_1}^2(v_\rho)}{Q^2} + m_{Z_2}^4(v_\rho) \ln \frac{m_{Z_2}^2(v_\rho)}{Q^2} \right) \\
& + 2m_W^4(v_\rho) \ln \frac{m_W^2(v_\rho)}{Q^2} + 2m_U^4(v_\rho) \ln \frac{m_U^2(v_\rho)}{Q^2} - 4m_t^4(v_\rho) \ln \frac{m_t^2(v_\rho)}{Q^2} \\
& + \frac{1}{64\pi^2} \left(m_{h_1}^4(v_\rho) \ln \frac{m_{h_1}^2(v_\rho)}{Q^2} - m_{h_2}^4(v_\rho) \ln \frac{m_{h_2}^2(v_\rho)}{Q^2} + 2m_{h^{++}}^4 \ln \frac{m_{h^{++}}^2(v_\rho)}{Q^2} \right) \\
& + \frac{T^4}{4\pi^2} \left[F_- \left(\frac{m_{h_1}(v_\rho)}{T} \right) - F_- \left(\frac{m_{h_2}(v_\rho)}{T} \right) + 2F_- \left(\frac{m_{h^{++}}(v_\rho)}{T} \right) \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{3T^4}{4\pi^2} \left[4F_+ \left(\frac{m_t(v_\rho)}{T} \right) + F_- \left(\frac{m_{Z_1}(v_\rho)}{T} \right) + F_- \left(\frac{m_{Z_2}(v_\rho)}{T} \right) \right. \\
& \left. + 2F_- \left(\frac{m_W(v_\rho)}{T} \right) + 2F_- \left(\frac{m_U(v_\rho)}{T} \right) \right],
\end{aligned}$$

where

$$\begin{aligned}
F_\mp \left(\frac{m_\phi}{T} \right) &= \int_0^{\frac{m_\phi}{T}} \alpha J_\mp^{(1)}(\alpha, 0) d\alpha, \\
J_\mp^{(1)}(\alpha, 0) &= 2 \int_\alpha^\infty \frac{(x^2 - \alpha^2)^{1/2}}{e^x \mp 1} dx.
\end{aligned}$$

The total effective potential in the RM331 model can be rewritten as follows

$$V_{eff}^{RM331} = V_{eff}(v_\chi) + V_{eff}(v_\rho).$$

IV. ELECTROWEAK PHASE TRANSITION

The symmetry breaking in the RM331 can take place sequentially: due to two scales of symmetry breaking which are very different, $v_{\chi_0} \gg v_{\rho_0}$ ($v_{\chi_0} \sim 4 - 5$ TeV [14, 18], $v_{\rho_0} = 246$ GeV) and because of the accelerating universe. The symmetry breaking $SU(3) \rightarrow SU(2)$ takes place before the symmetry breaking $SU(2) \rightarrow U(1)$.

The symmetry breaking $SU(3) \rightarrow SU(2)$ through χ_0 , generates masses of the heavy gauge bosons such as $U^{\pm\pm}$, V^\pm , Z_2 , and the exotic quarks. Therefore, the phase transition $SU(3) \rightarrow SU(2)$ only depends on v_χ . When our universe has been expanding and cooling due to v_{ρ_0} scale, the symmetry breaking or phase transition $SU(2) \rightarrow U(1)$ is turned on through ρ_0 , which generates masses of the SM particles and the last part of masses of $U^{\pm\pm}$. Therefore phase transition $SU(2) \rightarrow U(1)$ only depends on v_ρ . For the current universe having the baryon asymmetry, this asymmetry must exist from the initial conditions of the universe. If the early universe has not the baryon asymmetry, the current universe will not have it [1]. In other words, this asymmetry exists throughout periods of the universe to date. In order to describe baryogenesis, models must satisfy three conditions given by Sakharov. Especially in the third condition, the spontaneous symmetry breaking must be associated with the first order phase transition. In the RM331 model, the spontaneous symmetry breaking takes place in two different energy scales or other word speaking the electroweak phase transition is the combination of two different phase transitions. Therefore, if the RM331 model well describes this phenomenon, both two phase transitions are the first order phase transition.

In contrast, if one of two phase transitions is not the first order phase transition, the RM331 model will not fully describe this asymmetry, since this model does not ensure the continuity of baryogenesis in the universe.

Through the boson mass formulations in the above sections, we see that boson V^\pm only involves in the first phase transition $SU(3) \rightarrow SU(2)$. The gauge bosons Z_1 , W^\pm and h_1 only involve in the second phase transition $SU(2) \rightarrow U(1)$. However, $U^{\pm\pm}$, Z_2 and h^{--} involve in both two phase transitions.

With that structure of phase transition, we see that the mass of $U^{\pm\pm}$, is generated by both two phase transitions. When the universe at v_{χ_0} scale, the symmetry breaking $SU(3) \rightarrow SU(2)$ generates masses for the exotic quarks and a part of $U^{\pm\pm}$, or other word speaking, it eaten one Goldstone boson $\chi^{\pm\pm}$ of triplet χ . When the universe cool to v_{ρ_0} scale, the symmetry breaking $SU(2) \rightarrow U(1)$ is turned on, it generates masses for the SM particles and the last part of $U^{\pm\pm}$, or $U^{\pm\pm}$ eaten the other Goldstone boson $\rho^{\pm\pm}$ of triplet ρ .

A. Phase transition $SU(3) \rightarrow SU(2)$

This phase transition involves exotic quarks, heavy bosons, without involvement of the SM particles, the effective potential of the EWPT $SU(3) \rightarrow SU(2)$ is $V_{eff}(v_\chi)$.

$$\begin{aligned}
V_{eff}(v_\chi) = & V_0(v_\chi) + \frac{3}{64\pi^2} \left(m_{Z_2}^4(v_\chi) \ln \frac{m_{Z_2}^2(v_\chi)}{Q^2} - 12m_Q^4(v_\chi) \ln \frac{m_Q^2(v_\chi)}{Q^2} \right) \\
& + \frac{1}{64\pi^2} \left(m_{h_2}^4(v_\chi) \ln \frac{m_{h_2}^2(v_\chi)}{Q^2} + 2m_{h^{++}}^4(v_\chi) \ln \frac{m_{h^{++}}^2(v_\chi)}{Q^2} \right) \\
& + \frac{3}{64\pi^2} \left(2m_U^4(v_\chi) \ln \frac{m_U^2(v_\chi)}{Q^2} + 2m_V^4(v_\chi) \ln \frac{m_V^2(v_\chi)}{Q^2} \right) \\
& + \frac{T^4}{4\pi^2} \left[F_- \left(\frac{m_{h_2}(v_\chi)}{T} \right) + 2F_- \left(\frac{m_{h^{++}}(v_\chi)}{T} \right) + 12F_+ \left(\frac{m_Q(v_\chi)}{T} \right) \right] \\
& + \frac{3T^4}{4\pi^2} \left[F_- \left(\frac{m_{Z_2}(v_\chi)}{T} \right) + 2F_- \left(\frac{m_U(v_\chi)}{T} \right) + 2F_- \left(\frac{m_V(v_\chi)}{T} \right) \right].
\end{aligned}$$

The symmetry breaking scale is v_{χ_0} chosen to be 4 TeV [14, 18] and the masses of three exotic quarks are m_Q . Therefore, the effective potential can be rewritten as follows

$$V_{SU(3) \rightarrow SU(2)}^{eff} = D'(T^2 - T_0'^2)v_\chi^2 - E'Tv_\chi^3 + \frac{\lambda'_T}{4}v_\chi^4$$

The minimum conditions are

$$V_{eff}(v_{\chi_0}) = 0; \quad \frac{\partial V_{eff}(v_{\chi})}{\partial v_{\chi}}(v_{\chi_0}) = 0; \quad \frac{\partial^2 V_{eff}(v_{\chi})}{\partial v_{\chi}^2}(v_{\chi_0}) = m_{h_2}^2(v_{\chi})|_{v_{\chi}=v_{\chi_0}},$$

where

$$\begin{aligned} D' &= \frac{1}{24v_{\chi_0}^2} \left\{ 6m_U^2(v_{\chi}) + 3m_{Z_2}^2(v_{\chi}) + 6m_V^2(v_{\chi}) + 18m_Q^2(v_{\chi}) + 2m_{h_2}^2(v_{\chi}) + 2m_{h^\pm}^2(v_{\chi}) \right\}, \\ T_0'^2 &= \frac{1}{D} \left\{ \frac{1}{4}m_{h_2}^2(v_{\chi}) - \frac{1}{32\pi^2 v_{\chi_0}^2} \left(6m_U^4(v_{\chi}) + 3m_{Z_2}^4(v_{\chi}) + 6m_V^4(v_{\chi}) - 36m_Q^4(v_{\chi}) \right. \right. \\ &\quad \left. \left. + m_{h_2}^4(v_{\chi}) + 2m_{h^\pm}^4(v_{\chi}) \right) \right\}, \\ E' &= \frac{1}{12\pi v_{\chi_0}^3} (6m_U^3(v_{\chi}) + 3m_{Z_2}^3(v_{\chi}) + 6m_V^3(v_{\chi}) + m_{h_2}^3(v_{\chi}) + 2m_{h^\pm}^3(v_{\chi})), \\ \lambda_T' &= \frac{m_{h_2}^2(v_{\chi})}{2v_{\chi_0}^2} \left\{ 1 - \frac{1}{8\pi^2 v_{\chi_0}^2 m_{h_2}^2(v_{\chi})} \left[6m_V^4(v_{\chi}) \ln \frac{m_V^2(v_{\chi})}{bT^2} + 3m_{Z_2}^4(v_{\chi}) \ln \frac{m_{Z_2}^2(v_{\chi})}{bT^2} \right. \right. \\ &\quad \left. \left. + 6m_U^4(v_{\chi}) \ln \frac{m_U^2(v_{\chi})}{bT^2} - 36m_Q^4(v_{\chi}) \ln \frac{m_Q^2(v_{\chi})}{b_F T^2} + m_{h_2}^4(v_{\chi}) \ln \frac{m_{h_2}^2(v_{\chi})}{bT^2} \right. \right. \\ &\quad \left. \left. + 2m_{h^\pm}^4(v_{\chi}) \ln \frac{m_{h^\pm}^2(v_{\chi})}{bT^2} \right] \right\}. \end{aligned}$$

The values of $V_{eff}(v_{\chi})$ at the two minima become equal at the critical temperature

$$T_c' = \frac{T_0'}{\sqrt{1 - E'^2/D'\lambda_{T_c}'^2}}. \quad (14)$$

The problem here is that there are three variables: the masses of h_2 , h^{--} and Q . However, for simplicity, following the ansatz in [9], we assume $m_{h_2} = X$, $m_{h^{--}} = m_Q = K$ and $m_{Z_2}(v_{\chi}) = 4.821$ TeV. Note that the contributions from h_2 , h^{--} , Z_2 , in this phase transition, are X or K , different to their contributions in the phase transition $SU(2) \rightarrow U(1)$. In order to have the first phase transition, the phase transition strength must be larger than the unit, i.e., $\frac{v_{\chi c}}{T_c'} \geq 1$.

In figure 1, we have plotted K as a function of $m_{h_2}(v_{\chi})$, with $m_{h_2}(v_{\chi}) > 1$ TeV.

According to Fig. 1, if X is larger than 1 TeV, the heavy particle masses are in range of few TeVs in order to have the first order phase transition. In addition, this phase transition can be strong first order.

B. Phase transition $SU' \rightarrow U'$

This phase transition dose not involve the exotic quarks and boson V^\pm and the contribution from $U^{\mp\mp}$ is equal to W^\mp . The effective potential of the EWPT $SU(2) \rightarrow U(1)$ is




FIG. The minimum conditions are contours of $E'/\lambda'_{T'_c}$ as contours of $E'/\lambda'_{T'_c}$.

$V_{eff}(v_\rho)$:

$$\begin{aligned}
 V_{eff}(v_\rho) = & V_0(v_\rho) + \frac{3}{64\pi^2} \left(m_{Z_1}^4(v_\rho) \ln \frac{m_{Z_1}^2(v_\rho)}{Q^2} + m_{Z_2}^4(v_\rho) \ln \frac{m_{Z_2}^2(v_\rho)}{Q^2} \right. \\
 & + 2m_W^4(v_\rho) \ln \frac{m_W^2(v_\rho)}{Q^2} + 2m_U^4(v_\rho) \ln \frac{m_U^2(v_\rho)}{Q^2} - 4m_t^4(v_\rho) \ln \frac{m_t^2(v_\rho)}{Q^2} \left. \right) \\
 & + \frac{1}{64\pi^2} \left(m_{h_1}^4(v_\rho) \ln \frac{m_{h_1}^2(v_\rho)}{Q^2} + m_{h_2}^4(v_\rho) \ln \frac{m_{h_2}^2(v_\rho)}{Q^2} + 2m_{h^{++}}^4 \ln \frac{m_{h^{++}}^2(v_\rho)}{Q^2} \right) \\
 & + \frac{T^4}{4\pi^2} \left[F_- \left(\frac{m_{h_1}(v_\rho)}{T} \right) + F_- \left(\frac{m_{h_2}(v_\rho)}{T} \right) + 2F_- \left(\frac{m_{h^{++}}(v_\rho)}{T} \right) \right] \\
 & + \frac{3T^4}{4\pi^2} \left[4F_+ \left(\frac{m_t(v_\rho)}{T} \right) + F_- \left(\frac{m_{Z_1}(v_\rho)}{T} \right) + F_- \left(\frac{m_{Z_2}(v_\rho)}{T} \right) \right. \\
 & \left. + 2F_- \left(\frac{m_W(v_\rho)}{T} \right) + 2F_- \left(\frac{m_U(v_\rho)}{T} \right) \right].
 \end{aligned}$$

The minimum conditions are

$$V_{eff}(v_{\rho_0}) = 0; \quad \frac{\partial V_{eff}}{\partial v_\rho}(v_{\rho_0}) = 0; \quad \frac{\partial V_{eff}}{\partial v_\rho}(v_{\rho_0}) = m_{h_1}^2 + m_{h_2}^2(v_\rho)|_{v_\rho=v_{\rho_0}}.$$

From the above minimum conditions, we see that in this EWPT $m_{h_2}^2(v_\rho)$ plays the role to generate masses of the last heavy particles and $m_{h_1}^2$ generates masses of the SM particles.

With the symmetry breaking scale equal to $Q \equiv v_{\rho_0} = v_0 = 246$ GeV, the high-temperature expansion of this potential has the form

$$V_{eff}^{RM331} = D(T^2 - T_0^2) \cdot v_\rho^2 - ET|v_\rho|^3 + \frac{\lambda_T}{4} v_\rho^4,$$

where

$$\begin{aligned} D &= \frac{1}{24v_0^2} \left[6m_W^2(v_\rho) + 6m_U^2(v_\rho) + 3m_{Z_1}^2(v_\rho) + 3m_{Z_2}^2(v_\rho) \right. \\ &\quad \left. + 6m_t^2(v_\rho) + m_{h_1}^2(v_\rho) + m_{h_2}^2(v_\rho) + 2m_{h^\pm}^2(v_\rho) \right], \\ T_0^2 &= \frac{1}{D} \left\{ \frac{1}{4}(m_{h_1}^2(v_\rho) + m_{h_2}^2(v_\rho)) - \frac{1}{32\pi^2 v_0^2} \left(6m_W^4(v_\rho) + 6m_U^4(v_\rho) + 3m_{Z_1}^4(v_\rho) \right. \right. \\ &\quad \left. \left. + 3m_{Z_2}^4(v_\rho) - 12m_t^4(v_\rho) + m_{h_1}^4(v_\rho) + m_{h_2}^4(v_\rho) + 2m_{h^\pm}^4(v_\rho) \right) \right\}, \\ E &= \frac{1}{12\pi v_0^3} \left(6m_W^3(v_\rho) + 6m_U^3(v_\rho) + 3m_{Z_1}^3(v_\rho) + 3m_{Z_2}^3(v_\rho) \right. \\ &\quad \left. + m_{h_1}^3(v_\rho) + m_{h_2}^3(v_\rho) + 2m_{h^\pm}^3(v_\rho) \right), \\ \lambda_T &= \frac{m_{h_1}^2(v_\rho) + m_{h_2}^2(v_\rho)}{2v_0^2} \left\{ 1 - \frac{1}{8\pi^2 v_0^2 (m_{h_1}^2(v_\rho) + m_{h_2}^2(v_\rho))} \left[6m_W^4(v_\rho) \ln \frac{m_W^2(v_\rho)}{bT^2} \right. \right. \\ &\quad \left. \left. + 3m_{Z_1}^4(v_\rho) \ln \frac{m_{Z_1}^2(v_\rho)}{bT^2} + 3m_{Z_2}^4(v_\rho) \ln \frac{m_{Z_2}^2(v_\rho)}{bT^2} + 6m_U^4(v_\rho) \ln \frac{m_U^2(v_\rho)}{bT^2} \right. \right. \\ &\quad \left. \left. - 12m_t^4(v_\rho) \ln \frac{m_t^2(v_\rho)}{b_F T^2} + m_{h_1}^4(v_\rho) \ln \frac{m_{h_1}^2(v_\rho)}{bT^2} + m_{h_2}^4(v_\rho) \ln \frac{m_{h_2}^2(v_\rho)}{bT^2} \right. \right. \\ &\quad \left. \left. + 2m_{h^\pm}^4(v_\rho) \ln \frac{m_{h^\pm}^2(v_\rho)}{bT^2} \right] \right\} \end{aligned} \quad (15)$$

The effective potential has two minimum points, the first minimum at $v_\rho = 0$ and the second one at $v_{\rho_c} = \frac{2ET_c}{\lambda_{T_c}}$. In the limit $E \rightarrow 0$, we have the second order phase transition. In order to have the first order phase transition, the phase transition strength has to be larger than the unit, i.e., $\frac{v_{\rho_c}}{T_c} \geq 1$. The critical temperature T_c is given by

$$T_c = \frac{T_0}{\sqrt{1 - E^2/D\lambda_{T_c}}}. \quad (16)$$

The equation (16) is self-consistent with the critical temperature because λ_{T_c} is a function of T_c . According to the LHC, we take $m_{h_1} = 125$ GeV and put $m_{h_2}(v_\rho) = Z$, $m_{h^{--}}(v_\rho) = Y$ and $m_{Z_2}(v_\rho) = 14.53$ GeV.

In Fig.2, we show the masses region of h_1 and h_{++} where the necessary condition of the first order phase transition is imposed. According to Fig.2, and by the numerical evaluation, the strength of the EWPT is in range, $1 \leq 2E/\lambda_{T_c} < 5$. Therefore, in the RM331 we always have the first order phase transition but it is weak (at the v_ρ scale).




FIG. The plot shows the mass m_{h_2} [Gev] versus $m_{h^{--}}$ [Gev]. The curves represent different parameter spaces or theoretical models. The solid line represents the most massive state, while the dashed and dotted lines represent lighter states. The jagged lines indicate regions of instability or specific physical constraints.

The contributions from new particles (at the first symmetry breaking) make of the first order phase transition that the Standard Model cannot. However, there is one thing special, heavy particles as $U^{\pm\pm}$, h_2 , h^{--} , Z_2 that contribute only the little part in their total masses.




FIG. For a particular value of v_ρ at T_c , the potential V has a minimum at $v_\rho = 0$. For $T > T_c$, the minimum shifts to $v_\rho > 0$. For $T < T_c$, the minimum shifts to $v_\rho < 0$. The curves represent different parameter spaces or theoretical models.

In Fig. 3, when temperature go close to T_c , the second minimum slowly formed distinct, ie the phase transition nucleation appears. When temperature go to T_c , the symmetry breaking phase is turned on and temperature go down T_c , the system switches to the symmetry breaking phase.

To finish this section, we conclude that the effective potential of this model is different from that of the SM, and it has the contributions from heavy bosons, which plays role of the triggers for the first order phase transition with $m_{h_1} = 125$ GeV.

C. Constraint on mass of charged Higgs boson

In Section IV, from the phase transition $SU(2) \rightarrow U(1)$, we have derived

$$200 \text{ GeV} < Y = m_{h_{++}}(v_\rho) < 1200 \text{ GeV}$$

and

$$0 < Z = m_{h_2}(v_\rho) < 624 \text{ GeV}.$$

Therefore, we get

$$\begin{aligned} (200 \text{ GeV})^2 &< \frac{\lambda_4}{2} v_{\rho_0}^2 < (1200 \text{ GeV})^2, \\ (0 \text{ GeV})^2 &< \frac{\lambda_3^2}{4\lambda_2} v_{\rho_0}^2 < (624 \text{ GeV})^2. \end{aligned} \quad (17)$$

Taking into account of the recent Higgs boson mass (125 GeV), combining with the formula (5), we also obtain

$$\left(\lambda_1 - \frac{\lambda_3^2}{4\lambda_2} \right) v_{\rho_0}^2 = (125 \text{ GeV})^2. \quad (18)$$

Combination of the equations in (17) and (18) leads to

$$\begin{aligned} 0 &< \lambda_1 < 6.692, \\ 0 &< \frac{\lambda_3^2}{4\lambda_2} < 6.434, \\ 1.321 &< \lambda_4 < 47.59. \end{aligned}$$

From equation (19), leading to $\lambda_1, \lambda_2, \lambda_4$ that must be positive, satisfying the above bound conditions and $3.258 \text{ TeV} < m_{h_{++}} < 19.549 \text{ TeV}$. Thus the mass of the heavy Higgs must be some few TeVs, the electroweak phase transition in this model is the first order phase transition. We hope that the heavy particles will uncover many more new physics.

V. CONCLUSION AND OUTLOOKS

We have used the effective potential at finite temperature to study the structure of the electroweak phase transition in the RM331 model. This phase transition is separated into two phases. The first transition period is $SU(3) \rightarrow SU(2)$, or the symmetry breaking in the energy scale v_{χ_0} (in order to generate masses for the heavy particles and the exotic quarks). The second phase transition is $SU(2) \rightarrow U(1)$ at v_{ρ_0} , which generates masses for all usual fermions and the SM gauge bosons. The electroweak phase transition in this model (in the scale v_ρ) may be the weakly first order electroweak phase transition with $m_{h_1} = 125$ GeV if the heavy bosons masses are some few TeVs. So this is strong enough to study the baryon asymmetry.

If Z_2 exists, its mass is some few TeVs, so its contribution to the electroweak phase transition is very large. Therefore, the electroweak phase transition in this model is completely turned on. In other words, the baryon asymmetry problem in this model is directly related to the mass of Z_2 .

The self-interactions of Higgs in this model are more complicated than the SM, because heavy particles involve in both two phase transitions. Thus calculating the quantum corrections can reveal many new physical phenomena and opens up new relations further between Cosmology and Particle Physics. In addition, from the phase transitions, we can get some bounds on the Higgs self-couplings.

Although we only work on the RM331 model, but this calculation can still apply to other 3-3-1 models such as the recent supersymmetric reduced minimal 3-3-1 model [22] and the 3-3-1-1 model [23]. So the 3-3-1 models can have specific advantages more than the SM in explaining the baryon asymmetry problem.

Our next works will calculate the sphaleron rate and CP violations in 3-3-1 models, in order to analysis in details electroweak baryogenesis. In addition, by using the electroweak phase transition or baryogenesis problem, we can predict masses of the heavy particles beyond the SM such as 3-3-1 models.

Acknowledgment

The authors thank P. V. Dong and D. T. Huong for critical remarks and stimulating discussions. This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 103.01-2011.63.

u anov. *Physical Foundations of Cosmology*, Cambridge University Press, A
 D arov. *J Exp an or ys* **71**, **71**
 K Ka ant La n K u u a n n an apos n ov. *ys v L tt* **77**
 , F Cs or. Z Fo or. an J H t r. *ys v L tt* **82**, J Grant.
 H n ars *ys v D* **64**, D= no o. K u u a n n A ranb r.
 JHE **08**,
 Bast ro G C Hu on F K n. D oy an pat *ys L tt B* **489**.
 , A non. D E orr ss y. an C E a n r. *ys v D* **70**.
 Ha K C K E J Yoo. an D on. *ys v D* **70**.
 J Esp nosa. Konstan n an F va u *ys B*,
 Kan ura. E na a. n ou an Ya a a. JHE
 D J H C un. an A J Lon. *ys v D* **81**.
 J C n G Laport H Ya as ta. Kra. JHE **0907**
 Kan ura. Y a a E na a. *ys L tt B* **606**.
 Ha A an K *ys v D* **81**.
 G Bar nbo an us. *ys v D* **58**.
 Das. J Fox. A Ku ar. an n r. JHE **1011**, D C un. an A J
 Lon. *ys v D* **84**, Car na. a a. an C E a n r. *ys*
v D **85**, A A r an asr. *ys v D* **85**, D Bora
 an J C n *ys v D* **86**.
 w r suits n at t at part s ov r at CE s a H. s boson.
 ttp] pr ss w b rn
 F sano an t *ys v D*, H Fra pton. *ys v L tt*

Foot *et al*, ys v D 80

A x G D as t ys v D 80

n r J F a an J t r ys v D , Foot H Lon

an uan A ran ys v D arX v p p] J C ont ro F

sano an t ys v D , H Lon ys v D

H Lon ys v D H Lon o ys L tt A13

C A r s av n ys v D , A Do F sano o

ys L tt A ys v D Don H Lon Int J

o ys A

A on Y Gra o an L A an ys v D 67 Don H

Lon D un an D oa ys v D 73 Don an H

Lon A v H En r y ys 2008 arX v Don r

Huon D Huon an H Lon ys v D 74 Don H

Lon an D oa ys v D 73 Don H Lon an D

oa ys v D 75 Don H Hum an H Lon ys v

D 86

J G Frr ra Jr D n ro C A r s an o r u s a va ys

v D 84 Huy n La H Lon an on

arX v p p

D ys v D 49

art Data Group J B r n r et al ys v D 86

or xa p s D A Gut rr A on an L A an Eur ys J C 46

Y A Cout n o Gu ara s an A A po u no arX v

Duon an E ys L tt B J C ont ro t C

o r u ys v D A Da art n J A o r u ys

L tt B C ont ro t C o r u ys v D

Don D Huon C o r u H Lon u ys B

Don D Huon uy H Lon u ys B

Huon L Hu C o r u H Lon u ys B J G Frr ra

C A r s o r u s a va A a p r r nt arX v D B n

L Hu D Huon H Lon r nt arX v
 Don H Hun D a ys v D 87 ,
 At $K v_\rho \equiv v_{\rho_0}$ G an $v_\chi \equiv v_{\chi_0}$ \div In t s wor w oos v_{χ_0}