Put your logo here



Goshawk. Adult. Male (06-IV)

NORTHERN GOSHAWK (Accipiter gentilis)



Male: 48-61 cm; **female**: 58-64 cm. Short and rounded wings and long tail; yellow or orange iris; tail with 4-5 bands; grey brown upper-parts; pale and barred underparts. **Juveniles** are brown with pale underparts mottled brown.



Goshawk. Adult. Pattern of head and breast.





Goshawk. Juvenile. Pattern of head and breast.



SIMILAR SPECIES

Kestrel and Lesser Kestrel have long tails, but have dark iris and rufous upperparts. Female Sparrowhawk can be very similar to a male Goshawk but has a smaller size; juvenile Sparrowhawk has barred underparts.



Kestrel. Female



Sparrowhawk. Juvenile



Sparrowha wk. Female

SEXING

In adult birds, male with bluish grey upperparts; ear covers deep grey. Female with browish upperparts; ear-coverts light grey or brownish. Female with bigger size than male. In juveniles, plumage of both sexes alike but can be sexed by size.

CAUTION: head and upperparts pattern is individually very variable; size in this species is very different depending on its geographical location (northern birds bigger than southern ones), so it would be necessary to know the geographical origin to sex by size without errors.

Put your logo here



Goshawk Sexing. Pattern of head: top male; bottom female.







Goshawk Sexing. Pattern of upperparts: left male; right female.

AGEING

3 age groups can be recognized:

Juvenil with brown streaked head with only a slight supercilium; feathers of upperparts brown with pale edge; pale underparts with dark vertical streaks; pale yellow iris.


2nd year autumn/3rd year spring with grey brown upperparts with brown and worn retained juvenile feathers specially on rump and mantle; with some brown juvenile feathers retained on wing coverts, specially on lesser and median coverts; breast feathers with 3 or 4 transversal broad and brown lines, sometimes tinged rufous; sometimes with some narrow, brownish and worn juvenile secondaries retained; yellow or yellow-orange iris.

Adult with uniformly dark crown and earcoverts with clear white supercilium; uniform brown grey or slate-brown upperparts, sometimes with thinly white tipped feathers; breast feathers with 4 or 5 transversal narrow and grey -brown lines, the outermost narrower than the others; if retained secondaries then with adult pattern; orange to red iris.











Goshawk Ageing. Pattern of upperparts:left adult; right ju-

Write your website here


Page 2

Put your logo here

MOULT

Complete **postbreeding** moult but central secondaries often retained; usually finished in September. **Postjuvenile** moult begins in **2nd year** and is complete but occasionally some central secondaries, rump feathers and wing coverts retained.

PHENOLOGY



STATUS IN ARAGON

Resident, with contribution of wintering European birds. Breeds in all the Region in appropriated forests.



Goshawk. Adult. Female (12-XII).



Goshawk. 2nd year autumn. Male ()



Goshawk. 2nd year autumn. Female (20-XI)



Goshawk. Juvenile. Male (30-VII)



Goshawk. Juvenile. Female (06-X)

Put your logo here






Goshawk. Adult. Head pattern: top male (06-IV30-VII); bottom female (12-XII).





Goshawk. Juvenile. Head pattern: top male (30-VII); bottomfemale (06-X).





Goshawk. 2nd year autumn. Head pattern: top male (); bottom female (20-XI).



Goshawk. Adult. Crown pattern: left male (06-IV); right female (12-XII).



Goshawk. 2nd year autumn. Crown pattern: left male (); right female (20-XI).

Put your logo here



Goshawk. Juvenile. Crown pattern: left male (30-VII); right female (06-X).



Goshawk. Adult. Breast pattern: left male (06-IV); right female (12-XII).



Goshawk. 2nd year autumn. Breast pattern: left male (); right female (20-XI).



Goshawk. Juvenile. Breast pattern: left male (30-VII); right female (06-X).





Goshawk. Adult. Pattern of breast feathers: top male (06-IV); bottom female (12-XII).



Goshawk. 2nd year autumn. Pattern of breast feathers: top male (); bottom female (20-XI).








Goshawk. Juvenile. Pattern of breast feathers: top male (30-VII); bottom female (06-X).

Put your logo here





Goshawk. Adult. Rump pattern: left male (06-IV); right female (12-XII).



Goshawk. 2nd year autumn. Rump pattern: left male (); right female (20-XI).



Goshawk. Juvenile. Rump pattern: left male (30-VII); right female (06-X).



Goshawk. Adult. Upperparts pattern: left male (06-IV); right female (12-XII).





Goshawk. 2nd year autumn. Upperparts pattern: left male (); right female (20-XI).


Put your logo here



Goshawk. Juvenile. Upperparts pattern: left male (30 -VII); right female (06-X).



Goshawk. Adult. Tail pattern: left male (06-IV); right female (12-XII).



Goshawk. 2nd year autumn. Tail pattern: left male (); right female (20-XI).



Goshawk. Juvenile. Tail pattern: left male (30-VII); right female (06-X).





Goshawk. Adult. Pattern of primary coverts: top male (06-IV); bottom female (12-XII).

Put your logo here







Goshawk. 2nd year autumn. Pattern of primary coverts: top male (); bottom female (20-XI).



Goshawk. Adult. Female: pattern of wing coverts (12 -XII).





venile. Pattern of primaty coverts: top male (30-VII); bottom female (06-X).

Goshawk. Ju-

Goshawk. 2nd year autumn. Male: pattern of wing coverts ().





Goshawk. Adult. Male: pattern of wing coverts (06-IV).

Goshawk. 2nd year autumn. Female: pattern of wing coverts (20-XI).



Goshawk. Juvenile. Male: pattern of wing coverts (30-VII).

Put your logo here



Goshawk. Juvenile. Female: pattern of wing coverts (06-X).



Goshawk. Adult. Male: pattern of primaries (06-IV).




Goshawk. Adult. Female: pattern of primaries (12-XII).





Goshawk. 2nd year autumn. Male: pattern of primaries ().

Put your logo here





Goshawk. 2nd year autumn. Female: pattern of primaries (20-XI).



Goshawk. Juvenile. Male: pattern of primaries (30-VII).





Goshawk. Juvenile. Female: pattern of primaries (06-X).




Goshawk. Adult. Male: pattern of secondaries (06-IV).

Put your logo here

117 Northern Goshawk

<image>

Goshawk. Adult. Female: pattern of secondaries (12-XII).



Goshawk. 2nd year autumn. Male: pattern of secondaries ().



Goshawk. 2nd year autumn. Female: pattern of secondaries (20-XI).



Goshawk. Juvenile. Male: pattern of secundaries (30-VII).


<image>

Goshawk. Juvenile. Female: pattern of secundaries (06-X).






Goshawk. Adult. Male: pattern of wing (06-IV).



Goshawk. Adult. Female: pattern of wing (12-XII).



Goshawk. 2nd year autumn. Male: pattern of wing ().



Goshawk. 2nd year autumn. Female: pattern of wing (20-XI).

Put your logo here

Put your logo here

117 Northern Goshawk

<image>

Goshawk. Juvenile. Male: pattern of wing (30-VII).



Goshawk. Juvenile. Female: pattern of wing (06-X).