

# On 3-decomposable geometric drawings of $K_n$

B. Ábrego\*    S. Fernández-Merchant\*    J. Leaños†    G. Salazar†

November 13, 2018

## Abstract

The point sets of all known optimal rectilinear drawings of  $K_n$  share an unmistakable clustering property, the so-called *3-decomposability*. It is widely believed that the underlying point sets of all optimal rectilinear drawings of  $K_n$  are 3-decomposable. We give a lower bound for the minimum number of ( $\leq k$ )-sets in a 3-decomposable  $n$ -point set. As an immediate corollary, we obtain a lower bound for the crossing number  $\overline{cr}(\mathcal{D})$  of any rectilinear drawing  $\mathcal{D}$  of  $K_n$  with underlying 3-decomposable point set, namely  $\overline{cr}(\mathcal{D}) > \frac{2}{27} (15 - \pi^2) \binom{n}{4} + \Theta(n^3) \approx 0.380029 \binom{n}{4} + \Theta(n^3)$ . This closes this gap between the best known lower and upper bounds for the rectilinear crossing number  $\overline{cr}(K_n)$  of  $K_n$  by over 40%, under the assumption of 3-decomposability.

## 1 Introduction

Figure 1 shows the point set of an optimal (crossing minimal) rectilinear drawing of  $K_9$ , with an evident partition of the 9 vertices into 3 highly structured clusters of 3 vertices each:




Figure 1: The points in this optimal drawing of  $K_9$  are clustered into 3 sets.

A similar, natural, highly structured partition into 3 clusters of equal size is observed in *every* known optimal drawing of  $K_n$ , for every  $n$  multiple of 3 (see [4]). Even for those values of  $n$  (namely,  $n > 27$ ) for which the exact rectilinear crossing number  $\overline{cr}(K_n)$  of  $K_n$  is not known, the best available examples also share this property [4].

In all these examples, a set  $S$  of  $n$  points in general position is partitioned into sets  $A$ ,  $B$ , and  $C$ , with  $|A| = |B| = |C| = n/3$  with the following properties:

- (i) There is a directed line  $\ell_1$  such that, as we traverse  $\ell_1$ , we find the  $\ell_1$ -orthogonal projections of the points in  $A$ , then the  $\ell_1$ -orthogonal projections of the points in  $B$ , and then the  $\ell_1$ -orthogonal projections of the points in  $C$ ;

\*Department of Mathematics, California State University Northridge.

†Instituto de Física, Universidad Autónoma de San Luis Potosí, Mexico. Supported by FAI-UASLP and by CONACYT Grant 45903.

- (ii) there is a directed line  $\ell_2$  such that, as we traverse  $\ell_2$ , we find the  $\ell_2$ -orthogonal projections of the points in  $B$ , then the  $\ell_2$ -orthogonal projections of the points in  $A$ , and then the  $\ell_2$ -orthogonal projections of the points in  $C$ ; and
- (iii) there is a directed line  $\ell_3$  such that, as we traverse  $\ell_3$ , we find the  $\ell_3$ -orthogonal projections of the points in  $B$ , then the  $\ell_3$ -orthogonal projections of the points in  $C$ , and then the  $\ell_3$ -orthogonal projections of the points in  $A$ .

**Definition** A point set that satisfies conditions (i)–(iii) above is *3-decomposable*. We also say that the underlying rectilinear drawing of  $K_n$  is *3-decomposable*.

A possible choice of  $\ell_1, \ell_2$ , and  $\ell_3$  for the example in Figure 1 is illustrated in Figure 2.




Figure 2: The 9-point set  $S$  gets naturally partitioned into three clusters  $A, B$ , and  $C$ . The  $\ell_1$ -,  $\ell_2$ -, and  $\ell_3$ -orthogonal projections of  $A, B$ , and  $C$  satisfy conditions (i)–(iii), and so  $S = A \cup B \cup C$  is 3-decomposable.

## 1.1 The main result

It is widely believed that all optimal rectilinear drawings of  $K_n$  are 3-decomposable. One of our main results in this paper is the following lower bound for the number of crossings in all such drawings.

**Theorem 1** *Let  $\mathcal{D}$  be a 3-decomposable rectilinear drawing of  $K_n$ . Then the number  $\overline{\text{cr}}(\mathcal{D})$  of crossings in  $\mathcal{D}$  satisfies*

$$\overline{\text{cr}}(\mathcal{D}) \geq \frac{2}{27} (15 - \pi^2) \binom{n}{4} + \Theta(n^3) \approx 0.380029 \binom{n}{4} + \Theta(n^3).$$

The best known general lower and upper bounds for the rectilinear crossing number  $\overline{\text{cr}}(K_n)$  are  $0.37968 \binom{n}{4} + \Theta(n^3) \leq \overline{\text{cr}}(K_n) \leq 0.38054 \binom{n}{4} + \Theta(n^3)$  (see [3] and [2]). Thus the

bound given by Theorem 1 closes this gap by over 40%, under the (quite feasible) assumption of 3-decomposability.

To prove Theorem 1 (in Section 2), we exploit the close relationship between rectilinear crossing numbers and  $(\leq k)$ -sets, unveiled independently by Ábrego and Fernández-Merchant [1] and by Lovász et al. [8].

Recall that a  $(\leq k)$ -set of a point set  $S$  is a subset  $T$  of  $S$  with  $|T| \leq k$  such that some straight line separates  $T$  and  $S \setminus T$ . The number  $\chi_{\leq k}(S)$  of  $(\leq k)$ -sets of  $S$  is a parameter of independent interest in discrete geometry (see [6]), and, as we recall in Section 2, is closely related to the rectilinear crossing number of the geometric graph induced by  $S$ .

The main ingredient in the proof of Theorem 1 is the following bound (Theorem 2) for the number of  $(\leq k)$ -sets in 3-decomposable point sets. The bound is in terms of the following quantity (by convention,  $\binom{r}{s} = 0$  if  $r < s$ ),

$$Y(k, n) := 3 \binom{k+1}{2} + 3 \binom{k - n/3 + 1}{2} + 3 \sum_{j=2}^b j(j+1) \binom{k+1 - (\frac{1}{2} - \frac{1}{3j(j+1)})3n}{2} - \frac{1}{3}, \quad (1)$$

where  $b := b(k, n)$  is the unique integer such that  $\binom{b(k,n)+1}{2} < n/(n-2k-1) \leq \binom{b(k,n)+2}{2}$ .

**Theorem 2** *Let  $S$  be a 3-decomposable set of  $n$  points in general position, where  $n$  is a multiple of 3, and let  $k < n/2$ . Then*

$$\chi_{\leq k}(S) \geq Y(k, n).$$

The best general lower bound for  $\chi_{\leq k}(S)$  is the sum of the first two terms in (1) (see [3] and [2]). Thus the third summand in (1) is the improvement we report, under the assumption of 3-decomposability.

The proofs of Theorems 1 and 2 are in Sections 2 and 3, respectively. In Section 4 we present some concluding remarks and open questions.

## 2 Proof of Theorem 1

Let  $\mathcal{D}$  be a 3-decomposable rectilinear drawing of  $K_n$ , and let  $S$  denote the underlying  $n$ -point set, that is, the vertex set of  $\mathcal{D}$ . Besides Theorem 2, our main tool is the following relationship between  $(\leq k)$ -sets and the rectilinear crossing number (see [1] or [8]):

$$\overline{\text{cr}}(\mathcal{D}) = \sum_{1 \leq k \leq (n-2)/2} (n-2k-1) \chi_{\leq k}(S) + \Theta(n^3). \quad (2)$$

Combining Theorem 2 and Eq. (2), and noting that both the  $-1$  in the factor  $n-2k-1$  and the summand  $-1/3$  in (1) only contribute to smaller order terms, we obtain:

$$\begin{aligned}
\overline{\text{cr}}(\mathcal{D}_n) &\geq \sum_{k=1}^{\binom{n-2}{2}} (n-2k) \left( 3 \binom{k+1}{2} + 3 \binom{k-n/3+1}{2} + 3 \sum_{j=2}^b j(j+1) \binom{k+1 - (\frac{1}{2} - \frac{1}{3j(j+1)})3n}{2} \right) + \Theta(n^3) \\
&\geq 24 \binom{n}{4} \sum_{k=1}^{\binom{n-2}{2}} \frac{(1-2k/n)}{n} \left( 3 \binom{k+1}{2} + 3 \binom{k-n/3+1}{2} + 3 \sum_{j=2}^b j(j+1) \binom{k+1 - (\frac{1}{2} - \frac{1}{3j(j+1)})3n}{2} \right) / n^2 + \Theta(n^3) \\
&= 24 \binom{n}{4} \left( \sum_{k=1}^{\binom{n-2}{2}} \binom{3}{2} \frac{(1-2k/n)}{n} \left( \frac{k}{n} \right)^2 + \sum_{k=n/3}^{\binom{n-2}{2}} \binom{3}{2} \frac{(1-2k/n)}{n} \left( \frac{k}{n} - \frac{1}{3} \right)^2 \right. \\
&\quad \left. + \sum_{j=2}^{\infty} j(j+1) \sum_{k=(\frac{1}{2}-1/(3j(j+1)))n}^{\binom{n-2}{2}} \binom{3}{2} \frac{(1-2k/n)}{n} \left( \frac{k}{n} - \left( \frac{1}{2} - \frac{1}{3j(j+1)} \right) \right)^2 \right) + \Theta(n^3) \\
&= 24 \binom{n}{4} \left( \int_0^{1/2} (3/2)(1-2x)x^2 dx + \int_{1/3}^{1/2} (3/2)(1-2x)(x-1/3)^2 dx \right. \\
&\quad \left. + \sum_{j=2}^{\infty} (3/2)j(j+1) \int_{\frac{1}{2}-\frac{1}{3j(j+1)}}^{1/2} (1-2x)(x - (1/2 - 1/(3j(j+1))))^2 dx \right) + \Theta(n^3)
\end{aligned}$$

Elementary calculations show that  $\frac{3}{2} \int_0^{1/2} (1-2x)x^2 dx = 3/8$ ,  $\frac{3}{2} \int_{1/3}^{1/2} (1-2x)(x-1/3)^2 dx = 1/216$ , and  $\int_{\frac{1}{2}-\frac{1}{3j(j+1)}}^{1/2} (1-2x)(x - (1/2 - 1/(3j(j+1))))^2 dx = (1/486)j^4(1+j)^4$ .

Thus,

$$\overline{\text{cr}}(\mathcal{D}_n) \geq 24 \binom{n}{4} \left( 3/8 + 1/216 + (3/2) \sum_{j=2}^{\infty} \frac{1}{486j^3(j+1)^3} \right) + \Theta(n^3).$$

Since  $\sum_{j=2}^{\infty} \frac{1}{j^3(j+1)^3} = \sum_{j=2}^{\infty} \left( \frac{1}{j^3} - \frac{3}{j^2} + \frac{6}{j} - \frac{1}{(j+1)^3} - \frac{3}{(j+1)^2} - \frac{6}{j+1} \right) = \frac{79}{8} - \pi^2$ , then

$$\overline{\text{cr}}(\mathcal{D}_n) \geq \frac{2}{27} \left( 15 - \pi^2 \right) \binom{n}{4} + \Theta(n^3). \quad \blacksquare$$

## 3 Proof of Theorem 2

The first step to prove Theorem 2 is to obtain an equivalent (actually, more general) formulation in terms of circular sequences (namely Proposition 3 below).

### 3.1 Circular sequences: reducing Theorem 2 to Proposition 3

All the geometrical information of a point set  $S$  gets encoded in (any halfperiod of) the *circular sequence* associated to  $S$ . We recall that a circular sequence on  $n$  elements is a doubly infinite sequence  $\dots \pi_{-1}, \pi_0, \pi_1, \dots$  of permutations of the points in  $S$ , where consecutive permutations differ in a transposition of neighboring elements, and, for every  $i$ ,  $\pi_i$  is the reverse permutation of  $\pi_{i+\binom{n}{2}}$ . Thus a circular sequence on  $n$  elements has period  $2\binom{n}{2}$ , and all the information is encoded in an  $n$ -halfperiod, that is, a sequence of  $\binom{n}{2} + 1$  consecutive permutations.

Each  $n$ -point set  $S$  has an associated circular sequence  $\mathbf{\Pi}_S$ , which contains all the geometrical information of  $S$  [7]. As we observed above, any  $n$ -halfperiod  $\Pi$  of  $\mathbf{\Pi}_S$  contains all the information of  $\mathbf{\Pi}_S$ , and so  $n$ -halfperiods are usually the object of choice to work

with. In an  $n$ -halfperiod  $\Pi = \pi_0, \pi_1, \dots, \pi_{\binom{n}{2}}$ , the *initial* permutation is  $\pi_0$  and the *final* permutation is  $\pi_{\binom{n}{2}}$ .

Not every  $n$ -halfperiod  $\Pi$  arises from a point set  $S$ . We refer the reader to the seminal work by Goodman and Pollack [7] for further details.

Observe that if  $S$  is 3-decomposable, then there is an  $n$ -halfperiod  $\Pi$  of the circular sequence associated to  $S$ , whose points can be labeled  $a_1, \dots, a_{n/3}, b_1, \dots, b_{n/3}, c_1, \dots, c_{n/3}$ , so that:

- (i) The initial permutation  $\pi_0$  reads  $a_{n/3}, a_{n/3-1}, \dots, a_1, b_1, b_2, \dots, b_{n/3}, c_1, c_2, \dots, c_{n/3}$ ;
- (ii) there is an  $s$  such that in the  $(s+1)$ -st permutation first the  $b$ 's appear consecutively, then the  $a$ 's appear consecutively, and then the  $c$ 's appear consecutively; and
- (iii) there is a  $t$ , with  $t > s$ , such that in the  $(t+1)$ -st permutation first the  $b$ 's appear consecutively, then the  $c$ 's appear consecutively, and then the  $a$ 's appear consecutively.

**Definition** An  $n$ -halfperiod  $\Pi$  that satisfies properties (i)–(iii) above is 3-decomposable.

A transposition that occurs between elements in sites  $i$  and  $i+1$  is an  $(i, i+1)$ -transposition. An  $i$ -critical transposition is either an  $(i, i+1)$ -transposition or an  $(n-i, n-i+1)$ -transposition, and a  $(\leq k)$ -critical transposition is a transposition that is  $i$ -critical for some  $i \leq k$ . If  $\Pi$  is an  $n$ -halfperiod, then  $\eta_{\leq k}(\Pi)$  denotes the number of  $(\leq k)$ -critical transpositions in  $\Pi$ .

The key result is the following.

**Proposition 3** *Let  $\Pi$  be a 3-decomposable  $n$ -halfperiod, and let  $k < n/2$ . Then*

$$\eta_{\leq k}(\Pi) \geq Y(k, n).$$

**Proof of Theorem 2.** Let  $S$  be 3-decomposable, and let  $\Pi$  be an  $n$ -halfperiod of the circular sequence associated to  $S$ , that satisfies properties (i)–(iii) above. Then  $\Pi$  is 3-decomposable. Now, for any point set  $T$  and any halfperiod  $\Pi_T$  associated to  $T$ , the  $(\leq k)$ -critical transpositions of  $\Pi_T$  are in one-to-one correspondence with  $(\leq k)$ -sets of  $T$ . Applying this to  $\Pi$  and  $S$ , it follows that  $\chi_{\leq k}(S) = \eta_{\leq k}(\Pi)$ . Applying Proposition 3, Theorem 2 follows. ■

We devote the rest of this section to the proof of Proposition 3.

### 3.2 Proof of Proposition 3

Throughout this section,  $\Pi = (\pi_0, \pi_1, \dots, \pi_{\binom{n}{2}})$  is a 3-decomposable  $n$ -halfperiod, with initial permutation  $\pi_0 = (a_{n/3}, a_{n/3-1}, \dots, a_1, b_1, \dots, b_{n/3}, c_1, \dots, c_{n/3})$ .

In order to (lower) bound the number of  $(\leq k)$ -critical transpositions in 3-decomposable circular sequences, we distinguish between two types of transpositions. A transposition is *homogeneous* if it occurs between two  $a$ 's, between two  $b$ 's, or between two  $c$ 's; otherwise it is *heterogeneous*. We let  $\eta_{\leq k}^{\text{hom}}(\Pi)$  (respectively  $\eta_{\leq k}^{\text{het}}(\Pi)$ ) denote the number of homogeneous (respectively heterogeneous)  $(\leq k)$ -critical transpositions in  $\Pi$ , so that

$$\eta_{\leq k}(\Pi) = \eta_{\leq k}^{\text{hom}}(\Pi) + \eta_{\leq k}^{\text{het}}(\Pi). \quad (3)$$

### 3.2.1 Bounding (actually, calculating) $\eta_{\leq k}^{\text{het}}(\Pi)$

Let us call a transposition an *ab*-transposition if it involves one *a* and one *b*. We similarly define *ac*- and *bc*-transpositions. Thus, each heterogeneous transposition is either an *ab*- or an *ac*- or a *bc*-transposition.

Since in  $\Pi$  each *ab*-transposition moves the involved *a* to the right and the involved *b* to the left, then (a) for each  $i \leq n/3$ , there are *exactly*  $i$  *i*-critical *ab* transpositions; and (b) for each  $i$ ,  $n/3 < i < 2n/3$ , there are *exactly*  $2n/3 - i$  *i*-critical *ab*-transpositions. Since the same holds for *ac*- and *bc*-transpositions, it follows that for each  $i \leq n/3$ , there are *exactly*  $3i$  *i*-critical heterogeneous transpositions, and for each  $i$ ,  $n/3 < i < 2n/3$ , exactly  $2n - 3i$  *i*-critical heterogeneous transpositions.

Therefore, for each  $k \leq n/3$ , there are exactly  $\sum_{i \leq k} 3i = 3 \binom{k+1}{2}$  ( $\leq k$ )-critical transpositions, and if  $n/3 < k < n/2$ , then there are exactly  $\sum_{i \leq n/3} 3i + \sum_{n/3 < i \leq k} 2n - 3i + \sum_{n-k-1 < i \leq 2n/3-1} 2n - 3i = 3 \binom{n/3+1}{2} + (k - n/3)n$  ( $\leq k$ )-critical transpositions.

We now summarize these results.

**Proposition 4** *Let  $\Pi$  be a 3-decomposable  $n$ -halfperiod, and let  $k < n/2$ . Then*

$$\eta_{\leq k}^{\text{het}}(\Pi) = \begin{cases} 3 \binom{k+1}{2} & \text{if } k \leq n/3, \\ 3 \binom{n/3+1}{2} + (k - n/3)n & \text{if } n/3 < k < n/2, \end{cases}$$

### 3.2.2 Bounding $\eta_{\leq k}^{\text{hom}}(\Pi)$

Our goal here is to give a lower bound (see Proposition 10) for the number  $\eta_{\leq k}^{\text{hom}}(\Pi)$  of homogeneous ( $\leq k$ )-critical transpositions in a 3-decomposable  $n$ -halfperiod  $\Pi$ .

Our approach is to find an *upper* bound for  $\eta_{>k}^{aa}(\Pi)$ , which will denote the number of *aa*-transpositions that are *not* ( $\leq k$ )-critical ( $\eta_{>k}^{bb}(\Pi)$  and  $\eta_{>k}^{cc}(\Pi)$  are defined analogously). Since the total number of *aa*-transpositions is  $\binom{n/3}{2}$ , this will yield a lower bound for the contribution of *aa*-transpositions (and, by symmetry, for the contribution of *bb*-transpositions and of *cc*-transpositions) to  $\eta_{\leq k}^{\text{hom}}(\Pi)$ .

**Remark 5** *For every  $k \leq n/3$ , it is a trivial task to construct  $n$ -halfperiods  $\Pi$  for which  $\eta_{\leq k}^{\text{hom}}(\Pi) = 0$ . In view of this, we concentrate our efforts on the case  $k > n/3$ .*

A transposition between elements in positions  $i$  and  $i + 1$ , with  $k + 1 \leq i \leq n - k - 1$ , is *valid*. Thus our goal is to (upper) bound the number of valid *aa*-transpositions.

Let  $D_{\Pi}^{aa}$  be the digraph with vertex set  $a_1, \dots, a_{n/3}$ , and such that there is a directed edge from  $a_\ell$  to  $a_j$  if and only if  $\ell < j$  and the transposition that swaps  $a_\ell$  and  $a_j$  is valid. For  $j = 1, \dots, n/3$ , we let  $[a_j]_{\Pi}^+$  (respectively  $[a_j]_{\Pi}^-$ ) denote the outdegree (respectively indegree) of  $a_j$  in  $D_{\Pi}^{aa}$ . We define  $D_{\Pi}^{bb}$ ,  $D_{\Pi}^{cc}$ ,  $[b_j]_{\Pi}^+$ ,  $[b_j]_{\Pi}^-$ ,  $[c_j]_{\Pi}^+$  and  $[c_j]_{\Pi}^-$  analogously.

The inclusion of the symbol  $\Pi$  in  $D_{\Pi}^{aa}$ ,  $[a_i]_{\Pi}^+$ , etc., is meant to emphasize the dependence on the specific  $n$ -halfperiod  $\Pi$ . For brevity we will omit the reference to  $\Pi$  and simply write  $D^{aa}$ ,  $D^{bb}$ ,  $D^{cc}$ ,  $[a_i]^+$ ,  $[a_i]^-$ , and so on. No confusion will arise from this practice.

The importance of  $D^{aa}$ ,  $D^{bb}$ , and  $D^{cc}$  is clear from the following observation.

**Remark 6** *For each  $n$ -halfperiod  $\Pi$ , the number of edges of  $D^{aa}$  equals  $\eta_{>k}^{aa}(\Pi)$ . Indeed, to each valid *aa*-transposition, that is, each transposition that contributes to  $\eta_{>k}^{aa}(\Pi)$ , there corresponds a unique edge in  $D^{aa}$ . Analogous observations hold for  $D^{bb}$  and  $D^{cc}$ .*

In view of Remark 6, we direct our efforts to bounding the number of edges in  $D^{aa}$ . The essential observation to get this bound is the following:

$$[a_j]^- \leq \min\{n - 2k - 1 + [a_j]^+, (n/3) - j\}. \quad (4)$$

To see this, simply note that  $[a_j]^- \leq n - 2k - 1 + [a_j]^+$ , since  $n - 2k - 1 + [a_j]^+$  is clearly the maximum possible number of valid moves in which  $a_j$  moves right, and trivially  $[a_j]^- \leq (n/3) - j$ , since there are only  $(n/3) - j$   $a_\ell$ 's with  $\ell < j$ .

**Proposition 7** *If  $\Pi$  is a 3-decomposable  $n$ -halfperiod, and  $n/3 < k < n/2$ , then  $D^{aa}$  has at most  $\binom{n/3}{2} - (1/3) \left( Y(k, n) - 3\binom{n/3+1}{2} - (k - n/3)n \right)$  edges.*

*Proof.* Let  $\mathcal{D}_{k,n}$  denote the class of all digraphs with vertex set  $a_1, \dots, a_{n/3}$ , with every directed edge  $\overrightarrow{a_\ell a_j}$  satisfying  $\ell < j$  and  $[a_j]^- \leq \min\{n - 2k - 1 + [a_j]^+, n/3 - j\}$ .

We argue that any graph in  $\mathcal{D}_{k,n}$  has at most  $\binom{n/3}{2} - (1/3) \left( Y(k, n) - 3\binom{n/3+1}{2} - (k - n/3)n \right)$  edges. This clearly finishes the proof, since  $D^{aa} \in \mathcal{D}_{k,n}$ .

To achieve this, we note that it follows from the work in Section 2 in [5] that the maximum number of edges of such a digraph is attained in the digraph  $D_{k,n}$  recursively constructed as follows. First define that all the directed edges arriving at  $a_{n/3}$  are the edges  $\overrightarrow{a_j a_{n/3}}$  for  $j = (n/3) - 1, \dots, (n/3) - n - 2k - 2$ . Now, for  $j + 1 \leq n/3$ , once all the directed edges arriving at  $a_{j+1}$  have been determined, fix that (all) the directed edges arriving at  $a_j$  are  $\overrightarrow{a_\ell a_j}$ , for all those  $\ell$  that satisfy  $j - \ell \leq n - 2k - 1 + [a_j]^+$ .

Since no digraph in  $\mathcal{D}_{k,n}$  has more edges than  $D_{k,n}$ , to finish the proof it suffices to bound the number of edges of  $D_{k,n}$ . This is the content of Claim 8 below.

**Claim 8**  *$D_{k,n}$  has at most  $\binom{n/3}{2} - (Y(k, n) - 3\binom{n/3+1}{2} - (k - n/3)n)/3$  edges.*

*Sketch of proof.* Since we know the exact indegree of each vertex in  $D_{k,n}$ , we know the exact number of edges of  $D_{k,n}$ , and so the proof of Claim 8 is no more than a straightforward, but quite long and tedious, calculation. ■

**Corollary 9** *If  $\Pi$  is a 3-decomposable  $n$ -halfperiod, and  $n/3 < k < n/2$ , then each of  $D^{bb}$  and  $D^{cc}$  has at most  $\binom{n/3}{2} - (1/3) \left( Y(k, n) - 3\binom{n/3+1}{2} - (k - n/3)n \right)$  edges.*

*Proof.* In the proof of Proposition 7, the only relevant property about  $D^{aa}$  is that the  $a$ 's form a set of  $n/3$  points that in some permutation of  $\Pi$  (namely  $\pi_0$ ) appear all consecutively and at the beginning of the permutation. Since  $\Pi$  is 3-decomposable, this condition is also satisfied by the set of  $b$ 's and by the set of  $c$ 's. ■

We now summarize the results in the current subsection.

**Proposition 10** *If  $\Pi$  is a 3-decomposable  $n$ -halfperiod, and  $n/3 < k < n/2$ , then*

$$\eta_{\leq k}^{\text{hom}}(\Pi) \geq Y(k, n) - 3\binom{n/3+1}{2} - (k - n/3)n.$$

*Proof.* By Remark 6, the number  $\eta_{>k}^{aa}(\Pi)$  of  $aa$ -transpositions that are *not* ( $\leq k$ )-critical equals the number of edges in  $D^{aa}$ , which by Proposition 7 is at most  $(1/3) \left( Y(k, n) - 3\binom{n/3+1}{2} - (k - n/3)n \right)$ . Since the total number of  $aa$ -transpositions is  $\binom{n/3}{2}$ , then the number of  $aa$ -transpositions that contribute to  $\eta_{\leq k}(\Pi)$  is at least  $\binom{n/3}{2} - \left( \binom{n/3}{2} - (1/3) \left( Y(k, n) - 3\binom{n/3+1}{2} - (k - n/3)n \right) \right) = (1/3) \left( Y(k, n) - 3\binom{n/3+1}{2} - (k - n/3)n \right)$ . By Corollary 9,  $bb$ - and  $cc$ -transpositions contribute in at least the same amount to  $\eta_{\leq k}^{\text{hom}}(\Pi)$ , and so the claimed inequality follows. ■

### 3.2.3 Proof of Proposition 3

Proposition 3 follows immediately from Eq. (3) and Propositions 4 and 10. ■

## 4 Concluding remarks

All the lower bounds proved above remain true for point sets that satisfy conditions (i) and (ii) (and not necessarily condition (iii)) for 3-decomposability.

## References

- [1] B.M. Ábrego and S. Fernández–Merchant, A lower bound for the rectilinear crossing number, *Graphs and Comb.*, **21** (2005), 293–300.
- [2] B.M. Ábrego, J. Balogh, S. Fernández–Merchant, J. Leaños, and G. Salazar, An extended lower bound on the number of ( $\leq k$ )-edges to generalized configurations of points and the pseudolinear crossing number of  $K_n$ . Submitted (2007).
- [3] O. Aichholzer, J. García, D. Orden, and P. Ramos, New lower bounds for the number of ( $\leq k$ )-edges and the rectilinear crossing number of  $K_n$ , *Discr. Comput. Geom.*, to appear.
- [4] O. Aichholzer. On the rectilinear crossing number. Available online at <http://www.ist.tugraz.at/staff/aichholzer/crossings.html>.
- [5] J. Balogh and G. Salazar,  $k$ -sets, convex quadrilaterals, and the rectilinear crossing number of  $K_n$ , *Discr. Comput. Geom.* **35** (2006), 671–690.
- [6] P. Brass, W.O.J. Moser, and J. Pach, *Research Problems in Discrete Geometry*. Springer, New York (2005).
- [7] J. E. Goodman and R. Pollack, On the combinatorial classification of nondegenerate configurations in the plane, *J. Combin. Theory Ser. A* **29** (1980), 220–235.
- [8] L. Lovász, K. Vesztergombi, U. Wagner, and E. Welzl, Convex Quadrilaterals and  $k$ -Sets. *Towards a Theory of Geometric Graphs*, (J. Pach, ed.), Contemporary Math., AMS, 139–148 (2004).



## Appendix: Proof of Claim 8

Since  $D_{k,n}$  is a well-defined digraph, and we know the exact indegree of each of its vertices, Claim 8 is no more than long and tedious, yet elementary, calculation.

The purpose of this Appendix is to give the full details of this calculation.

We prove Claim 8 in two steps. First we obtain an expression for the *exact* value of the number of edges of  $D_{k,n}$ , and then we show that this exact value is upper bounded by the expression in Claim 8.

### 1 The exact number of edges in $D_{k,n}$

The exact number of edges in  $D_{k,n}$  is a function of the following parameters. Let  $i, j$  be positive integers with  $i \leq j$ . Then:

- $b(i, j)$  is the (unique) nonnegative integer such that  $\binom{b(i,j)+1}{2} < j/i \leq \binom{b(i,j)+2}{2}$ ; and
- $q(i, j)$  and  $r(i, j)$  are the (unique) integers satisfying  $0 \leq q(i, j) < i, 1 \leq r(i, j) \leq b(i, j) + 1$  and such that

$$j = i \binom{b(i,j)+1}{2} + q(i, j)(b(i, j) + 1) + r(i, j) \quad (\text{A-1})$$

For brevity, in the rest of the section we let  $s := n/3$  and  $m := n - 2k - 1$ .

The key observation is that we know the indegree of each vertex in  $D_{k,n}$ :

**Proposition 1 (Proposition 19 in [5])** *For each integer  $1 \leq i \leq s$ , and each vertex  $a_i$  of  $D_{k,n}$ ,  $[a_i]^- = b(i, s)m + q(i, s)$ .*

The number of edges of  $D_{k,n}$  equals the sum of the indegrees over all vertices in  $D_{k,n}$ . Thus our main task is to find a closed expression for the sum  $\sum_{1 \leq i \leq s} [a_i]^-$ . This is the content of our next statement.

**Proposition 2 (Exact number of edges of  $D_{k,n}$ )** *The number  $\sum_{1 \leq i \leq s} [a_i]^-$  of edges of  $D_{k,n}$  is*

$$\begin{aligned} E(k, n) := & 2m^2 \binom{b(m, s) + 1}{3} + \binom{b(m, s) + 1}{2} \binom{m}{2} + 2m \cdot q(m, s) \binom{b(m, s) + 1}{2} + \\ & \binom{q(m, s)}{2} \binom{b(m, s) + 1}{1} + r(m, s) (m \cdot b(m, s) + q(m, s)). \end{aligned}$$

*Proof.* We break the index set of the summation  $\sum_{1 \leq i \leq s} [a_i]^-$  into three parts, in terms of  $\alpha := m \binom{b(m,s)+1}{2}$  and  $\beta := q(m, s)(b(m, s) + 1)$ . We let  $A = \sum_{1 \leq i \leq \alpha} [a_i]^-$ ,  $B = \sum_{\alpha+1 \leq i \leq \alpha+\beta} [a_i]^-$ , and  $C = \sum_{\alpha+\beta+1 \leq i \leq s} [a_i]^-$  so that

$$\sum_{1 \leq i \leq s} [a_i]^- = A + B + C. \quad (\text{A-2})$$

We calculate each of  $A$ ,  $B$ , and  $C$  separately.

*Calculating  $A$*

If  $\ell, j$  are integers such that  $0 \leq \ell \leq m - 1$  and  $0 \leq j \leq b(m, s)$ , we define  $S_j := \{i : b(i, m) = j\}$  and  $T_{j,\ell} := \{i : b(i, m) = j, q(i, m) = \ell\}$ . Note that  $S_1, S_2, \dots, S_{b(m,s)}$  is a

partition of  $\{1, 2, \dots, n\}$  and that for each  $j \leq b(m, s) - 1$ ,  $T_{j,0}, T_{j,1}, \dots, T_{j,m-1}$  is a partition of  $S_j$ .

Note that  $A$  can be rewritten as  $\sum_{0 \leq j \leq b(m,s)-1} \sum_{i \in S_j} [a_i]^-$ . By Proposition 1 this equals  $\sum_{0 \leq j \leq b(m,s)-1} \sum_{i \in S_j} (m \cdot b(i, m) + q(i, m))$ . That is,

$$A = \sum_{0 \leq j \leq b(m,s)-1} \left( m \sum_{i \in S_j} b(i, m) + \sum_{i \in S_j} q(i, m) \right).$$

Since  $0 \leq q(i, m) \leq m - 1$  for all  $i$ , and  $T_{j,0}, T_{j,1}, \dots, T_{j,m-1}$  is a partition of  $S_j$ , then  $\sum_{i \in S_j} q(i, m) = \sum_{0 \leq l \leq m-1} \sum_{i \in T_{j,l}} q(i, m)$ . Thus,

$$A = \sum_{0 \leq j \leq b(m,s)-1} \left( m \sum_{i \in S_j} b(i, m) + \sum_{0 \leq l \leq m-1} \sum_{i \in T_{j,l}} q(i, m) \right). \quad (\text{A-3})$$

On other hand, for  $0 \leq j \leq b(m, s) - 1$  and  $0 \leq \ell \leq m - 1$ , it is not difficult to verify that  $|T_{j,\ell}| = j + 1$ . This implies that  $|S_j| = m(j + 1)$ .

By definition of  $S_j$  we have

$$\sum_{i \in S_j} b(i, m) = \sum_{i \in S_j} j = j |S_j| = jm(j + 1). \quad (\text{A-4})$$

By definition of  $T_{j,\ell}$  we have

$$\sum_{i \in T_{j,\ell}} q(i, m) = \sum_{i \in T_{j,\ell}} \ell = \ell |T_{j,\ell}| = \ell(j + 1). \quad (\text{A-5})$$

Substituting (A-4) and (A-5) into (A-3) we obtain

$$\begin{aligned} A &= \sum_{0 \leq j \leq b(m,s)-1} \left( m(jm(j + 1)) + \sum_{0 \leq \ell \leq m-1} \ell(j + 1) \right) \\ &= \sum_{0 \leq j \leq b(m,s)-1} \left( 2m^2 \binom{j+1}{2} + (j + 1) \sum_{0 \leq \ell \leq m-1} \ell \right) \\ &= \sum_{0 \leq j \leq b(m,s)-1} \left( 2m^2 \binom{j+1}{2} + (j + 1) \binom{m}{2} \right) \\ &= 2m^2 \binom{b(m,s)+1}{3} + \binom{b(m,s)+1}{2} \binom{m}{2}. \end{aligned} \quad (\text{A-6})$$

### Calculating $B$

Since  $b(i, m) = b(m, s)$  for each  $i \geq \alpha + 1$ , and  $[a_i]^- = m \cdot b(i, m) + q(i, m)$ , then  $B = \sum_{\alpha+1 \leq i \leq \alpha+\beta} [a_i]^- = \sum_{\alpha+1 \leq i \leq \alpha+\beta} (m \cdot b(i, m) + q(i, m))$ .

Therefore

$$\begin{aligned}
B &= \sum_{\alpha+1 \leq i \leq \alpha+\beta} m \cdot b(m, s) + \sum_{\alpha+1 \leq i \leq \alpha+\beta} q(i, m) \\
&= m \cdot b(m, s) \sum_{\alpha+1 \leq i \leq \alpha+\beta} 1 + \sum_{\alpha+1 \leq i \leq \alpha+\beta} q(i, m) \\
&= m \cdot b(m, s)\beta + \sum_{\alpha+1 \leq i \leq \alpha+\beta} q(i, m) \\
&= m \cdot b(m, s)q(m, s)(b(m, s) + 1) + \sum_{\alpha+1 \leq i \leq \alpha+\beta} q(i, m).
\end{aligned}$$

On other hand it is easy to check that  $|T_{b(m,s),k}| = b(m, s) + 1$  for every  $k$  such that  $0 \leq k \leq q(m, s) - 1$ . Since  $0 \leq q(i, m) \leq q(m, s) - 1$  for every  $i$  such that  $\alpha + 1 \leq i \leq \alpha + \beta$ , then  $T_{b(m,s),0}, T_{b(m,s),1}, \dots, T_{b(m,s),q(m,s)-1}$  is a partition of  $\{\alpha + 1, \alpha + 2, \dots, \alpha + \beta\}$ . Thus,

$$\sum_{\alpha+1 \leq i \leq \alpha+\beta} q(m, s) = \sum_{0 \leq \ell \leq q(m,s)-1} \sum_{i \in T_{b(m,s),\ell}} q(i, m). \quad (\text{A-7})$$

We note that  $\sum_{i \in T_{b(m,s),\ell}} q(i, m) = \ell |T_{b(m,s),k}| = \ell(b(m, s) + 1)$ . Using this fact in (A-7) we obtain

$$\sum_{\alpha+1 \leq i \leq \alpha+\beta} q(i, m) = \sum_{0 \leq \ell \leq q(m,s)-1} \ell(b(m, s) + 1) = \binom{q(m,s)}{2}(b(m, s) + 1).$$

Thus,

$$\begin{aligned}
B &= m \cdot b(m, s)q(m, s)(b(m, s) + 1) + \binom{q(m,s)}{2}(b(m, s) + 1) \\
&= 2m \cdot q(m, s) \binom{b(m,s)+1}{2} + \binom{q(m,s)}{2}(b(m, s) + 1). \quad (\text{A-8})
\end{aligned}$$

### Calculating $C$

Since  $b(i, m) = b(m, s)$ ;  $q(i, m) = q(m, s)$  for each  $i$  such that  $i \geq \alpha + \beta + 1$ ; and  $[a_i]^- = m \cdot b(i, m) + q(i, m)$ , it follows that

$$\begin{aligned}
C &= \sum_{\alpha+\beta+1 \leq i \leq s} [a_i]^- = \sum_{\alpha+\beta+1 \leq i \leq s} m \cdot b(i, m) + q(i, m) \\
&= \sum_{\alpha+\beta+1 \leq i \leq s} m \cdot b(m, s) + q(m, s) \\
&= (s - \alpha - \beta) \left( m \cdot b(m, s) + q(m, s) \right)
\end{aligned}$$

From (A-1) it follows that  $r(m, s) = (s - \alpha - \beta)$ , and so

$$C = r(m, s)(m \cdot b(m, s) + q(m, s)). \quad (\text{A-9})$$

Now from (A-6), (A-8), and (A-9), it follows that  $E(k, n) = A + B + C$ , and so Proposition 2 follows from (A-2). ■

## 2 Upper bound for number of edges in $D_{k,n}$ : Proof of Claim 8

First we bound the number of  $(\leq k)$ -edges in 3-decomposable  $n$ -halfperiods in terms of the expression  $E(k, n)$  in Proposition 2.

**Proposition 3** *Let  $\Pi$  be a 3-decomposable  $n$ -halfperiod, and let  $k < n/2$ . Then*

$$\eta_{\leq k}(\Pi) \geq \begin{cases} 3\binom{k+1}{2} & \text{if } k \leq n/3, \\ 3\binom{n/3+1}{2} + (k - n/3)n + 3\left(\binom{n/3}{2} - E(k, n)\right) & \text{if } n/3 < k < n/2. \end{cases}$$

*Proof.* Obviously,  $\eta_{\leq k}(\Pi) \geq \eta_{\leq k}^{\text{het}}(\Pi)$  and so the case  $k \leq n/3$  follows from Proposition 4. Now suppose that  $n/3 < k < n/2$ . Recall that  $\eta_{\leq k}^{\text{hom}}(\Pi) = \eta_{>k}^{aa}(\Pi) + \eta_{>k}^{bb}(\Pi) + \eta_{>k}^{cc}(\Pi)$ . Now the total number of  $aa$ - (and  $bb$ -, and  $cc$ -) transpositions is exactly  $\binom{n/3}{2}$ , and so  $\eta_{\leq k}^{\text{hom}}(\Pi) = 3\binom{n/3}{2} - \eta_{>k}^{aa}(\Pi) - \eta_{>k}^{bb}(\Pi) - \eta_{>k}^{cc}(\Pi)$ . Thus it follows from Remark 6 and Proposition 2 that  $\eta_{\leq k}^{\text{hom}}(\Pi) \geq 3\left(\binom{n/3}{2} - E(k, n)\right)$ . This fact, together with Proposition 4, implies that  $\eta_{\leq k}(\Pi) = \eta_{\leq k}^{\text{het}}(\Pi) + \eta_{\leq k}^{\text{hom}}(\Pi) \geq 3\binom{n/3+1}{2} + (k - n/3)n + 3\left(\binom{n/3}{2} - E(k, n)\right)$ , as claimed. ■

Must explain that for  $bb$  and  $cc$  same digraph works

**Proof of Claim 8.** Recall that  $s := n/3$  and  $m := n - 2k - 1$ . By Remark 5 we know that  $k > n/3$ , and so  $s \geq m$ . From (A-1), it follows that

$$q(m, s) = \frac{s - m\binom{b(m,s)+1}{2} - r(m, s)}{b(m, s) + 1}. \quad (\text{A-10})$$

Now by Proposition 3,  $\eta_{\leq k}(\Pi) \geq L(k, n)$ , where

$$L(k, n) := 3\binom{s+1}{2} + (k - s)n + 3\left(\binom{s}{2} - E(k, n)\right). \quad (\text{A-11})$$

Substituting in  $E(k, n)$  the value of  $q(m, s)$  given in (A-10), a (long and tedious yet) totally elementary simplification yields

$$L(n, k) - Y(k, n) - 1/3 = \frac{1}{8(b(m, s) + 1)} \left( 5b(m, s)^2 + 4b(m, s)^3 + b(m, s)^4 + b(m, s)(-12r(m, s) + 2) + 12(r(m, s) - 1)r(m, s) \right).$$

Define  $f(b(m, s), r(m, s)) := L(n, k) - Y(k, n) - 1/3$ . An elementary calculation shows that  $r_0(m, s) := (b(m, s) + 1)/2$  minimizes  $f(b(m, s), r(m, s))$ . Thus  $f(b(m, s), r(m, s)) \geq f(b(m, s), r_0(m, s)) = (b(m, s) + 3)(b(m, s) + 1)(b(m, s) - 1)/8$ . Since  $b(m, s)$  is a nonnegative integer, it follows that  $f(b(m, s), r(m, s)) \geq -1/3$  and therefore  $L(n, k) - Y(k, n) \geq 0$ .

By (A-11),  $E(k, n) = (1/3)\left(3\binom{s+1}{2} + (k - s)n + 3\binom{s}{2} - L(k, n)\right)$ . Since  $-L(n, k) \leq -Y(k, n)$ , then  $E(k, n) \leq (1/3)\left(3\binom{s+1}{2} + (k - s)n + 3\binom{s}{2} - Y(k, n)\right) = \binom{s}{2} - (1/3)(Y(k, n) - 3\binom{s+1}{2} - (k - s)n)$ . This proves Claim 8, since  $E(k, n)$  is the total number of edges in  $\mathcal{D}_{k,n}$ . ■