

LETTER TO THE EDITOR

The seeds of star formation in the filamentary infrared-dark cloud G011.11–0.12^{*}

Th. Henning, H. Linz, O. Krause, S. Ragan, H. Beuther, R. Launhardt, M. Nielbock, and T. Vasyunina

Max-Planck-Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany
e-mail: [henning, linz, krause, ragan, beuther, rl, nielbock, vasyunina]@mpia.de

Received 31 March 2010 ; accepted 28 April 2010

ABSTRACT

Context. Infrared-dark clouds (IRDCs) are the precursors to massive stars and stellar clusters. G011.11–0.12 is a well-studied filamentary IRDC, though, to date, the absence of far-infrared data with sufficient spatial resolution has limited the understanding of the structure and star-formation activity.

Aims. We use *Herschel* to study the embedded population of young pre- and protostellar cores in this IRDC.

Methods. We examine the cloud structure, which appears in absorption at short wavelength and in emission at longer wavelength. We derive the properties of the massive cores from the spectral energy distributions of bright far-infrared point sources detected with the PACS instrument aboard *Herschel*.

Results. We report on the detection and characterization of pre- and protostellar cores in a massive filamentary infrared-dark cloud G011.11–0.12 using PACS. We characterize 18 cores directly associated with the filament, two of which have masses over $50 M_{\odot}$, making them the best candidates to become massive stars in G011.11–0.12. These cores are likely at various stages of protostar formation, showing elevated temperature ($\langle T \rangle \sim 22$ K) with respect to the ambient gas reservoir. The core masses ($\langle M \rangle \sim 24 M_{\odot}$) are small compared to that in the cold filament. The mean core separation is 0.9 pc, well in excess of the Jeans length in the filament.

Conclusions. We confirm that star formation in IRDCs is underway and diverse, and IRDCs have the capability of forming massive stars and clusters.

Key words. stars: formation - stars: protostars - techniques: photometric - ISM: individual: G011.11-0.12

1. Introduction

Star formation occurs in cold, dense molecular cores. In local low-mass star-forming regions, this process takes place in relatively isolated environments and is better understood compared to the distant regions in which massive stars form. Specifically, it is the identification of the cold precursors to stars - the so-called “pre-stellar cores” that has led to a greater understanding of the initial conditions of low-mass star formation; however, the high-mass counterpart has been more difficult to identify. Thus it is not clearly understood how the early stages of massive star formation compare to that of low-mass stars.

Infrared-dark clouds (IRDCs) are complexes of cold ($T < 20$ K), dense ($n > 10^4 \text{ cm}^{-3}$) molecular gas, now believed to be the precursors to massive stars and star clusters (e.g. Rathborne et al. 2006), although there is diversity among this class of objects. Early observations revealed that the cold dust in IRDCs absorbs the bright Galactic background emission at $8 \mu\text{m}$ (Egan et al. 1998; Hennebelle et al. 2001) and emits at sub-mm wavelengths (e.g. Carey et al. 2000). Later, high-resolution studies found that IRDCs exhibit significant substructure (e.g. Butler & Tan 2009; Ragan et al. 2009), and (sub-)millimeter studies show that they host massive embedded star formation (e.g. Rathborne et al. 2007; Beuther & Steinacker 2007; Hennemann et al. 2009).

The spectral energy distribution (SED) of massive embedded protostars is expected to peak in the far-infrared but is usually very faint in the mid-infrared. The coldest cores are not readily detectable with *Spitzer* because of sensitivity limitations at $24 \mu\text{m}$ and confusion at longer wavelengths due to limited spatial resolution. With *Herschel* (Pilbratt et al. 2010), we now have the spatial resolution and the capability to perform large-area maps to search for these deeply embedded objects and to characterize the SEDs.

The large, filamentary IRDC G011.11–0.12 was selected as a target in the Early Phases of Star Formation (EPoS) *Herschel* key program (PI: O. Krause). Together with complementary *Spitzer* data, we use PACS (Poglitsch et al. 2010) and SPIRE (Griffin et al. 2010) observations to constrain the properties of the massive embedded cores within this IRDC.

Due to its high absorbing contrast against the Galactic background emission (particularly at $8 \mu\text{m}$), the IRDC G011.11–0.12 ($d = 3.6$ kpc) has been subject to detailed studies. Carey et al. (1998) confirm the presence of dense molecular gas based on H_2CO observations. Carey et al. (2000) report 850 and $450 \mu\text{m}$ continuum observations of the thermal dust emission which precisely follows the structure seen in absorption in the mid-infrared, and the column density reaches up to 10^{23} cm^{-2} (see also Parsons et al. 2009). The so-called P1 position is seen in emission in both 8 and $850 \mu\text{m}$ images (Fig. 1), a likely indicator of embedded star formation.

Pillai et al. (2006a) observe G011.11–0.12 as part of an ammonia survey, including the (1,1) and (2,2) inversion transitions with $40''$ spatial resolution. The masses of four sub-

^{*} *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. It is open for proposals for observing time from the worldwide astronomical community.




Fig. 1. The IRDC G011.11–0.12 from *Spitzer* to SCUBA wavelengths. (a) *Spitzer* IRAC $8\mu\text{m}$ image. (b) *Spitzer* MIPS $24\mu\text{m}$ image. The peak SCUBA positions (P1, P2,...) as assigned by Johnstone et al. (2003) are indicated. (c) *Herschel* PACS $70\mu\text{m}$ image. (d) *Herschel* PACS $100\mu\text{m}$ image with the positions of analyzed sources (see Sect. 4) indicated. Sources with no $24\mu\text{m}$ counterpart are circled in cyan, sources with $24\mu\text{m}$ counterparts are circled in green. (e) *Herschel* PACS $160\mu\text{m}$ image. (f) *Herschel* SPIRE $250\mu\text{m}$ image. (g) *Herschel* SPIRE $350\mu\text{m}$ image. (h) $8\mu\text{m}$ extinction map (derived from *Spitzer* data) with (blue) SCUBA $850\mu\text{m}$ continuum contours (levels are $0.4\ 0.8\ 1.2\ \text{Jy beam}^{-1}$). Source labeling same as in panel (d). The approximate beam size is indicated in the lower-left corner of each panel.

regions (P1, P2, P6, and P7 in the Johnstone et al. (2003) nomenclature) were found to be several hundred M_{\odot} each, though other mass-tracing techniques find higher values for the tot

Table 1. Photometric Properties of Protostars

Src. Num.	RA (J2000)	Dec. (J2000)	Temp. (K)	Lum. (L_{\odot})	Mass (M_{\odot})	Notes
1	18:10:47.0	-19:10:09	21	6	3	
2	18:10:40.3	-19:11:34	23	26	9	P3, e
3	18:10:39.9	-19:16:39	18	5	5	
4	18:10:38.8	-19:19:01	24	8	1	e
5	18:10:38.2	-19:19:50	25	66	10	e
6	18:10:45.9	-19:20:46	16	15	40	
7	18:10:46.3	-19:21:46	20	6	4	
8	18:10:33.7	-19:22:08	21	25	9	P2, e
9	18:10:28.4	-19:22:29	24	1346	240	P1, e
10	18:10:03.8	-19:23:06	23	27	7	e
11	18:10:16.4	-19:24:19	19	9	7	
12	18:10:19.4	-19:24:24	21	11	5	e
13	18:10:09.2	-19:25:55	26	23	2	P2, e
14	18:10:06.9	-19:26:00	20	9	5	
15	18:10:05.5	-19:26:34	24	58	11	P7, e
16	18:10:04.7	-19:27:43	24	25	5	P6, e
17	18:10:04.0	-19:28:16	23	28	6	P6
18	18:10:07.4	-19:29:15	23	139	82	P6, e
19	18:10:25.1	-19:17:49	24	478	79	e
20	18:10:27.1	-19:18:12	23	45	11	e
21	18:10:07.7	-19:18:39	21	19	9	
22	18:10:42.7	-19:19:58	21	11	5	e
23	18:10:03.8	-19:21:04	20	8	5	e
24	18:10:13.8	-19:22:04	17	9	15	e


Notes – “P” labels indicate spatial correspondence to regions identified by Johnstone et al. (2003). “e” indicates where we infer the presence of an embedded protostar based on a $24 \mu\text{m}$ detection. Objects above the horizontal line are located “on” the IRDC filament; sources below are “off.”

3. Results and discussion

In *Spitzer* images, G011.11–0.12 appears mostly dark at both 8 and $24 \mu\text{m}$, as shown in Fig. 1, though there are some indications of point sources at $24 \mu\text{m}$. However, the nature of these objects remains undefined because they lack counterparts at shorter wavelengths. The IRDC filament is dark in the PACS 70 and $100 \mu\text{m}$ filter images, but shows little evidence of absorption at $160 \mu\text{m}$. In the continuum maps, starting with $250 \mu\text{m}$ SPIRE map to the SCUBA $850 \mu\text{m}$ map (Di Francesco et al. 2008), shown in contours in Fig. 1, the extinction filament seen at shorter wavelengths now appears in emission.

3.1. Characterization of embedded cores

Herschel provides both the ability to make large-scale maps and the spatial resolution to isolate point sources from bright background emission for the first time. At the given scales between 18000 and 43000 AU (our resolution limit from the 5 to $12''$ PACS beam), we probe at the size scale of “cores,” as defined by Bergin & Tafalla (2007). We detect cores



- Parsons, H., Thompson, M. A., & Chrysostomou, A. 2009, MNRAS, 399, 1506
- Pilbratt, M., , I., Author, I.-I., et al. 2010, A&A, this volume
- Pillai, T., Wyrowski, F., Carey, S. J., & Menten, K. M. 2006a, A&A, 450, 569
- Pillai, T., Wyrowski, F., Menten, K. M., & Krügel, E. 2006b, A&A, 447, 929
- Poglitsch, M., , I., Author, I.-I., et al. 2010, A&A, this volume
- Ragan, S. E., Bergin, E. A., & Gutermuth, R. A. 2009, ApJ, 698, 324
- Rathborne, J. M., Jackson, J. M., & Simon, R. 2006, ApJ, 641, 389
- Rathborne, J. M., Simon, R., & Jackson, J. M. 2007, ApJ, 662, 1082