

Stability Constraints on Classical de Sitter Vacua

Gary Shiu^{1,2} and Yoske Sumitomo³

¹ Department of Physics, University of Wisconsin, Madison, WI 53706, USA

² Institute for Advanced Study, Hong Kong University of Science and Technology, Hong Kong

³ Tata Institute of Fundamental Research, Mumbai 400005, India

Email: shiu at physics.wisc.edu, yoskesumitomo at gmail.com

Abstract

We present further no-go theorems for classical de Sitter vacua in Type II string theory, i.e. de Sitter constructions that do not invoke non-perturbative effects or explicit supersymmetry-breaking localized sources. By analyzing the stability of the 4D potential arising from compactification on manifolds with curvature fluxes and orientifold planes, we found that additional ingredients beyond the minimal ones presented so far are necessary to avoid the presence of unstable modes. We enumerate the minimal setups for (meta-)stable de Sitter vacua to arise in this context.

Contents

1	Introduction	1
2	No-go Theorems for de Sitter Extrema	3
3	No-go Theorems for the Stability of de Sitter Extrema	5
4	Evading the no-goes	8
5	Discussions	9
A	Instability Analysis for Other Examples	11
A	R_6, H_3, F_0 , O4–O6 in IIA	
A	R_6, H_3, F_1, F_3 , O in IIB	
A	R_6, H_3, F_1 , O–O5 in IIB	
A 4	R_6, H_3, F_5 , O–O5 in IIB	
A 5	R_6, H_3, F_1 , O–O7 in IIB	
A 6	R_6, H_3, F_3 , O–O7 in IIB	
A 7	R_6, H_3, F_5 , O–O7 in IIB	

1 Introduction

The observational evidence for an accelerating universe adds another serious wrinkle to the already vexing issue of moduli stabilization in string theory. Besides having to stabilize a myriad of moduli self consistently at values that give physically acceptable couplings, requiring the vacuum solution to have positive energy introduces a new layer of complications. Unlike their anti de Sitter and Minkowskian counterparts, de Sitter solutions are much harder to construct as they are not amenable to the powerful tools of supersymmetry. Furthermore, the de Sitter solutions one might hope to obtain from a string compactification are at best metastable. There exist generically supersymmetric vacua in the decompactified limit and any candidate de Sitter vacua are subject to all kinds of perturbative and non perturbative instabilities.

In the past decade, motivated partly by the increasing observational support for dark energy, various proposals for constructing metastable de Sitter vacua from string theory have been suggested. These proposals can be broadly divided into two types: those that hinges on non perturbative effects and/or explicit SUSY breaking localized sources (most notably fluxes and variations thereof) and those that do not [6]. We refer to the latter as *classical de Sitter solutions* as their constructions involve only classical ingredients such as internal curvature, fluxes and orientifold planes. In contrast to non perturbative effects in string compactifications which

are difficult to compute in full detail the simplicity of these classical de Sitter solutions allows for *explicit* models to be constructed. Among these models some of them can be shown to be genuine solutions of the D equations of motion as well as critical points of the dimensionally reduced theory. The classical de Sitter solutions explored so far are admittedly far from realistic e.g. the Standard Model sector has not yet been implemented and an exponentially small cosmological constant is difficult to achieve. Nonetheless they serve to illustrate the general issues one may encounter in constructing explicit de Sitter vacua from string theory. Furthermore explicit constructions of such solutions though not fully realistic may shed light on conceptual issues of de Sitter space such as a microscopic understanding of its entropy and holography etc.

If one may draw hints from the aforementioned attempts in constructing explicit de Sitter vacua a recurrent lesson seems to be the ubiquity of tachyons. While anti de Sitter flux vacua are abundant [7–1] de Sitter solutions are hard to come by. Even if de Sitter extrema are found they are plagued with one or more unstable modes. Similar searches for de Sitter vacua within 4D supergravity also seems to suggest that tachyonic modes are omnipresent [2–4] though the tachyons found in explicit reductions of D backgrounds (e.g. [6–5] are not necessarily in the “sGoldstino” direction [6]. Taking cues from these earlier attempts we set out to prove some no go theorems for the existence of stable de Sitter vacua. Of course no go theorems always come with assumptions and there are ways around them. Our work thus helps in sharpening the requirements needed for constructing explicit de Sitter vacua from string theory.

We found that some necessary constraints on the absence of tachyonic modes in classical de Sitter solutions can be stated in a surprisingly clean and simple way. From the scalings of various contributions to the potential with respect to the universal moduli (i.e. dilaton and the breathing mode) and upon simplifications around a positive potential extremum we analyzed the conditions under which the moduli mass matrix contains necessarily an unstable mode or a flat direction upon diagonalization. The conditions we found supplemented with earlier no go theorems on the existence of de Sitter extrema [4–5–6] therefore provide us with a more refined guide to search for (meta) stable de Sitter vacua in classical supergravity. As we shall see the conditions on the stability of de Sitter extrema are simple but yet powerful enough to show that the minimal setups evading the no goes in [4–5–6] turn out to all suffer from perturbative instabilities. To obtain stable de Sitter vacua additional ingredients (e.g. more types of fluxes and/or O planes) have to be introduced. Our approach further allows us to enumerate the *minimal* ingredients needed for constructing classical de Sitter vacua.

Although our focus is on finding classical vacua with positive cosmological constant in this paper we should mention here some recent related attempts in clarifying the possibilities of realizing a time varying dark energy in string theory. A shift symmetry is often invoked to

¹The non-perturbative (in)stability of AdS vacua were discussed in [18, 19]. See also the non-perturbative instability argument [20] for AdS_5 through a bubble of nothing.

prevent any undesired couplings between the heavy modes and the quintessence field. There are two strands of approaches to weakly break this shift symmetry in order to obtain a time varying dark energy: one with a non perturbative potential [5, 6] and the other with a classical potential from an NS5 brane [7] (based on [8]). The latter class of models are similar in spirit to the classical de Sitter vacua considered here as only perturbative ingredients are introduced.

This paper is organized as follows. In Section 2 we revisit the no go theorems for de Sitter extrema. Some form of these results were already obtained previously in the literature [4, 5, 6] but our approach is more systematic and suited for our subsequent discussions about the stability of these extrema. In Section 3 we generalize the setups in Section 2 and derive some new no go theorems for (meta) stable de Sitter vacua. We enumerate several “minimal” setups necessary to evade these no goes. We end with some discussions in Section 5. Some details are relegated to an appendix.

2 No-go Theorems for de Sitter Extrema

Here we analyze the conditions for de Sitter extrema to arise in classical supergravity (with localized sources). Though some forms of these results were previously obtained [4, 5, 6] our result here is more complete and our presentation will streamline our subsequent discussion in Section 3 on more complicated set ups and on the stability of classical de Sitter vacua.

Our analysis applies to both Type IIA and Type IIB string theories. Consider compactification to 4D with the following ansatz for the metric:

$$ds_{10}^2 = \tau^{-2} ds_4^2 + \rho ds_6^2 \quad (2.1)$$

where we took the Weyl factor to be $\tau = e^{-\phi} \rho^{3/2}$ such that the kinetic terms for the universal moduli ρ and τ in the 4D Einstein frame do not mix.

Various fluxes H_3, F_p localized q brane sources and the 6D curvature contribute to the 4D potential in some specific way:

$$V_{H_3} = A_{H_3} \tau^{-2} \rho^{-3}, \quad V_{F_p} = A_{F_p} \tau^{-4} \rho^{3-p}, \quad V_q = A_q \tau^{-3} \rho^{(q-6)/2}, \quad V_{R_6} = A_{R_6} \tau^{-2} \rho^{-1}. \quad (2.2)$$

The coefficients A_{H_3} and A_{F_q} of the flux potentials are defined to be positive while the coefficients A_q of the p branes contributions (including D branes and O planes) and A_{R_6} for the curvature contribution can be either positive or negative. Note that all these potentials go to zero when we take $\tau \rightarrow \infty$ while keeping the others finite. Therefore there always exist a Minkowski vacuum asymptotically.

Here we follow more closely the discussions in [6] which analyze the conditions for a de Sitter extremum to arise in this context. The idea is that if we can find an inequality of the

following form:

$$\begin{aligned} D &\equiv -a\tau\partial_\tau - b\rho\partial_\rho, \\ DV &\geq cV, \end{aligned} \quad ($$

with non trivial real constants a, b and $c > 0$ then a positive energy extremum of the potential is excluded. To evade this no go one can enumerate a set of minimal ingredients needed

As a simple example let's consider an effective 4D potential receiving contributions from R_6, H_3 and two more components taken from the set of F_p, O_q . D branes typically introduce additional open string moduli and thus as a first pass we do not include them for simplicity. Here we analyze all possibilities which can evade the condition (4.1) and then confirm that all candidates of this type were tabulated in Table 6. From now on we restrict ourselves to the following ingredients for simplicity:

$$\begin{aligned} &F_0, F_2, F_4, F_6, O4, O6 \text{ in IIA} \\ &F_1, F_3, F_5, O_-, O5, O7 \text{ in IIB} \end{aligned} \quad (4.2)$$

More complicated setups give us more examples some excluded by the no goes and some evading them. However since we will find examples of each type within this setup we will concentrate on the above limited components.

First we focus on the case with R_6, H_3, F_p, O_q (same as in Table 6). If the following conditions are satisfied:

$$\begin{aligned} DV &= (a - b)V - bV_{H_3} - (a - (p - 4)b)V_{F_p} - \left(a - \left(-\frac{q}{2}\right)b\right)V_q, \\ a - b &> 0, \quad b \geq 0, \quad a - (p - 4)b \geq 0, \quad a - \left(-\frac{q}{2}\right)b \leq 0, \end{aligned} \quad (4.3)$$

we see that $DV \geq (a - b)V$ and this leads to a no go theorem for de Sitter extrema. However the inequality can be violated if we have the following sources:

$$\begin{aligned} \text{IIA} : & (F_0, O4), (F_0, O6), (F_2, O4), \\ \text{IIB} : & (F_1, O_-), (F_1, O5), (F_3, O_-), (F_5, O_-) \end{aligned} \quad (4.4)$$

with suitable sign of R_6 for each case which we omit here but will be clarified later. A nontrivial H_3 is required in some cases but we do not specify the details here.

Upon a similar analysis but with $R_6, H_3, F_{p_1}, F_{p_2}$ and also $R_6, H_3, O_{q_1}, O_{q_2}$ we are left with two additional cases i.e. $(O_-, O5), (O_-, O7)$ which can potentially evade the no go. However a more detailed analysis showed that these two cases can evade the no go only if the number of $O5$ (respectively $O7$) is zero. Therefore we have in total 7 cases listed in Table 6 where the no go does not apply. Note that evading the no go here only means that the 4D potential can admit a de Sitter extremum but does not guarantee the stability of such extremum.

Now if we include the contributions to the potential from the O8–O9 planes the constraints for evading the no goes leave us only with (O–O9). However again this situation is possible only when the number of O9 is zero. Thus we can conclude that the result obtained in (6) is most general with two additional ingredients beyond R_6 and H_3 .

In the next section we will proceed further to analyze the stability of the candidate de Sitter extrema. Although the ingredients presented in (6) are minimal in terms of evading the no go (for de Sitter extrema) all these minimal scenarios turn out to give only unstable extrema. Since the more complicated setups we consider in the next section subsume the simpler cases enumerated here we will relegate our discussion of the no go for stability to the next section.

3 No-go Theorems for the Stability of de Sitter Extrema

We now generalize the setup in the previous section to include three additional components beyond R_6 and H_3 and reanalyze the no go theorems for de Sitter extrema presented in (6). The three components are taken from the set: F_p, O_q in particular those listed in (4). Then evading the no go for extremal (6) leaves us with the following possibilities:

$$\begin{aligned} \text{IIA} : & (F_0, F_2, O4), (F_0, F_4, O4), (F_0, F_6, O4), (F_2, F_4, O4), (F_2, F_6, O4), \\ & (F_0, F_2, O6), (F_0, F_4, O6), (F_0, F_6, O6), (F_0, O4, O6), (F_2, O4, O6), \\ \text{IIB} : & (F_1, F_3, O), (F_1, F_5, O), (F_3, F_5, O), (F_1, F_3, O5), (F_1, F_5, O5), \\ & (F_1, O, O5), (F_1, O, O7), (F_3, O, O5), (F_3, O, O7), (F_5, O, O5), \\ & (F_5, O, O7), (F_1, O5, O7), \end{aligned} \quad (7)$$

We will analyze the stability of de Sitter extrema arising from all these cases. Again the sign of the 6D curvature should be chosen appropriately in each case and to avoid cluttering our discussion we omit such details here.

Let us examine the (in)stability of the candidates listed in (7). We restrict our analysis to the universal moduli subspace as the positivity of the mass matrix of this D subspace is a necessity condition for the absence of unstable modes in the full moduli space according to the *Sylvester's criterion* in linear algebras². The criterion can be stated as follows (see e.g. [10]):

An $N \times N$ Hermitian (e.g. real symmetric) matrix is positive definite if and only if the determinants of the upper left $n \times n$ submatrices ($n \leq N$) are all positive or more mathematically precise all of the leading principle minors are positive.

For instance let us apply this criterion first to the 6×6 mass matrix M of the (ρ, τ) subspace. The positivity of the determinant of M requires the diagonal components to be both positive

²We are grateful to Thomas van Riet, Timm Wrase, and especially Marco Zagermann for bringing our attention this point.

or both negative. In addition, the positivity of the upper left most component is also required (by Sylvester's criterion for M to be positive definite thus $\text{tr}(M) > 0$ and $\det(M) > 0$). Now applying Sylvester's criterion to the mass matrix of the full moduli space, we see that $\text{tr}(M) > 0$ together with $\det(M) > 0$ are necessary conditions for the full moduli mass matrix to be positive definite.

The eigenvalues of the two by two mass matrix $M \equiv \partial_{\rho_i} \partial_{\rho_j} V|_{\text{ext}}$ can be easily calculated:

$$\text{eigenvalues}(M) = - \left(\text{tr}(M) \pm \sqrt{(\text{tr}(M))^2 - 4 \det(M)} \right). \quad (3.10)$$

So a stable minimum can exist if both $\text{tr}(M) > 0$ and $\det(M) \leq (\text{tr}(M))^2/4$. We also consider the case in which we have a zero eigenvalue separately. In the following, we will show that in the minimal scenarios enumerated in (3.6), there is at least one tachyonic or flat direction. Furthermore, in the remainder of this section, we will show some no go examples where $\text{tr}(M) > 0$ and $\det(M) > 0$ cannot be simultaneously satisfied. Therefore, owing to the Sylvester's criterion, the full moduli mass matrix must necessarily contain at least an unstable mode.

We now analyze the stability of the setups enumerated in (3.6), which were already shown to evade the no go for de Sitter extrema. Since going through the details case by case is not very illuminating, we will work out one case in detail here and relegate the details of all other cases to appendix A. We will further enumerate in the next section the minimal ingredients for evading the *refined* no go for de Sitter vacua we found here.

Let's consider classical de Sitter solutions in Type IIA string theory with R_6, H_3, F_0, F_2 and O4, as these ingredients were shown to evade the no go for de Sitter extrema. As the critical point, one finds the following two constraints between the coefficients of the potential and the values of the moduli fields

$$A_{H_3} = - \frac{A_{R_6} \rho^2}{7} - \frac{A_{F_0} \rho^6}{7\tau^2} - \frac{A_{F_2} \rho^4}{\tau^2}, \quad A_4 = - \frac{4A_{R_6} \tau}{7} - \frac{8A_{F_0} \rho^4}{7\tau} - \frac{A_{F_2} \rho^2}{\tau}. \quad (3.11)$$

Usually, we solve for the moduli given the coefficients; here, however, we use the constraint equations to replace the coefficients with moduli fields instead. This does not mean that these coefficients are functions of the universal moduli, as these equalities only hold on shell. Upon substituting (3.11) into the potential at the extremum, we get:

$$V_{\text{ext}} = \frac{A_{F_0} \rho^3}{7\tau^4} - \frac{A_{R_6}}{7\rho\tau^2}. \quad (3.12)$$

Here we see that a positive A_{R_6} corresponding to a negatively curved manifold helps to obtain a positive extremum of the potential, otherwise A_{F_0} should be sufficiently large to compensate for the contribution from a positive (or zero) internal curvature.

We proceed to analyze the stability of the extremum following the argument around (3.12). Actually, we do not need to know the trace of the mass matrix in this example to show the no go

for its stability. Instead, we focus on the determinant at the extrema which can be written as:

$$\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) = \frac{4}{56\rho^4\tau^{10}} \left[(4A_{R_6}\tau^2 - 87A_{F_0}\rho^4 - 5A_{F_2}\rho^2)^2 - 5(5A_{F_2}^2\rho^4 - 8A_{F_0}A_{F_2}\rho^6 - 4A_{F_0}^2\rho^8) \right]. \quad (5)$$

Since we are interested in $V_{\text{ext}} > 0$, $A_{H_3} \geq 0$ we are limited to the range: $-A_{F_0}\rho^4 < A_{R_6}\tau^2 \leq 7A_{F_2}\rho^2 - A_{F_0}\rho^4$. From the explicit expression of the determinant and the above inequalities we found that the determinant is bounded from above by:

$$\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) < -\frac{6A_{F_0}^2\rho^4}{\tau^{10}}. \quad (6)$$

Therefore we see that the constraints $V_{\text{ext}} > 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $A_{H_3} \geq 0$ cannot be satisfied simultaneously regardless of the sign of R_6 .

Let us analyze the case in which $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) = 0$ while $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$ meaning that at least one eigenvalue of the mass matrix is zero. However we exclude the situations which can satisfy $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) > 0$ since then the examples have stable dS vacua in some region. Actually there is only one situation which have a zero eigenvalue not positive which is $(R_6, F_5, 0)$ without any other sources. In this case the potential can be rewritten as

$$V(\sigma) = \frac{A_{F_5}}{\sigma^4} - \frac{A_3}{\sigma^3} - \frac{A_{R_6}}{\sigma^2}, \quad (7)$$

with the definition $\sigma = \tau\rho^{1/2}$. Since the non trivial potential is generated only for the direction of σ we have one remaining flat direction defined by $\delta = \tau\rho^{-1/2}$ in 4D moduli space. Therefore one modulus field δ cannot be stabilized by the classical ingredients we consider.

Applying similar analysis to all the cases in (4) we found that Type II string theories in the presence of R_6, H_3 and only the following sources:

$$\begin{aligned} \text{IIA} : & (F_0, F_2, 0, 4), (F_0, 0, 4, 0, 6), \\ \text{IIB} : & (F_1, F_3, 0), (F_1, 0, 0, 5), (F_1, 0, 0, 7), (F_3, 0, 0, 7), (F_5, 0, 0, 5), (F_5, 0, 0, 7) \end{aligned} \quad (8)$$

do not lead to metastable de Sitter vacua. For the cases in Type IIA listed above in (8) $V_{\text{ext}} > 0$ and $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$ cannot be satisfied simultaneously. While in Type IIB we cannot satisfy $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$ with $V_{\text{ext}} > 0$ except for the case $(R_6, F_5, 0)$ in which we have one flat direction. Thus we found a new no go theorem which states that for the cases enumerated in (8) when $V_{\text{ext}} > 0$ there is at least one tachyonic or flat direction. We also see that the “minimal” setups (enumerated in (6) designed to evade the no go for de Sitter extrema turn out to be all unstable. Even if we generalized these “minimal” setups in (6) to include more contributions from fluxes and orientifold planes (see (3)) some of these generalized setups still forbid stable positive minima.

4 Evading the no-goes

Next we consider the cases in () other than the ones in (8) which were already excluded in the previous section based on stability. Among these many cases we would like to classify what kinds of sources are essential in evading the no go theorems for de Sitter vacua. Just like the previous section we will work out one case in detail and then simply provide a list of cases which can evade all the no goes which we found after a thorough analysis.

Let's consider the setup with R_6, H_3, F_0, F_2 O6. At the extremum the coefficients A_{H_3}, A_6 and the potential go like:

$$\begin{aligned} A_{H_3} &= -\frac{A_{R_6}\rho^2}{\tau^2} - \frac{A_{F_2}\rho^4}{\tau^2} - \frac{A_{F_0}\rho^6}{\tau^2}, \quad A_6 = -\frac{4A_{R_6}\tau}{9\rho} - \frac{4A_{F_2}\rho}{9\tau} - \frac{A_{F_0}\rho^3}{\tau}, \\ V_{\text{ext}} &= -\frac{A_{F_2}\rho}{9\tau^4} - \frac{A_{R_6}}{9\rho\tau^2}. \end{aligned} \quad (4)$$

Then we can analyze the stability of the de Sitter critical point from the trace and the determinant of the mass matrix at the extrema:

$$\begin{aligned} \text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= \frac{1}{\rho^3\tau^6} (5A_{F_2}\rho^4 - A_{F_0}\rho^6 - 6A_{F_2}\rho^2\tau^2 - A_{R_6}\rho^2\tau^2 - 7A_{F_0}\rho^4\tau^2 - A_{R_6}\tau^4), \\ \det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= \frac{4}{\rho^4\tau^{10}} (7A_{F_2}^2\rho^4 - A_{F_0}A_{F_2}\rho^6 - 9A_{F_2}A_{R_6}\rho^2\tau^2 - A_{F_0}A_{R_6}\rho^4\tau^2 - A_{R_6}^2\tau^4). \end{aligned} \quad (4)$$

To obtain the validity range for de Sitter minima we need to satisfy $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) > 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) > 0$ simultaneously with $V_{\text{ext}} > 0$, $A_{H_3} \geq 0$, $A_6 < 0$. After some simplifications these inequalities leave us with the following range of parameters:

$$\frac{A_{F_2}\rho^2}{\tau^2} < A_{R_6} < \frac{A_{F_2}\rho^2}{7\tau^2}, \quad A_{F_0} > \frac{-7A_{F_2}^2\rho^4 - 9A_{F_2}A_{R_6}\rho^2\tau^2 - A_{R_6}^2\tau^4}{A_{F_2}\rho^6 - A_{R_6}\rho^4\tau^2}, \quad (4)$$

which can be satisfied and thus the no go theorem for the stability of de Sitter extrema can be evaded. Note that $A_{H_3} > 0$ for the parameter region above.

Since we are interested in finding the parameter regime which gives stable de Sitter vacua we omitted the case when $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) = 0$ which can be satisfied non trivially in this example if the second inequality in (4) becomes an equality. If an eigenvalue of the mass matrix is zero we need to check higher order terms for stability.

One can repeat this analysis for all the other cases. Since the details are not very illuminating we simply state our results as follows:

(i) In Type IIA string theory we can satisfy the conditions for the existence of de Sitter critical points and the requirements of stability in the following setups:

- R_6, H_3, F_0, F_2 O6 with $A_{R_6} > 0$, $A_{H_3} > 0$

- R_6, H_3, F_0, F_4 O6 with $A_{R_6} > , A_{H_3} \geq$
- R_6, H_3, F_0, F_6 O6 with $A_{R_6} > , A_{H_3} \geq$
- R_6, H_3, F_0, F_4 O4 with $A_{R_6} \geq , A_{H_3} >$ or $A_{R_6} > , A_{H_3} =$
- R_6, H_3, F_0, F_6 O4 with $A_{R_6} \geq , A_{H_3} >$ or $A_{R_6} > , A_{H_3} =$
- R_6, H_3, F_2, F_4 O4 with $A_{R_6} > , A_{H_3} \geq$
- R_6, H_3, F_2, F_6 O4 with $A_{R_6} > , A_{H_3} \geq$
- R_6, H_3, F_2 O4 O6 with $A_{R_6} > , A_{H_3} >$

(ii) In Type IIB string theory the following setups can admit stable de Sitter minimum:

- R_6, H_3, F_3, F_5 O with $A_{R_6} > , A_{H_3} >$
- R_6, H_3, F_1, F_5 O with $A_{R_6} \geq , A_{H_3} >$
- R_6, H_3, F_1, F_5 O5 with $A_{R_6} > , A_{H_3} \geq$
- R_6, H_3, F_1, F_3 O5 with $A_{R_6} > , A_{H_3} >$
- R_6, H_3, F_3 O O5 with $A_{R_6} > , A_{H_3} >$
- R_6, H_3, F_1 O5 O7 with $A_{R_6} > , A_{H_3} >$

As we increase the number of ingredients (fluxes and localized sources) we are likely to find more and more examples that evade the no goes (for both the existence of an extremum and its stability) but the above list constitutes the “minimal” setups for classical de Sitter vacua. Interestingly the no goes can be evaded with compactifications on a *positively curved* or a *flat* manifold in the presence of H_3 flux even though in most cases a negatively curved manifold is preferred.

5 Discussions

In this paper we have presented several new no go theorems for classical de Sitter vacua i.e. de Sitter constructions using only 6D curvature fluxes and O planes in Type IIA and IIB string theories. In addition to the no goes for de Sitter extrema previously derived in this context [4, 5, 6] we found that constraints on the stability of these extrema can further eliminate a significant portion of the landscape. We enumerate the *minimal* ingredients needed to evade these no goes. Most of these minimal setups we found involve negatively curved 6D manifolds as originally suggested in [1] though there still remain several interesting possibilities with positively curved 6D manifolds. It would be interesting to see if such minimal setups can indeed be realized in terms of explicit models.

Recently it was argued that warping and/or stringy corrections are necessarily important for compactifications on manifolds whose curvature is everywhere negative [17]. This is because the equations of motion cannot be satisfied pointwise in an everywhere negatively curved internal space if the only negative tension objects at our disposal – i.e. the orientifold planes – are localized. The universal Kahler moduli defined in (2.1) is that of an unwarped case, but is modified in the presence of warping [18, 4] (see in particular [4] which is more suited for our present discussion). In this work we sidestep these issues of warping by implicitly smearing the orientifold planes, which can be thought of as an approximation before the fully backreacted solutions are found.³ Furthermore, since our analysis is carried out in the 4D effective field theory where the internal space is integrated out, it would also apply to manifolds which are not everywhere negatively curved, but with an overall negative 6D curvature. We should emphasize that while we found the necessary constraints for stable classical dS minima, there is no guarantee that explicit backgrounds satisfying the requirements exist. Moreover, the full backreaction of several localized sources remains an open challenging issue.

We focus our search for de Sitter vacua whose constructions do not invoke explicit SUSY breaking localized sources. Introducing sources such as anti-branes would certainly enlarge the list of possibilities. For example, the no-go theorem for the stability of de Sitter extremum constructed from IIB string theory with R_6 , H_3 , F_0 , O4 and O6 can be evaded by replacing O4 with D4 $\overline{D4}$ pairs. In fact, the no-go theorems can similarly be evaded for R_6 , H_3 , F_1 , D $\overline{D}$, O5, also for R_6 , H_3 , F_5 , O $\overline{D5}$ $\overline{D5}$ and R_6 , H_3 , F_5 , O $\overline{D7}$ $\overline{D7}$. Pairs of localized D $\overline{D}$ 4 were also used for uplifting the AdS vacua to dS in [19]. Recently the backreaction of such D $\overline{D}$ pairs in the Klebanov-Strassler throat were discussed in [4, 45] (see also further discussions in [46–48]). Related studies on the backreaction of localized sources in simpler setups (though more closely related to the classical de Sitter vacua discussed here) were considered in [49–5].

Acknowledgment

We have benefited from discussions with Ulf Danielsson, Hirotaka Hayashi, Masazumi Honda, Shajid Haque, Paul Koerber, Thomas Van Riet, S. H. Henry Tye, Sandip P. Trivedi, Timm Wrase and Marco Zagermann. GS and YS thank the Hong Kong Institute for Advanced Study greatly for their hospitality. GS is supported in part by a DOE grant under contract DE-FG 95ER4-896 and a Cottrell Scholar Award from Research Corporation.

³We consider smearing just on a base manifold transverse to the Oq -planes, such that the Oq -planes do not acquire additional moduli dependences as compared to a localized one.

A Instability Analysis for Other Examples

This appendix is devoted to complete the stability analysis for all the cases in (8) which were skipped in section 4 and demonstrate that they all lead to a no go for stable de Sitter vacua

A.1 R_6, H_3, F_0 , O4, O6 in IIA

The requirements of an extremum lead to the following two conditions:

$$A_{H_3} = -\frac{A_{R_6}\rho^6}{\tau^2} - \frac{A_{F_0}\rho^6}{\tau^2} - \frac{A_4\rho^2}{\tau}, \quad A_6 = -\frac{A_{F_0}\rho^3}{\tau} - \frac{4A_{R_6}\tau}{9\rho} - \frac{7A_4}{9\rho}. \quad (\text{A } 3)$$

Then the potential at its extremum upon substituting A_{H_3} and A_6 becomes

$$V_{\text{ext}} = \frac{A_{R_6}}{9\rho\tau^2} - \frac{A_4}{9\rho\tau^3}. \quad (\text{A } 4)$$

Therefore we see that a negatively curved manifold together with a number of O4 planes or with a certain amount of F_0 flux help to obtain dS extrema

The determinant of the mass matrix at the extremum becomes

$$\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) = \frac{1}{\rho^4\tau^9} (6 A_4 A_{F_0} \rho^4 - 7 A_4^2 \tau - 84 A_{F_0} A_{R_6} \rho^4 \tau - 4 A_4 A_{R_6} \tau^2 - 8 A_{R_6}^2 \tau^3). \quad (\text{A } 5)$$

Thus we cannot satisfy $V_{\text{ext}} > 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $A_{H_3} \geq 0$, $A_4 \leq 0$ simultaneously. This means there is at least one tachyonic direction at the positive extremum

A.2 R_6, H_3, F_1, F_3 , O3 in IIB

The conditions on extrema lead to the following potential at the critical point:

$$A_{H_3} = \frac{A_{F_3}\rho^3}{\tau^2} - \frac{A_{F_1}\rho^5}{\tau^2}, \quad A_3 = -\frac{A_{F_3}\rho^{3/2}}{\tau} - \frac{8A_{F_1}\rho^{7/2}}{\tau} - A_{R_6}\rho^{1/2}\tau, \\ V_{\text{ext}} = \frac{A_{F_1}\rho^2}{\tau^4} - \frac{A_{R_6}}{\rho\tau^2}. \quad (\text{A } 6)$$

The trace and the determinant of the mass matrix go like

$$\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) = \frac{1}{\rho^3\tau^6} (4A_{F_3}\rho^3 - 9A_{F_3}\rho\tau^2 - 4A_{R_6}\rho^2\tau^2 - A_{R_6}\tau^4 - A_{F_1}\rho^3\tau^2), \\ \det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) = -\frac{6}{\rho^3\tau^{10}} (4A_{F_1}^2\rho^5 - A_{F_3}A_{R_6}\tau^2 - A_{F_1}\rho^2(A_{F_3}\rho - A_{R_6}\tau^2)), \quad (\text{A } 7)$$

We see that the conditions $V_{\text{ext}} > 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $A_{H_3} \geq 0$ cannot be satisfied simultaneously

A.3 R_6, H_3, F_1 , O3, O5 in IIB

At the extremum of the potential:

$$\begin{aligned} A_{H_3} &= \frac{A_{F_1}\rho^5}{\tau^2} - \frac{A_5\rho^{5/2}}{\tau}, \quad A_3 = -\frac{8A_{F_1}\rho^{7/2}}{\tau} - \frac{A_{R_6}\rho^{1/2}\tau}{\tau} - \frac{4A_5\rho}{\tau}, \\ V_{\text{ext}} &= \frac{A_{F_1}\rho^2}{\tau^4} - \frac{A_{R_6}}{\rho\tau^2} - \frac{A_5}{6\rho^{1/2}\tau^3}, \end{aligned} \quad (\text{A } 6)$$

while the trace and the determinant of the mass matrix are found to be

$$\begin{aligned} \text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= \frac{1}{4\rho^3\tau^5} \begin{pmatrix} -4A_5\rho^{5/2} & 7A_5\rho^{1/2}\tau^2 - 8A_{R_6}\rho^2\tau - A_{R_6}\tau^3 & 64A_{F_1}\rho^3\tau \end{pmatrix}, \\ \det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= -\frac{1}{\rho^{7/2}\tau^{10}} \begin{pmatrix} 8A_{F_1}\rho^{5/2} & A_5\tau & 4A_{F_1}\rho^3 & A_5\rho^{1/2}\tau & A_{R_6}\tau^2 \end{pmatrix}. \end{aligned} \quad (\text{A } 7)$$

Again we cannot satisfy $V_{\text{ext}} > 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $A_{H_3} \geq 0$, $A_3 \leq 0$ simultaneously regardless of the sign of R_6

A.4 R_6, H_3, F_5 , O3, O5 in IIB

At the extremum of the potential:

$$\begin{aligned} A_{H_3} &= \frac{A_5\rho^{5/2}}{\tau}, \quad A_3 = -\frac{4A_{F_5}}{\rho^{1/2}\tau} - \frac{A_{R_6}\rho^{1/2}\tau}{\tau} - \frac{4A_5\rho}{\tau}, \\ V_{\text{ext}} &= -\frac{A_{F_5}}{\rho^2\tau^4} - \frac{A_{R_6}}{\rho\tau^2} - \frac{A_5}{6\rho^{1/2}\tau^3}. \end{aligned} \quad (\text{A } 8)$$

then the trace and the determinant of the mass matrix becomes

$$\begin{aligned} \text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= \frac{1}{4\rho^4\tau^6} \begin{pmatrix} 6A_{F_5}\rho^2 & 4A_{F_5}\tau^2 - 4A_5\rho^{7/2}\tau & 7A_5\rho^{3/2}\tau^3 - 8A_{R_6}\rho^3\tau^2 - A_{R_6}\rho\tau^4 \end{pmatrix}, \\ \det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= -\frac{A_5(-6A_{F_5} - A_{R_6}\rho\tau^2 - A_5\rho^{3/2}\tau)}{\rho^{9/2}\tau^9}. \end{aligned} \quad (\text{A } 9)$$

In this case we cannot satisfy $V_{\text{ext}} > 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) > 0$, $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $A_{H_3} \geq 0$, $A_5 \leq 0$ simultaneously. When we satisfy $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) = 0$, $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$ the system is reduced to (R_6, F_5, O) which was showed to have one flat direction in (7)

A.5 R_6, H_3, F_1 , O3, O7 in IIB

At the extremum of the 4D potential:

$$\begin{aligned} A_{H_3} &= \frac{A_{F_1}\rho^5}{\tau^2} - \frac{A_7\rho^{7/2}}{\tau}, \quad A_3 = -\frac{8A_{F_1}\rho^{7/2}}{\tau} - \frac{A_{R_6}\rho^{1/2}\tau}{\tau} - \frac{5A_7\rho^2}{\tau}, \\ V_{\text{ext}} &= \frac{A_{F_1}\rho^2}{\tau^4} - \frac{A_7\rho^{1/2}}{\tau^3} - \frac{A_{R_6}}{\rho\tau^2}. \end{aligned} \quad (\text{A } 10)$$

Then the trace and the determinant of the mass matrix go like

$$\begin{aligned}\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= \frac{1}{\rho^3\tau^5} \begin{pmatrix} A_{F_1}\rho^3\tau - 4A_{R_6}\rho^2\tau - A_{R_6}\tau^3 - 4A_7\rho^{7/2} & A_7\rho^{3/2}\tau^2 \end{pmatrix}, \\ \det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= \frac{-4}{\rho^{5/2}\tau^{10}} \begin{pmatrix} 6A_{F_1}^2\rho^{9/2} & A_7\tau^2(5A_7\rho^{3/2} & 4A_{R_6}\tau & 4A_{F_1}(5A_7\rho^3\tau & A_{R_6}\rho^{3/2}\tau^2 \end{pmatrix}.\end{aligned}\tag{A}$$

It can be shown that $V_{\text{ext}} > 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $A_{H_3} \geq 0$ cannot be satisfied

A.6 R_6, H_3, F_3 , **O3, O7** in IIB

At the extremum of the 4D potential:

$$\begin{aligned}A_{H_3} &= \frac{A_{F_3}\rho^3}{\tau^2} - \frac{A_7\rho^{7/2}}{\tau}, \quad A_3 = -\frac{A_{F_3}\rho^{3/2}}{\tau} - \frac{A_{R_6}\rho^{1/2}\tau}{\tau} - \frac{5A_7\rho^2}{\tau}, \\ V_{\text{ext}} &= \frac{A_7\rho^{1/2}}{\tau^2} - \frac{A_{R_6}}{\rho\tau^2}\end{aligned}\tag{A}$$

Then the determinant of the mass matrix becomes

$$\begin{aligned}\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= \frac{1}{\rho^3\tau^6} \begin{pmatrix} 4A_{F_3}\rho^3 & 9A_{F_3}\rho\tau^2 - 4A_7\rho^{7/2}\tau & A_7\rho^{3/2}\tau^3 - 4A_{R_6}\rho^2\tau^2 - A_{R_6}\tau^4 \end{pmatrix}, \\ \det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) &= -\frac{4}{\rho^3\tau^9} \begin{pmatrix} 4A_{F_3}A_{R_6}\tau & 5A_7^2\rho^2\tau & 4A_7(A_{F_3}\rho^{3/2} & A_{R_6}\rho^{1/2}\tau^2 \end{pmatrix}.\end{aligned}\tag{A}$$

Thus we cannot satisfy $V_{\text{ext}} > 0$, $\det(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $\text{tr}(\partial_{\rho_i}\partial_{\rho_j}V|_{\text{ext}}) \geq 0$, $A_{H_3} \geq 0$ simultaneously

A.7 R_6, H_3, F_5 , **O3, O7** in IIB

At the extremum of the 4D potential:

$$\begin{aligned}A_{H_3} &= \frac{A_7\rho^{7/2}}{\tau}, \quad A_3 = -\frac{4A_{F_5}}{\rho^{1/2}\tau} - \frac{A_{R_6}\rho^{1/2}\tau}{\tau} - \frac{5A_7\rho^2}{\tau}, \\ V_{\text{ext}} &= -\frac{A_{F_5}}{\rho^2\tau^4} - \frac{A_{R_6}}{\rho\tau^2} - \frac{A_7\rho^{1/2}}{\tau^3}.\end{aligned}\tag{A 4}$$

We see immediately that the only allowed situation is $A_{H_3} = A_7 = 0$ for O7 a case subsumed in the analysis around (7 which has at least one flat direction

References

- S Kachru R Kallosh A D Linde and S P Trivedi “De Sitter vacua in string theory ” *Phys. Rev.* **D68** (46 5 arXiv:hep-th/0301240
- M P Hertzberg S Kachru W Taylor and M Tegmark “Inflationary Constraints on Type IIA String Theory ” *JHEP* **12** (7 95 arXiv:0711.2512 [hep-th]
- E Silverstein “Simple de Sitter Solutions ” *Phys. Rev.* **D77** (8 6 6 arXiv:0712.1196 [hep-th]
- 4 S S Haque G Shiu B Underwood and T Van Riet “Minimal simple de Sitter solutions ” *Phys. Rev.* **D79** (9 86 5 arXiv:0810.5328 [hep-th]
- 5 U H Danielsson S S Haque G Shiu and T Van Riet “Towards Classical de Sitter Solutions in String Theory ” *JHEP* **09** (9 4 arXiv:0907.2041 [hep-th]
- 6 U H Danielsson *et al.* “De Sitter hunting in a classical landscape ” arXiv:1103.4858 [hep-th]
- 7 C Caviezel *et al.* “On the Cosmology of Type IIA Compactifications on SU(3) Structure Manifolds ” *JHEP* **04** (9 arXiv:0812.3551 [hep-th]
- 8 R Flauger S Paban D Robbins and T Wrase “Searching for slow roll moduli in inflation in massive type IIA supergravity with metric fluxes ” *Phys. Rev.* **D79** (9 86 arXiv:0812.3886 [hep-th]
- 9 B de Carlos A Guarino and J M Moreno “Flux moduli stabilisation Supergravity algebras and no go theorems ” *JHEP* **01** (arXiv:0907.5580 [hep-th]
- B de Carlos A Guarino and J M Moreno “Complete classification of Minkowski vacua in generalised flux models ” *JHEP* **02** (76 arXiv:0911.2876 [hep-th]
- C Caviezel T Wrase and M Zagermann “Moduli Stabilization and Cosmology of Type IIB on SU(3) Structure Orientifolds ” *JHEP* **04** (arXiv:0912.3287 [hep-th]
- X Dong B Horn E Silverstein and G Torroba “Micromanaging de Sitter holography ” *Class. Quant. Grav.* **27** (May 45 1005.5403
- D Andriot E Goi R Minasian and M Petrini “Supersymmetry breaking branes on solvmanifolds and de Sitter vacua in string theory ” *JHEP* **05** (8 arXiv:1003.3774 [hep-th]
- 4 G Dibitetto R Linares and D Roest “Flux Compactifications Gauge Algebras and De Sitter ” *Phys. Lett.* **B688** (96 arXiv:1001.3982 [hep-th]
- 5 U H Danielsson P Koerber and T Van Riet “Universal de Sitter solutions at tree level ” *JHEP* **05** (9 arXiv:1003.3590 [hep-th]

- 6 T Wrase and M Zagermann
“On Classical de Sitter Vacua in String Theory ” *Fortschr. Phys.* **58** (Mar 9 6 9 1003.0029
- 7 O DeWolfe A Giryavets S Kachru and W Taylor “Type IIA moduli stabilization ” *JHEP* **07** (5 66 hep-th/0505160
- 8 P Narayan and S P Trivedi “On The Stability Of Non Supersymmetric AdS Vacua ” *JHEP* **07** (89 arXiv:1002.4498 [hep-th]
- 9 D Harlow “Metastability in Anti de Sitter Space ” 1003.5909
G T Horowitz J Orgera and J Polchinski “Nonperturbative Instability of AdS₅ x S⁵ Zk ” *Phys. Rev.* **D77** (8 4 4 arXiv:0709.4262 [hep-th]
L Covi *et al.*
“de Sitter vacua in no scale supergravities and Calabi Yau string models ” *JHEP* **06** (Apr 8 57 0804.1073
M Gomez Reino J Louis and C A Scrucca
“No metastable de Sitter vacua in N= supergravity with only hypermultiplets ” *JHEP* **02** (Dec 9 0812.0884
A Borghese and D Roest “Metastable supersymmetry breaking in extended supergravity ” *JHEP* **05** (arXiv:1012.3736 [hep-th]
- 4 L Brizi and C A Scrucca “The lightest scalar in theories with broken supersymmetry ” 1107.1596
- 5 K Choi “String or M theory axion as a quintessence ” *Phys. Rev.* **D62** (4 5 9 hep-ph/9902292
- 6 P Svrcek “Cosmological constant and axions in string theory ” arXiv:hep-th/0607086
- 7 S Panda Y Sumitomo and S P Trivedi “Axions as Quintessence in String Theory ” *Phys. Rev.* **D83** (8 5 6 arXiv:1011.5877 [hep-th]
- 8 E Silverstein and A Westphal “Monodromy in the CMB: Gravity Waves and String Inflation ” *Phys. Rev.* **D78** (8 6 arXiv:0803.3085 [hep-th]
- 9 L McAllister E Silverstein and A Westphal “Gravity Waves and Linear Inflation from Axion Monodromy ” *Phys. Rev.* **D82** (46 arXiv:0808.0706 [hep-th]
R Flauger L McAllister E Pajer A Westphal and G Xu
“Oscillations in the CMB from Axion Monodromy Inflation ” *JCAP* **1006** (July 9 0907.2916
G T Gilbert “Positive definite matrices and sylvester’s criterion ” *The American Mathematical Monthly* **98** (99 no pp 44 46
<http://www.jstor.org/stable/2324036>

- M R Douglas and R Kallosh
 “Compactification on negatively curved manifolds” *JHEP* **06** (Jan 2006) 040
 1001.4008
- S B Giddings and A Maharana “Dynamics of warped compactifications and the shape of the warped landscape” *Phys. Rev.* **D73** (2006) 065010 [arXiv:hep-th/0507158]
- 4 H Kodama and K Uzawa “Comments on the four dimensional effective theory for warped compactification” *JHEP* **03** (2006) 055 [arXiv:hep-th/0512104]
- 5 H Kodama and K Uzawa “Moduli instability in warped compactification”
 arXiv:hep-th/0601100
- 6 P Koerber and L Martucci
 “From ten to four and back again: how to generalize the geometry” *JHEP* **08** (July 2008) 077
 0707.1038
- 7 G Shiu G Torroba B Underwood and M R Douglas “Dynamics of Warped Flux Compactifications” *JHEP* **06** (2008) 084 [arXiv:0803.3068 [hep-th]]
- 8 M R Douglas and G Torroba “Kinetic terms in warped compactifications”
JHEP **05** (2009) 085 [arXiv:0805.3700 [hep-th]]
- 9 L Martucci
 “On moduli and effective theory of $N=1$ warped flux compactifications” *JHEP* **05** (Feb 2006) 092
 0902.4031
- 4 H Y Chen Y Nakayama and G Shiu “On D brane Dynamics at Strong Warping”
Int. J. Mod. Phys. **A25** (2010) 4905 [arXiv:0905.4463 [hep-th]]
- 4 A R Frey G Torroba B Underwood and M R Douglas “The Universal Kaehler Modulus in Warped Compactifications” *JHEP* **01** (2010) 096 [arXiv:0810.5768 [hep-th]]
- 4 S Kachru J Pearson and H L Verlinde “Brane Flux Annihilation and the String Dual of a Non Supersymmetric Field Theory” *JHEP* **06** (2006) 063 [arXiv:hep-th/0112197]
- 4 O DeWolfe S Kachru and M Mulligan “A Gravity Dual of Metastable Dynamical Supersymmetry Breaking” *Phys. Rev.* **D77** (2008) 065018 [arXiv:0801.1520 [hep-th]]
- 44 P McGuirk G Shiu and Y Sumitomo “Non supersymmetric infrared perturbations to the warped deformed conifold” *Nucl. Phys.* **B842** (2010) 84 [arXiv:0910.4581 [hep-th]]
- 45 I Bena M Grana and N Halmagyi “On the Existence of Meta stable Vacua in Klebanov Strassler” *JHEP* **09** (2009) 087 [arXiv:0912.3519 [hep-th]]
- 46 I Bena G Giecold M Grana and N Halmagyi “On The Instanton Potential From Antibrane in Warped Throats” arXiv:1011.2626 [hep-th]

- 47 A Dymarsky “On gravity dual of a metastable vacuum in Klebanov–Strassler theory ”
JHEP **05** (5 arXiv:1102.1734 [hep-th]
- 48 I Bena G Giecold M Grana N Halmagyi and S Massai “The backreaction of anti D
branes on the Klebanov–Strassler geometry ” arXiv:1106.6165 [hep-th]
- 49 J Blaback *et al.* “Smeared versus localised sources in flux compactifications ”
JHEP **12** (4 arXiv:1009.1877 [hep-th]
- 5 J Blaback *et al.* “The problematic backreaction of SUSY breaking branes ”
arXiv:1105.4879 [hep-th]