

Yohji Akama · Shinji Iizuka

Random fields on model sets with localized dependency and their diffraction

Received: date / Accepted: date

Abstract For a random field on a general discrete set, we introduce a condition that the range of the correlation from each site is within a predefined compact set D . For such a random field ω defined on the model set Λ that satisfies a natural geometric condition, we develop a method to calculate the diffraction measure of the random field. The method partitions the random field into a finite number of random fields, each being independent and admitting the law of large numbers. The diffraction measure of ω consists almost surely of a pure-point component and an absolutely continuous component. The former is the diffraction measure of the expectation $E[\omega]$, while the inverse Fourier transform of the absolutely continuous component of ω turns out to be a weighted Dirac comb which satisfies a simple formula. Moreover, the pure-point component will be understood quantitatively in a simple exact formula if the weights are continuous over the internal space of Λ . Then we provide a sufficient condition that the diffraction measure of a random field on a model set is still pure-point.

Keywords Diffraction · Pure point spectrum · Absolutely continuous spectrum · Quasicrystal · Model set

PACS 61.05.cc · 61.05.cp

Mathematics Subject Classification (2000) MSC primary 52C23; · secondary 37B50

The preliminary version is presented at the fifth Asian International Workshop on Quasicrystals, 2009, Tokyo Japan.

Yohji Akama
Mathematical Institute
Tel.: +81-(0)22-795-7708
Fax: +81-(0)22-795-6400
E-mail: akama@math.tohoku.ac.jp

Shinji Iizuka
Research and Development Section, Hitachi East Japan Solutions, Ltd.

1 Introduction

A physical quasicrystal is a material which has (1) a diffraction pattern with Bragg peaks and (2) a symmetry that ordinary crystals cannot have. The set of the atomic positions in a quasicrystal is mathematically modelled by a *model set* [18], which is defined by introducing an extra space (“internal space”), a relatively compact subset (“*window*”) of the internal space, and a so-called *star map* $(-)^*$ such that for each site s of a model set the point s^* belongs to the window. The topological properties of the window cause the pure-point diffraction measure (see [23] and [7] for example) of the model set, which explain the aforementioned properties (1) of the quasicrystal.

Although model sets are proved to have necessarily a pure-point diffraction measure, real quasicrystals have diffraction measures with not only Bragg peaks (pure-point component) but also diffuse scattering (absolutely continuous component). The phenomenon is explained from a physical point of view with a probabilistic effect in [14], and in [5] where to the sites of the model set associated are *independent* random variables. In [12], Hof regarded the thermal motion of atoms as i.i.d. random displacements, and then studied the influence on the diffraction measure of aperiodic monoatomic crystals. Since correlations ([10]) are present in a quasicrystal, we equip model sets with a localized probabilistic dependency, to quantitatively study the ability of diffuse scattering to characterize local structures and defects in materials. In [16], Lenz employed a dynamical system of point sets to study the diffraction measures of percolation and the random displacement models based on aperiodic order. Recently, in [21], Müller and Richard also made a rigorous approach on these models by using sets of σ -algebras.

For a model set Λ , we consider a complex-valued random field $\{X_s\}_{s \in \Lambda}$ with dependency localized as follows: there is a finite patch D such that each site s has correlation on, at most, sites belonging to the patch D relative to s . This localized dependency condition seems essentially the same as the so-called “finite range condition” of stochastic analysis. We call a random field on a model set subject to the localized dependency condition a *finitely randomized model set*. We develop a method to calculate the diffraction measure of such complex-valued random field (Section 3). For the diffraction measures of finitely randomized model sets, we determine quantitatively the pure-point component (in Section 5) and the absolutely continuous component (in Section 4). Our approach is mostly based on the *finite local complexity* of model sets, as in [23]. As a consequence, if the fourth noncentral moments $\{E[|X_s|^4]\}_{s \in \Lambda}$ is bounded, and if the expectation $E[X_s]$ at each site s as well as the covariance X_s and X_{s-g} of sites s and $s-g$ in the finitely randomized model set are given by continuous functions $e(s^*)$ and $c_g(s^*)$, where s^* is the value of s by the star map, then

1. the inverse Fourier transform of the absolutely continuous component is a Dirac comb such that the support $\{g_1, \dots, g_n\}$ is the smallest D and the weight of the δ_{g_i} is the average of the covariances $\text{Cov}[X_s, X_{s-g_i}]$ ($s, s-g_i \in \Lambda$); and
2. the pure-point component is the diffraction measure of a Dirac comb $\sum_{s \in \Lambda} E[X_s] \delta_s$, the expectation of the random field.

This type of theorems are also seen in some other models with i.i.d. conditions. See [2] and references therein.

On the other hand, from the viewpoint of stochastic processes, we provide a sufficient condition for a randomly weighted Dirac comb on a model set to have diffraction measure whose expectation is still pure-point. The sufficient condition is satisfied when the set of the weights X_s forms a Wiener stochastic process $\{X_{\star^{-1}(y)}\}_y$ parametrized by the window $W \subset \mathbb{R}$. We draw this observation by providing quantitatively the diffraction measure of the deterministically weighted Dirac comb on a model set. The quantitative estimate will be established with the help of the so-called torus parametrization which was introduced in [3], then was extended in [23] to the locally compact, σ -compact Abelian Hausdorff groups (LCAG for short), and was finally fully exploited in [4].

2 Basic properties of model sets: review

Throughout this paper, $\mathfrak{G}$ and $\mathfrak{G}_{\text{int}}$ are locally compact, σ -compact Abelian Hausdorff groups (LCAG for short).

Definition 1 A *cut-and-project scheme* (c.-p. scheme, for short) is a triple $\mathcal{S} = (\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$ such that (1) $\mathfrak{G}$ and $\mathfrak{G}_{\text{int}}$ are called a *physical space* and an *internal space* respectively; (2) $\tilde{L}$ is a lattice of $\mathfrak{G} \times \mathfrak{G}_{\text{int}}$, that is, a discrete subgroup of $\mathfrak{G} \times \mathfrak{G}_{\text{int}}$ with $(\mathfrak{G} \times \mathfrak{G}_{\text{int}})/\tilde{L}$ being compact; (3) The canonical projection $\Pi : \mathfrak{G} \times \mathfrak{G}_{\text{int}} \rightarrow \mathfrak{G}$ is injective on $\tilde{L}$, and the image of $\tilde{L}$ by the other canonical projection $\Pi_{\text{int}} : \mathfrak{G} \times \mathfrak{G}_{\text{int}} \rightarrow \mathfrak{G}_{\text{int}}$ is dense in the internal space $\mathfrak{G}_{\text{int}}$. For each $s \in \Pi(\tilde{L})$, we write s^* for $\Pi_{\text{int}} \circ (\Pi|_{\tilde{L}})^{-1}(s)$, L for $\Pi(\tilde{L})$, and L^* for $(L)^*$. The $(-)^*$ is called the *star map*. Define

$$\Lambda(W) := \{\Pi(x) ; x \in \tilde{L}, \Pi_{\text{int}}(x) \in W\}.$$

For any LCAG G , any subsets $A, B \subset G$, and any $x \in G$, let $A \pm B$ be $\{a \pm b ; a \in A, b \in B\}$ and $x + A$ be $\{x + a ; a \in A\}$. A set $A \subset \mathfrak{G}$ is said to be *uniformly discrete*, if $(A - A) \cap U = \{0\}$ for some open neighborhood U of 0, while A is said to be *relatively dense*, if $\mathfrak{G} = A + K$ for some compact set K . The interior, the closure and the boundary of A in LCAG G are denoted by $\text{Int}_G(A)$, $\text{Cl}_G(A)$ and $\partial_G A$, respectively. Here the subscript G is omitted if no confusion occurs.

Lattices in $G \times H$ with G, H being LCAG's are written as $\tilde{L}, \tilde{M}, \dots$, and their canonical projections to G are written as $L, M, \dots$, respectively.

Lemma 1 *If $\Lambda \subset \mathfrak{G}$ is uniformly discrete, there is a compact neighborhood U of 0 such that $U = -U$ and $(s + U) \cap (s' + U) = \emptyset$ for all distinct $s, s' \in \Lambda$.*

Definition 2 (Model set) Let $(\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$ be a c.-p. scheme. By a *window*, we mean a nonempty, measurable, relatively compact subset of the internal space $\mathfrak{G}_{\text{int}}$. If $W \subset \mathfrak{G}_{\text{int}}$ is a window, $\Lambda(W)$ is called a *model set*.

It is well-known that any model set is uniformly discrete. See [20, Proposition 2], for example. Every LCAG has a unique Haar measure up to positive real multiples. Throughout this paper, we fix Haar measures of the LCAG's $\mathfrak{G}$ and $\mathfrak{G}_{\text{int}}$. The Haar measure of $\mathfrak{G}_{\text{int}}$ is denoted by θ , and the integration of a function with respect to the Haar measure of $\mathfrak{G}$ ($\mathfrak{G}_{\text{int}}$ resp.) is denoted by $\int \cdots dx$ ($\int \cdots dy$ resp.) as usual. The Haar measure of a set A is just denoted by $|A|$ if no confusion occurs. By a *van Hove sequence* of $\mathfrak{G}$, we mean an increasing sequence $\{D_n\}_{n \in \mathbb{N}}$ of compact subsets of $\mathfrak{G}$ such that $|D_n| > 0$ for every $n \in \mathbb{N} = \{1, 2, 3, \dots\}$ and for every compact subset $K \subset \mathfrak{G}$, $\lim_{n \leq} |\partial^K(D_n)|/|D_n| = 0$, where for a compact set $A \subset \mathfrak{G}$,

$$\partial^K(A) := ((A + K) \setminus \text{Int}(A)) \cup ((Cl(\mathfrak{G} \setminus A) - K) \cap A). \quad (1)$$

In [23], the existence of a van Hove sequence for any LCAG is derived from

Proposition 1 ([11, Theorem 9.8]) *For every locally compact, compactly generated Abelian Hausdorff group H , there are $l, m \in \mathbb{Z}_{\geq 0}$, a compact Abelian Hausdorff group K , and an isomorphism φ from H to $\mathbb{R}^l \times \mathbb{Z}$*

Actually we have $\gamma_\omega^{(n)} = \sum_{g \in S} \eta_\omega^{(n)}(g) \delta_g$ where $\eta_\omega^{(n)}(g)$ is the summation of $w_s \overline{w_t} |D_n|^{-1}$ over $s, t \in S \cap D_n$ such that $s - t = g$. Since S is discrete and D_n is compact, $\eta_\omega^{(n)}$ is well-defined, and $\gamma_\omega^{(n)}$ is so.

Let $C_c(\mathfrak{G})$ be the set of complex continuous functions on $\mathfrak{G}$ with compact support in $\mathfrak{G}$. Let the *autocorrelation measure* of ω be the limit γ_ω of $\gamma_\omega^{(n)}$ in the vague topology. Then γ_ω is written as

$$\gamma_\omega = \sum_{g \in S} \eta_\omega(g) \delta_g, \quad \eta_\omega(g) = \lim_{n \leq} \eta_\omega^{(n)}(g).$$

For any LCAG H , the *dual group* of H is denoted by $\widehat{H}$. Fix a Haar measure h of H . The Fourier transform of a function $f : H \rightarrow \mathbb{C}$ is $\widehat{f} : \widehat{H} \rightarrow \mathbb{C}$ such that $\widehat{f}(\chi) := \int_H f(x) \chi(x) dh(x)$. The *diffraction measure* of ω is, by definition, the Fourier transform $\widehat{\gamma_\omega}$ of the autocorrelation measure γ_ω . A measure on $\mathfrak{G}$ has Fourier transform as a measure on $\widehat{\mathfrak{G}}$ as follows:

Proposition 3 ([1, Theorem 2.1, Theorem 4.1]) *Suppose λ is a measure on $\mathfrak{G}$. The Fourier transform (if it exists) of λ is a unique measure $\widehat{\lambda}$ on $\widehat{\mathfrak{G}}$ such that for all $\varphi \in C_c(\mathfrak{G})$*

$$\int_{\mathfrak{G}} (\varphi * \tilde{\varphi})(x) d\lambda(x) = \int_{\widehat{\mathfrak{G}}} |\widehat{\varphi}(-\chi)|^2 d\widehat{\lambda}(\chi).$$

Moreover if λ is positive definite (i.e., $\int_{\mathfrak{G}} (\varphi * \tilde{\varphi})(x) d\lambda(x) \geq 0$ for all $\varphi \in C_c(\mathfrak{G})$), then λ indeed has the Fourier transform $\widehat{\lambda}$ which is positive. Here $\tilde{\varphi} : \mathfrak{G} \rightarrow \mathbb{C}$ is a function $x \in \mathfrak{G} \mapsto \overline{\varphi(-x)}$.

A Haar measure on $\widehat{\mathfrak{G}}$ is $\widehat{\delta}_0$ where δ_0 is the Dirac measure at 0 on $\mathfrak{G}_{\text{int}}$. Then the equation of Proposition 3 amounts to a Plancherel formula. The integral of a function with respect to the Haar measure of $\widehat{\mathfrak{G}}$ is denoted by $\int \cdots d\chi$. Because $\widehat{\eta_\omega} = \gamma_\omega$, we have $\widehat{\gamma_\omega} = \gamma_\omega$ and thus $\widehat{\gamma_\omega}$ is positive by [1, Proposition 4.1].

A measure μ on $\mathfrak{G}$ is said to be *translationally bounded*, if for every $\varphi \in C_c(\mathfrak{G})$ the set $\{\int_{\mathfrak{G}} \varphi(x+t) d\mu(x) ; t \in \mathfrak{G}\}$ is bounded.

Baake-Moody established the pure-point diffraction of weighted Dirac combs on model sets, by using Weyl's theorem ([6, Proposition 6.2]) for model sets and an ingenious topological space.

Proposition 4 ([7, Theorem 2]) *Suppose that $\Lambda(W)$ is a model set, and the window $W \subset \mathfrak{G}_{\text{int}}$ satisfies that the Haar measure $\theta(\partial W)$ is 0. Then for any $f : \mathfrak{G}_{\text{int}} \rightarrow \mathbb{C}$ supported and continuous on $Cl(W)$, the diffraction measure $\widehat{\gamma_\omega}$ of the Dirac comb $\omega = \sum_{s \in \Lambda(W)} f(s^*) \delta_s$ is translationally bounded, nonnegative and pure-point.*

3 Finitely randomized model sets

We need Weyl's theorem for random fields on model sets, in Section 4. So for a model set $\Lambda_{\mathcal{O}}(W)$ over a c.-p. scheme $\mathcal{O}$, we wish to decompose a random field $\{X_s\}_{s \in \Lambda_{\mathcal{O}}(W)}$ into a finite number of *independent* random fields, each being on a model set over the same c.-p. scheme $\mathcal{O}$. For a sufficient condition of such a decomposition, we consider a geometric condition (Subsection 3.2) on model sets and a condition for random fields (Subsection 3.1). Hereafter, a “random variable” is abbreviated as “RV,” and the cardinality of a finite set A is denoted by $\#A$.

3.1 Independence in random field

Definition 3 (Dependency set) Let $\{X_s\}_{s \in S}$ be a random field on a discrete set S . A *dependency set* (*d-set* for short) is a set $D = -D \subset S - S$ such that for any finite sets $P, Q \subset S$, if a set $(P - Q)$ is disjoint from D , then a pair of a $\#P$ -dimensional random vector $(X_s)_{s \in P}$ and a $\#Q$ -dimensional random vector $(X_s)_{s \in Q}$ is independent. A d-set necessarily has 0 as an element. If a random field has a dependency set, we can replace it with an arbitrary superset of it.

A d-set is a patch such that each site s has correlation on, at most, sites belonging to the patch relative to s . Recall that a sequence $\{X_1, X_2, \dots\}$ of RV's is independent, if so are any finite subsequences. We set the product $\prod_{i=1}^{\infty} (\dots)$ to be 1.

Lemma 2 (Independence) Let D be a d-set of a random field $\{X_s\}_{s \in S}$ on an FLC subset S of an LCAG, and let N be any subset of the LCAG. If $D \cap ((s + N) - (t + N)) = \emptyset$ for any distinct $s, t \in S$, then a sequence $\left\{ \prod_{t \in (s+N) \setminus S} X_t ; s \in S \right\}$ of RV's is independent. Furthermore, the random field is independent, if and only if the random field has $\{0\}$ as a d-set.

Proof We show that a sequence $\left\{ \prod_{t \in (s+N) \setminus S} X_t ; s \in S \right\}$ of RV's is independent for any finite subset $S' = \{s_1, \dots, s_{\nu}\}$ of S . The proof is by induction on $\nu = \#S'$. When $\nu = 1$, it is trivial, so assume $\nu > 1$. A set $(\{s_1, \dots, s_{\nu-1}\} + N) - (s_{\nu} + N) = \bigcup_{i=1}^{\nu-1} (s_i + N - (s_{\nu} + N))$ is disjoint from D by the premise. So, a random vector $(X_t ; t \in (\{s_1, \dots, s_{\nu-1}\} + N) \cap S)$ is independent from a random vector $(X_t ; t \in (s_{\nu} + N) \cap S)$. Thus a $(\nu - 1)$ -dimensional random vector $(\prod_{t \in (s_i+N) \setminus S} X_t ; 1 \leq i \leq \nu - 1)$ is independent from an RV $\prod_{t \in (s_{\nu}+N) \setminus S} X_t$. Because a sequence $\left\{ \prod_{t \in (s_i+N) \setminus S} X_t ; 1 \leq i \leq \nu - 1 \right\}$ of RV's has length $\nu - 1$ and is independent by the induction hypothesis on ν , we are done. The if-part of the last sentence of the statement is proved by taking $N = \{0\}$, while the only-if part is immediate. $\square$

Definition 4 A *finitely randomized model set* (FRMS for short) on a model set $\Lambda(W)$ is a random field $\{X_s\}_{s \in \Lambda(W)}$ with a *finite* d-set D .

The FRMS can be regarded as a Dirac comb with random weights. When each X_s is an indicator (i.e., a $\{0, 1\}$ -valued RV), we intend that $X_s > 0$ if and only if $s \in \Lambda(W)$ indeed appears.

Example 1 Suppose $\Gamma = \Lambda(W)$ is a model set such that the star map $(-)^*$ is injective. Suppose a set C is $\{p_1, \dots, p_n\} \subset L$ such that $(W + p_i^*) \cap (W + p_j^*) \cap L^* = \emptyset$ ($i \neq j$). Then $\Lambda = \Lambda(W + C^*)$ is also a model set, and for every $s \in \Lambda$ there are unique $t \in \Gamma$ and unique $i \in \{1, \dots, n\}$ such that

$$s = t + p_i. \quad (2)$$

Let $\{\mathbf{Y}_t\}_{t \in \Gamma}$ be an infinite independent sequence of n -dimensional random vectors taking values in $\{0, 1\}^n$. By using (2), define a random field $\{X_s\}_{s \in \Lambda}$ so that X_s is the i -th component of $\mathbf{Y}_t$. Then we can prove that $\{X_s\}_{s \in \Lambda}$ is indeed a random field by Kolmogorov's consistency theorem [24, p.129]. Below we explain that $D := C - C$ is a d-set of this random field. Let P, Q be finite subsets of Λ such that the pair of a $\#P$ -dimensional random vector $(X_s)_{s \in P}$ and a $\#Q$ -dimensional random vector $(X_s)_{s \in Q}$ is not independent. Take minimal subsets P', Q' of Γ such that $P \subset P' + C$ and $Q \subset Q' + C$. Because of the choice of the finite sets P, Q , the random vector $(X_s)_{s \in P}$ is not independent from the random vector $(X_s)_{s \in Q}$. So the pair of random vectors $(\mathbf{Y}_t)_{t \in P'}$ and $(\mathbf{Y}_t)_{t \in Q'}$ is not independent. But the set $\{\mathbf{Y}_t\}_{t \in \Gamma}$ is independent, so there is $t \in P' \cap Q'$. Then, by the minimality of P', Q' , there are $p, q \in C$ such that $t + p \in P$ and $t + q \in Q$. Therefore $p - q \in P - Q$. Thus $(P - Q) \cap D \neq \emptyset$.

Example 2 (FRMS caused by random shift of windows) For a model set $\Lambda(W)$ with both of $\mathfrak{G}$ and $\mathfrak{G}_{\text{int}}$ being Euclidean vector spaces, physicists often associate to each site $s \in L$ its own window $W_s = W + y_s$ where the “shift” y_s is an RV ranging over a window $R \subset \mathfrak{G}_{\text{int}}$ with nonempty interior. Then $W + R$ is again a window. For $s \in \Lambda(W + R)$, define an RV X_s such that $X_s = 1$ whenever the RV y_s takes a value in $s^* - W$, in other words, $s^* \in W_s = W + y_s$, while $X_s = 0$ otherwise. If the sequence $\{y_s\}_{s \in \Lambda(W)}$ of the RV's is independent, then the random field $\{X_s\}_{s \in \Lambda(W+R)}$ is independent, so it is an FRMS on a model set $\Lambda(W + R)$.

However, if $y_s(\omega) = y_t(\omega)$ for any $s, t \in L$ and for any ω of the probability space, then the random field is not an FRMS, because no finite d-set can be taken owing to the existence of a relatively dense subset $\Gamma := \Lambda((W + R) \setminus W) \subset \Lambda(W + R)$ such that a sequence $\{X_s\}_{s \in \Gamma}$ of the RV's is not independent. Here the relative density follows from $\text{Int}((W + R) \setminus W) \neq \emptyset$, and the proof is in the appendix.

3.2 Finitely periodic model sets and internal space

Roughly speaking, we say a subset A of H is a finitely periodic, if and only if for each $x \in (A - A) \setminus \{0\}$ there is a positive integer ℓ such that every arithmetical progression with the common difference being x has length smaller than ℓ . The precise definition is as follows:

Definition 5 For any LCAG H , any set $A \subset H$, any $x \in H$ and any $s \in A$, $\ell_A(x; s)$ is 0 if $x = 0$, or else is the maximum $k \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$ such that $\{s - nx; 0 \leq n \leq k\} \subset A$. So if $x \notin A - A$, then $\ell_A(x; s)$ is 0 for any $s \in A$. Set

$$\ell_A(x) := \max_{s \in A} \ell_A(x; s).$$

If $\ell_A(x)$ is finite for any $x \in A - A$, we say A is *finitely periodic*. We say an FRMS $\{X_s\}_{s \in A}$ is *finitely periodic*, if so is A .

Lemma 3 1. *If a model set is finitely periodic, the star map is injective.*
 2. *A model set is finitely periodic, if the star map is injective and the internal space is isomorphic to $\mathbb{R}^l \times \mathbb{Z}^m \times F$ for some $l, m \in \mathbb{Z}_{\geq 0}$ and some finite Abelian group F .*

Proof (1) If the kernel of the star map has nonzero element x , then for any site s in the model set, we have an infinite arithmetical progression $(s + nx)_{n \in \mathbb{N}}$.
 (2) Assume the model set $\Lambda(W)$ over a c.-p. scheme $(\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$ is not finitely periodic. Let $N = \#F$. Then there is $x \in (\Lambda(W) - \Lambda(W)) \setminus \{0\}$ satisfies that for all $\ell \in \mathbb{N}$ there is $s_\ell \in \Lambda(W)$ such that

$$\{s_\ell - nx; 0 \leq n < N\ell\} \subseteq \Lambda(W).$$

Let φ be the isomorphism from $\mathfrak{G}_{\text{int}}$ to the $\mathbb{R}^l \times \mathbb{Z}^m \times F$ and let ψ_1 be a homomorphism $L \xrightarrow{*} L^* \hookrightarrow \mathfrak{G}_{\text{int}} \xrightarrow{\varphi} \mathbb{R}^l \times \mathbb{Z}^m \times F \xrightarrow{\pi} \mathbb{R}^l \times \mathbb{Z}^m$ where π is the canonical projection, and ψ_2 be the other homomorphism from L to F . Since F is a finite group, $N\psi_2(x) = 0$. The injectivity of the star map implies $\psi_1(Nx) \neq 0$.

Then

$$\{\psi_1(s_\ell) - n\psi_1(x); 0 \leq n < N\ell\} \subseteq \pi(\varphi(W)).$$

We can take the integer ℓ greater than d/d where d is the diameter of the compact set $\pi(\varphi(Cl(W))) \subseteq \mathbb{R}^l \times \mathbb{Z}^m$ and d is the norm of $\psi_1(Nx)$. But this is a contradiction. $\square$

Example 3 1. The vertex sets of the rhombic Penrose tilings are finitely periodic model sets with $\mathfrak{G}_{\text{int}} = \mathbb{C} \times (\mathbb{Z}/5\mathbb{Z})$ and injective star maps. See [19, Section 3.2].

2. If an internal space is the compact Abelian group of p -adic integers, we can find a model set [8] which is not finitely periodic.

3.3 Decomposition of finitely randomized model sets

Definition 6 A *cut-and-project subscheme* of a c.-p. scheme $(\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$ is a c.-p. scheme $(\mathfrak{G}, \mathfrak{H}_{\text{int}}, \tilde{M})$ such that $\tilde{M} \subset \tilde{L}$, $\mathfrak{H}_{\text{int}} \subset \mathfrak{G}_{\text{int}}$, and the topology of $\mathfrak{H}_{\text{int}}$ is the relative topology induced from $\mathfrak{G}_{\text{int}}$.

Lemma 4 *If an LCAG $\mathfrak{H}_{\text{int}}$ is a subgroup of an LCAG $\mathfrak{G}_{\text{int}}$, then the Haar measure θ of $\mathfrak{G}_{\text{int}}$ restricted to $\mathfrak{H}_{\text{int}}$ is a Haar measure ϑ of $\mathfrak{H}_{\text{int}}$ such that*

$$\theta(\partial_{\mathfrak{G}_{\text{int}}}(B)) = 0 \Rightarrow \vartheta(\partial_{\mathfrak{H}_{\text{int}}}(B \cap \mathfrak{H}_{\text{int}})) = 0.$$

Proof It is immediate that ϑ is a Haar measure of $\mathfrak{H}_{\text{int}}$. Note $\partial_{\mathfrak{H}_{\text{int}}}(B \cap \mathfrak{H}_{\text{int}})$ is $(Cl(B \cap \mathfrak{H}_{\text{int}}) \cap \mathfrak{H}_{\text{int}}) \setminus (Int(B \cap \mathfrak{H}_{\text{int}}) \cap \mathfrak{H}_{\text{int}})$ contained by $(Cl(B) \cap \mathfrak{H}_{\text{int}}) \setminus Int(B) = \partial_{\mathfrak{G}_{\text{int}}}(B) \cap \mathfrak{H}_{\text{int}}$. So, if $\theta(\partial_{\mathfrak{G}_{\text{int}}}(B)) = 0$ then $\theta(\partial_{\mathfrak{H}_{\text{int}}}(B \cap \mathfrak{H}_{\text{int}})) = 0$, which implies $\vartheta(\partial_{\mathfrak{H}_{\text{int}}}(B \cap \mathfrak{H}_{\text{int}})) = 0$ since ϑ is the restriction of $\mathfrak{G}_{\text{int}}$'s Haar measure θ to $\mathfrak{H}_{\text{int}}$. $\square$

Lemma 5 *For any c.-p. scheme $(\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$ with L being finitely generated and for any finite subset $D \subset L$ there are a c.-p. subscheme $\mathcal{S} = (\mathfrak{G}, \mathfrak{H}_{\text{int}}, \tilde{M})$ of $(\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$ and a finite complete representation system R of L/M with $D \subset R$.*

Proof By the structure theorem of a finitely generated Abelian group, $\tilde{L}$ is isomorphic to $\mathbb{Z}^v \times \mathbb{Z}/\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}/\mathbb{Z}_{n_u}$ for some $u, v \in \mathbb{Z}_{\neq 0}$ and for some integers $n_1, \dots, n_u \geq 2$. Because D is regarded as a finite subset of $\mathbb{Z}^v \times \mathbb{Z}/\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}/\mathbb{Z}_{n_u}$, there is an integer k greater than the “first components” of any elements of D . Without loss of generality, each n_i divides k . Then put $M := kL$. We can find a finite complete representation system $R \supseteq D$. Let $\tilde{M} := \{(t, t^*) ; t \in M\}$ and $\mathfrak{H}_{\text{int}} := Cl(M^*) \subset \mathfrak{G}_{\text{int}}$. Then $\mathfrak{H}_{\text{int}}$ is a σ -compact LCAG with the relative topology induced from $\mathfrak{G}_{\text{int}}$, and $(\mathfrak{G}, \mathfrak{H}_{\text{int}}, \tilde{M})$ is a c.-p. scheme with the star map being a restriction of the star map of $(\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$. $\square$

We consider the following condition on FRMS's:

Condition 1 $\Lambda_{\mathcal{O}}(W)$ is a finitely periodic model set over a c.-p. scheme $\mathcal{O} = (\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$ with L being finitely generated.

Recall that θ is a Haar measure of $\mathfrak{G}_{\text{int}}$. The subscheme $\mathcal{S}$ given in Lemma 5 is used in the following theorem:

Theorem 1 (Decomposition) *Suppose an FRMS $\{X_s\}_s$ $\Lambda_{\mathcal{O}}(W)$ satisfies Condition 1. Then there are a c.-p. subscheme $\mathcal{S} = (\mathfrak{G}, \mathfrak{H}_{\text{int}}, \tilde{M})$ of the c.-p. scheme $\mathcal{O}$ and a finite complete representation system $R = \{r_C \in L ; C \in L/M\}$ of L/M such that following holds:*

for each $g \in \Lambda_{\mathcal{O}}(W) - \Lambda_{\mathcal{O}}(W)$, each $C \in L/M$, each $r \in R$ with $r \equiv g \pmod{M}$, and each nonnegative integer $k \leq \ell := \ell_{\Lambda_{\mathcal{O}}(W)}(g - r)$, there is a relatively compact set $V_{C,k} \subseteq W$ such that for

$$S_{C,k} := r_C + \Lambda(V_{C,k}), \quad (3)$$

we have

1. *the sequence $\{X_s \overline{X_s - g} ; s \in S_{C,k}, s - g \in \Lambda_{\mathcal{O}}(W)\}$ is independent; and*
2. *if the Haar measure $\theta(\partial W)$ is 0, then $\text{dense}_{D_n - n}(S_{C,k})$ exists.*

Furthermore $\Lambda_{\mathcal{O}}(W)$ is the disjoint union of all the $S_{C,k}$'s.

Proof Let the subscheme $\mathcal{S}$ and the finite complete representation system R be as in Lemma 5 applied for the c.-p. scheme $\mathcal{O}$ and a d-set D of the FRMS $\{X_s\}_{s \in \Lambda_{\mathcal{O}}(W)}$. Since $0 \in D \subseteq R$ by Lemma 5, if $0 = g \equiv r \pmod{M}$ then $r = 0$. Because $\Lambda_{\mathcal{O}}(W)$ is finitely periodic, ℓ is finite, and we have relatively compact sets

$$W_k := \{y \in W ; \ell_W(g^* - r^* ; y) = k\} \quad (0 \leq k \leq \ell).$$

Then $\Lambda_{\mathcal{O}}(W) = \coprod_{k=0}^{\ell} \{s \in L ; s^* \in W_k\}$ where $\coprod$ is a disjoint union. Since $L = \coprod_{C \in L/M} (r_C + M)$, we have $\Lambda_{\mathcal{O}}(W) = \coprod_{k=0}^{\ell} \coprod_{C \in L/M} \{s \in r_C + M ; s^* - r_C^* \in (W_k - r_C^*) \cap \mathfrak{H}_{\text{int}}\}$. Therefore $\Lambda_{\mathcal{O}}(W) = \coprod_{k=0}^{\ell} \coprod_{C \in L/M} \left(r_C + M \cap (W_k - r_C^*) \cap \mathfrak{H}_{\text{int}} \right)$. So define

$$V_{C,k} := (W_k - r_C^*) \cap \mathfrak{H}_{\text{int}}. \quad (4)$$

Now we prove the assertion (1) of Theorem 1 below. By Lemma 2 with $N = \{-g, 0\}$, it is sufficient to prove the following claim : for any distinct $s, t \in S_{C,k}$, $(\{s, s - g\} - \{t, t - g\}) \cap D = \emptyset$. In other words, for any $g \in D$ we have (i) $s - t = (s - g) - (t - g) \neq g$; (ii) $s - (t - g) \neq g$; and (iii) $(s - g) - t \neq g$.

It is proved as follows: By Lemma 5, $M \cap D = \{0\}$, which derives the assertion (i) from $s - t \in M \setminus \{0\}$. If $g = 0$, then (ii) and (iii) follow from (i). So let $g \neq 0$. Assume (ii) is false. Then $g - g = s - t \in M$, so $g \equiv r \pmod{M}$ for a unique $r \in R$. Then $g = r$, because $g \in D$ belongs to R . Therefore $t - s = g - r$. But $t^* - (g^* - r^*) = s^* \in W_k \ni t^*$ contradicts against the definition of W_k . Therefore (ii) holds. The assertion (iii) follows from (ii) since $D = -D$.

We prove the assertion (2) of Theorem 1. Recall that the LCAG $\mathfrak{H}_{\text{int}}$ of the subscheme $\mathcal{S}$ is a subgroup of the LCAG $\mathfrak{G}_{\text{int}}$. As in Lemma 4, we write ϑ for a Haar measure of $\mathfrak{H}_{\text{int}}$. The premise $\theta(\partial_{\mathfrak{G}_{\text{int}}} W) = 0$ implies

$$\vartheta(\partial_{\mathfrak{H}_{\text{int}}} V_{C,k}) = 0. \quad (5)$$

To see it first observe that $W_k = W \cap (W + g - r) \cap \cdots \cap (W + k(g - r)) \setminus (W + (k + 1)(g - r))$. So the premise implies $\theta(\partial_{\mathfrak{G}_{\text{int}}} (W_k - r_C^*)) = 0$ because of fact $\partial_{\mathfrak{G}_{\text{int}}} (A \cap B) \subset \partial_{\mathfrak{G}_{\text{int}}} (A) \cup \partial_{\mathfrak{G}_{\text{int}}} (B)$ and fact $\partial_{\mathfrak{G}_{\text{int}}} (A) = \partial_{\mathfrak{G}_{\text{int}}} (\mathfrak{G}_{\text{int}} \setminus A)$. Thus (5) follows from Lemma 4.

Since $\{D_n - r_C\}_n$ is a van Hove sequence too, $\text{dense}_{D_n - r_C}(\Lambda(V_{C,k}))$ converges by Proposition 2. But it is $\text{dense}_{D_n - r_C}(S_{C,k})$ by the definition (3) of $S_{C,k}$. $\square$

4 Absolutely continuous component of diffraction and covariance

If a complex-valued FRMS $\omega = \{X_s\}_{s \in \Lambda}$ satisfies Condition 1 and all of the expectation $E[X_s]$ and the covariances between X_s and X_{s-g} are “continuous” with respect to $s^* \in \mathfrak{G}_{\text{int}}$ for any $g \in \Lambda - \Lambda$, then we quantitatively give the diffraction measure of ω as follows:

- the inverse Fourier transform of the absolutely continuous component is a Dirac comb whose support is the smallest d-set; and
- the pure-point component is the diffraction measure of a Dirac comb which is the expectation of the FRMS ω .

Here

Definition 7 The *expectation* of an FRMS $\omega = \{X_s\}_{s \in \Lambda}$ is, by definition, a Dirac comb $E[\omega] = \{E[X_s]\}_{s \in \Lambda}$, that is, $\sum_{s \in \Lambda} E[X_s] \delta_s$.

We use Kolmogorov's strong law of large numbers. By the variance of an RV X , we mean $V[X] = E[|X - E[X]|^2] = E[|X|^2] - E[X]E[\overline{X}]$.

Proposition 5 ([24, Corollary 1.4.9]) *Suppose that $\{b_m; m \in \mathbb{N}\}$ is a nondecreasing sequence of positive numbers which tends to infinity, and that a set $\{X_n\}_{n \in \mathbb{N}}$ of square integrable RV's is independent. If $\sum_{m=1}^{\infty} V[X_m] b_m^{-2} < \infty$, then*

$$\lim_{m \leq} \frac{1}{b_m} \sum_{i=1}^m (X_i - E[X_i]) = 0 \quad (\text{almost surely}).$$

Lemma 6 *If a set $A \subset \mathfrak{G}$ is a nonempty and discrete and $\{Y_s\}_{s \in A}$ is an independent set of RV's with the variances $V[Y_s]$ bounded uniformly from above, then for any van Hove sequence $\{D_n\}_n$*

$$\lim_{n \leq} \frac{1}{\#(A \cap D_n)} \sum_{s \in A \cap D_n} (Y_s - E[Y_s]) = 0 \quad (\text{almost surely}).$$

Proof There is an enumeration $\{s_i\}_{i \in \mathbb{N}}$ of A without repetition which exhausts $A \cap D_1$ first then $A \cap D_2$, $A \cap D_3$, and so on, because $A \subseteq \bigcup_n D_n$ and because $\#(A \cap D_n) < \infty$ follows from the discreteness of A and the compactness of D_n . For $m \in \mathbb{N}$ let b_m be $\#(A \cap D_n)$ where n is the smallest integer such that $s_m \in A \cap D_n$. If $m = \#(A \cap D_n)$ for such n , then $b_m = m$. In general, we have $b_m \geq m$, because the van Hove sequence $\{D_n\}_n$ is increasing. So $\sum_{m=1}^{\infty} b_m^{-2} \leq \sum_{m=1}^{\infty} m^{-2} < \infty$. Because $V[Y_s]$ is uniformly bounded from above by the premise, we have $\sum_{m=1}^{\infty} V[Y_{s_m}] m^{-2} < \infty$. Since the sequence $\{Y_{s_i}\}_{i \in \mathbb{N}}$ is independent from the premise, Proposition 5 implies that the sequence $\left\{ \sum_{i=1}^m (Y_{s_i} - E[Y_{s_i}]) / b_m \right\}_{m \in \mathbb{N}}$ converges to 0 almost surely. Hence a subsequence $\left\{ \sum_{s \in A \cap D_n} (Y_s - E[Y_s]) / \#(A \cap D_n) \right\}_{n \in \mathbb{N}}$ does so almost surely. $\square$

By the *covariance* between complex-valued RV's X_s and X_t , we mean

$$\text{Cov}[X_s, X_t] = E[(X_s - E[X_s])(\overline{X_t - E[X_t]})] = E[X_s \overline{X_t}] - E[X_s]E[\overline{X_t}].$$

Condition 2 *An FRMS $\{X_s\}_{s \in \Lambda(W)}$ has functions $e, c_g \in C_c(\mathfrak{G}_{\text{int}})$ such that*

$$\begin{aligned} E[X_s] &= e(s^*), & (s \in \Lambda(W)); \\ \text{Cov}[X_s, X_{s+g}] &= c_g(s^*), & (g \in \Lambda(W) - \Lambda(W), s \in \Lambda(W) \cap (\Lambda(W) + g)). \end{aligned}$$

Example 4 In Example 1, suppose that there are $m \in C(\mathfrak{G}_{\text{int}}, \mathbb{R}^n)$ and $S \in C(\mathfrak{G}_{\text{int}}, \mathbb{R}^{n \times n})$ such that $m(t^*)$ is the mean $E[\mathbf{Y}_t]$ and $S(t^*)$ is the covariance matrix $E[(\mathbf{Y}_t - E[\mathbf{Y}_t])(\mathbf{Y}_t - E[\mathbf{Y}_t])^\top]$ for all $t \in \Gamma$. Then functions e, c_g ($g \in \Lambda - \Lambda$) indeed belong to $C_c(\mathfrak{G}_{\text{int}})$. For the FRMS of Example 2, assume further that each shift y_s is subject to a continuous probabilistic density function h with $\text{supp } h = R$. Then, for $s \in L$, $(1_W * h)(s^*) = \int_{\mathfrak{G}_{\text{int}}} 1_W(s^* - y)h(y)dy = \int_{W - s^*} h(y)dy = P(s^* \in W + y_s)$, the probability for $s \in \Lambda(W + R)$ to indeed appear. So $e = 1_W * h$.

Theorem 2 Let $\omega = \{X_s\}_{s \in \Lambda(W)}$ be an FRMS such that Condition 2 holds, $\{E[|X_s|^4]\}_{s \in \Lambda(W)}$ is bounded, and W is compact but the Haar measure $\theta(\partial W)$ of $\mathfrak{G}_{\text{int}}$ is 0. Then the diffraction measure $\widehat{\gamma_\omega}$ of ω is almost surely $\widehat{\gamma_{E[\omega]}} + A$, where

1. $\widehat{\gamma_{E[\omega]}}$ is a pure-point diffraction measure.
2. A is an absolutely continuous, real-valued measure on $\mathfrak{G}$. In fact, there is some d -set D of the FRMS ω such that the Radon-Nikodým derivative of A with respect to the Haar measure $d\chi$ of $\mathfrak{G}$ is $\sum_{g \in D} A_g \chi(-g)$ where

$$A_g = \int_{W_{\uparrow}(W+g^*)} c_g(y)dy = \lim_{n \leq s \in \Lambda(W)_{\uparrow}} \sum_{(A(W)+g)_{\uparrow} \cap D_n} \frac{\text{Cov}[X_s, X_{s-g}]}{|D_n|}.$$

Proof We use the notation of Theorem 1. Let $g \in \Lambda_{\mathcal{O}}(W) - \Lambda_{\mathcal{O}}(W)$. Because $\Lambda_{\mathcal{O}}(W)$ is the disjoint union of $S_{C,k}$'s over $(C, k) \in (L/M) \times \{0, \dots, \ell\}$, we have $s, s-g \in \Lambda_{\mathcal{O}}(W) \cap D_n$ if and only if there is a unique $(C, k) \in (L/M) \times \{0, \dots, \ell\}$ such that $s \in S_{C,k} \cap (\Lambda_{\mathcal{O}}(W) + g) \cap D_n \cap (D_n + g)$. Furthermore $r_C - g \in R - M$ and

$$S_{C,k} \cap (\Lambda_{\mathcal{O}}(W) + g) = r_C + \Lambda(W_{C,k}), \quad (6)$$

where

$$W_{C,k} := V_{C,k} \cap ((W - r_C^* + g^*) \cap \mathfrak{H}_{\text{int}}). \quad (7)$$

Thus,

$$\lim_{n \leq s \in \Lambda_{\mathcal{O}}(W)_{\uparrow}} \sum_{D_n} \frac{(\dots)}{|D_n|} = \sum_{\substack{C \\ 0 \leq k \leq \ell}} \lim_{n \leq s \in \substack{(r_C + \Lambda_S(W_{C,k})) \\ D_n \cap (D_n + g)}} \sum_{D_n \cap (D_n + g)} \frac{(\dots)}{|D_n|}. \quad (8)$$

Suppose $C \in L/M$, $0 \leq k \leq \ell$ and $S_{C,k} \neq \emptyset$.

Claim 1 For any $f \in C_c(\mathfrak{G}_{\text{int}})$, as $n \rightarrow \infty$, both of the summation of $f(s^*)|D_n|^{-1}$ over $s \in (r_C + \Lambda(W_{C,k})) \cap D_n \cap (D_n + g)$ and the summation of $f(s^*)|D_n|^{-1}$ over $s \in (r_C + \Lambda(W_{C,k})) \cap D_n$ converge to the same value.

Proof The absolute value Δ of the difference between the two summations is not greater than $\sum_{s \in (r_C + \Lambda_S(W_{C,k})) \setminus (D_n \setminus (D_n + g))} |f(s^*)| / |D_n|$. Here, by (6), a set $r_C + \Lambda_S(W_{C,k})$ is a subset of $\Lambda_\phi(W) + g$, and is uniformly discrete. Thus by Lemma 1, there is a compact neighborhood U of 0 such that for all n , $\#((r_C + \Lambda_S(W_{C,k})) \cap (D_n \setminus (D_n + g))) \leq |D_n \setminus (D_n + g)| / |U|$. By the premise $f \in C_c(\mathfrak{G}_{\text{int}})$, there is $b \geq 0$ such that $|f(y)| \leq b$ for all $y \in \mathfrak{G}_{\text{int}}$. Thus $\Delta \leq |D_n \setminus (D_n + g)| \cdot b |U|^{-1} / |D_n| = b |U|^{-1} |D_n \setminus (D_n + g)| / |D_n| \leq b |U|^{-1} |\mathfrak{P}^{0, -g}(D_n)| / |D_n| \rightarrow 0$. Moreover, by Proposition 2, the summation of $f(s^*) / |D_n|$ over $s \in (r_C + \Lambda_S(W_{C,k})) \cap D_n$ converges as n goes to infinity. To see it, the definition (7) of $W_{C,k}$ implies $\partial_{\mathfrak{H}_{\text{int}}}(W_{C,k})$ is contained by $\partial_{\mathfrak{H}_{\text{int}}}(V_{C,k}) \cup \partial_{\mathfrak{H}_{\text{int}}}((W - r_C^* + g^*) \cap \mathfrak{H}_{\text{int}})$ which has null measure with respect to the Haar measure ϑ of $\mathfrak{H}_{\text{int}}$, where $\vartheta(\partial_{\mathfrak{H}_{\text{int}}}(V_{C,k})) = 0$ by (5), while the premise $\theta(\partial W) = 0$ of Theorem 2 implies that $(W - r_C^* + g^*) \cap \mathfrak{H}_{\text{int}}$ has ϑ -null boundary by Lemma 4. $\square$

Claim 2 It holds almost surely that as n goes to ∞ ,

$$\sum_{s \in (r_C + \Lambda_S(W_{C,k})) \setminus (D_n \setminus (D_n + g))} \frac{X_s \overline{X_{s-g}}}{|D_n|} - \sum_{s \in (r_C + \Lambda_S(W_{C,k})) \cap (D_n \setminus (D_n + g))} \frac{\mathbb{E}[X_s \overline{X_{s-g}}]}{|D_n|} \rightarrow 0. \quad (9)$$

Proof Note that the left-hand side of (9) is the product of the following two:

$$u_n := \sum_{s \in (r_C + \Lambda_S(W_{C,k})) \setminus (D_n \setminus (D_n + g))} \frac{X_s \overline{X_{s-g}} - \mathbb{E}[X_s \overline{X_{s-g}}]}{\# \{(r_C + \Lambda_S(W_{C,k})) \cap D_n \cap (D_n + g)\}},$$

$$v_n := \# \{(r_C + \Lambda_S(W_{C,k})) \cap D_n \cap (D_n + g)\} / |D_n|.$$

Here u_n tends to 0 almost surely, because by the premise the variances $V[X_s \overline{X_{s-g}}] \leq \mathbb{E}[|X_s \overline{X_{s-g}}|^2] \leq \mathbb{E}[|X_s|^4]/2 + \mathbb{E}[|X_{s-g}|^4]/2$ are uniformly bounded, from which Theorem 1 (1), the equality (8) and Lemma 6 imply that $\lim_n u_n = 0$ almost surely. On the other hand, v_n tends to dense $\lim_n \# \{(r_C + \Lambda_S(W_{C,k})) \cap D_n \cap (D_n + g)\} < \infty$ by Claim 1. $\square$

The second sum in the left-hand side of (9) has the following limit in the limit of n .

Claim 3 As n goes to infinity, $\sum_{s \in (r_C + \Lambda_S(W_{C,k})) \cap (D_n \setminus (D_n + g))} \mathbb{E}[X_s \overline{X_{s-g}}] / |D_n|$ converges to

$$\lim_n \sum_{s \in (r_C + \Lambda_S(W_{C,k})) \setminus (D_n \setminus (D_n + g))} \frac{e(s^*) \overline{e(s^* - g)}}{|D_n|} + \lim_n \sum_{s \in (r_C + \Lambda_S(W_{C,k})) \cap D_n} \frac{c_g(s^*)}{|D_n|}. \quad (10)$$

Proof By Condition 2, Claim 1 and Proposition 2, the two limits in (10) are convergent, and (10) is

$$\lim_{n \leq} \frac{1}{|D_n|} \sum_{s \in (r_C + A_S(W_{C,k})) \setminus D_n \setminus (D_n + g)} \left(e(s^*) \overline{e(s^* - g^*)} + c_g(s^*) \right).$$

But the summand is $E[X_s \overline{X_{s-g}}]$ by Condition 2. $\square$

By taking the summation (9) and (10) respectively over any $(C, k) \in (L/M) \times \{0, \dots, \ell\}$ such that $S_{C,k} \neq \emptyset$, by (8), we have almost surely $\eta_\omega(g) = \eta_{E[\omega]}(g) + A_g$, and thus $\widehat{\gamma}_\omega = \widehat{\gamma_{E[\omega]}} + \sum_g A_g \chi(-g)$.

We continue the proof of Theorem 2.

The assertion (1) of Theorem 2 holds because the pure-pointness of $E[\omega]$ follows from Proposition 4 and Condition 2.

The assertion (2) is proved as follows: The Radon-Nikodým derivative $A(\chi) = \sum_g A_g \chi(-g)$ is actually $A(\chi) = \sum_g A_g \chi(-g)$ for any d-set D of the FRMS ω . Indeed $\text{Cov}[X_s, X_{s-g}]$ is equal to 0 for any $g \in (S - S) \setminus D$ and for any $s \in \Lambda_O(W) \cap (\Lambda_O(W) + g)$, because a pair of X_s and X_{s-g} is independent for this g by Definition 3. This completes the proof of Theorem 2. $\square$

Corollary 1 *Under the same assumption as Theorem 2, the smallest d-set of the FRMS is the support of the Dirac comb which is obtained by the inverse Fourier transform of the absolutely continuous component A of the diffraction measure $\widehat{\gamma}_\omega$.*

Proof The inverse Fourier transform of A is $\int_{\widehat{\mathfrak{G}}} \sum_g A_g \chi(-g) \chi(x) d\chi$ which is $\sum_g A_g \int_{\widehat{\mathfrak{G}}} \chi(-g+x) d\chi = \sum_g A_g \delta_g(x)$. $\square$

Example 5 If the FRMS in Theorem 2 is independent, it has a d-set $D = \{0\}$ by Lemma 2 and the absolutely continuous component of the diffraction measure $\widehat{\gamma}_\omega$ is $A = \lim_n \sum_{s \in \Lambda(W) \setminus D_n} V[X_s] / |D_n|$.

If we add a mild condition “ $Cl(Int(W)) = W$ ” to the theorem, we can quantitatively provide the pure-point component $\widehat{\gamma_{E[\omega]}}$ by using a following theorem (Theorem 3) (and can dispense with Proposition 4.)

5 Diffraction of weighted Dirac comb and torus parametrization

Let ω be a weighted Dirac comb on $\Lambda(W)$ where $\Lambda(W)$ is a c.-p. set over a c.-p. scheme $\mathcal{S} = (\mathfrak{G}, \widehat{\mathfrak{G}_{\text{int}}}, \tilde{L})$. To describe the support of $\widehat{\gamma}_\omega$, we use the dual c.-p. scheme $(\widehat{\mathfrak{G}}, \widehat{\mathfrak{G}_{\text{int}}}, \tilde{\mathcal{L}})$ for the c.-p. scheme $\mathcal{S}$. Here the lattice $\tilde{\mathcal{L}}$ is the *annihilator* of $\tilde{L}$ in $\widehat{\mathfrak{G}} \times \widehat{\mathfrak{G}_{\text{int}}} \simeq \widehat{\mathfrak{G}} \times \widehat{\mathfrak{G}_{\text{int}}}$. That is $\tilde{\mathcal{L}}$ is the LCAG of $(\chi, \eta) \in \widehat{\mathfrak{G}} \times \widehat{\mathfrak{G}_{\text{int}}}$ such that $\chi(s)\eta(s^*) = 1$ for all $(s, s^*) \in \tilde{L}$. Then $(\widehat{\mathfrak{G}}, \widehat{\mathfrak{G}_{\text{int}}}, \tilde{\mathcal{L}})$ is indeed a c.-p. scheme. See [19, Section 5] for the proof. Let $\mathcal{L}$ ($\mathcal{L}^*$ resp.)

stand for the canonical projection of $\tilde{\mathcal{L}}$ to $\widehat{\mathfrak{G}}$ ($\widehat{\mathfrak{G}_{\text{int}}}$ resp.), and let the star map be $(-)^* : \mathcal{L} \rightarrow \mathcal{L}^*$. Then

$$\chi(s) = \overline{\chi^*(s^*)} \quad (\chi \in \mathcal{L}, s \in L). \quad (11)$$

Let $\mathfrak{T}$ be a compact Abelian group $(\mathfrak{G} \times \mathfrak{G}_{\text{int}})/\tilde{L}$. Then the lattice $\tilde{\mathcal{L}}$ of the c.-p. scheme satisfies

$$\tilde{\mathcal{L}} \simeq \widehat{\mathfrak{T}}, \quad (12)$$

and the composite $\varpi : \widehat{\mathfrak{T}} \rightarrow \mathcal{L}$ of the isomorphism and the projection restricted to $\tilde{\mathcal{L}}$ is indeed a bijection:

$$\varpi(\xi) := \xi((\bullet, 0) + \tilde{L}) \in \mathcal{L}. \quad (13)$$

Recall that $|\tilde{L}| = \int_{\mathfrak{T}} dx dy$.

We say an FLC set $P \subset \mathfrak{G}$ is *repetitive* if for every compact set $K \subset \mathfrak{G}$ there exists compact set $K' \subset \mathfrak{G}$ such that for all $t_1, t_2 \in \mathfrak{G}$ there exists $s \in K'$ such that $(t_1 + P) \cap K = (s + t_2 + P) \cap K$. According to [23], we say a window $W \subset \mathfrak{G}_{\text{int}}$ has *no nontrivial translation invariance* if $\{c \in \mathfrak{G}_{\text{int}} ; c + W = W\} = \{0\}$.

Theorem 3 *Assume that $\Lambda(W)$ is a repetitive model set over a c.-p. scheme $(\mathfrak{G}, \mathfrak{G}_{\text{int}}, \tilde{L})$ where $\Lambda(W) - \Lambda(W)$ generates L , $W = Cl(Int(W))$, W has no non-trivial translation invariance, and the Haar measure $\theta(\partial W)$ is 0. Assume furthermore that*

$$b \in C_c(\mathfrak{G}_{\text{int}}), \text{ supp } b \subset W, \omega = \sum_{s \in \Lambda(W)} b(s^*) \delta_s. \quad (14)$$

Then the diffraction measure of ω is $\widehat{\gamma_\omega} = \sum_{\chi \in \mathcal{L}} \frac{|\hat{b}(\chi^)|^2}{|\tilde{L}|^2} \delta_\chi$.*

The theorem with the physical space $\mathfrak{G}$ being $\mathbb{R}^n$ was proved by Hof [13], and that for Dirac combs with constant weights was by Schlottmann [23]. A general theorem for weighted Dirac comb with the weights arising from an “admissible” Radon-Nikodým derivative of the $\tilde{L}$ -invariant measure was studied in Lenz-Richard [17, Theorem 3.3]. Our theorem is another form of a weaker version of Lenz-Richard’s theorem.

In order to prove the theorem, we employ a uniquely ergodic dynamical system $X_{\Lambda(W)}$ made from $\Lambda(W)$, and connection of the autocorrelation measure γ_ω to a complex Hilbert space over $X_{\Lambda(W)}$. Then we prove lemmas about the so-called *torus parametrization* of $\Lambda(W)$, introduced in [3] and generalized by [23].

Proposition 6 ([23])

1. *For every FLC set $\Lambda \subset \mathfrak{G}$, the closure X_Λ of the $\mathfrak{G}$ -orbit $\{\Lambda + g ; g \in \mathfrak{G}\}$ of Λ with some uniform topology is a complete, compact Hausdorff space.*

2. Suppose a model set $\Lambda := \Lambda(W)$ satisfies the same assumptions as Theorem 3. Then X_Λ will be a minimal and uniquely ergodic dynamical system, with the $\mathfrak{G}$ -action $(x, P) \in \mathfrak{G} \times X_\Lambda \mapsto P + x \in X_\Lambda$.

Hereafter, the uniquely ergodic probability measure of X_Λ will be denoted by ν , and the complex Hilbert space over X_Λ with the inner product

$$\langle \Phi_1, \Phi_2 \rangle_\nu := \int_{X_\Lambda} \Phi_1(P) \overline{\Phi_2(P)} d\nu(P) \quad (\Phi_i \in L^2(X_\Lambda, \nu))$$

will be denoted by $L^2(X_\Lambda, \nu)$.

Proposition 7 (Torus parametrization) Assume the same assumptions as Theorem 3, and let $\mathfrak{T}$ be a compact Abelian group $(\mathfrak{G} \times \mathfrak{G}_{\text{int}})/\tilde{L}$ with the Haar probability measure τ , and let $\mathfrak{G}$ act on $\mathfrak{T}$ by $(x, t) \in \mathfrak{G} \times \mathfrak{T} \mapsto t + (x, 0) \in \mathfrak{T}$. Then there are a continuous surjection

$$\beta : X_\Lambda \rightarrow \mathfrak{T}$$

and a full measure subset X'_Λ of X_Λ such that (1) β preserves the $\mathfrak{G}$ -action, (2) $\beta(\Lambda) = \tilde{L}$, and (3) $\beta|_{X'_\Lambda}$ is injective with the range $\beta(X'_\Lambda)$ disjoint from $\beta(X_\Lambda \setminus X'_\Lambda)$.

From the proposition, we can derive a following:

Lemma 7 Let $L^2(\mathfrak{T}, \tau)$ be a complex Hilbert space with the inner product

$$\langle \alpha_1, \alpha_2 \rangle_\tau := \int_{\mathfrak{T}} \alpha_1(t) \overline{\alpha_2(t)} d\tau(t) \quad (\alpha_1, \alpha_2 \in L^2(\mathfrak{T}, \tau)).$$

Then we have a bijective isometry

$$\iota : L^2(X_\Lambda, \nu) \rightarrow L^2(\mathfrak{T}, \tau) \quad ; \quad \Theta \mapsto \Theta \circ (\beta)^{-1}.$$

By using [1, Proposition 1.4], we can prove the following:

Fact 1 If λ and μ are translationally bounded, nonnegative measures on $\mathfrak{G}$ and $\{D_n\}_n$ is a van Hove sequence on $\mathfrak{G}$, then in the vague topology

$$\lim_{n \leq} \frac{(\widetilde{\lambda|_{D_n}}) * \mu|_{D_n} - \tilde{\lambda} * (\mu|_{D_n})}{|D_n|} = \lim_{n \leq} \frac{(\widetilde{\lambda|_{D_n}}) * \mu - \tilde{\lambda} * (\mu|_{D_n})}{|D_n|} = 0.$$

In the following Lemma, recall that ω is a weighted Dirac comb mentioned in Theorem 3 and the weight depends on a function $b \in C_c(\mathfrak{G}_{\text{int}})$.

Lemma 8 Suppose the same assumptions as Theorem 3. Let Λ be $\Lambda(W)$. Then, for all $\varphi_1, \varphi_2 \in C_c(\mathfrak{G})$, there are unique $\Phi_1, \Phi_2 \in L^2(X_\Lambda, \nu)$ such that for any $x \in \mathfrak{G}$,

$$\Phi_i(\Lambda - x) = (\varphi_i * \omega)(x), \quad (15)$$

$$(\varphi_1 * \widetilde{\varphi_2} * \gamma_\omega)(0) = \langle \Phi_1, \Phi_2 \rangle_\nu. \quad (16)$$

Proof Let $b \in C_c(\mathfrak{G}_{\text{int}})$ be as in Theorem 3 and $\beta : X_\Lambda \rightarrow \mathfrak{T}$ be as in Proposition 7. Define $g_i \in C_c(\mathfrak{G} \times \mathfrak{G}_{\text{int}})$ by $g_i(x, y) := \varphi_i(-x)b(y)$. Put $K_i := \text{supp } g_i$. Let $P \in X_\Lambda$. Because $\beta(P) \in X_\Lambda$ is a discrete set, $\beta(P) \cap K_i$ is finite. Thus $\Phi_i(P) := \sum_{(x,y) \in \beta(P)} g_i(x, y)$ is indeed a finite sum and well-defined.

To establish $\Phi_i \in L^2(X_\Lambda, \nu)$, it is sufficient to verify that a function $f_i : \mathfrak{T} \rightarrow \mathbb{C} ; t \mapsto \sum_{(x,y) \in t} g_i(x, y)$ is continuous, because Φ_i is the composition of the continuous function β and f_i .

Because $g_i \in C_c(\mathfrak{G} \times \mathfrak{G}_{\text{int}})$, for all $\varepsilon > 0$ there exists a compact neighborhood $U \subset \mathfrak{G} \times \mathfrak{G}_{\text{int}}$ of 0 such that for all $z, z' \in \mathfrak{G} \times \mathfrak{G}_{\text{int}}$ with $z' \in z + U$, we have $|g_i(z') - g_i(z)| < \varepsilon$. Then $\{z + \tilde{L} ; z \in U\}$ is a compact neighborhood of 0 in $\mathfrak{T}$. Let $t, t' \in \mathfrak{T} = (\mathfrak{G} \times \mathfrak{G}_{\text{int}})/\tilde{L}$ such that $t' - t$ belongs to the compact neighborhood $\{z + \tilde{L} ; z \in U\}$. Then $t' = t + (u, v)$ for some $(u, v) \in U$. So $|f_i(t') - f_i(t)| \leq \sum_{z \in t} |g_i(z + (u, v)) - g_i(z)|$. We show it is less than $\#((K_i - U) \cup K_i) \cap t \times \varepsilon$. If $z \in t$ contributes to the summand, then $z + (u, v) \in K_i$ or $z \in K_i$, so $z \in ((K_i - U) \cup K_i) \cap t$. Here $((K_i - U) \cup K_i) \cap t$ is finite since t is a translation of the lattice $\tilde{L}$ and $(K_i - U) \cup K_i$ is compact. Thus f_i is continuous.

To prove (15), observe

$$\Phi_i(\Lambda - x) = \sum_{s \in L} \varphi_i(-s + x)b(s^*) \quad (17)$$

follows from $\beta(\Lambda - x) = \tilde{L} - (x, 0)$. By the condition $\text{supp } b \subset W$ of (14), the weight $b(s^*)$ vanishes for any $s \notin \Lambda(W)$. So the range L of s in the summation (17) can be replaced with Λ . Thus (15) holds.

The left-hand side of (16) is $\lim_n \left(\varphi_1 * \widetilde{\varphi_2 * \omega}|_{D_n} * \widetilde{\omega}|_{D_n} \right) (0) \cdot |D_n|^{-1}$ by definition of γ_ω . But by Fact 1, it is $\lim_n \left(\widetilde{(\varphi_2 * \omega)} * (\varphi_1 * \omega)|_{D_n} \right) (0) \cdot |D_n|^{-1}$, which is equal to $\lim_n \int_{D_n} \overline{\Phi_2(\Lambda - x)} \Phi_1(\Lambda - x) dx \cdot |D_n|^{-1}$ by (15). By the pointwise ergodic theorem [22], it is $\int_{X_\Lambda} \overline{\Phi_2(P)} \Phi_1(P) d\nu(P) = \langle \Phi_1, \Phi_2 \rangle_\nu$. $\square$

Here is a technical lemma concerning van Hove sequences and uniformly discrete sets.

Lemma 9 *For any uniformly discrete subset Λ of $\mathfrak{G}$, any bounded complex sequence $\{w_s\}_{s \in \Lambda}$, any $\chi \in \widehat{\mathfrak{G}}$, any $\varphi \in C_c(\mathfrak{G})$, and any van Hove sequence $\{D_n\}_n$ of $\mathfrak{G}$,*

$$\frac{1}{|D_n|} \left| \sum_{s \in \Lambda} w_s \int_{D_n} \chi(x) \varphi(x - s) dx - \sum_{s \in \Lambda \setminus D_n} w_s \int_{\mathfrak{G}} \chi(x) \varphi(x - s) dx \right|$$

tends to 0, as n goes to ∞ .

Proof The numerator is bounded from above by

$$\sum_{s \in \bigcup_{\Lambda} D_n} |w_s| \int_{\mathfrak{G} \setminus D_n} |\varphi(x-s)| dx + \sum_{s \in \bigcup_{\Lambda} D_n} |w_s| \int_{D_n} |\varphi(x-s)| dx. \quad (18)$$

The summands in the former summation and the latter summation are bounded, because so are the sequence $\{w_s\}_{s \in \Lambda}$ and $\varphi \in C_c(\mathfrak{G})$. So it is sufficient to show that the set of s that “contributes” to (18) has density 0 with respect to $\{D_n\}_{n \in \mathbb{N}}$.

Let $s \in \Lambda$ “contribute” to (18). If $s \in \Lambda$ “contributes” to the former summation, then $s \in D_n$ for some $x \in \mathfrak{G} \setminus D_n$ such that $x-s \in \text{supp } \varphi$. So $s \in [Cl(\mathfrak{G} \setminus D_n) - \text{supp } \varphi] \cap D_n \subset \partial^{\text{supp } \varphi}(D_n)$, by the definition (1) of $\partial^{\text{supp } \varphi}(D_n)$. Similarly, if $s \in \Lambda$ “contributes” to the latter summation then $s \notin D_n$ for some $x \in D_n$ such that $x-s \in \text{supp } \varphi$, so $s \in (D_n - \text{supp } \varphi) \setminus \text{Int}(D_n) \subset \partial^{-\text{supp } \varphi}(D_n)$. Then $s \in \partial^{\text{supp } \varphi}(D_n) \cup \partial^{-\text{supp } \varphi}(D_n) \subset \partial^K(D_n)$ for some compact set $K \subset \mathfrak{G}$ such that $K = -K \supseteq \text{supp } \varphi$.

Thus we have only to verify $\lim_n \frac{\#\{s \in \Lambda; s \in \partial^K(D_n)\}}{|D_n|} = 0$. This is proved as follows: By Lemma 1, there is a compact set $U = -U$ such that $(s+U) \cap (t+U) = \emptyset$ for any distinct points $s, t \in \Lambda$. Thus $U + \{s \in \Lambda; s \in \partial^K(D_n)\} \subset \partial^{K+U}(D_n)$, and the Haar measure of the left-hand side of the inclusion is $|U| \cdot \#\{s \in \Lambda; s \in \partial^K(D_n)\}$. Thus $|U| \cdot \lim_n \frac{\#\{s \in \Lambda; s \in \partial^K(D_n)\}}{|D_n|} \leq \lim_n \frac{|\partial^{K+U}(D_n)|}{|D_n|} = 0$ by the definition of van Hove sequence. $\square$

Lemma 10 *Suppose the same assumptions as in Theorem 3. Then*

1. *For a complex Hilbert space $L^2(\widehat{\mathfrak{T}})$ with the inner product*

$$\langle \kappa_1, \kappa_2 \rangle_{\widehat{\mathfrak{T}}} := \sum_{\xi \in \widehat{\mathfrak{T}}} \kappa_1(\xi) \overline{\kappa_2(\xi)} \quad (\kappa_1, \kappa_2 \in \widehat{\mathfrak{T}}),$$

we have a bijective isometry $L^2(X_\Lambda, \nu) \xrightarrow{\iota} L^2(\mathfrak{T}, \tau) \xrightarrow{\widehat{\iota}} L^2(\widehat{\mathfrak{T}})$.

2. *For φ_1, Φ_1 of Lemma 8, and the projection ϖ given in (13), we have*

$$\widehat{(\iota\Phi_1)}(\xi) = \frac{\widehat{\varphi_1}(-\varpi(\xi))}{|\tilde{L}|} \widehat{b}(\varpi(\xi)^*) \quad (\xi \in \widehat{\mathfrak{T}}).$$

Proof (1) By (12), $\widehat{\mathfrak{T}}$ is discrete, so the inner product of the Hilbert space $L^2(\widehat{\mathfrak{T}})$ is in fact a summation. Recall that Fourier transform is a bijective isometry. So the conclusion follows from Lemma 7.

(2) Put $\chi := \varpi(\xi)$. The left-hand side $\widehat{(\iota\Phi_1)}(\xi)$ of the equation is a Fourier transform $\int_{\widehat{\mathfrak{T}}} \xi(t) (\iota\Phi_1)(t) d\tau(t)$, which is $\int_{X_\Lambda} \xi \circ \beta(P) \Phi_1(P) d\nu(P)$ by Lemma 7. By Proposition 6 (2) and the pointwise ergodic theorem [22], the integral is $\lim_n \frac{1}{|D_n|} \int_{D_n} \xi(\beta(\Lambda - x)) \Phi_1(\Lambda - x) dx \cdot |D_n|^{-1}$. Since $\beta(\Lambda - x) = \tilde{L} - (x, 0)$ by Proposition 7, we have $\xi(\beta(\Lambda - x)) = \xi(\tilde{L} - (x, 0)) =$

$\varpi(\xi)(-x) = \overline{\chi}(x)$. So $(\widehat{\iota\Phi_1})(\xi) = \lim_n \int_{D_n} \chi(x) \Phi_1(\Lambda - x) dx \cdot |D_n|^{-1}$. By (15) of Lemma 8 and the premise $\omega = \sum_{s \in \Lambda(W)} b(s^*) \delta_s$, we have

$$(\widehat{\iota\Phi_1})(\xi) = \lim_{n \leq} \frac{1}{|D_n|} \int_{D_n} \chi(x) \sum_{s \in \Lambda(W)} \varphi_1(x - s) b(s^*) dx.$$

We can apply Lemma 9 to above, since $\Lambda(W)$ is uniformly discrete, the sequence $\{b(s^*)\}_{s \in \Lambda(W)}$ of weights is bounded, and $\varphi_1 \in C_c(\mathfrak{G})$. Thus

$$(\widehat{\iota\Phi_1})(\xi) = \lim_{n \leq} \frac{1}{|D_n|} \sum_{s \in \Lambda(W) \cap D_n} b(s^*) \int_{\mathfrak{G}} \chi(x) \varphi_1(x - s) dx.$$

Here $\int_{\mathfrak{G}} \chi(x) \varphi_1(x - s) dx = \chi(s) \widehat{\varphi_1}(-\chi)$ is $\overline{\chi^*(s^*)} \widehat{\varphi_1}(-\chi)$ by (11). Therefore

$$(\widehat{\iota\Phi_1})(\xi) = \widehat{\varphi_1}(-\chi) \lim_n \frac{1}{|D_n|} \sum_{s \in \Lambda(W) \cap D_n} b(s^*) \overline{\chi^*(s^*)}. \quad (19)$$

Since $b \in C_c(\mathfrak{G}_{\text{int}})$, the Fourier transform $\hat{b}$ of b is well-defined. So Proposition 2 implies that the limit in (19) is $\int_W b(y) \overline{\chi^*}(y) dy / |\tilde{L}|$, which is $\hat{b}(\chi^*) / |\tilde{L}|$ by $\text{supp } b \subset W$. Therefore (19) implies the desired consequence. $\square$

Proof of Theorem 3 By Proposition 3, it is sufficient to verify

$$\int_{\mathfrak{G}} (\varphi * \tilde{\varphi})(x) d\gamma_{\omega}(x) = \int_{\mathfrak{G}} |\widehat{\varphi}(-\gamma)|^2 d \left(\sum_{\chi \in \mathcal{L}} \frac{|\hat{b}(\chi^*)|^2}{|\tilde{L}|^2} \delta_{\chi} \right) (\gamma) \geq 0. \quad (20)$$

Since $\tilde{\gamma}_{\omega} = \gamma_{\omega}$, the leftmost integral is $((\varphi * \tilde{\varphi}) * \gamma_{\omega})(0)$, which is $\langle \Phi, \Phi \rangle_{\nu}$ by (16). Because each Φ corresponds uniquely to φ by Lemma 8 and because of Lemma 10, it is $\sum_{\chi \in \mathcal{L}} |\widehat{\varphi}(-\chi) \hat{b}(\chi^*)|^2 \cdot |\tilde{L}|^{-2}$, the right-hand side of (20). $\square$

By the Theorem we have proved, we can see that the pure-point diffraction is still observed as long as the sample path of random weights is continuous on the internal space of the model set. The condition is comparable Baake-Moody's sufficient condition for deterministic model sets to have pure-point diffraction; their condition demands the continuity with respect to the internal space.

Theorem 4 *Suppose the same assumptions as in Theorem 3. If a complex-valued stochastic process $\{B_y(\omega)\}_{y \in W}$ is such that the sample path $b_{\omega}(y) = 1_W(y) B_y(\omega)$ is continuous on W almost every ω , then a Dirac comb $\pi(\omega) = \sum_{s \in \Lambda(W)} B_{s^*}(\omega) \delta_s$ has a diffraction measure $\widehat{\gamma}_{\pi}(\omega)$ which expectation is pure-point*

$$\mathbb{E}_{\omega}[\widehat{\gamma}_{\pi}(\omega)] = \sum_{\chi \in \mathcal{L}} \frac{\mathbb{E}[|\hat{b}_{\omega}(-\chi)|^2]}{|\tilde{L}|^2} \delta_{\chi}.$$

For example, if $W = [0, 1]$, $\{B_y(\omega)\}_y$ is an Ornstein-Uhlenbeck process.

In [9], for particle gases over FLC sets with Gibbs random field under a suitable interaction potential restrictions, Baake-Zint proved that the diffraction measures do not have singular continuous components and explicitly described the pure-point component and the absolutely continuous component by using the covariance of the random field.

Weak dependence of a random field over a point set S is studied typically with *Dobrushin interdependence matrix* [15, p. 32] $(C_{x,y})_{x,y \in S}$. For example, Külske [15] derived that if a Gibbs field has a Dobrushin interdependence matrix with the largest row-sum and the largest column-sum both less than 1, then it satisfies a concentration inequality for functions. We see that every finitely randomized model set has a Dobrushin interdependence matrix $(C_{x,y})_{x,y \in S}$ with each row and each column having bounded number of nonzero entries, because a mass at a point x is independent from that of any point y whenever $x - y$ is not in the dependency set D . So, it is natural to generalize finitely randomized model sets S to randomized model sets S having a finite set D with $\sup_{x \in S} \sum_{y \in S \setminus (D+x)} C_{x,y} < 1$, for example. We hope such randomized model sets with weak dependence satisfy our results and a concentration inequality for functions.

Acknowledgements The first author thanks Prof. Michael Baake and anonymous referee.

References

1. Argabright, L., Gil de Lamadrid, J.: Fourier analysis of unbounded measures on locally compact abelian groups. American Mathematical Society, Providence, R.I. (1974). Memoirs of the American Mathematical Society, No. 145
2. Baake, M., Birkner, M., Moody, R.V.: Diffraction of stochastic point sets: Explicitly computable examples. *Communications in Mathematical Physics* **293**, 611–660 (2010)
3. Baake, M., Hermisson, J., Pleasants, P.A.B.: The torus parametrization of quasiperiodic LI-classes. *Journal of Physics A: Mathematical and General* **30**(9), 3029–3056 (1997).
4. Baake, M., Lenz, D., Moody, R.V.: Characterization of model sets by dynamical systems. *Ergodic Theory Dynam. Systems* **27**(2), 341–382 (2007).
5. Baake, M., Moody, R.V.: Diffractive point sets with entropy. *J. Phys. A: Math. Gen* **31**, 9023–9039 (1998)
6. Baake, M., Moody, R.V.: Self-similar measures for quasicrystals. In: *Directions in mathematical quasicrystals, CRM Monogr. Ser.*, vol. 13, pp. 1–42. Amer. Math. Soc., Providence, RI (2000)
7. Baake, M., Moody, R.V.: Weighted Dirac combs with pure point diffraction. *J. Reine Angew. Math.* **573**, 61–94 (2004)
8. Baake, M., Moody, R.V., Schlottmann, M.: Limit-(quasi)periodic point sets as quasicrystals with p -adic internal spaces. *J. Phys. A* **31**(27), 5755–5765 (1998).
9. Baake, M., Zint, N.: Absence of singular continuous diffraction for discrete multi-component particle models. *J. Stat. Phys.* **130**(4), 727–740 (2008).
10. Barabash, R., Ice, G., Turchi, P. (eds.): *Diffuse Scattering and the Fundamental Properties of Materials*. Momentum press, New York (2009)
11. Hewitt, E., Ross, K.A.: Abstract harmonic analysis. Vol. I, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 115, second edn. Springer-Verlag, Berlin (1979). Structure of topological groups, integration theory, group representations

12. Hof, A.: Diffraction by aperiodic structures at high temperatures. *J. Phys. A* **28**(1), 57–62 (1995).
13. Hof, A.: On diffraction by aperiodic structures. *Comm. Math. Phys.* **169**(1), 25–43 (1995).
14. Janot, C.: Quasicrystals: A primer, *Monographs on the physics and chemistry of materials*, vol. 50, 2nd edn. Oxford University Press, Oxford (1994)
15. Külske, C.: Concentration inequalities for functions of Gibbs fields with application to diffraction and random Gibbs measures. *Comm. Math. Phys.* **239**(1-2), 29–51 (2003).
16. Lenz, D.: Continuity of eigenfunctions of uniquely ergodic dynamical systems and intensity of Bragg peaks. *Comm. Math. Phys.* **287**(1), 225–258 (2009).
17. Lenz, D., Richard, C.: Pure point diffraction and cut and project schemes for measures: the smooth case. *Math. Z.* **256**(2), 347–378 (2007).
18. Meyer, Y.: Algebraic numbers and harmonic analysis. North-Holland Publishing Co., Amsterdam (1972). North-Holland Mathematical Library, Vol. 2
19. Moody, R.V.: Meyer sets and their duals. In: The mathematics of long-range aperiodic order (Waterloo, ON, 1995), *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, vol. 489, pp. 403–441. Kluwer Acad. Publ., Dordrecht (1997)
20. Moody, R.V.: Uniform distribution in model sets. *Canad. Math. Bull.* **45**(1), 123–130 (2002)
21. Müller, P., Richard, C.: Ergodic properties of randomly coloured point sets. *Canadian Journal of Mathematics* (2012). DOI 10.4153/CJM-2012-009-7. On-Line First
22. Nevo, A.: Pointwise ergodic theorems for actions of groups. In: Handbook of dynamical systems. Vol. 1B, pp. 871–982. Elsevier B. V., Amsterdam (2006)
23. Schlottmann, M.: Generalized model sets and dynamical systems. In: Directions in mathematical quasicrystals, *CRM Monogr. Ser.*, vol. 13, pp. 143–159. Amer. Math. Soc., Providence, RI (2000)
24. Stroock, D.W.: Probability theory, an analytic view. Cambridge University Press, Cambridge (1993)

A Proof of “If $\text{Int}(W) \neq \emptyset$ then $\Lambda(W)$ is relatively dense.”

Variants of this assertion have already appeared in Schlottmann, Lagarias and so on.

Since $\tilde{L}$ is a lattice of $\mathfrak{G} \times \mathfrak{G}_{\text{int}}$, there is a complete system C of representatives of a compact, quotient set $(\mathfrak{G} \times \mathfrak{G}_{\text{int}})/\tilde{L}$ such that C is relatively compact. Then $\mathfrak{G} \times \mathfrak{G}_{\text{int}} = \tilde{L} + C$. Since the window W has nonempty interior and L^* is dense in $\mathfrak{G}_{\text{int}}$, it follows that $\mathfrak{G}_{\text{int}} = L^* - W$. Moreover, since $\Pi_{\text{int}}(\text{Cl}(C))$ is compact, there is a finite subset F of L such that $\Pi_{\text{int}}(\text{Cl}(C)) \subset F^* - W$. Let $K := \Pi(\text{Cl}(C)) - F$, which is compact in $\mathfrak{G}$. For each $x \in \mathfrak{G}$, there are $(t, t^*) \in \tilde{L}$ and $(c, d) \in C$ such that $(x, 0) = (t, t^*) + (c, d)$. Since $\Pi_{\text{int}}(C) \subset F^* - W$, there are $f \in F$ and $w \in W$ such that $d = f^* - w$. Then

$$(x, 0) = (t, t^*) + (c, f^* - w) = (t + c, t^* + f^* - w).$$

So $t^* + f^* = w \in W$, and thus $t + f \in \Lambda(W)$. Since $x = t + c = (t + f) + (c - f)$ and $c - f \in \Pi(C) - F \subset K$, we have $x \in \Lambda(W) + K$. Therefore $\mathfrak{G} = \Lambda(W) + K$.