

# QUASI-CIRCLES THROUGH PRESCRIBED POINTS

JOHN M. MACKAY

ABSTRACT. We show that in an  $L$ -annularly linearly connected,  $N$ -doubling, complete metric space, any  $n$  points lie on a  $\lambda$ -quasi-circle, where  $\lambda$  depends only on  $L, N$  and  $n$ . This implies, for example, that if  $G$  is a hyperbolic group that does not split over any virtually cyclic subgroup, then any geodesic line in  $G$  lies in a quasi-isometrically embedded copy of  $\mathbb{H}^2$ .

*Version accepted by Indiana University Mathematics Journal*<sup>1</sup>

## 1. INTRODUCTION

Menger’s theorem for graphs extends to the following topological result, known as the “ $n$ -Bogensatz” or  $n$ -arc connectedness theorem.

**Theorem 1.1** ([Nöb32, Zip33, Why48]). *If  $X$  is a connected, locally connected, locally compact metric space that cannot be disconnected by removing any  $n - 1$  points, then any two points in  $X$  can be joined by  $n$  arcs, pairwise disjoint apart from their endpoints.*

A well known corollary of this result is that any  $n$  points in  $X$  lie on a simple closed curve (see Theorem 4.1 and remark in [TV08]).

In this paper, we prove analogues of these theorems for quasi-arcs and quasi-circles using quantitative topological arguments. Quasi-circles arise naturally in geometric function theory and in the study of boundaries of hyperbolic groups, and our results have consequences for the geometry of such groups.

**1.1. Statement of results.** In 1963, Ahlfors [Ahl63] showed that a Jordan curve  $\gamma \subset \mathbb{R}^2$  is the image of  $\mathbb{S}^1 \subset \mathbb{R}^2$  under some quasiconformal homeomorphism of  $\mathbb{R}^2$  if and only if  $\gamma$  is *linearly connected*:

---

*Date:* March 12, 2013.

*2010 Mathematics Subject Classification.* Primary 30L10; Secondary 30C65, 51F99.

*Key words and phrases.* Quasi-circle, quasi-arc, linearly connected, bounded turning,  $n$ -Bogensatz.

JMM was supported by EPSRC grant “Geometric and analytic aspects of infinite groups”.

<sup>1</sup><http://www.iumj.indiana.edu/IUMJ/Preprints/5211.pdf>

**Definition 1.2.** A complete metric space  $(X, d)$  is  $L$ -linearly connected, for some  $L \geq 1$ , if for every  $x, y \in X$ , there exists a continuum  $J$  containing  $x$  and  $y$ , so that  $\text{diam}(J) \leq Ld(x, y)$ .

(This is also called  $L$ -bounded turning. Note that in the above definition, we may assume  $J$  is an arc.)

More generally, a *quasi-circle* (respectively *quasi-arc*) is a quasisymmetric image of the standard Euclidean circle (respectively interval). Tukia and Väisälä showed that a metric Jordan curve (i.e., a metric space homeomorphic to  $\mathbb{S}^1$ ) is a quasi-circle if and only if it is doubling and linearly connected [TV80], and likewise for metric Jordan arcs. Recall that a metric space  $(X, d)$  is  $N$ -doubling, for some  $N \in \mathbb{N}$ , if any ball of radius  $r > 0$  in  $X$  can be covered by  $N$  balls of radius  $r/2$ .

In the remainder of this paper, we define a  $\lambda$ -quasi-circle (or  $\lambda$ -quasi-arc) to be a metric Jordan curve (or metric Jordan arc) that is doubling and  $\lambda$ -linearly connected.

The spaces we study have the property that they have no local cut points, in the following quantitatively controlled sense.

**Definition 1.3.** Let  $(X, d)$  be a metric space. The annulus around  $x$  between radii  $r$  and  $R$  is denoted by  $A(x, r, R) = \overline{B}(x, R) \setminus B(x, r)$ .

A metric space  $(X, d)$  is  $L$ -annularly linearly connected, for some  $L \geq 1$ , if it is connected, and given  $r > 0$ ,  $p \in X$ , any two points  $x, y \in A(p, r, 2r)$  lie in an arc  $J$  so that  $x, y \in J \subset A(p, r/L, 2Lr)$ .

(We may assume (on replacing  $L$  by  $8L$ ) that  $X$  is also  $L$ -linearly connected.)

Our main theorems are quantitative versions of Theorems 1.1 and 4.1.

**Theorem 1.4.** Let  $X$  be a  $N$ -doubling,  $L$ -annularly linearly connected, and complete metric space. For any  $n \in \mathbb{N}$ , there exists  $\lambda = \lambda(L, N, n)$  so that any distinct  $x, y \in X$  can be joined by  $n$  different  $\lambda$ -quasi-arcs, where the concatenation of any two forms a  $\lambda$ -quasi-circle.

**Theorem 1.5.** Suppose  $(X, d)$  is a non-trivial,  $N$ -doubling,  $L$ -annularly linearly connected, complete metric space. Then any finite set  $T \subset X$  lies on a  $\lambda$ -quasi-circle  $\gamma \subset X$ , where  $\lambda = \lambda(L, N, |T|)$ . Moreover, if  $|T| \geq 2$  we can ensure that  $\text{diam}(\gamma) \leq \lambda \text{diam}(T)$ .

Note that these results apply to the boundaries of many hyperbolic groups. For example, we observe the following.

**Corollary 1.6.** Suppose  $G$  is a  $\delta$ -hyperbolic group, which does not virtually split over any finite or two-ended subgroup. Then any  $n$  geodesics

in  $G$  lie in the image of an  $(L, C)$ -quasi-isometry  $f : \mathbb{H}^2 \rightarrow G$ , where  $L$  and  $C$  depend only on  $G$  and  $n$ .

*Proof.* The boundary  $\partial_\infty G$ , given some fixed visual metric, is doubling and annularly linearly connected [Mac10, Proof of Corollary 1.2]. Let  $x_1, \dots, x_{2n} \in \partial_\infty G$  be the endpoints of the geodesics.

We apply Theorem 1.5 to find a quasi-circle through  $x_1, \dots, x_{2n}$ , and extend this to find a quasi-isometrically embedded hyperbolic plane  $f : \mathbb{H}^2 \rightarrow G$  [BS00, Theorems 7.4, 8.2]. Up to modifying  $f$  by a finite distance, we may assume that the geodesics lie in the image of  $f$ .  $\square$

**1.2. Background and remarks.** Any two points in a connected, locally connected, locally compact metric space can be joined by an arc. The analogous statement for quasi-arcs was proved by Tukia, and is a key tool in this paper.

**Theorem 1.7** ([Tuk96, Theorem 1A],[Mac08, Corollary 1.2]). *Suppose  $(X, d)$  is an  $N$ -doubling,  $L$ -linearly connected, complete metric space. Then there exists  $\lambda = \lambda(N, L)$  so that any two points in  $X$  can be connected by a  $\lambda$ -quasi-arc.*

For quasi-circles, as far as the author knows, the only non-trivial existence result known prior to this paper is the following result of Bonk and Kleiner. (For an analogous statement for certain relatively hyperbolic groups, see [MS11, Theorem 1.3].)

**Theorem 1.8** ([BK05, Theorem 1]). *If  $G$  is a hyperbolic group and its boundary  $\partial_\infty G$  is not totally disconnected, then  $\partial_\infty G$  contains a quasi-circle.*

Theorem 1.8 is motivated by the problem of finding surface subgroups in hyperbolic groups: undistorted surface subgroups give quasi-isometric embeddings of  $\mathbb{H}^2$  in the group, and quasi-isometric embeddings of  $\mathbb{H}^2$  exactly correspond to quasi-circles in the boundary. Theorem 1.8 showed that there is no geometric obstruction to finding such a surface subgroup once the group is not virtually free, answering a question of Papasoglu.

This result is proved by using Theorem 1.7, a dynamical argument, and Arzelà-Ascoli; the indirectness involved means that this method cannot show that every point in  $\partial_\infty G$  lies in a quasi-circle.

We now consider Theorems 1.4 and 1.5. In these results, we cannot weaken the annular linear connectedness condition to the topological condition of Nöbeling. For example, the set  $X = \{(x, y) : 0 \leq x \leq 1, |y| \leq x^2\}$  is doubling, linearly connected, and has no local cut points, but there is no quasi-circle in  $X$  that contains the point  $(0, 0)$ .

One might hope for a stronger result than Theorem 1.5, where rather than a quantitative no local cut points condition, we merely assume a quantitative version of “cannot be disconnected by removing  $N$  points.” For example, perhaps in a doubling, LLC, complete metric space, any two points lie on a quasi-circle.

However, our arguments fail in this case, as we strongly use rescaling and Gromov-Hausdorff limits of sequences of spaces. The LLC(2) condition need not be preserved under such limits: consider a sequence of larger and larger circles converging to a line.

On the other hand, Theorem 1.5 is sharp in the following two senses. First, the hypotheses of this theorem do not suffice to ensure that  $x_1, \dots, x_n$  lie on  $\gamma$  in the cyclic order given. (Consider the closed unit square, which is doubling and annularly linearly connected, and label the four corners clock-wise  $x_1, x_2, x_3, x_4$ . There is no topologically embedded circle containing these points in cyclic order  $x_1, x_3, x_2, x_4$ .)

Second,  $\lambda$  must depend on  $n$ , otherwise one could take increasingly dense subsets of the sphere and find uniform quasi-circles through these sets. In the limit, this gives a contradiction.

The key technical tool that we use in this paper is a new method of joining two quasi-arcs together to make a quasi-arc. This is described in Section 2. The “quasi-arc  $n$ -Bogensatz” Theorem 1.4 is proved in Section 3. Finally, we prove Theorem 1.5 in Section 4.

**1.3. Notation.** We denote balls in a metric space  $(X, d)$  by  $B(x, r) = \{y \in X : d(x, y) < r\}$ . The open neighbourhood of  $A \subset X$  of size  $r$  is  $N(A, r) = \{y \in X : d(y, A) < r\}$ . If  $B = B(x, r)$ , and  $t > 0$ , then  $tB = B(x, tr)$ . Similarly, if  $V = N(A, r)$ , and  $t > 0$ , then  $tV = N(A, tr)$ .

If  $C$  is a constant depending only on  $C_1, C_2$ , then we write  $C = C(C_1, C_2)$ .

**1.4. Acknowledgements.** I thank Daniel Groves for bringing this question to my attention, and Bruce Kleiner and Alessandro Sisto for helpful comments.

## 2. JOINING TOGETHER QUASI-ARCS

Any arc in a doubling, linearly connected space can be straightened into a quasi-arc.

**Theorem 2.1** ([Tuk96, Theorem 1B], [Mac08, Theorem 1.1]). *Suppose  $(X, d)$  is a  $L$ -linearly connected,  $N$ -doubling, complete metric space. For every arc  $A$  in  $X$  and every  $\epsilon > 0$ , there is an arc  $J$  that  $\epsilon$ -follows  $A$ , has the same endpoints as  $A$ , and is an  $\alpha\epsilon$ -local  $\lambda$ -quasi-arc, where*

$\lambda = \lambda(L, N) \geq 1$  and  $\alpha = \alpha(L, N) > 0$ . That is, for any  $x, y \in J$  with  $d(x, y) \leq \alpha\epsilon$ , we have  $\text{diam}(J[x, y]) \leq \lambda d(x, y)$ .

The notation we use for arcs is described below.

**Definition 2.2** ([Mac08]). For any  $x$  and  $y$  in an embedded arc  $A$ , let  $A[x, y]$  be the closed, possibly trivial, subarc of  $A$  that lies between them. Let  $A[x, y) = A[x, y] \setminus \{y\}$ , and so on.

An arc  $B$   $\iota$ -follows an arc  $A$  if there exists a (not necessarily continuous) map  $p : B \rightarrow A$ , sending endpoints to endpoints, such that for all  $x, y \in B$ ,  $B[x, y]$  is in the  $\iota$ -neighbourhood of  $A[p(x), p(y)]$ ; in particular,  $p$  displaces points at most  $\iota$ .

The goal of this section is to refine Theorem 2.1 to the following situation. Suppose an arc  $I$  is formed from two quasi-arcs joined by an arc  $I' \subset I$ . We show how to modify  $I$  only near  $I'$  to create a quasi-arc.

**Theorem 2.3.** Let  $X$  be an  $N$ -doubling,  $L$ -linearly connected, complete metric space. Let  $A$  be an arc formed of three consecutive subarcs  $A_1 = A[a_0, a_1]$ ,  $A_2 = A[a_1, a_2]$  and  $A_3 = A[a_2, a_3]$ . Suppose that  $A_1$  and  $A_3$  are  $\epsilon$ -local  $\lambda$ -quasi-arcs, and  $d(A_1, A_3) \geq 2\epsilon$ .

Then there exists an  $\alpha\epsilon$ -local  $\lambda'$ -quasi-arc  $J$  that  $\alpha\epsilon$ -follows  $A$ , for  $\alpha = \alpha(L, N, \lambda) > 0$  and  $\lambda' = \lambda'(L, N, \lambda) > 1$ . Moreover,  $J$  contains the initial and final connected components of  $A \setminus N(A_2, 2\epsilon)$ .

This theorem follows the proof of Theorem 2.1 given in [Mac08] verbatim, once we establish the following modified version of [Mac08, Proposition 2.1].

**Proposition 2.4.** We assume the hypotheses of Theorem 2.3. There exists constants  $s = s(L, N, \lambda) > 0$  and  $S = S(L, N, \lambda) > 0$  with the following property: for each  $\iota \in (0, \epsilon)$  there exists an arc  $J$  that  $\iota$ -follows  $A$ , contains the initial and final connected components of  $A \setminus N(A_2, 2\iota)$ , and satisfies

$$(*) \quad \forall x, y \in J, d(x, y) < s\iota \implies \text{diam}(J[x, y]) < S\iota.$$

*Proof.* We modify the proof of [Mac08, Proposition 2.1]. To simplify notation, we replace  $L$  by  $\max\{L, \lambda\}$ .

Let  $r = \iota/20L$ , and let  $\mathcal{N}$  be a maximal  $r$ -separated net in  $X$  containing  $a_0$  and  $a_3$ . Then there exists  $\delta = \delta(L, N, \lambda) \in (0, 1)$  and a collection of sets  $\{V_x\}_{x \in \mathcal{N}}$  so that each  $V_x$  is a union of finitely many (closed) arcs in  $X$ , and for all  $x, y \in \mathcal{N}$ :

- (1)  $d(x, y) \leq 2r \implies y \in V_x$ .
- (2)  $\text{diam}(V_x) \leq 5Lr$ .
- (3)  $V_x \cap V_y = \emptyset \implies d(V_x, V_y) > \delta r$ .

$$(4) \quad B(x, r) \cap (A_1 \cup A_3) \subset V_x.$$

To show this, we follow the proof of [Mac08, Lemma 3.1], with the exception that when we construct  $V_x^{(0)}$ , we also add closed arcs from  $B(x, 2r) \cap (A_1 \cup A_3)$  so that the hypotheses of (4) are satisfied, and arcs joining them to  $x$  in  $B(x, 2rL)$ . Observe that  $\text{diam}(V_x^{(0)}) \leq 4Lr$ , so the rest of the proof of [Mac08, Lemma 3.1] follows unchanged.

Now cover  $A_2$  by connected open arcs which lie in some  $B(z, r)$ ,  $z \in \mathcal{N}$ , and take a finite subcover of  $A_2$ . Let  $y_1, y_2, \dots, y_m$  be points in  $A_2$  lying in the arcs corresponding to this cover, in the order given by  $A_2$ , and let  $z_1, z_2, \dots, z_m \subset \mathcal{N}$  be the centres of the associated balls.

The collection of sets  $\{V_x\}$  is locally finite, and each  $V_x$  is compact, so there exists a point  $q_0 \in A_1$  that is the first point in  $A_1$  to be contained in some  $V_{w_0}$  which meets  $\bigcup_j V_{z_j}$ .

Let  $K$  be the union of  $V_x$  so that  $V_x \cap A_3 \neq \emptyset$ . Define  $w_i$  inductively as follows, for  $i > 0$ . If  $V_{w_{i-1}} \cap K \neq \emptyset$ , set  $n = i$ , and stop. Otherwise, let  $k_i = \max\{j : V_{w_{i-1}} \cap V_{z_j} \neq \emptyset\}$ , set  $w_i = z_{k_i}$ , and continue.

Finally, let  $q_{n+1}$  be the last point in  $A_3$  to be contained in some  $V_{w_n}$  meeting  $V_{w_{n-1}}$ . By (1), this process is well defined.

We use this sequence to build our path  $J$  in stages. Set  $J_{-1} = A_1[a_0, q_0]$ . Let  $J_0$  be an arc in  $V_{w_0}$  that joins  $q_0$  to  $q_1 \in V_{w_1}$ , where  $J_0 \cap A[a_0, q_0] = \{q_0\}$  and  $J_0 \cap V_{w_1} = \{q_1\}$ . Now for  $i$  from 1 to  $n-1$ , let  $J_i$  be an arc in  $V_{w_i}$  that joins  $q_i$  in  $V_{w_i}$  to some  $q_{i+1} \in V_{w_{i+1}}$ , where  $q_{i+1}$  is the first point of  $J_i$  to meet  $V_{w_{i+1}}$ . Let  $J_n$  be an arc in  $V_{w_n}$  that joins  $q_n$  to  $q_{n+1} \in A[q_{n+1}, a_3]$ , where  $J_n \cap A[q_{n+1}, a_3] = \{q_{n+1}\}$ . To finish, we set  $J_{n+1} = A[q_{n+1}, a_3] = A_3[q_{n+1}, a_3]$ .

We claim that the arc  $J = J_{-1} \cup J_0 \cup \dots \cup J_{n+1}$  satisfies our conclusions, for suitable  $s$  and  $S$ .

To show that  $J$   $\iota$ -follows  $A$ , define a coarse map  $f : J \rightarrow A$  as follows. If  $x \in J_{-1} \cup J_{n+1}$ , let  $f(x) = x$ . If  $x \in J[q_0, q_1]$ , set  $f(x) = q_0$ , and if  $x \in J[q_n, q_{n+1}]$ , set  $f(x) = q_{n+1}$ . For  $x \in J[q_i, q_{i+1}]$ ,  $i = 1, \dots, n-1$ , set  $f(x) = y_{k_i} \in A_2 \cap B(z_{k_i}, r)$ .

It is straightforward to check that  $f$  satisfies the definition of  $\iota$ -following. For example, suppose  $y \in J_i, y' \in J_{i'}$  and  $1 \leq i \leq i' \leq n-1$ . Then

$$\begin{aligned} J[y, y'] &\subset J[q_i, q_{i'+1}] \subset N(\{w_i, \dots, w_{i'}\}, 5Lr) \\ &\subset N(\{y_{k_i}, \dots, y_{k_{i'}}\}, 5Lr + r) \subset N(A[y_{k_i}, y_{k_{i'}}], 5Lr + r) \\ &\subset N(A[f(y), f(y')], \iota). \end{aligned}$$

The other cases follow in similar fashion.

As  $f$  is the identity on  $J_{-1} \cup J_{n+1}$ , and  $d(q_0, A_2), d(q_{n+1}, A_2) < \iota$ ,  $J$  contains the required components of  $A_1$  and  $A_3$ .

All that remains is to show that  $J$  satisfies (\*). Suppose that  $y \in J_i, y' \in J_{i'}$ , with  $y < y'$  in  $J$ , and  $d(y, y') < r\delta$ . There are four cases.

(i) If  $i = i' = -1$  or  $i = i' = n + 1$ , we have  $\text{diam}(J[y, y']) \leq \lambda d(y, y') \leq Lr\delta$ .

(ii) If  $0 \leq i, i' \leq n$ , then  $d(V_{w_i}, V_{w_{i'}}) < r\delta$  so by construction and (3) we have  $|i - i'| \leq 1$  and thus by (2),  $\text{diam}(J[y, y']) \leq 10Lr$ .

(iii) If  $i = -1, i' \geq 0$  then by (4),  $y$  lies in some  $V_x$ , and  $d(V_x, V_{w_{i'}}) \leq d(y, y') < \delta r$ , so by (3) and construction,  $i' = 0$ . Thus  $d(y', q_0) \leq \text{diam}(V_{w_0}) \leq 5Lr$ , and  $d(y, q_0) \leq 5Lr + d(y, y') < (5L + \delta)r$ . So

$$\text{diam}(J[y, y']) = \text{diam}(J[y, q_0] \cup J[q_0, y']) \leq \lambda(5L + \delta)r + 5Lr \leq 11L^2r.$$

(iv) The case  $i \leq n, i' = n + 1$  follows similarly to (iii).

We let  $s = \delta r/\iota = \delta/20L$  and  $S = \max\{Lr\delta, 10Lr, 11L^2r\}/\iota = 11L/20$ , and have proven the proposition.  $\square$

### 3. MANY QUASI-ARCS BETWEEN TWO POINTS

Our goal in this section is the following theorem.

**Theorem 1.4** *Let  $X$  be a  $N$ -doubling,  $L$ -annularly linearly connected, and complete metric space. For any  $n \in \mathbb{N}$ , there exists  $\lambda = \lambda(L, N, n)$  so that any distinct  $x, y \in X$  can be joined by  $n$  different  $\lambda$ -quasi-arcs, so that the concatenation of any two forms a  $\lambda$ -quasi-circle.*

The key part of this theorem is the following proposition that splits a quasi-arc into two relatively close and separated quasi-arcs. This uses arguments similar to [Mac10, Section 3].

**Proposition 3.1.** *Given  $\lambda_0 \geq 1, \epsilon > 0$ , there exists  $\lambda = \lambda(L, N, \lambda_0, \epsilon) \geq 1$  and  $\eta = \eta(L, N, \lambda_0, \epsilon) > 0$  with the following property:*

*For any  $\lambda_0$ -quasi-arc  $A = A[a, b]$  in an  $N$ -doubling,  $L$ -annularly linearly connected, complete metric space  $X$ , there exist two  $\lambda$ -quasi-arcs  $J = J[a, b]$  and  $J' = J'[a, b]$  with the following properties:*

*For all  $z \in (J \cup J') \setminus \{a, b\}$ ,*

$$(3.2) \quad d(z, A) \leq \epsilon d(z, \{a, b\}), \text{ and}$$

$$(3.3) \quad \max\{d(z, J), d(z, J')\} \geq \eta d(z, \{a, b\}).$$

*Proof.* We may rescale so that  $d(a, b) = 1$ , and assume that  $\epsilon < 1$ . Let  $\delta = 1/10\lambda_0$ .

We consider  $A$  in the natural order from  $a$  to  $b$ . For each  $i \in \mathbb{N}$ , let  $x_{-i}$  be the first point in  $A$  at distance  $\delta^i$  from  $a$ , and let  $x_i$  be the last point in  $A$  at distance  $\delta^i$  from  $b$ .

Let  $D_1 = \epsilon\delta/3\lambda_0$ , and for  $i \in \mathbb{Z} \setminus \{0\}$ , let  $B_i = B(x_i, D_1\delta^{|i|})$ .




FIGURE 1. Splitting a quasi-arc into a quasi-circle

For  $i < 0$  let  $A_i = A[x_{i-1}, x_i]$ , let  $A_0 = A[x_{-1}, x_1]$ , and for  $i > 0$  let  $A_i = A[x_i, x_{i+1}]$ . Set  $D_2 = D_1\delta/10\lambda_0L$ , and for  $i \in \mathbb{Z}$  let  $V_i = N(A_i, D_2\delta^{|i|})$ .

**Lemma 3.4.** *These neighbourhoods have the following properties:*

- (1) If  $i \neq j$ , then  $B_i \cap B_j = \emptyset$ .
- (2) If  $i < 0$  and  $j < i$ , then  $V_{i+1} \cap B_j = V_{i+1} \cap V_j = B_i \cap V_j = \emptyset$ .
- (3) If  $i < 0$ , then  $(V_i \cap V_{i-1}) \subset B_{i-1}$ .

*Proof.* (1) This is immediate.

(2) This follows from the following claim. If for some  $i < 0$ , we have  $z \in A[a, x_{i-1}]$ ,  $z' \in A[x_i, x_{i+1}]$  then  $d(z, z') \geq D_1\delta^{|i-1|} + D_1\delta^{|i+1|}$ , for otherwise

$$\text{diam}(A[z, z']) \leq \lambda_0(D_1\delta^{|i-1|} + D_2\delta^{|i+1|}) \leq \frac{1}{2}\delta^{|i|},$$

but

$$\text{diam}(A[z, z']) \geq \text{diam}(A[x_{i-1}, x_i]) \geq \delta^{|i|} - \delta^{|i-1|} \geq \frac{2}{3}\delta^{|i|}.$$

(3) If not, there exist  $z \in A_{i-1}$ ,  $z' \in A_i$  outside  $\frac{1}{2}B_{i-1}$ , so that  $d(z, z') \leq D_2\delta^{|i-1|} + D_2\delta^{|i|}$ . Therefore

$$\text{diam}(A[z, z']) \leq \lambda_0 D_2 \delta^{|i-1|} (1 + \delta^{-1}) \leq \frac{1}{10} D_1 \delta^{|i-1|}$$

but  $A[z, z']$  must pass through the centre of  $\frac{1}{2}B_{i-1}$ , so  $\text{diam}(A[z, z']) \geq \frac{1}{2}D_1\delta^{|i-1|}$ , a contradiction.  $\square$

We now split  $A$  into two disjoint arcs along the subarcs  $A_{2i}$ ,  $i \in \mathbb{Z}$ . See Figure 1.

**Lemma 3.5.** *For  $i \in \mathbb{Z}$  we can find two  $\lambda_1$ -quasi-arcs  $J_{2i}, J'_{2i}$  that  $\frac{1}{2}D_2\delta^{|2i|}$ -follow  $A_{2i}$ , and are  $\eta_1\delta^{|2i|}$  separated, where  $\eta_1 = \eta_1(L, N, \lambda_0, \epsilon) > 0$  and  $\lambda_1 = \lambda_1(L, N, \lambda_0, \epsilon) > 1$ .*

*Proof.* Using [Mac10, Lemma 3.3] with “ $\epsilon$ ” equal to  $\frac{1}{4}D_2\delta^{|2i|}$ , we split  $A_{2i}$  into two  $2\eta_1\delta^{|2i|}$  separated arcs that  $\frac{1}{4}D_2\delta^{|2i|}$ -follow  $A_{2i}$ , for some  $\eta_1 = \eta_1(L, N, \epsilon, \lambda_0) \in (0, \frac{1}{4}D_2)$ . We then apply Theorem 2.1 to these



arcs with “ $\epsilon$ ” equal to  $\frac{1}{2}\eta_1\delta^{|2i|}$ , to get two  $\frac{1}{2}\alpha\eta_1\delta^{|2i|}$ -local  $\lambda'_1$ -quasi-arcs  $J_{2i}, J'_{2i}$  that  $\frac{1}{2}D_2\delta^{|2i|}$ -follow  $A[x_1, y_1]$  and are  $\eta_1\delta^{|2i|}$ -separated, for suitable  $\alpha = \alpha(L, N)$  and  $\lambda'_1 = \lambda'_1(L, N)$ .

Every  $\beta$ -local  $\mu$ -quasi-arc of diameter  $D$  is a  $\max\{\mu, D/\beta\}$ -quasi-arc, and these arcs have diameter at most  $2\lambda_0\delta^{|2i|}$ , so  $J_{2i}, J'_{2i}$  are  $\lambda_1$ -quasi-arcs, for  $\lambda_1 = \lambda_1(\alpha, \eta_1, \lambda_0, \lambda'_1)$ .  $\square$

Now, following the proof of [Mac10, Lemma 3.5], for each  $i \in \mathbb{Z}$ , one can join the pair of arcs  $J_{2i}, J'_{2i}$  to the arcs  $J_{2i-2}, J'_{2i-2}$  inside  $\frac{1}{2}B_{2i-1} \cup \frac{1}{2}V_{2i-1} \cup \frac{1}{2}B_{2i-2}$ , with control on the separation between the resulting arcs. We prove this in the case  $i \leq 0$ ;  $i > 0$  is handled similarly.

The separation properties of Lemma 3.4 ensure that the following process can be applied independently in each location.

*Topological joining:* Join the endpoints of  $J_{2i}, J'_{2i}$  to  $A$  inside the ball  $B(x_{2i-1}, \frac{1}{2}LD_2\delta^{|2i|}) = (LD_2/2D_1\delta)B_{2i-1}$ . Similarly, join the endpoints of  $J_{2i-2}, J'_{2i-2}$  to  $A$  inside  $(LD_2/2D_1)B_{2i-2}$ .

We “unzip”  $A$  along this segment to join the two pairs of arcs  $J_{2i}, J'_{2i}$  and  $J_{2i-2}, J'_{2i-2}$  by two disjoint arcs  $\tilde{J}_{2i-1}, \tilde{J}'_{2i-1}$  in  $\frac{1}{4}V_{2i-1}$  (see [Mac10, Lemma 3.1, Lemma 3.5]). This involves discarding the ends of the four given arcs, but all changes take place inside  $\frac{1}{4}B_{2i-2} \cup \frac{1}{4}V_{2i-1} \cup \frac{1}{4}B_{2i-1}$ .

*Quantitative control:* As in [Mac10, Lemma 3.5], compactness arguments ensure that  $d(\tilde{J}_{2i-1}, \tilde{J}'_{2i-1}) \geq 2\eta_2\delta^{|2i-1|}$ , for some value  $\eta_2 = \eta_2(L, N, \epsilon, \lambda_0, \lambda_1) > 0$ .

We now straighten these separated arcs into quasi-arcs.

*Straightening:* We assume, after swapping  $J_*, J'_*$  if necessary, that  $\tilde{J}_{2i-1}$  joins  $J_{2i}$  and  $J_{2i-2}$ , and that  $\tilde{J}'_{2i-1}$  joins  $J'_{2i}$  and  $J'_{2i-2}$ .

We apply Theorem 2.3 to  $J_{2i-2} \cup \tilde{J}_{2i-1} \cup J_{2i}$  with “ $\epsilon$ ” equal to  $\frac{1}{2}\eta_2\delta^{|2i-1|}$ , to straighten the arc into a  $\lambda_2$ -quasi-arc  $J_{2i-2} \cup J_{2i-1} \cup J_{2i}$ , making changes only in  $\frac{1}{2}B_{2i-2} \cup \frac{1}{2}V_{2i-1} \cup \frac{1}{2}B_{2i-1}$ . Here  $\lambda_2 = \lambda_2(L, N, \eta_2, \lambda_1) \geq \lambda_1$ . Again, we may discard ends of  $J_{2i}, J'_{2i}, J_{2i-2}, J'_{2i-2}$  in  $\frac{1}{2}B_{2i-2} \cup \frac{1}{2}B_{2i}$ .

We claim that the arcs  $J = \bigcup_{i \in \mathbb{Z}} J_i$  and  $J' = \bigcup_{i \in \mathbb{Z}} J'_i$  satisfy our requirements.

**Lemma 3.6.**  *$J$  and  $J'$  are  $\lambda$ -quasi-arcs, for  $\lambda = \lambda(L, N, \lambda_0, \epsilon)$ .*

*Proof.* Suppose  $x, y \in J$ , where  $x \in J_i$  and  $y \in J_j$ ,  $i \leq j$ . It suffices to consider the following three cases.

If  $|i - j| \leq 1$ , then  $\text{diam}(J[x, y]) \leq \lambda_2 d(x, y)$ .

If  $i < 0 < j$  then  $d(a, x), d(b, y) \leq \frac{1}{5}$ , so  $\text{diam}(J[x, y]) \leq \text{diam}(J) \leq 2\lambda_0 \leq 4\lambda_0 d(x, y)$ .

If  $i+1 < j \leq 0$ , then  $x \in \frac{1}{2}B_{i-1} \cup \frac{1}{2}V_i \cup \frac{1}{2}B_i$ , and  $y \in \frac{1}{2}B_{j-1} \cup \frac{1}{2}V_j \cup \frac{1}{2}B_j$  (where  $B_0 = B_1$ ). Thus by Lemma 3.4,  $d(x, y) \geq \frac{1}{2}D_2\delta^{|j|}$ , so

$$\text{diam}(J[x, y]) \leq \text{diam}(J[a, x_j]) \leq 2\lambda_0\delta^{|j|} \leq \frac{4\lambda_0}{D_2}d(x, y).$$

We set  $\lambda = \max\{\lambda_2, 4\lambda_0/D_2\}$ , and are done.  $\square$

It remains to check the neighbourhood and separation conditions.

**Lemma 3.7.** *J and J' satisfy (3.2).*

*Proof.* It suffices to consider  $z \in J_i$ ,  $i \leq 0$ . If  $z \in \frac{1}{2}B_{i-1}$ , then  $d(z, A) < \frac{1}{2}D_1\delta^{|i-1|}$ , and  $d(z, a) > \delta^{|i-1|}(1 - \frac{1}{2}D_1) > \frac{1}{2}\delta^{|i-1|}$ . Therefore, as  $D_1 < \epsilon$ , (3.2) holds. Similarly, if  $z \in \frac{1}{2}B_i$ , (3.2) holds.

It remains to check when  $z \in \frac{1}{2}V_i$ . Then there exists some  $z' \in A_i$  so that  $d(z, z') \leq \frac{1}{2}D_2\delta^{|z|}$ . Thus as  $d(a, A_i) \geq \delta^{|i-1|}/\lambda_0$ ,

$$d(z, a) \geq d(z', a) - \frac{1}{2}D_2\delta^{|z|} \geq \delta^{|z|}\left(\frac{\delta}{\lambda_0} - \frac{D_2}{2}\right) \geq \frac{1}{20\lambda_0^2}\delta^{|z|}.$$

Since  $d(z, A) \leq \frac{1}{2}D_2\delta^{|z|}$ , and  $20\lambda_0^2D_2/2 \leq \epsilon$ , we are done.  $\square$

**Lemma 3.8.** *J and J' satisfy (3.3), for some  $\eta = \eta(L, N, \epsilon, \lambda_0)$ .*

*Proof.* Suppose  $z \in J_i \subset J$ , for  $i \leq 0$ . Let  $z' \in J'$  be the closest point to  $z$ . If  $z' \in J'_{i-1} \cup J'_i \cup J'_{i+1}$ , then  $d(z, z') \geq \min\{\eta_1\delta^{|z|}, \eta_2\delta^{|i-1|}\}$ . Otherwise, by Lemma 3.4,  $d(z, z') \geq \frac{1}{2}D_2\delta^{|z|}$ .  $\square$

This completes the proof of Proposition 3.1.  $\square$

Observe that the relative separation condition (3.3) proven above suffices to show that we have a quasi-circle.

**Lemma 3.9.** *If J, J' are two  $\lambda$ -quasi-arcs with the same endpoints a, b, and satisfying (3.3) for some  $\eta \in (0, 1)$ , then  $\gamma = J \cup J'$  is a  $6\lambda/\eta$ -quasi-circle.*

*Proof.* Clearly  $\gamma$  is a topological circle. Let  $x, y \in \gamma$  be two points we wish to check for linear connectivity. The only non-trivial case is when (up to relabelling)  $x \in J \setminus \{a, b\}$  and  $y \in J' \setminus \{a, b\}$ .

First suppose that  $d(\{x, y\}, \{a, b\}) \geq \frac{1}{2}d(a, b)$ . Then by (3.3), we have  $d(x, y) \geq \eta \cdot \frac{1}{2}d(a, b)$ , so

$$(3.10) \quad \text{diam}(\gamma[x, y]) \leq \text{diam}(\gamma) \leq 2\lambda d(a, b) \leq \frac{4\lambda}{\eta}d(x, y),$$

where  $\gamma[x, y]$  denotes an appropriate subarc of  $\gamma$ .

Otherwise, we may suppose that  $d(x, a) \leq \frac{1}{2}d(a, b)$ , and so  $d(x, y) \geq \eta \cdot d(x, a)$ . If  $d(x, y) \geq \frac{1}{3}d(a, b)$  then as in (3.10) we have  $\text{diam}(\gamma) \leq$

$6\lambda d(x, y)$ . So we assume that  $d(x, y) \leq \frac{1}{3}d(a, b)$ , giving that  $d(y, a) \leq \frac{5}{6}d(a, b)$ , hence  $d(y, b) \geq \frac{1}{5}d(y, a)$ . Thus  $d(x, y) \geq \eta \cdot \frac{1}{5}d(y, a)$ . Therefore,

$$\begin{aligned} \text{diam}(\gamma[x, y]) &\leq \text{diam}(J[a, x]) + \text{diam}(J'[a, y]) \leq \lambda d(a, x) + \lambda d(a, y) \\ &\leq \frac{\lambda}{\eta} d(x, y) + \frac{5\lambda}{\eta} d(x, y) = \frac{6\lambda}{\eta} d(x, y). \quad \square \end{aligned}$$

We now complete the proof of the “quasi-arc  $n$ -Bogensatz.”

*Proof of Theorem 1.4.* We may assume that  $n = 2^m$ .

We claim that by induction on  $m$ , we can find  $\kappa_m = \kappa_m(L, N) \geq 1$  and  $\eta_m = \eta_m(L, N) \in (0, 1)$  so that there are  $2^m$  different  $\kappa_m$ -quasi-arcs from  $x$  to  $y$  that pairwise satisfy (3.3) with  $\eta = \eta_m$ .

The  $m = 0$  case follows from Theorem 1.7, finding a  $\kappa_0$ -quasi-arc between  $x$  and  $y$ , where  $\kappa_0 = \kappa_0(L, N)$ . We set  $\eta_0 = 1$ .

For the induction step with  $m \geq 1$ , apply Proposition 3.1 to each of the  $2^{m-1}$  previous  $\kappa_{m-1}$ -quasi-arcs, with  $\epsilon_m = \frac{1}{4}\eta_{m-1}$ . This results in  $2^m$  different  $\kappa_m$ -quasi-arcs, pairwise satisfying (3.3) for some value of  $\eta$  which we denote by  $\eta_m$ . (Here  $\kappa_m = \kappa_m(L, N, \kappa_{m-1}, \epsilon_m) > 1$ , and  $\eta_m = \eta_m(L, N, \kappa_{m-1}, \epsilon_m) > 0$ .)

Finally, Lemma 3.9 completes the proof. □

#### 4. QUASI-CIRCLES THROUGH $n$ POINTS

The following corollary of the  $n$ -Bogensatz is well known. To motivate the proof of Theorem 1.5, we include a short proof.

**Theorem 4.1.** *Let  $X$  be a connected, locally connected and locally compact metric space. If  $n \geq 2$  and  $X$  is not disconnected by the removal of any  $n - 1$  points, then any  $n$  points in  $X$  lie on a simple closed curve.*

*Proof.* The  $n = 2$  case is just a restatement of Theorem 1.1.

We prove the  $n > 2$  case by induction. Suppose  $x_1, \dots, x_n$  are given. By induction, we can find a simple closed curve  $\gamma$  containing  $x_1, \dots, x_{n-1}$ ; we relabel so that they are in the cyclic order  $x_1, \dots, x_{n-1}$ . Let  $D$  be a closed disc with centre labelled  $x_*$ , and choose subsets  $\{y_1, \dots, y_n\} \subset \partial D$  and  $\{y'_1, \dots, y'_n\} \subset \gamma$ . Let  $Y$  be the topological space formed from  $D$  and  $X$  by gluing together  $y_i$  and  $y'_i$  for each  $1 \leq i \leq n$ .

The space  $Y$  is connected, locally connected, locally compact, and cannot be disconnected by the removal of any  $n - 1$  points. Thus Theorem 1.1 gives  $n$  disjoint arcs  $\alpha_1, \dots, \alpha_n$  from  $x_*$  to  $x_n$  in  $Y$ . For each  $i = 1, \dots, n$ , let  $\beta_i$  be the closed, connected subarc of  $\alpha_i$  which contains  $x_n$  and exactly one point  $z_i$  of  $\gamma$ . Each point  $z_i$  lies in one of  $\gamma[x_1, x_2), \gamma[x_2, x_3), \dots, \gamma[x_{n-1}, x_1)$ . By the pigeonhole principle, two of

the points lie in the same interval, and so we use these two  $\beta$  arcs to find a simple closed curve containing  $x_1, \dots, x_n$ .  $\square$

This proof cannot be used directly in the quasi-arc case: the space  $Y$  has local cut points. Moreover, to apply the straightening techniques of Theorem 2.3 we need a quasi-arc of controlled size through each  $x_i$ . In adapting this proof, the following corollary of the  $n$ -Bogensatz, due to Zippin, will be useful.

**Theorem 4.2** ([Zip33, Corollary 9]). *Let  $X$  be a connected, locally connected, locally compact, separable metric space. If  $A, B \subset X$  are compact subsets of size at least  $n$ , and there is no subset  $S \subset X$  of size at most  $n - 1$  so that  $A \setminus S$  and  $B \setminus S$  lie in different components of  $X \setminus S$ , then there exists  $n$  disjoint arcs joining  $A$  and  $B$ .*

*Proof of Theorem 1.5.* The  $n = 1$  and  $n = 2$  cases follow from Theorem 1.4. We prove the  $n > 2$  case by strong induction.

By induction, there exists  $\lambda_1 = \lambda_1(L, N, n - 1)$  so that any set  $T$  of at most  $n - 1$  points in an  $N$ -doubling,  $L$ -annularly linearly connected, complete metric space  $X$  must lie in a  $\lambda_1$ -quasi-circle  $\gamma$  with  $\text{diam}(\gamma) \leq \lambda_1 \text{diam}(T)$ .

Suppose  $x_1, \dots, x_n$  are given. Without loss of generality, we may assume that  $d(x_1, x_2) \leq d(x_i, x_j)$  for all  $i \neq j$ , and that  $d(x_1, x_i) \leq d(x_1, x_{i+1})$  for  $i = 2, \dots, n - 1$ . We rescale so that  $d(x_1, x_n) = 1$ .

Let  $s = d(x_1, x_2)$ ,  $S = \text{diam}(\{x_1, \dots, x_n\}) \in [1, 2]$ , and set  $\delta = 1/200L^2\lambda_1^3$ .

The proof splits into two cases.

*Case 1:* Suppose  $s \geq \delta^{n-1}$ .

By induction, there exists a  $\lambda_1$ -quasi-circle  $\alpha_1$  through  $x_2, \dots, x_n$  of diameter at most  $2\lambda_1$ , and at least  $s$ . We relabel  $x_2, \dots, x_n$  so that they lie in  $\alpha_1$  in this cyclic order.

Now suppose  $d(x_1, \alpha_1) \leq s/10L\lambda_1$ . Then one can alter  $\alpha_1$  using a detour in  $A(x_1, s/10L^2\lambda_1, s/5\lambda_1)$  to find a simple closed curve  $\alpha_2$  which does not meet  $B(x_1, s/10L^2\lambda_1)$ . Since this only cuts out loops of  $\alpha_1$  in  $B(x_1, s/5)$ ,  $\alpha_2$  agrees with  $\alpha_1$  outside  $B(x_1, s/5)$ , and is a  $\lambda_1$ -quasi-arc there. Therefore we can apply Theorem 2.3 with  $\epsilon = s/100L^2\lambda_1$  to straighten  $\alpha_2$  into a  $\lambda_2$ -quasi-circle  $\beta_1$ , which passes through  $x_2, \dots, x_n$ , and does not meet  $B(x_1, s/20L^2\lambda_1)$ , for  $\lambda_2 = \lambda_2(L, N, \lambda_1, s/S) \geq \lambda_1$ .

If  $d(x_1, \alpha_1) \geq s/10L\lambda_1$ , then we set  $\beta_1 = \alpha_1$  and continue.

By the  $n = 2$  case of the theorem, we find a  $\lambda_1$ -quasi-circle  $\beta_2$  through  $x_1$  of diameter at least  $s/50L^2\lambda_1^2$ , inside  $B(x_1, s/40L^2\lambda_1)$ .

As  $X$  has no local cut points, no two disjoint compacta can be separated by removing any finite number of points. Therefore, Theorem 4.2

implies that we can join  $\beta_1$  to  $\beta_2$  by  $2n$  disjoint arcs inside  $B(x_1, 4LS)$ . We can control the separation of these arcs.

**Lemma 4.3** (Cf. [Mac10, Lemma 3.3]). *We can join  $\beta_1$  to  $\beta_2$  by  $2n$  arcs in  $B(x_1, 4\lambda_1 LS)$  that are  $\delta_* S$ -separated, for  $\delta_* = \delta_*(L, N, \lambda_1, \lambda_2, s/S) > 0$ .*

*Proof.* This follows from a compactness argument: if not, there is a sequence of configurations giving counterexamples. To be precise, we can find (on rescaling to  $S = 1$ ), a sequence

$$\{\mathcal{C}^i = (X^{(i)}, x_1^{(i)}, \beta_1^{(i)}, \beta_2^{(i)})\}_{i \in \mathbb{N}}$$

so that for each  $i \in \mathbb{N}$ ,  $X^{(i)}$  is an  $L$ -annularly linearly connected,  $N$ -doubling, complete metric space with base point  $x_1^{(i)}$ , and  $\beta_1^{(i)}$  and  $\beta_2^{(i)}$  are  $\lambda_2$ -quasi-circles in  $B(x_1^{(i)}, 2\lambda_1 S)$ , with uniformly controlled diameter and separation. Moreover, there do not exist  $2n$  disjoint arcs connecting  $\beta_1^{(i)}$  to  $\beta_2^{(i)}$  which are  $1/i$  separated.

Such configurations have a subsequence that converges to a limit configuration  $(X^\infty, x_1^\infty, \beta_1^\infty, \beta_2^\infty)$  in the Gromov-Hausdorff topology. We apply Theorem 4.2 to the limit space  $X^\infty$  to find  $2n$  disjoint arcs joining  $\beta_1^\infty$  to  $\beta_2^\infty$  inside  $B(x_1^\infty, 3\lambda_1 LS)$ . As these arcs are disjoint, they are separated by some definite distance. These arcs will then lift back to  $\mathcal{C}^i$  for sufficiently large  $i$  to give a contradiction.  $\square$

Now of these  $2n$  arcs, at most  $n$  of them can be  $\frac{1}{2}\delta_* S$  close to any of the  $n$  different points  $x_1, \dots, x_n$ . Therefore, we can find  $n$  arcs  $\gamma'_1, \dots, \gamma'_n$  which join  $\beta_1$  to  $\beta_2$ , are  $\delta_* S$ -separated, and have distance at least  $\frac{1}{2}\delta_* S$  from any  $x_i$ .

By the pigeonhole principle, two of the arcs in  $\{\gamma'_j\}$  must have endpoints that lie in the same arc out of  $\beta_1(x_2, x_3), \dots, \beta_1(x_{n-1}, x_n)$  and  $\beta_1(x_n, x_2)$ . Let us call these arcs  $\gamma_1 = \gamma_1[y_1, z_1]$  and  $\gamma_2 = \gamma_2[y_2, z_2]$ , where  $y_1, y_2 \in \beta_1$ , and  $z_1, z_2 \in \beta_2$ .

Let  $\gamma_3$  be the simple closed curve formed from  $\beta_1[y_1, y_2]$  (containing  $x_2, \dots, x_n$ ),  $\gamma_1, \gamma_2$ , and  $\beta_2[z_1, z_2]$  (containing  $x_1$ ). As  $\beta_1, \beta_2$  are quasi-arcs, and we have control on the distance of  $\gamma_1, \gamma_2$  from  $x_1, \dots, x_n$ , we can apply Theorem 2.3 to straighten  $\gamma_2$  into a  $\lambda$ -quasi-circle  $\gamma$ , where  $\lambda = \lambda(L, N, \lambda_1, \lambda_2, \delta_*, s/S) = \lambda(L, N, n)$ . Moreover,  $\text{diam}(\gamma) \leq 4\lambda_1 S$  as desired.

*Case 2:* Suppose  $s < \delta^{n-1}$ .

This case is similar to Case 1, except now  $s$  may be arbitrarily small, so we replace  $\beta_2$  by a quasi-circle through  $x_1$  and all points close to it.




FIGURE 2. Joining two quasi-circles in case 2

Consider the set  $U = \{d(x_1, x_i)\}_{i=3}^{n-1}$  of size  $n - 3$ . One of the intersections  $U \cap [\delta^{n-1}, \delta^{n-2}), \dots, U \cap [\delta^2, \delta^1)$  is empty. Thus there exists  $m \in \{2, \dots, n - 1\}$  so that  $d(x_1, x_m) \leq \delta d(x_1, x_{m+1})$ .

Let  $\alpha_1$  be a  $\lambda_1$ -quasi-circle through  $\{x_1, x_{m+1}, x_{m+2}, \dots, x_n\}$ . Similarly to case 1, use the  $L$ -annularly linearly connected property for  $A(x_1, 4\lambda_1 Ld(x_1, x_m), 8\lambda_1 Ld(x_1, x_m))$  to find a circle  $\alpha_2$  that detours  $\alpha_1$  around  $B(x_1, 4\lambda_1 d(x_1, x_m))$ , while only cutting out loops in

$$B(x_1, 8\lambda_1^2 L^2 d(x_1, x_m)) \subset B(x_1, \frac{4}{5}d(x_1, x_{m+1})).$$

In particular,  $\alpha_2$  contains  $\{x_{m+1}, \dots, x_n\}$ , and we relabel so they are in this cyclic order.

We use Theorem 2.3 with  $\epsilon = \lambda_1 d(x_1, x_m)$  to straighten  $\alpha_2$  into a quasi-circle  $\beta_1$  which remains outside  $B(x_1, 3\lambda_1 d(x_1, x_m))$ . Moreover,  $\beta_1$  will  $9\lambda_1 Ld(x_1, x_m)$ -follow  $\alpha_1$ . Inside  $B(x_1, 9\lambda_1^2 L^2 d(x_1, x_m))$ ,  $\beta_1$  is a  $\lambda_3$ -quasi-arc, where  $\lambda_3 = \lambda_3(L, N, \lambda_1)$  is independent of  $d(x_1, x_m)$ .

Let  $\beta_2$  be a  $\lambda_1$ -quasi-circle through  $\{x_1, \dots, x_m\}$ , relabelled so they are in this cyclic order, of diameter at most  $2\lambda_1 d(x_1, x_m)$  (see Figure 2).

As in Case 1, by Theorem 4.2 we can join  $\beta_1$  to  $\beta_2$  by  $2n$  disjoint arcs inside  $B(x_1, 10\lambda_1^2 L^2 d(x_1, x_m))$ . Inside this ball we have control on the diameter of  $\beta_2$ , and the quasi-arc constants of  $\beta_1, \beta_2$ . Therefore, a similar argument to Lemma 4.3 gives that these arcs are  $\delta_* d(x_1, x_m)$  separated, where  $\delta_* = \delta_*(L, N, \lambda_1, \lambda_3)$ .

As before,  $n$  of these arcs, let us call them  $\gamma'_1, \dots, \gamma'_n$ , will join  $\beta_1$  to  $\beta_2$ , be  $\delta_* d(x_1, x_m)$ -separated, and have distance at least  $\frac{1}{2}\delta_* d(x_1, x_m)$  from any  $x_i$ .

By the pigeonhole principle, two of the arcs in  $\{\gamma'_j\}$  must have endpoints that lie in the same arc out of  $\beta_2(x_1, x_2), \dots, \beta_2(x_{m-1}, x_m)$  and  $\beta_2(x_m, x_1)$ . Let us call these arcs  $\gamma_1 = \gamma_1[y_1, z_1]$  and  $\gamma_2 = \gamma_2[y_2, z_2]$ , where  $y_1, y_2 \in \beta_1$ , and  $z_1, z_2 \in \beta_2$ . (Again, see Figure 2.)

Using the fact that  $\beta_1$  follows  $\gamma_1$ , we see that the diameter of the smaller arc  $\beta_1[y_1, y_2]$  is at most  $100\lambda_1^3 L^2 d(x_1, x_m) < \frac{1}{2}d(x_1, x_{m+1})$ . Therefore, there is a subarc  $\beta'_1[y_1, y_2] \subset \beta_1$  containing  $x_{m+1}, \dots, x_n$ .

Let  $\gamma_3$  be the simple closed curve formed from  $\beta'_1, \gamma_1, \gamma_2$ , and  $\beta_2[z_1, z_2]$  (containing  $x_1, \dots, x_m$ ). As  $\beta_1, \beta_2$  are quasi-arcs, and we have control on the distance of  $\gamma_1, \gamma_2$  from  $x_1, \dots, x_n$ , we can apply Theorem 2.3 with  $\epsilon = \frac{1}{2}\delta_* d(x_1, x_m)$  to straighten  $\gamma_2$  into a quasi-circle  $\gamma$ .

Let us show that  $\gamma$  is a quasi-circle with controlled constant. Observe that  $\gamma$   $D$ -follows  $\alpha_1$ , where  $D = 10\lambda_1^2 L^2 d(x_1, x_m)$ . Let  $f : \gamma \rightarrow \alpha_1$  be the associated map. Consider the following three cases.

(i) From Theorem 2.3, there exists  $\lambda_4 = \lambda_4(L, N, \lambda_1, \delta_*)$  so that if  $z, z' \in \gamma \cap B(x_1, 10\lambda_1 D)$ , then  $\text{diam}(\gamma[z, z']) \leq \lambda_4 d(z, z')$ .

(ii) If  $\gamma[z, z'] \cap B(x_1, 2\lambda_1 D) = \emptyset$ , then  $\gamma[z, z'] = \alpha_1[z, z']$ , so we have  $\text{diam}(\gamma[z, z']) \leq \lambda_1 d(z, z')$ .

(iii) Otherwise, we know that  $\text{diam}(\gamma[z, z']) \geq 8\lambda_1 D$ , so

$$\begin{aligned} \text{diam}(\gamma[z, z']) &\leq 2D + \text{diam}(\alpha_1[f(z), f(z')]) \leq 2D + \lambda_1 d(f(z), f(z')) \\ &\leq 2D + 2\lambda_1 D + \lambda_1 d(z, z'), \end{aligned}$$

so  $\frac{1}{2} \text{diam}(\gamma[z, z']) \leq \lambda_1 d(z, z')$ , thus  $\text{diam}(\gamma[z, z']) \leq 2\lambda_1 d(z, z')$ .

Therefore,  $\gamma$  is a  $\lambda$ -quasi-circle, where  $\lambda = \max\{\lambda_4, 2\lambda_1\}$  depends only on  $L, N, n$ . Observe that  $\text{diam}(\gamma) \leq 4\lambda_1 S$ , as desired.  $\square$

## REFERENCES

- [Ahl63] L. V. Ahlfors. Quasiconformal reflections. *Acta Math.*, 109:291–301, 1963.
- [BK05] M. Bonk and B. Kleiner. Quasi-hyperbolic planes in hyperbolic groups. *Proc. Amer. Math. Soc.*, 133(9):2491–2494 (electronic), 2005.
- [BS00] M. Bonk and O. Schramm. Embeddings of Gromov hyperbolic spaces. *Geom. Funct. Anal.*, 10(2):266–306, 2000.
- [Mac08] J. M. Mackay. Existence of quasi-arcs. *Proc. Amer. Math. Soc.*, 136(11):3975–3981, 2008.
- [Mac10] J. M. Mackay. Spaces and groups with conformal dimension greater than one. *Duke. Math. J.*, 153(2):211–227, 2010.
- [MS11] J. M. Mackay and A. Sisto. Quasi-hyperbolic planes in relatively hyperbolic groups. *Preprint*, arxiv:1111.2499, 2011.
- [Nöb32] G. Nöbeling. Eine Verschärfung des  $n$ -Beinsatzes. *Fund. Math.*, 18:23–38, 1932.
- [Tuk96] P. Tukia. Spaces and arcs of bounded turning. *Michigan Math. J.*, 43(3):559–584, 1996.
- [TV80] P. Tukia and J. Väisälä. Quasisymmetric embeddings of metric spaces. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 5(1):97–114, 1980.

- [TV08] C. Thomassen and A. Vella. Graph-like continua, augmenting arcs, and Menger's theorem. *Combinatorica*, 28(5):595–623, 2008.
- [Why48] G. T. Whyburn. On  $n$ -arc connectedness. *Trans. Amer. Math. Soc.*, 63:452–456, 1948.
- [Zip33] L. Zippin. Independent arcs of a continuous curve. *Ann. of Math. (2)*, 34(1):95–113, 1933.

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, OXFORD, UK.  
*E-mail address:* john.mackay@maths.ox.ac.uk