

Thin-shell concentration for convex measures

Matthieu Fradelizi, Olivier Guédon and Alain Pajor

Abstract

We prove that for $s < 0$, s -concave measures on \mathbb{R}^n satisfy a thin-shell concentration similar to the log-concave case. It leads to a Berry-Esseen type estimate for most of their one dimensional marginal distributions. We also establish sharp reverse Hölder inequalities for s -concave measures.

1 Introduction

For any subsets $A, B \subset \mathbb{R}^n$, the Minkowski sum is defined by

$$A + B = \{a + b : a \in A, b \in B\}.$$

Let $s \in [-\infty, 1]$. A measure μ on \mathbb{R}^n is called s -concave whenever

$$\mu((1 - \lambda)A + \lambda B) \geq ((1 - \lambda)\mu(A)^s + \lambda\mu(B)^s)^{1/s},$$

for every $\lambda \in [0, 1]$ and every compact subsets $A, B \subset \mathbb{R}^n$ such that $\mu(A)\mu(B) > 0$. When $s = 0$, this inequality should be read as

$$\mu((1 - \lambda)A + \lambda B) \geq \mu(A)^{1-\lambda}\mu(B)^\lambda$$

2010 *Mathematics Subject Classification*: Primary 60E15, 60F10, 52A23; Secondary 52A40, 46B09.

Key words and phrases: Isotropic, convex measure, concentration inequalities, thin-shell, large-deviation, KLS conjecture.

and it defines μ as a log-concave measure. When $s = -$, the measure is said to be convex and the inequality is replaced by

$$\mu((1 - \lambda)A + \lambda B) \geq \min(\mu(A), \mu(B)).$$

Notice that the class of s -concave measures on \mathbb{R}^n is decreasing in s so that any s -concave measure is a convex measure. Any s -concave measure with $s \geq 0$ is log-concave and the thin-shell concentration for log-concave measures has been studied in [16, 17, 19, 22, 23]. The purpose of this paper is to prove a thin-shell concentration for s -concave measures in the case $s < 0$, which we consider from now on. By measure, we always mean probability measure.

The class of s -concave measures was introduced and studied in [10, 11], where a complete characterization was established. An s -concave measure is supported on some convex subset of an affine subspace where it has a density (see Section 2 for more details). When the support of an s -concave measure μ generates the whole space, we say that μ is full-dimensional.

A random vector with an s -concave distribution is called s -concave. The linear image of an s -concave random vector is also s -concave. We say that a random vector is full-dimensional if its distribution is full-dimensional. It is known that any semi-norm of an s -concave random vector with $s < 0$ has moments of all order $p \in (0, -1/s)$ (see [10] and [1]). The Euclidean norm of an s -concave random vector X has a finite moment of order 2 if and only if $s > -1/2$. Since we are interested in comparison of moments of the Euclidean norm with the moment of order 2, we will always assume that $-1/2 < s < 0$.

Let $n \geq 1$ be an integer. The Euclidean space \mathbb{R}^n is equipped with its Euclidean norm $\|\cdot\|_2$ and its scalar product $\langle \cdot, \cdot \rangle$. Its unit sphere is denoted by S^{n-1} and its unit ball by B_2^n . We say that a random vector X is isotropic if $\mathbb{E}X = 0$ and for every $\theta \in S^{n-1}$, $\mathbb{E}\langle X, \theta \rangle^2 = 1$. Observe that if X is an s -concave full-dimensional random vector and $-1/2 < s$, we can always find an affine transformation A such that AX is isotropic.

Let $p \in \mathbb{R}$ and $X \in \mathbb{R}^n$ be a random vector. Assume that $\|X\|_2$ has finite moments of order 2 and p with the convention that $(\mathbb{E}\|X\|_2^p)^{1/p} = \exp(\mathbb{E} \ln \|X\|_2)$ for $p = 0$. We define

$$\alpha_p(X) := \left| \frac{(\mathbb{E}\|X\|_2^p)^{1/p}}{(\mathbb{E}\|X\|_2^2)^{1/2}} - 1 \right|.$$

Our main result is the following

Theorem 1. *Let $r > 2$. Let $X \in \mathbb{R}^n$ be a full-dimensional $(-1/r)$ -concave random vector.*

If X is isotropic, then for any p such that $p \leq c \min(r, n^{1/3})$, we have

$$\alpha_p(X) \leq \frac{C p - 2}{r} + \left(\frac{C p - 2}{n^{1/3}} \right)^{3/5},$$

where C and c are universal constants.

In the general case (when X is not isotropic), let A be an affine transformation such that AX is full-dimensional and isotropic. Then for any $p \in \mathbb{R}$ such that $p \leq c \min\left(r, \frac{n^{1/3}}{\|A\|^{2/3}\|A^{-1}\|^{2/3}}\right)$, we have

$$\alpha_p(X) \leq \frac{C p - 2}{r} + \left(\frac{C p - 2 (\|A\| \|A^{-1}\|)^{2/3}}{n^{1/3}} \right)^{3/5},$$

where C and c are universal constants.

We also show (see Remark 15) that for $r > n + \sqrt{n}$, the estimate of $\alpha_p(X)$ in Theorem 1 can be improved and recovers the estimate of the log-concave case from [19].

To present the connections between moment inequalities, concentration in a thin-shell and the Berry-Esseen theorem for one dimensional marginals, let us introduce some notations.

Let $X \in \mathbb{R}^n$ be an isotropic random vector. Thus $\mathbb{E} X^2 = n$. Define $\varepsilon(X)$ to be the smallest number $\varepsilon > 0$ such that

$$\mathbb{P} \left(\left| \frac{X^2}{\sqrt{n}} - 1 \right| \geq \varepsilon \right) \leq \varepsilon. \quad (1)$$

If $\varepsilon(X) = o(1)$ with respect to the dimension n , we say that X is concentrated in a thin-shell. This is the usual jargon of the subject. More rigorously, it suggests that we are considering a sequence of random vectors (X_n) with $X_n \in \mathbb{R}^n$ and that $\varepsilon(X_n) = o(1)$ as n goes to ∞ . It was shown in [2] (see also [13, 14]) that if an isotropic random vector X uniformly distributed on a convex body in \mathbb{R}^n is such that $\varepsilon(X) = o(1)$, then almost all one dimensional marginal distributions of X satisfy a Berry-Esseen theorem. More generally, let $X \in \mathbb{R}^n$ be an isotropic random vector, it was proved in [7] that

$$\sigma_{n-1} \left(\theta \in S^{n-1} : \sup_{t \in \mathbb{R}} \mathbb{P}(X, \theta \leq t) - \Phi(t) \geq 4\varepsilon(X) + \delta \right) \leq 4n^{3/8} e^{-cn\delta^4},$$

where σ_{n-1} denotes the rotation invariant probability measure on the unit sphere S^{n-1} , Φ is the standard normal distribution function and $c > 0$ is a universal constant. It is worth noticing that the result from [7] does not assume log-concavity. Assuming only that X is isotropic, we get that if $\varepsilon(X)$ is $o(1)$ then almost all the one dimension marginal distributions of X are approximately Gaussian. The fact that indeed for all log-concave random vector $\varepsilon(X) = o(1)$ was proved later in [22, 17] and the best estimate at this date [19] is that

$$\varepsilon(X) = O(n^{-1/6} \log n).$$

Now let $p > 2$ and assume that X is isotropic and that X_2 has a finite moment of order p . Then $\varepsilon(X)$ is $o(1)$ if and only if $\alpha_p(X)$ is $o(1)$, see Remark 4 below. Hence Theorem 1 ensures that if $r \rightarrow +\infty$ with the dimension n then any isotropic $(-1/r)$ -concave random vector satisfies a thin-shell concentration and therefore almost all its one dimensional marginals verify a Berry-Esseen theorem. As a matter of fact, this condition on r is necessary. If r is fixed and does not depend on the dimension n , Proposition 5 gives an example of an isotropic $(-1/r)$ -concave random vector $X \in \mathbb{R}^n$ which does not satisfy a thin-shell concentration. Remark 6 also shows the asymptotic sharpness of Theorem 1, since for this example, for a fixed $p > 2$, $\alpha_p(X) \geq C(p-2)/r$ for r and n large enough, where $C > 0$ is a universal constant.

To build the proof of Theorem 1, we need to extend to the case of s -concave measures several tools coming from the study of log-concave measures. This is the purpose of Section 2. Some of them were already achieved by Bobkov [8], like analog of the Ball's bodies [5] in the s -concave setting. Some others were also noticed previously (see e.g. [8], [1]) but not with the most accurate point of view. These new ingredients are analog to the results of [12] in the log-concave setting and are at the heart of our proof. As in the approach of [16] or [19], an important ingredient is the log-Sobolev inequality on $SO(n)$. It follows e.g. from the work of Bakry and Émery [4] and the calculus of the Ricci curvature of $SO(n)$ (see [21, Formula (F6)] for example) that for any Lipschitz function $f : SO(n) \rightarrow \mathbb{R}^+$ (see sections 3 and 4 for definitions)

$$\mathbb{E}(f(U) \log f(U)) - \mathbb{E}f(U) \log(\mathbb{E}f(U)) \leq \frac{c}{n} \mathbb{E}(\log f(U)^2 f(U)), \quad (2)$$

where U is uniformly distributed on $SO(n)$. It allows to get reverse Hölder inequalities (see inequality (15) in [16]): for every $f : SO(n) \rightarrow \mathbb{R}$, let L be

the log-Lipschitz constant of f (that is the Lipschitz constant of $\log f$), then for every $q > r > 0$,

$$(\mathbb{E} f(U)^q)^{1/q} \leq \exp\left(\frac{cL^2}{n}(q-r)\right) (\mathbb{E} f(U)^r)^{1/r}, \quad (3)$$

where U is uniformly distributed on $SO(n)$.

Let X be a $(-1/r)$ -concave random vector in \mathbb{R}^n with full-dimensional support and distributed according to a measure with a density function $w : \mathbb{R}^n \rightarrow \mathbb{R}_+$. For any linear subspace E , denote by P_E the orthogonal projection onto E and for any $x \in E$ denote by

$$\pi_E w(x) = \int_{x+E^\perp} w(y) dy$$

the marginal of w on E . Given an integer k between 1 and n , a real number $p \in (-k, r)$, a linear subspace E_0 of \mathbb{R}^n of dimension k and $\theta_0 \in S(E_0)$, where $S(E_0)$ denotes the unit sphere of E_0 , we define the function $h_{k,p} : SO(n) \rightarrow \mathbb{R}_+$ by

$$h_{k,p}(u) := S^{k-1} \int_0^\infty t^{p+k-1} \pi_{u(E_0)} w(tu(\theta_0)) dt, \quad (4)$$

for every $u \in SO(n)$, where S^{k-1} denotes the area of the sphere.

Following the approach of [23, 16], we observe that for any $p \in (-k, r)$

$$\mathbb{E} X^p = \frac{\Gamma((p+n)/2)\Gamma(k/2)}{\Gamma(n/2)\Gamma((p+k)/2)} \mathbb{E} h_{k,p}(U), \quad (5)$$

where U is uniformly distributed on $SO(n)$. In view of (5) and the definition of $h_{k,p}$, we notice that it is of importance to work with family of measures which are stable after taking the marginals and it is clear from the definition that for any subspace E , if X is $(-1/r)$ -concave, then $P_E X$ is also $(-1/r)$ -concave.

In the next section 2, we first introduce more notation and recall important facts concerning convex measures. Then we give an example of an isotropic $(-1/r)$ -concave random vector $X \in \mathbb{R}^n$ that does not satisfy a thin-shell concentration, when r is **kn**

these two sections are either classical or variation of known results; their proofs are shifted to the appendix.

Acknowledgement. We deeply and warmly thank the referee for his constructive comments on the first version of the paper. It forced us to clarify several main points and we hope that it improved the presentation of the paper.

2 Preliminary results for s -concave measures

We first recall some properties of s -concave measures and their relation to β -concave functions.

The class of s -concave measures was introduced and studied in [10, 11], where the following complete characterization was established. An s -concave measure μ on \mathbb{R}^n is supported on some convex subset of an affine subspace where it has a density. When this subspace is the whole space, we say that μ is full-dimensional. In this case, its density w is β -concave with $\beta = s/(1 - ns)$. Recall that a function $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ is called β -concave whenever

$$f((1 - \lambda)x + \lambda y) \geq ((1 - \lambda)f(x)^\beta + \lambda f(y)^\beta)^{1/\beta}$$

for every $\lambda \in [0, 1]$ and every $x, y \in \mathbb{R}^n$ such that $f(x)f(y) > 0$, where the right hand side is replaced by $f(x)^{1-\lambda}f(y)^\lambda$ for $\beta = 0$. Note that when $\beta < 0$ which will be the case below, β -concavity means that f^β is convex on its convex support $\bullet \bullet f > 0$.

We will use a similar language for probability measure, random vector and function which are related here as distribution, law of a random vector and density of probability. It is important to remember that when $X \in \mathbb{R}^n$ is $(-1/r)$ -concave full-dimensional, then the result recalled above states that its distribution has a support that generates \mathbb{R}^n and has a density which is $(-1/(n + r))$ -concave.

Recall that for every $x > 0$, $\Gamma(x) = \int_0^\infty u^{x-1}e^{-u} du$ and for every $x, y > 0$, $B(x, y) = \int_0^1 u^{x-1}(1 - u)^{y-1} du = \int_0^{+\infty} u^{x-1}(u + 1)^{-(x+y)} du$.

The following inequality of Paley-Zygmund type is well known.

Lemma 2. *Let $2 < p < s$. Let Y be a non-negative random variable with finite s -moment. Then for every $0 \leq t \leq (\mathbb{E}Y^p)^{1/p}$ we have*

$$\mathbb{P}(Y \geq t) \geq \left(\frac{\mathbb{E}Y^p - t^p}{(\mathbb{E}Y^s)^{p/s}} \right)^{s/(s-p)}.$$

Proof. Using Hölder inequality, we have

$$\mathbb{E}Y^p = \mathbb{E}Y^p 1_{Y < t} + \mathbb{E}Y^p 1_{Y \geq t} \leq t^p + (\mathbb{E}Y^s)^{p/s} \mathbb{P}(Y \geq t)^{1-p/s}.$$

Thus

$$\mathbb{P}(Y \geq t) \geq \left(\frac{\mathbb{E}Y^p - t^p}{(\mathbb{E}Y^s)^{p/s}} \right)^{s/(s-p)}.$$

□

Proposition 3. *Let $2 < p < s$. Let $X \in \mathbb{R}^n$ be an isotropic random vector such that X_2 has a finite s -moment. Then*

$$\min \left(\frac{\alpha_p(X)}{2}, \left(\frac{p\alpha_p(X)/2}{(\alpha_s(X) + 1)^p} \right)^{s/(s-p)} \right) \leq \varepsilon(X) \leq ((\alpha_p(X) + 1)^p - 1)^{1/3}.$$

Proof. Let $\varepsilon > 0$. Applying Lemma 2 to $Y = X_2 / (\mathbb{E} X_2^2)^{1/2}$, $t = \varepsilon + 1$ and noticing that $\mathbb{E}Y^p = (\alpha_p(X) + 1)^p$, $\mathbb{E}Y^s = (\alpha_s(X) + 1)^s$, we get that

$$\mathbb{P} \left(\frac{X_2}{(\mathbb{E} X_2^2)^{1/2}} \geq 1 + \varepsilon \right) \geq \left(\frac{(\alpha_p(X) + 1)^p - (\varepsilon + 1)^p}{(\alpha_s(X) + 1)^p} \right)^{s/(s-p)}$$

whenever $0 < \varepsilon \leq \alpha_p(X)$. Since for $p \geq 1$ and $x \geq y \geq 1$, $x^p - y^p \geq p(x - y)$, we have

$$\mathbb{P} \left(\frac{X_2}{(\mathbb{E} X_2^2)^{1/2}} \geq 1 + \varepsilon \right) \geq \left(\frac{p(\alpha_p(X) - \varepsilon)}{(\alpha_s(X) + 1)^p} \right)^{s/(s-p)}.$$

Therefore

$$\mathbb{P} \left(\frac{X_2}{(\mathbb{E} X_2^2)^{1/2}} \geq 1 + \varepsilon \right) \geq \left(\frac{p\alpha_p(X)/2}{(\alpha_s(X) + 1)^p} \right)^{s/(s-p)}$$

whenever $0 < \varepsilon \leq \alpha_p(X)/2$. The left-hand side inequality follows.

Since for $q \geq 1$, $x - 1 \leq x^q - 1$ for every $x \geq 0$, Markov inequality gives

$$\mathbb{P} \left(\left| \frac{X_2}{(\mathbb{E} X_2^2)^{1/2}} - 1 \right| \geq \varepsilon \right) \leq \mathbb{P} \left(\left| \frac{X_2^q}{(\mathbb{E} X_2^2)^{q/2}} - 1 \right| \geq \varepsilon \right) \leq \frac{\mathbb{E} \left| \frac{|X_2|^q}{(\mathbb{E} |X_2|^2)^{q/2}} - 1 \right|^2}{\varepsilon^2}.$$

To conclude the right-hand side inequality, take $q = p/2$ and observe that

$$\mathbb{E} \left| \frac{X_2^q}{(\mathbb{E} X_2^2)^{q/2}} - 1 \right|^2 = (\alpha_{2q}(X) + 1)^{2q} + 1 - 2(\alpha_q(X) + 1)^q \leq (\alpha_{2q}(X) + 1)^{2q} - 1.$$

□

Remark 4. Let $2 < p < s$. Let $X \in \mathbb{R}^n$ be an isotropic random vector such that X_2 has a finite s -moment. Proposition 3 shows that $\varepsilon(X)$ is $o(1)$ if and only if $\alpha_p(X)$ is $o(1)$ when $n \rightarrow \infty$.

Now we estimate $\varepsilon(X)$ for an example which shows that an isotropic $(-1/r)$ -concave random vector $X \in \mathbb{R}^n$ may not satisfy a thin-shell concentration. In the proposition below, the notation $\underline{\lim}$ refers to the limit inferior.

Proposition 5. Let $r > 2$. There exists a sequence $(X_n)_n$ of isotropic $(-1/r)$ -concave random vectors $X_n \in \mathbb{R}^n$ such that

$$\underline{\lim}_{n \rightarrow \infty} \varepsilon(X_n) \geq c(r) > 0,$$

where $c(r) > 0$ depends only on r .

Proof. Let $r > 2$ and $2 < p < r$ and let $X_n \in \mathbb{R}^n$ be an isotropic random vector with density

$$f_{n,r}(x) = \frac{c_1}{(1 + c_2 \|x\|^2)^{r+n}},$$

where c_1 and c_2 are normalization factors. From [10, 11], such a random vector is $(-1/r)$ -concave. An immediate computation gives that

$$\frac{(\mathbb{E} X_n^p)^{1/p}}{(\mathbb{E} X_n^2)^{1/2}} = \left(\frac{B(n+p, r-p)}{B(n, r)} \right)^{1/p} \left(\frac{B(n+2, r-2)}{B(n, r)} \right)^{-1/2}.$$

For fixed r and $2 < p < r$, we have

$$\lim_{n \rightarrow +\infty} \frac{(\mathbb{E} X_n^p)^{1/p}}{(\mathbb{E} X_n^2)^{1/2}} = \left(\frac{\Gamma(r-p)}{\Gamma(r)} \right)^{1/p} \left(\frac{\Gamma(r-2)}{\Gamma(r)} \right)^{-1/2} \quad (6)$$

and by the strict log-convexity of the Gamma function, we have

$$\lim_{n \rightarrow +\infty} (\alpha_p(X_n) + 1) = \lim_{n \rightarrow +\infty} \frac{(\mathbb{E} X_n^p)^{1/p}}{(\mathbb{E} X_n^2)^{1/2}} > 1.$$

As a consequence for any $2 < p < r$, $\lim_{n \rightarrow +\infty} \alpha_p(X_n) > 0$.

Now let $2 < p < s < r$. From Proposition 3, we get

$$\underline{\lim}_{n \rightarrow +\infty} \varepsilon(X_n) \geq \lim_{n \rightarrow \infty} \min \left(\frac{\alpha_p(X_n)}{2}, \left(\frac{p\alpha_p(X_n)/2}{(\alpha_s(X_n) + 1)^p} \right)^{s/(s-p)} \right) > 0. \quad (7)$$

Choose $p = (2+r)/2$ and $s = (p+r)/2$ for which $2 < p < s < r$ and note that the right hand side term in equation (7) depends only on r . This concludes the proof. □

Remark 6. Let $2 < p < r$ and let $r \rightarrow \infty$. Applying Stirling formula in (6) when $r \rightarrow \infty$, a calculation gives that

$$\lim_{r \rightarrow \infty} r \lim_{n \rightarrow \infty} \alpha_p(X_n) = (p - 2)/2.$$

This asymptotic estimate shows that for a fixed $p > 2$ and r and n large enough, then $\alpha_p(X_n) \geq C(p - 2)/r$ where $C > 0$ is a universal constant. This proves the sharpness of Theorem 1 under these conditions.

We now prove some inequalities for s -concave measures that will be useful tools in the next section.

Theorem 7. (1) Let $f : [0, \infty) \rightarrow [0, \infty)$ be a measurable function such that $f_\infty > 0$. Then

$$p \rightarrow \left(\int_0^\infty pt^{p-1} f(t) dt / f_\infty \right)^{1/p}$$

is non-decreasing on its domain of definition.

(2) Let $\alpha > 0$ and $f : [0, \infty) \rightarrow [0, \infty)$ be $(-1/\alpha)$ -concave, continuous and integrable. Define $H_f : [0, \alpha) \rightarrow \mathbb{R}_+$ by

$$H_f(p) = \begin{cases} \frac{1}{B(p, \alpha - p)} \int_0^{+\infty} t^{p-1} f(t) dt & \text{for } 0 < p < \alpha \\ f(0) & \text{for } p = 0. \end{cases}$$

Then H_f is log-concave on $[0, \alpha)$.

The proof of the first part may be treated as in Lemma 2.1 in [25] and the proof of the second part is identical to the well known $(1/n)$ -concave case [12]. We postpone the proof of Theorem 7 to the appendix.

We present several consequences of this result such as some reverse Hölder inequalities with sharp constants in the spirit of Borell's [12] and Berwald's [6] inequalities.

Corollary 8. Let $r > 0$ and μ be a $(-1/r)$ -concave measure on \mathbb{R}^n . Let $\phi : \mathbb{R}^n \rightarrow \mathbb{R}_+$ such that $\mu_{\bullet\bullet}(\phi > 0)$ is convex and ϕ is concave on $\mu_{\bullet\bullet}(\phi > 0)$. Then the function

$$p \rightarrow \begin{cases} \frac{1}{pB(p, r-p)} \int \phi(x)^p d\mu(x) & \text{for } 0 < p < r \\ \mu_{\bullet\bullet}(\phi > 0) & \text{for } p = 0 \end{cases}$$

is log-concave on $[0, r)$.

Moreover, if $\mu_{\bullet\bullet}(\phi > 0) > 0$ then for any $0 < p \leq q < r$,

$$\left(\int_{\mathbb{R}^n} \phi(x)^q \frac{d\mu(x)}{\mu_{\bullet\bullet}(\phi > 0)} \right)^{1/q} \leq \frac{(qB(q, r-q))^{1/q}}{(pB(p, r-p))^{1/p}} \left(\int_{\mathbb{R}^n} \phi(x)^p \frac{d\mu(x)}{\mu_{\bullet\bullet}(\phi > 0)} \right)^{1/p}.$$

Proof. By the concavity of ϕ , for every $u, v \geq 0$ and every $\lambda \in [0, 1]$

$$(1 - \lambda)\mu_{\bullet\bullet}(\phi > u) + \lambda\mu_{\bullet\bullet}(\phi > v) \subset \mu_{\bullet\bullet}(\phi > (1 - \lambda)u + \lambda v).$$

By the $(-1/r)$ -concavity of μ , the function $f(t) = \mu_{\bullet\bullet}(\phi > t)$ is $(-1/r)$ -concave and it is clearly continuous on \mathbb{R}_+ . Observe by Fubini that for any $p > 0$,

$$\int_{\mathbb{R}^n} \phi(x)^p d\mu(x) = \int_0^{+\infty} pt^{p-1} f(t) dt.$$

The result follows from the part (2) of Theorem 7. The moreover part follows from the log-concavity since then $p \rightarrow (H_f(p)/f(0))^{1/p}$ is a non-increasing function. \square

The second corollary concerns the function $h_{k,p}$ defined in (4).

Corollary 9. Let $r > 0$ and $u \in SO(n)$. For any $(-1/(r+n))$ -concave function $w : \mathbb{R}^n \rightarrow \mathbb{R}_+$ and any subspace E_0 of dimension $k \leq n$, the function

$$p \rightarrow \begin{cases} \frac{h_{k,p}(u)}{B(p+k, r-p)} & \text{for } p > -k+1 \\ S^{k-1} \pi_{u(E_0)} w(0) & \text{for } p = -k+1 \end{cases}$$

is log-concave on $[-k+1, r)$.

Proof. Since w is $(-1/(r+n))$ -concave, we note that $t \rightarrow \pi_{U(E_0)}w(tu(\theta_0))$ is $(-1/(r+k))$ -concave and it is clearly continuous on \mathbb{R}_+ . Theorem 7 proves the result. \square

We finish with some geometric properties of a family of bodies introduced by K. Ball in [5] in the log-concave case.

Corollary 10. *Let $\alpha > 0$. Let $w : \mathbb{R}^n \rightarrow \mathbb{R}_+$ be a $(-1/\alpha)$ -concave function such that $w(0) > 0$. For $0 < a < \alpha$ let*

$$K_a(w) = \left\{ x \in \mathbb{R}^n; a \int_0^{+\infty} t^{a-1} w(tx) dt \geq w(0) \right\}.$$

Then for any $0 < a \leq b < \alpha$

$$\left(\frac{w(0)}{w_\infty} \right)^{\frac{1}{a}-\frac{1}{b}} K_a(w) \subset K_b(w) \subset \frac{(bB(b, \alpha - b))^{1/b}}{(aB(a, \alpha - a))^{1/a}} K_a(w).$$

Proof. Notice that the sets K_a are star-shaped with respect to the origin, that is for every $x \in K_a$ and every $\lambda \in [0, 1]$, $\lambda x \in K_a$. The radial function of K_a is

$$\rho_{K_a}(x) := \sup_{r \geq 0} r : rx \in K_a = \left(a \int_0^{+\infty} t^{a-1} \frac{w(tx)}{w(0)} dt \right)^{\frac{1}{a}}.$$

For any $x \in \mathbb{R}^n$, let f be the continuous $(-1/\alpha)$ -concave function defined on \mathbb{R}^+ by $f(t) = w(tx)/w(0)$. By (1) of Theorem 7, the function $a \rightarrow \left(\int_0^{+\infty} t^{a-1} \frac{f(t)}{\|f\|_\infty} dt \right)^{\frac{1}{a}}$ is non-decreasing. The left hand side inclusion follows. Moreover, from (2) of Theorem 7, the function $H_f : [0, \alpha) \rightarrow \mathbb{R}_+$ is log-concave on $[0, \alpha)$ with $H_f(0) = 1$. For $0 < a \leq b < \alpha$, we have thus $H_f(b)^{1/b} \leq H_f(a)^{1/a}$. The right hand side inclusion follows. \square

3 Thin shell for convex measures

The purpose of this section is to prove Theorem 1. We follow the strategy of the log-concave case initiated in [22, 17, 23] and further developed in [16, 19]. The support function h_K of a non-empty compact set $K \subset \mathbb{R}^n$ is defined by

$$h_K(\theta) = \sup_{x \in K} x, \theta \quad , \quad \theta \in \mathbb{R}^n.$$

To any random vector X in \mathbb{R}^n and any $p \geq 1$, we associate its Z_p^+ -body defined by its support function

$$h_{Z_p^+(X)}(\theta) = (\mathbb{E} \|X, \theta\|_+^p)^{1/p}.$$

When the distribution of X has a density g , we write $Z_p^+(g) = Z_p^+(X)$. Extending a theorem of Ball [5] for log-concave functions, Bobkov proved in [8, Remark 2.6] (see also [15, Theorem 3.1]) that if w is $(-1/(r+n))$ -concave on \mathbb{R}^n such that $w(0) > 0$, then

$$K_a(w) \text{ is convex and compact for any } 0 < a \leq r + n - 1. \quad (8)$$

In the case of log-concave measures [26, 27, 19, 20], several relations between the Z_p^+ bodies and the convex sets K_a are known. We need their analogue in the setting of s -concave measures for negative s . We start with two technical lemmas. We postpone their proofs to the appendix.

Lemma 11. *Let $x, y \geq 1$, then*

$$c \frac{x}{x+y} \leq (xB(x, y))^{1/x} \leq C \frac{x}{x+y}, \quad (9)$$

where c, C are positive universal constants. Moreover, for $k, r > 1$, the extension by continuity at 0 of the function $p \rightarrow \frac{1}{p} \log \frac{B(k+p, r-p)}{B(k, r)}$ is differentiable on $[-(\frac{k-1}{2}), \frac{r-1}{2}]$ and satisfies

$$0 \leq \frac{d}{dp} \left(\frac{1}{p} \log \frac{B(k+p, r-p)}{B(k, r)} \right) \leq \frac{1}{r-1} + \frac{1}{k-1} \quad (10)$$

for $p \in [-(\frac{k-1}{2}), \frac{r-1}{2}]$.

In this paper, we use the notion of geometric distance between sets, defined for every compact subsets $K, L \subset \mathbb{R}^n$ containing 0 in their interior by

$$d(K, L) = \inf_{\bullet\bullet} t_2/t_1 : t_1 L \subset K \subset t_2 L, t_1, t_2 > 0.$$

Let $n \geq 1$, $r \geq 2$ and w be the $(-1/(r+n))$ -concave density of a probability measure μ on \mathbb{R}^n . Then by Corollary 8 and Lemma 11, for $1 \leq p \leq q \leq r-1$, one has

$$Z_p^+(w) \subset Z_q^+(w) \subset c \frac{q}{p} \left(\inf_{\theta \in S^{n-1}} \mu(\bullet\bullet x : x, \theta > 0) \right)^{\frac{1}{q} - \frac{1}{p}} Z_p^+(w).$$

Fix $\theta \in S^{n-1}$ and define $F(t) = \mu_{\bullet\bullet}(x : x, \theta \leq t)$, for $t \in \mathbb{R}$. One has $\int_{\mathbb{R}} tF'(t)dt = \int_{\mathbb{R}^n} x, \theta w(x)dx = 0$ and F is $(-1/r)$ -concave. Using Jensen's inequality, we get

$$F(0)^{-\frac{1}{r}} = F \left(\int_{\mathbb{R}} tF'(t)dt \right)^{-\frac{1}{r}} \leq \int_{\mathbb{R}} F(t)^{-\frac{1}{r}} F'(t)dt = \left[\frac{F(t)^{1-\frac{1}{r}}}{1-\frac{1}{r}} \right]_{-\infty}^{+\infty} = \frac{1}{1-\frac{1}{r}}.$$

Hence $\mu_{\bullet\bullet}(x : x, \theta > 0) \geq (1 - \frac{1}{r})^r \geq 1/4$ for $r \geq 2$. We have recovered here in a simple way a Grünbaum's type inequality for convex measures due to Bobkov [8, Theorem 5.2]. We deduce that, for $1 \leq p \leq q \leq r - 1$,

$$Z_p^+(w) \subset Z_q^+(w) \subset C \frac{q}{p} Z_p^+(w) \quad \text{and} \quad d(Z_p^+(w), Z_q^+(w)) \leq C \frac{q}{p}. \quad (11)$$

Lemma 12. *Let r, m and p be such that m is a positive integer, $r \geq m + 1$ and $-\frac{m}{2} \leq p \leq r - 1$. Let F be a subspace of \mathbb{R}^n of dimension m and let g be a $(-1/(r + m))$ -concave density of a probability measure on F such that $\int_F xg(x)dx = 0$. Then we have*

$$d(K_{m+p}(g), Z_{\max(m,p)}^+(g)) \leq c,$$

where c is a universal constant.

As in [19], an important ingredient in the proof of the thin-shell concentration inequality is an estimate from above of the log-Lipschitz constant of the map on $SO(n) : u \rightarrow h_{k,p}(u)$. Let $\mathcal{M}_n(\mathbb{R})$ be the set of square $n \times n$ matrices. We equip

$$SO(n) =_{\bullet\bullet} u \in \mathcal{M}_n(\mathbb{R}) : u^t u = Id, \det(u) = 1$$

with its standard invariant Riemannian metric, which we specify for concreteness on $T_{Id}SO(n)$, the tangent space at the identity element $Id \in SO(n)$. Since $u^t u = Id$, this tangent space may be identified with the set of anti-symmetric matrices $_{\bullet\bullet} B \in \mathcal{M}_n(\mathbb{R}) : B^t + B = 0$. We define the scalar product $\langle B, B \rangle = \frac{1}{2} \text{tr}(B^t B)$ on $T_{Id}SO(n)$.

Proposition 13. *Let $n \geq 1$, $r > 10$ and w be the $(-1/(r + n))$ -concave density of a probability measure on \mathbb{R}^n such that $\int_{\mathbb{R}^n} xw(x)dx = 0$. Let k be an integer such that $k \geq 2$, $2k - 1 \leq n$ and $2k \leq r$. Let p such that*

$-\frac{k}{2} \leq p \leq r - 1$. Denote by $L_{k,p}$ the log-Lipschitz constant of the map on $SO(n) : u \rightarrow h_{k,p}(u)$. Then

$$L_{k,p} \leq C \max(k, p) d(Z_{\max(k,p)}^+(w), B_2^n),$$

where C is a universal constant.

Proof. For any subspace F of dimension m , the marginal $\pi_F(w)$ is a $(-1/(r+m))$ -concave function on F and from (8), for any $a \in [0, r+m-1]$, we associate the convex body $K_a(\pi_F(w))$ in F . Then the proof of Theorem 2.1 in [19, section 2.2] gives the upper bound:

$$L_{k,p} \leq \max_F (m+p) d(K_{m+p}(\pi_F(w)), B_2(F))$$

over all subspaces F of dimension $m = k, k+1, 2k-1$, where $B_2(F)$ is the Euclidean unit ball in F . By assumptions on k , we get that for these values of m , $m \leq 2k-1 \leq n$ and $r \geq 2k \geq m+1$ and $p \geq -k/2 \geq -m/2$. Hence from Lemma 12, we have

$$d(K_{m+p}(\pi_F(w)), B_2(F)) \leq c d(Z_{\max(m,p)}^+(\pi_F(w)), B_2(F)).$$

By definition, if X is the random vector with density w on \mathbb{R}^n , the marginal $\pi_F(w)$ is the density of the projection of X onto F , namely $P_F X$. By identification of the support functions, we have that, for any $\theta \in F$,

$$h_{Z_p^+(\pi_F(w))}^p(\theta) = \mathbb{E} P_F X, \theta \Big|_+^p = \mathbb{E} X, \theta \Big|_+^p.$$

This means that $Z_p^+(\pi_F(w)) = P_F(Z_p^+(w))$. Since the distance to the Euclidean ball cannot increase after projections, we conclude that

$$d(K_{m+p}(\pi_F(w)), B_2(F)) \leq c d(Z_{\max(m,p)}^+(w), B_2^n).$$

By equation (11), for $m = k, k+1, 2k-1$, one has

$$d(Z_{\max(m,p)}^+(w), Z_{\max(k,p)}^+(w)) \leq c.$$

This finishes the proof. \square

We define the q -condition number of a random vector X to be

$$\rho_q(X) = \frac{\sup_{|\theta|_2=1} (\mathbb{E} X, \theta \Big|_+^q)^{1/q}}{\inf_{|\theta|_2=1} (\mathbb{E} X, \theta \Big|_+^q)^{1/q}}.$$

Obviously, if w is the density of a full-dimensional random vector X in \mathbb{R}^n then $\rho_q(X) = d(Z_q^+(w), B_2^n)$.

Proposition 14. *With the same assumptions as in Proposition 13, if a random vector X with density w is isotropic then*

$$L_{k,p} \leq C \max(k, p)^2.$$

More generally if A is such that AX is isotropic then

$$L_{k,p} \leq C \max(k, p)^2 \|A\| \|A^{-1}\|. \quad (12)$$

Proof. Let $q = \max(k, p)$, then one has $1 \leq q \leq r - 1$. Using the triangular inequality we get

$$\rho_q(X) = d(Z_q^+(w), B_2^n) \leq d(Z_q^+(w), Z_2^+(w)) d(Z_2^+(w), B_2^n).$$

From equation (11) we deduce that $d(Z_q^+(w), Z_2^+(w)) \leq cq$. For any $\theta \in S^{n-1}$, $\mathbb{E} X, \theta = 0$, hence $\mathbb{E} X, \theta_+ = \mathbb{E} -X, \theta_+$. Using this equality and equation (11) we deduce that

$$(\mathbb{E} -X, \theta_+^2)^{\frac{1}{2}} \leq c \mathbb{E} -X, \theta_+ = c \mathbb{E} X, \theta_+ \leq c (\mathbb{E} X, \theta_+^2)^{\frac{1}{2}}.$$

Thus

$$\mathbb{E} X, \theta_+^2 \leq \mathbb{E} X, \theta^2 = \mathbb{E} X, \theta_+^2 + \mathbb{E} -X, \theta_+^2 \leq C \mathbb{E} X, \theta_+^2.$$

Hence if X is isotropic we deduce that $d(Z_2^+(w), B_2^n) \leq c'$. We conclude that

$$\rho_q(X) = d(Z_q^+(w), B_2^n) \leq C'q.$$

The conclusion follows from Proposition 13. In the general case, notice that $Z_q^+(AX) = AZ_q^+(X)$ and $d(AB_2^n, B_2^n) = \|A\| \|A^{-1}\|$, thus

$$\rho_q(X) \leq \rho_q(AX) \|A\| \|A^{-1}\|.$$

□

Proof of Theorem 1. Without loss of generality, we can assume that $r > 32$. Indeed, if $r \leq 32$ then the statement in Theorem 1 is valid for $p \leq cr$ and it gives only a comparison of $(\mathbb{E} X_2^p)^{1/p}$ with $(\mathbb{E} X_2^2)^{1/2}$ up to a constant factor. The result is a consequence of Theorem 5.2 in [1].

From now on, we assume that $r > 32$ and that $p \leq \frac{r}{8}$. We start by presenting a complete argument following [16]. This will give a complete

proof of Theorem 1 with a slightly weaker result. In the second part, we just indicate the needed modifications of the argument of [19] to get the complete conclusion.

In this first part, we will prove that for any $p \in [\frac{1}{\sqrt{n}}, \min(cn^{1/8}, \frac{r}{8})]$

$$(\mathbb{E} X_2^p \mathbb{E} X_2^{-p})^{1/p} \leq 1 + \frac{Cp}{r} + \left(\frac{Cp}{n^{1/3}}\right)^{3/5}. \quad (13)$$

Assuming (13), few elementary steps are needed to prove that for any p such that $p \leq \min(cn^{1/8}, \frac{r}{8})$,

$$\left| \frac{(\mathbb{E} X_2^p)^{1/p}}{(\mathbb{E} X_2^2)^{1/2}} - 1 \right| \leq \frac{C(1+p)}{r} + \left(\frac{C(1+p)}{n^{1/3}}\right)^{3/5}, \quad (14)$$

which is already enough to get a thin-shell concentration. Indeed, for $p \geq 2$, by Hölder inequality, we have

$$0 \leq \frac{(\mathbb{E} X_2^p)^{1/p}}{(\mathbb{E} X_2^2)^{1/2}} - 1 \leq \frac{(\mathbb{E} X_2^p)^{1/p}}{(\mathbb{E} X_2^{-p})^{-1/p}} - 1$$

and we conclude by (13). For $p \leq -2$, we have $p = -p \geq 2$ and from Hölder inequality and (13),

$$0 \leq \frac{(\mathbb{E} X_2^2)^{1/2}}{(\mathbb{E} X_2^p)^{1/p}} - 1 \leq \frac{(\mathbb{E} X_2^{|p|})^{1/|p|}}{(\mathbb{E} X_2^{-|p|})^{-1/|p|}} - 1 \leq \frac{Cp}{r} + \left(\frac{Cp}{n^{1/3}}\right)^{3/5}.$$

An elementary computation shows that

$$\left| \frac{(\mathbb{E} X_2^p)^{1/p}}{(\mathbb{E} X_2^2)^{1/2}} - 1 \right| \leq \frac{Cp}{r} + \left(\frac{Cp}{n^{1/3}}\right)^{3/5}.$$

For $p \in [-2, 2]$, by Hölder inequality, we have

$$0 \leq 1 - \frac{(\mathbb{E} X_2^p)^{1/p}}{(\mathbb{E} X_2^2)^{1/2}} \leq 1 - \frac{(\mathbb{E} X_2^{-2})^{-1/2}}{(\mathbb{E} X_2^2)^{1/2}}$$

and we conclude by the previous estimate for $p = -2$. This concludes the proof of (14).

Let us start the proof of (13). Let $p \in [\frac{1}{\sqrt{n}}, \min(cn^{1/8}, \frac{r}{8})]$ and k be an integer greater or equal than 2 such that $p < k \leq n$. We will optimize the choice of k at the end of the proof. Recall that by (5),

$$\mathbb{E} X_{\frac{p}{2}} = \frac{\Gamma((p+n)/2)\Gamma(k/2)}{\Gamma(n/2)\Gamma((p+k)/2)} \mathbb{E} h_{k,p}(U),$$

where U is uniformly distributed on $SO(n)$. Using that the function $\frac{d}{dp} \log \Gamma(p)$ is concave (see for example the proof of Lemma 11 in the appendix), we deduce that

$$\frac{d}{dp} \left(\frac{1}{p} \log \frac{\Gamma((p+n)/2)\Gamma(k/2)}{\Gamma((p+k)/2)\Gamma(n/2)} \right) \leq 0. \quad (15)$$

It follows that for any $0 < p < k$,

$$\frac{\Gamma((p+n)/2)\Gamma(k/2)}{\Gamma(n/2)\Gamma((p+k)/2)} \frac{\Gamma((-p+n)/2)\Gamma(k/2)}{\Gamma(n/2)\Gamma((-p+k)/2)} \leq 1.$$

Then for all $0 < p < r$ and $n \geq k > p$ we have

$$\mathbb{E} X_{\frac{p}{2}} \mathbb{E} X_{\frac{-p}{2}} \leq \mathbb{E} h_{k,p}(U) \mathbb{E} h_{k,-p}(U). \quad (16)$$

Applying log-Sobolev inequality (3) to $h_{k,p}$ and $h_{k,-p}$ we get

$$\mathbb{E} h_{k,p}(U)^2 \leq e^{\frac{cL_{k,p}^2}{n}} (\mathbb{E} h_{k,p}(U))^2, \quad \mathbb{E} h_{k,-p}(U)^2 \leq e^{\frac{cL_{k,-p}^2}{n}} (\mathbb{E} h_{k,-p}(U))^2. \quad (17)$$

Since $\text{Var} f = \mathbb{E} f^2 - (\mathbb{E} f)^2$ we deduce that

$$\begin{cases} \text{Var} h_{k,p}(U) \leq \left(e^{\frac{cL_{k,p}^2}{n}} - 1 \right) (\mathbb{E} h_{k,p}(U))^2, \\ \text{Var} h_{k,-p}(U) \leq \left(e^{\frac{cL_{k,-p}^2}{n}} - 1 \right) (\mathbb{E} h_{k,-p}(U))^2. \end{cases} \quad (18)$$

By Corollary 9, we know that $p \rightarrow h_{k,p}(u)/B(k+p, r-p)$ is log-concave on $[-k+1, r)$ hence

$$h_{k,p}(u) h_{k,-p}(u) \leq \left(\frac{B(k+p, r-p) B(k-p, r+p)}{B(k, r)} \right) h_{k,0}^2(u).$$

Taking the expectation with respect to $SO(n)$, we get that

$$\mathbb{E} h_{k,p}(U) h_{k,-p}(U) \leq \left(\frac{B(k+p, r-p) B(k-p, r+p)}{B(k, r)} \right) \mathbb{E} h_{k,0}^2(U).$$

Since $\mathbb{E}h_{k,0}(U) = 1$ we deduce from (17) that

$$\mathbb{E}h_{k,0}^2(U) \leq e^{\frac{cL_{k,0}^2}{n}}.$$

Assume that k is such that $k \leq r$ then by (10), we know that for $p \leq (k-1)/2$,

$$\left(\frac{B(k+p, r-p) B(k-p, r+p)}{B(k, r)} \right)^{1/p} \leq e^{2p(\frac{1}{k-1} + \frac{1}{r-1})} \leq e^{4p(\frac{1}{k} + \frac{1}{r})}$$

since $k, r \geq 2$. Hence

$$\mathbb{E}h_{k,p}(U)h_{k,-p}(U) \leq e^{\frac{cL_{k,0}^2}{n} + 4p^2(\frac{1}{k} + \frac{1}{r})}. \quad (19)$$

Moreover

$$\begin{aligned} \mathbb{E}h_{k,p}(U)h_{k,-p}(U) &= \mathbb{E}h_{k,p}(U)\mathbb{E}h_{k,-p}(U) + \text{Cov}(h_{k,p}(U), h_{k,-p}(U)) \\ &\geq \mathbb{E}h_{k,p}(U)\mathbb{E}h_{k,-p}(U) - \sqrt{\text{Var}h_{k,p}(U)\text{Var}h_{k,-p}(U)} \\ &\geq \mathbb{E}h_{k,p}(U)\mathbb{E}h_{k,-p}(U) \left(1 - \sqrt{\left(e^{\frac{cL_{k,p}^2}{n}} - 1 \right) \left(e^{\frac{cL_{k,-p}^2}{n}} - 1 \right)} \right) \end{aligned} \quad (20)$$

where the last inequality follows from (18). Assume moreover that k is such that $2k-1 \leq n$ and $2k \leq r$ then for $p \leq (k-1)/2$, we can evaluate $L_{k,p}$, $L_{k,-p}$ and $L_{k,0}$ from Proposition 14 since the assumptions are fulfilled. We get that if X is isotropic then $\max(L_{k,p}, L_{k,-p}, L_{k,0}) \leq Ck^2$. If $k \leq c_0n^{1/4}$ for a small enough numerical constant c_0 , we have

$$\sqrt{\left(e^{\frac{cL_{k,p}^2}{n}} - 1 \right) \left(e^{\frac{cL_{k,-p}^2}{n}} - 1 \right)} \leq c' \frac{k^4}{n} \leq \frac{1}{10}.$$

Combining this estimate with (20) and (19), we have proved that if k is an integer such that $k \geq 2$, $2k-1 \leq n$, $2k \leq r$, $k \leq c_0n^{1/4}$ and $2p+1 \leq k$ (this set of integers is not empty since $r > 32$ and $p \leq r/8$) then

$$\mathbb{E}h_{k,p}(U)\mathbb{E}h_{k,-p}(U) \leq \frac{e^{4p^2(\frac{1}{k} + \frac{1}{r}) + c\frac{k^4}{n}}}{1 - c' \frac{k^4}{n}} \leq e^{4p^2(\frac{1}{k} + \frac{1}{r}) + C\frac{k^4}{n}}.$$

For $p \leq 1$, we also force k to satisfy $k \leq C_0 p^{1/4} n^{1/4}$. Hence taking the power $1/p$ in the last expression, we conclude from (16) that

$$(\mathbb{E} X_2^p \mathbb{E} X_2^{-p})^{1/p} \leq e^{4p(\frac{1}{k} + \frac{1}{r}) + C \frac{k^4}{pn}} \leq 1 + cp \left(\frac{1}{k} + \frac{1}{r} \right) + c \frac{k^4}{pn},$$

since $p/k, p/r$ and k^4/pn are bounded by universal constants. It remains to optimize the choice of k . Let $p_0 = n^{-1/2}$. In this case we choose $k = 2$ and get

$$(\mathbb{E} X_2^{p_0} \mathbb{E} X_2^{-p_0})^{1/p_0} \leq 1 + \frac{C}{\sqrt{n}}. \quad (21)$$

If $p \geq n^{-1/2}$ we choose k to be an integer such that $\min(r/4, (p^2 n)^{1/5}) \leq k \leq 2 \min(r/4, (p^2 n)^{1/5})$ with the restriction $2p + 1 \leq k \leq cn^{1/4}$ and that $k \leq cp^{1/4} n^{1/4}$. For any p such that $p_0 \leq p \leq \min(cn^{1/8}, r/8)$, the integer k satisfies $k \geq 2$, $2k - 1 \leq n$, $2k \leq r$, $k \leq c_0 n^{1/4}$ and $2p + 1 \leq k$ and we get that

$$(\mathbb{E} X_2^p \mathbb{E} X_2^{-p})^{1/p} \leq 1 + \frac{Cp}{r} + \left(\frac{Cp}{n^{1/3}} \right)^{3/5}.$$

This ends the proof of (13).

In the second part, we follow the argument developed in [19] to get a better estimate. We deal now with the case of p being positive or negative and, as already said, we can assume without loss of generality that $r > 34$ and $p \leq r/8$. As in [19], our goal is to estimate

$$\frac{d}{dp} \log((\mathbb{E} X_2^p)^{\frac{1}{p}}) = \frac{d}{dp} \log((\mathbb{E} h_{k,p}(U))^{\frac{1}{p}}) + \frac{d}{dp} \left(\frac{1}{p} \log \frac{\Gamma((p+n)/2) \Gamma(k/2)}{\Gamma(n/2) \Gamma((p+k)/2)} \right).$$

Most of the computation of section 3.2 in [19] is identical. All the ingredients needed for the proof have been established and, adapting the argument done in section 3.2 in [19], we get

$$\frac{d}{dp} \log((\mathbb{E} X_2^p)^{\frac{1}{p}}) \leq \frac{c}{p^2 n} (2L_{k,p}^2 + 3L_{k,0}^2) + \frac{C}{k-1} + \frac{C}{r-1}. \quad (22)$$

For convenience of the reader, we will shortly reproduce the proof of (22) in the appendix.

Assume that X is isotropic. For any $2p \leq k \leq r/2$ (this set of integers is not empty since $r > 32$ and $p \leq r/8$), we know by Proposition 14, that

$L_{k,p}$ and $L_{k,0}$ are smaller than Ck^2 . We get that

$$\frac{d}{dp} \log((\mathbb{E} X \frac{p}{2})^{\frac{1}{p}}) \leq C \left(\frac{k^4}{p^2 n} + \frac{1}{k} + \frac{1}{r} \right).$$

We have to minimize this expression for k being an integer greater or equal than 2 and $k \in [2p, r/2]$. For $p \in [n^{-1/2}, cn^{1/3}]$, we set k being an integer such that $\min(r/4, 2(p^2 n)^{1/5}) \leq k \leq 2 \min(r/4, 2(p^2 n)^{1/5})$. Therefore k satisfies the restrictions and we get for any p such that $p \in [n^{-1/2}, cn^{1/3}]$,

$$\frac{d}{dp} \log((\mathbb{E} X \frac{p}{2})^{\frac{1}{p}}) \leq C \left(\frac{1}{(p^2 n)^{1/5}} + \frac{1}{r} \right). \quad (23)$$

After integration over p , we get that for all $p \in [n^{-1/2}, c \min(r, n^{1/3})]$

$$\left| \log \frac{(\mathbb{E} X \frac{p}{2})^{1/p}}{(\mathbb{E} X \frac{2}{2})^{1/2}} \right| \leq \frac{C p - 2}{r} + \frac{C p^{3/5} - 2^{3/5}}{n^{1/5}}.$$

Since $p^{3/5} - 2^{3/5} \leq p - 2^{3/5}$ and all terms in the right hand side of the inequality are bounded by a universal constant, we conclude by adjusting that

$$\left| \frac{(\mathbb{E} X \frac{p}{2})^{1/p}}{(\mathbb{E} X \frac{2}{2})^{1/2}} - 1 \right| \leq \frac{C p - 2}{r} + \left(\frac{C p - 2}{n^{1/3}} \right)^{3/5}, \quad p \in [n^{-1/2}, c \min(r, n^{1/3})].$$

Since (23) holds only for $p \geq n^{-1/2}$, we use (21) to bridge the gap between $-n^{-1/2}$ and $n^{-1/2}$. Indeed, from (21), the previous inequality for $p_0 = n^{-1/2}$ and using that $p_0 - 2 = 2 - p_0 \leq 2$, we get that for $p \in [-p_0, p_0]$,

$$\begin{aligned} (\mathbb{E} X \frac{p}{2})^{1/p} &\geq (\mathbb{E} X \frac{-p_0}{2})^{-1/p_0} \geq \frac{1}{1 + \frac{C}{\sqrt{n}}} (\mathbb{E} X \frac{p_0}{2})^{1/p_0} \\ &\geq \frac{1 - \frac{2C}{r} - (\frac{2C}{n^{1/3}})^{3/5}}{1 + \frac{C}{n^{1/5}}} (\mathbb{E} X \frac{2}{2})^{1/2}. \end{aligned}$$

An easy adaptation of the constants leads to the conclusion of Theorem 1 for all $p \in [-n^{-1/2}, n^{-1/2}]$.

Integrating again (23), we get, for $p \in [-c \min(r, n^{1/3}), -n^{-1/2}]$,

$$\frac{(\mathbb{E} X \frac{p}{2})^{1/p}}{(\mathbb{E} X \frac{-p_0}{2})^{-1/p_0}} \geq 1 - \frac{C p + p_0}{r} - \left(\frac{C p + p_0}{n^{1/3}} \right)^{3/5}.$$

Using that $p + p_0 \leq p - 2$ and the previous comparison of the moment of order $-p_0$ with the moment of order 2 and adjusting the constants, this proves that for all $p \in [-c \min(r, n^{1/3}), -n^{-1/2}]$,

$$\left| \frac{(\mathbb{E} X^p)^{1/p}}{(\mathbb{E} X^2)^{1/2}} - 1 \right| \leq \frac{C(p-2)}{r} + \left(\frac{C(p-2)}{n^{1/3}} \right)^{3/5}.$$

This concludes the proof of the first part of Theorem 1.

If X is such that AX is isotropic, we know from Proposition 14 that for any integer k such that $2p \leq k \leq r/2$,

$$\max(L_{k,p}, L_{k,0}) \leq Ck^2 \|A\| \|A^{-1}\|.$$

The proof is identical to the previous one replacing n by $\frac{n}{\|A\|^2 \|A^{-1}\|^2}$. \square

Remark 15. In [19], a preprocessing step consisted in adding a Gaussian isotropic vector to the random vector X in order to start at the very beginning with a better information on the Z_p^+ bodies associated to the measure. In [23, 16], this convolution argument played a role of regularization. It is natural to ask if such a process could be done in the situation of s -concave measure. Nothing is doable by adding a Gaussian vector because for $s < 0$, the new vector does not belong to any class of s -concave vectors. However, for $r > n$, we can build a similar argument, adding to X a random vector Z uniformly distributed on the Euclidean ball, see also [9]. Since Z is $(1/n)$ -concave and X is $(-1/r)$ -concave, the new vector $Y = \frac{X+Z}{\sqrt{2}}$ will be $(-1/(r-n))$ -concave. For any $p \geq 1$, we have (see inequality (4.7) in [19])

$$\alpha_p(X) \leq \alpha_{2p}(Y) (2 + \alpha_{2p}(Y))$$

so that it remains to bound $\alpha_{2p}(Y)$. It is easy to see that Y is such that for every $q \geq 2$ and every $\theta \in S^{n-1}$, $(\mathbb{E} Y, \theta^q)^{1/q} \geq c\sqrt{q}$. Adapting the proof of Proposition 14, we get $L_{k,p} \leq C \max(k, p)^{3/2}$. As in [19], this improvement leads to the following estimate: if $r - n > 2$, then for any p such that $1 \leq p \leq c \min(r - n, \sqrt{n})$

$$\alpha_{2p}(Y) \leq \frac{C(2p-2)}{r-n} + \left(\frac{C(2p-2)}{\sqrt{n}} \right)^{1/2}.$$

For $r > n + \sqrt{n}$, we recover the same thin-shell concentration as in the log-concave case. It would be interesting to understand in which precise

sense the s -concave measures are close to the log-concave measures for $s \in (-1/n, 1/n)$. Another question is to know what kind of preprocessing argument like in [24] would enable to recover the small ball estimates from [1].

4 Appendix

Proof of Theorem 7. (1) This result is classical. In the symmetric case, it follows from Lemma 2.1 in [25]. The general case is similar. We provide its proof for completeness. We may assume, without loss of generality, that $f|_{\infty} = 1$. Denote $I_p(f) = \int_0^{+\infty} t^{p-1} f(t) dt$. From Hölder inequality, the function $p \rightarrow \log(I_p(f))$ is convex on its convex support, thus the domain of definition of $I_p(f)$ is an interval. Let $0 < p < q$ be fixed such that $I_p(f) < +\infty$ and $I_q(f) < +\infty$. Let $a = (pI_p(f))^{1/p}$ and $\varphi(t) = t^{p-1}(f(t) - 1_{[0,a]}(t))$. Notice that $\varphi \leq 0$ on $[0, a]$, $\varphi \geq 0$ on $[a, +\infty)$ and $\int_0^{+\infty} \varphi(t) dt = 0$. Thus

$$I_q(f) - I_q(1_{[0,a]}) = \int_0^{+\infty} t^{q-p} \varphi(t) dt = \int_0^{+\infty} (t^{q-p} - a^{q-p}) \varphi(t) dt \geq 0,$$

since the integrand is non negative on \mathbb{R}_+ . We conclude that

$$I_q(f) \geq I_q(1_{[0,a]}) = \frac{a^q}{q} = \frac{1}{q} (pI_p(f))^{\frac{q}{p}}.$$

(2) Since f is $(-1/\alpha)$ -concave, there exists a convex function $\varphi : [0, +\infty) \rightarrow (0, +\infty)$ such that $f = \varphi^{-\alpha}$. Since f is integrable it follows that φ tends to $+\infty$ at $+\infty$. From the convexity of φ , one deduces that for some constant $c > 0$, $\varphi(t) \geq c(1+t)$. Thus $f(t) \leq (c+ct)^{-\alpha}$, for every $t \geq 0$. Therefore, $t^{p-1}f$ is integrable for every $p < \alpha$, which means that $H_f(p) < +\infty$ for every $0 < p < \alpha$. Let $p \in (0, \alpha)$ and $m, M > 0$. Define $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ by $g(t) = m \left(1 + \frac{t}{M}\right)^{-\alpha}$. Then

$$\int_0^{+\infty} t^{p-1} g(t) dt = mM^p \int_0^{+\infty} v^{p-1} (1+v)^{-\alpha} dv = mM^p B(p, \alpha - p).$$

Thus $H_g(p) = mM^p$, which implies that $\log(H_g)$ is affine on $(0, \alpha)$. Take $0 < a < b < c < \alpha$. Let $\lambda \in [0, 1]$ be such that $b = (1-\lambda)a + \lambda c$. Choose m and M such that $mM^a = H_f(a)$ and $mM^b = H_f(b)$ so that $H_g(a) = H_f(a)$ and $H_g(b) = H_f(b)$. If we prove that

$$\int_0^{+\infty} t^{c-1} (g - f)(t) dt \geq 0, \tag{24}$$

that is $H_g(c) \geq H_f(c)$, then using that $\log(H_g)$ is affine, we will deduce that

$$H_f(b) = H_g(b) = H_g(a)^{1-\lambda} H_g(c)^\lambda \geq H_f(a)^{1-\lambda} H_f(c)^\lambda$$

and this will prove the log-concavity of H on $(0, \alpha)$. If $f = g$ then (24) is satisfied so that in the following we assume that the function $h := g - f \equiv 0$. Let

$$H_1(t) = \int_t^{+\infty} s^{a-1} h(s) ds \quad \text{and} \quad H_2(t) = \int_t^{+\infty} s^{b-a-1} H_1(s) ds.$$

Since $h(t) = O(t^{-\alpha})$ at infinity, we deduce that $H_1(t) = O(t^{a-\alpha})$ and $H_2(t) = O(t^{b-\alpha})$. We have $\int_0^{+\infty} t^{a-1} h(t) dt = 0$ thus $H_1(+\infty) = H_1(0) = 0$. Obviously $H_2(+\infty) = 0$. We also observe

$$\begin{aligned} 0 &= \int_0^{+\infty} t^{b-1} h(t) dt = \int_0^{+\infty} t^{b-a} t^{a-1} h(t) dt = - \int_0^{+\infty} t^{b-a} H_1'(t) dt \\ &= [t^{b-a} H_1(t)]_0^{+\infty} + (b-a) \int_0^{+\infty} t^{b-a-1} H_1(t) dt = (b-a) H_2(0), \end{aligned}$$

whence $H_2(+\infty) = H_2(0) = 0$. Since $\int_0^{+\infty} t^{b-a-1} H_1(t) dt = 0$ and $H_1 \equiv 0$, the function H_1 has at least one change of sign. Moreover, using that $H_1(0) = H_1(+\infty) = 0$, we deduce that H_1' and therefore h has at least two sign changes. Since $h = g - f$ has the same sign as $f^{-\alpha} - g^{-\alpha}$ which is convex, it cannot have more than two sign changes. Thus it has exactly two sign changes at some $0 < t_1 < t_2$. Moreover, from the convexity of $f^{-\alpha} - g^{-\alpha}$, the sign of h has to be negative on (t_1, t_2) and positive on $(0, t_1)$ and $(t_2, +\infty)$. From an easy study of the function H_2 , we deduce that $H_2 \geq 0$. Therefore, using that $H_1(0) = H_1(+\infty) = H_2(0) = H_2(+\infty) = 0$, we get

$$\begin{aligned} \int_0^{+\infty} t^{c-1} h(t) dt &= \int_0^{+\infty} t^{c-a} t^{a-1} h(t) dt = - \int_0^{+\infty} t^{c-a} H_1'(t) dt \\ &= [-t^{c-a} H_1(t)]_0^{+\infty} + (c-a) \int_0^{+\infty} t^{c-a-1} H_1(t) dt \\ &= (c-a) \int_0^{+\infty} t^{c-b} t^{b-a-1} H_1(t) dt \\ &= (c-a) [-t^{c-b} H_2(t)]_0^{+\infty} + (c-a)(c-b) \int_0^{+\infty} t^{c-b-1} H_2(t) dt \\ &= (c-a)(c-b) \int_0^{+\infty} t^{c-b-1} H_2(t) dt \geq 0. \end{aligned}$$

This proves (24) and establish the log-concavity of H_f on $(0, \alpha)$. To get it on $[0, \alpha)$, it is enough to prove that H_f is continuous at 0. This follows from the observation that

$$B(p, \alpha - p) \underset{p \rightarrow 0}{\sim} \Gamma(p) \underset{p \rightarrow 0}{\sim} \frac{1}{p} \quad \text{thus} \quad H_f(p) \underset{p \rightarrow 0}{\sim} p \int_0^{+\infty} t^{p-1} f(t) dt.$$

And it is classical that, for a continuous function f , the right hand side term tends to $f(0)$ when $p \rightarrow 0$. \square

Proof of Lemma 11. Equation (9) follows easily from the classical bounds for the Gamma function (see [3]), valid for $x \geq 1$:

$$\sqrt{2\pi} x^{x-\frac{1}{2}} e^{-x} \leq \Gamma(x) \leq \sqrt{2\pi} x^{x-\frac{1}{2}} e^{-x+\frac{1}{12}}.$$

For equation (10), we write that

$$\frac{B(k+p, r-p)}{B(k, r)} = \frac{\Gamma(k+p)\Gamma(r-p)}{\Gamma(k)\Gamma(r)}.$$

Denote $G(p) = \log \Gamma(p)$, for $p > 0$. We know that $G''(p) = \sum_{i \geq 0} 1/(p+i)^2$ hence G'' is non-increasing and $0 \leq G''(p) \leq 1/(p-1)$, for $p > 1$. Denote $F_k(p) = \frac{G(k+p)-G(k)}{p}$, for $k > 0$ and $p > -k$. We have $F_k(p) = \int_0^1 G'(k+up) du$. Using that G'' is non-increasing, we get that for $k > 1$ and $p \geq -(k-1)/2$,

$$F'_k(p) = \int_0^1 G''(k+up) u du \leq G''\left(\frac{k+1}{2}\right) \int_0^1 u du = \frac{1}{2} G''\left(\frac{k+1}{2}\right) \leq \frac{1}{k-1}$$

and $F'_k(p) \geq 0$. Therefore, if $k > 1$, $r > 1$ and $-\frac{k-1}{2} \leq p \leq \frac{r-1}{2}$ then

$$\begin{aligned} 0 \leq \frac{d}{dp} \left(\frac{1}{p} \log \frac{B(k+p, r-p)}{B(k, r)} \right) &= \frac{d}{dp} (F_k(p) - F_r(-p)) \\ &= F'_k(p) + F'_r(-p) \leq \frac{1}{k-1} + \frac{1}{r-1}. \end{aligned}$$

\square

Proof of Lemma 12. We present here a similar proof than in the appendix of [19]. Applying Corollary 10 to $w = g$, $n = m$, $\alpha = r + m$, we deduce that, for $\frac{m}{2} \leq a \leq b \leq r + m - 1$, one has

$$\left(\frac{g(0)}{g \infty} \right)^{\frac{1}{a}-\frac{1}{b}} K_a(g) \subset K_b(g) \subset \frac{(bB(b, r+m-b))^{1/b}}{(aB(a, r+m-a))^{1/a}} K_a(g).$$

From Lemma 11, we have

$$\frac{(bB(b, r + m - b))^{1/b}}{(aB(a, r + m - a))^{1/a}} \leq c \frac{b}{a}$$

Moreover since $\int xg(x)dx = 0$, from Lemma 7.2 of [1], one has

$$\frac{g(0)}{g_\infty} \geq \left(\frac{r-1}{r+m-1} \right)^{r+m} \geq e^{-2m}.$$

Using that $\frac{1}{a} - \frac{1}{b} \leq \frac{1}{a} \leq \frac{2}{m}$, we deduce that $\left(\frac{g(0)}{\|g\|_\infty} \right)^{\frac{1}{a} - \frac{1}{b}} \geq e^{-4}$. We conclude that for $\frac{m}{2} \leq a \leq b \leq r + m - 1$, one has

$$e^{-4}K_a(g) \subset K_b(g) \subset c \frac{b}{a} K_a(g). \quad (25)$$

By integration in polar coordinates, it is well known [26] (see also [20]) that we have the following relation between the Z_q^+ -bodies associated with g and the Z_q^+ -bodies associated with one of the convex bodies $K_a(g)$: for any $0 < q < r$

$$Z_q^+(g) = g(0)^{1/q} Z_q^+(K_{m+q}(g)), \quad (26)$$

where for any body K , $Z_q^+(K)$ denotes the convex body whose support function is defined by

$$h_{Z_q^+(K)}(\theta) = \left(\int_K x, \theta_+^q dx \right)^{\frac{1}{q}}.$$

Let $\theta \in \mathbb{R}^m$ and K be a convex body containing 0. From Berwald's inequalities [6] applied to K $\bullet\bullet x, \theta \geq 0$ and the function $x \rightarrow x, \theta_+$ which is concave on K $\bullet\bullet x, \theta \geq 0$, the function

$$p \rightarrow \left(\frac{\int_K x, \theta_+^p dx}{mB(p+1, m) \text{Vol}(K \bullet\bullet x, \theta \geq 0)} \right)^{\frac{1}{p}}$$

is decreasing. Observe that for every $\theta \in \mathbb{R}^m$, $\lim_{p \rightarrow \infty} \left(\int_K x, \theta_+^p dx \right)^{\frac{1}{p}} = h_K(\theta)$ and that

$$(mB(p+1, m))^{\frac{1}{p}} = \left(m \int_0^1 u^p (1-u)^{m-1} du \right)^{\frac{1}{p}} \xrightarrow{p \rightarrow +\infty} 1.$$

We deduce that

$$\left(\frac{\int_K x, \theta \binom{q}{+} dx}{mB(q+1, m) \text{Vol}(K \bullet\bullet x, \theta \geq 0)} \right)^{\frac{1}{q}} \geq h_K(\theta).$$

Note also that $\int_K x, \theta \binom{q}{+} dx \leq h_K(\theta)^q \text{Vol}(K \bullet\bullet x, \theta \geq 0)$ and that $mB(q+1, m) = qB(q, m+1)$. Therefore

$$h_K(\theta) \geq \frac{h_{Z_q^+(K)}(\theta)}{\text{Vol}(K \bullet\bullet x, \theta \geq 0)^{1/q}} \geq (qB(q, m+1))^{1/q} h_K(\theta). \quad (27)$$

Now we establish that for $q = \max(p, m)$

$$d(K_{m+q}(g), Z_q^+(g)) \leq c. \quad (28)$$

Take $K = K_{m+q}(g)$. By Lemma 11, for any $q \geq m \geq 1$, $(qB(q, m+1))^{1/q} \geq cq/(m+q+1) \geq c/3$ and we deduce from (27) that for every $\theta \in \mathbb{R}^n$,

$$h_{K_{m+q}(g)}(\theta) \geq \frac{h_{Z_q^+(K_{m+q}(g))}(\theta)}{\text{Vol}(K_{m+q}(g) \bullet\bullet x, \theta \geq 0)^{1/q}} \geq \frac{c}{3} h_K(\theta).$$

where c is a universal constant. Together with (26), we conclude that

$$\begin{aligned} d(K_{m+q}(g), Z_q^+(g)) &= d(K_{m+q}(g), Z_q^+(K_{m+q}(g))) \\ &\leq c \frac{\sup_{\theta \in \mathbb{R}^n} \text{Vol}(K_{m+q}(g) \bullet\bullet x, \theta \geq 0)^{1/q}}{\inf_{\theta \in \mathbb{R}^n} \text{Vol}(K_{m+q}(g) \bullet\bullet x, \theta \geq 0)^{1/q}} \end{aligned} \quad (29)$$

for a universal constant c . Applying (25) for $a = m+1$ and $b = m+q$, we get

$$e^{-4} K_{m+1}(g) \subset K_{m+q}(g) \subset c \frac{m+q}{m+1} K_{m+1}(g).$$

Since $q \geq m$ and $\left(\frac{m+q}{m+1}\right)^{m/q} \leq e$, we get from (29)

$$d(K_{m+q}(g), Z_q^+(g)) \leq C \frac{\sup_{\theta \in \mathbb{R}^n} \text{Vol}(K_{m+1}(g) \bullet\bullet x, \theta \geq 0)^{1/q}}{\inf_{\theta \in \mathbb{R}^n} \text{Vol}(K_{m+1}(g) \bullet\bullet x, \theta \geq 0)^{1/q}}$$

for a universal constant C . Since g has its barycenter at the origin then $K_{m+1}(g)$ has also its barycenter at the origin and we deduce from a classical result of Grünbaum [18] that there exists a universal constant c for which

$$\frac{\sup_{\theta \in \mathbb{R}^n} \text{Vol}(K_{m+1}(g) \bullet\bullet x, \theta \geq 0)^{1/q}}{\inf_{\theta \in \mathbb{R}^n} \text{Vol}(K_{m+1}(g) \bullet\bullet x, \theta \geq 0)^{1/q}} \leq (e-1)^{1/q} \leq e-1.$$

And (28) is proved.

To conclude the proof of the Lemma, it is enough to establish that $d(K_{m+q}, K_{m+p}) \leq c$, where $q = \max(m, p)$. For $q = p$, this is obvious so we may assume that $q = m \geq p$. Then $\frac{m}{2} \leq m + p \leq m + q = 2m$ and using equation (25) for $a = m + p \leq b = 2m$, we deduce that

$$d(K_{m+p}(g), K_{2m}(g)) \leq ce^4 \frac{2m}{m+p} \leq 4ce^4.$$

□

Proof of inequality (22). Our goal is to estimate

$$\frac{d}{dp} \log \left((\mathbb{E} X^{\frac{p}{2}})^{\frac{1}{p}} \right) = \frac{d}{dp} \log \left((\mathbb{E} h_{k,p}(U))^{\frac{1}{p}} \right) + \frac{d}{dp} \left(\frac{1}{p} \log \frac{\Gamma((p+n)/2)\Gamma(k/2)}{\Gamma(n/2)\Gamma((p+k)/2)} \right).$$

As already mentioned in (15), by concavity of $p \rightarrow \frac{d}{dp} \log \Gamma(p)$, we have

$$\frac{d}{dp} \left(\frac{1}{p} \log \frac{\Gamma((p+n)/2)\Gamma(k/2)}{\Gamma(n/2)\Gamma((p+k)/2)} \right) \leq 0.$$

We use the following convention: let (Ω, μ) be a measurable space, for any measurable function $f : \Omega \rightarrow \mathbb{R}^+$, we set

$$\mathbb{E}_\mu(f) = \int f d\mu \quad \text{and} \quad \text{Ent}_\mu(f) = \mathbb{E}_\mu(f \log f) - \mathbb{E}_\mu(f) \log(\mathbb{E}_\mu(f)).$$

Let w be the density of the distribution of X on \mathbb{R}^n . Since X is $(-1/r)$ -concave, w is $(-1/(r+n))$ -concave on \mathbb{R}^n . To any fixed $u \in SO(n)$, we associate the measure μ_u on \mathbb{R}^+ with density

$$t \rightarrow S^{k-1} t^{k-1} \pi_{u(E_0)} w(tu(\theta_0))$$

so that

$$h_{k,p}(u) = S^{k-1} \int_0^\infty t^{p+k-1} \pi_{u(E_0)} w(tu(\theta_0)) dt = \mathbb{E}_{\mu_u}(t^p).$$

Define also $\mu_{k,p}$ the measure on \mathbb{R}^+ with density

$$t \rightarrow S^{k-1} t^{k-1} \mathbb{E} \pi_{U(E_0)} w(tU(\theta_0)).$$

Then $\mathbb{E}h_{k,p}(U) = \mathbb{E}_U \mathbb{E}_{\mu_U}(t^p) = \mathbb{E}_{\mu_{k,p}}(t^p)$. Since w is a density of probability, $\mu_{k,p}$ is a probability measure on \mathbb{R}^+ . A classical fact, verified by direct computation, is that

$$\frac{d}{dp} \log \left((\mathbb{E}_{\mu}(f^p))^{1/p} \right) = \frac{1}{p^2} \frac{\text{Ent}_{\mu}(f^p)}{\mathbb{E}_{\mu}(f^p)}.$$

Therefore

$$\begin{aligned} \frac{d}{dp} \log \left((\mathbb{E}h_{k,p}(U))^{1/p} \right) &= \frac{d}{dp} \log \left((\mathbb{E}_{\mu_{k,p}}(t^p))^{1/p} \right) \\ &= \frac{1}{p^2} \frac{\text{Ent}_{\mu_{k,p}}(t^p)}{\mathbb{E}_{\mu_{k,p}}(t^p)} = \frac{1}{p^2} \frac{\text{Ent}_{\mu_{k,p}}(t^p)}{\mathbb{E}h_{k,p}(U)}. \end{aligned} \quad (30)$$

The numerator can be decomposed into two terms:

$$\text{Ent}_{\mu_{k,p}}(t^p) = \mathbb{E}_U \text{Ent}_{\mu_U}(t^p) + \text{Ent}_U \mathbb{E}_{\mu_U}(t^p) = \mathbb{E}_U \text{Ent}_{\mu_U}(t^p) + \text{Ent}_U h_{k,p}(U).$$

To control the second term, we use the log-Sobolev inequality (2):

$$\frac{1}{p^2} \frac{\text{Ent}_U h_{k,p}(U)}{\mathbb{E}h_{k,p}(U)} \leq \frac{c}{p^2 n} \frac{\mathbb{E} \left(\log h_{k,p}^2(U) h_{k,p}(U) \right)}{\mathbb{E}h_{k,p}(U)} \leq \frac{cL_{k,p}^2}{p^2 n}. \quad (31)$$

To control the first term, we start by observing that for a fixed $u \in SO(n)$,

$$\begin{aligned} \frac{1}{p^2} \frac{\text{Ent}_{\mu_u}(t^p)}{\mathbb{E}_{\mu_u}(t^p)} &= \frac{d}{dp} \log \left((\mathbb{E}_{\mu_u}(f^p))^{1/p} \right) = \frac{d}{dp} \left(\frac{1}{p} \log h_{k,p}(u) \right) \\ &= \frac{d}{dp} \frac{1}{p} \left(\log \frac{h_{k,p}(u)}{B(p+k, r-p)} - \log \frac{h_{k,0}(u)}{B(k, r)} + \log \frac{B(p+k, r-p)}{B(k, r)} + \log h_{k,0}(u) \right). \end{aligned}$$

By Corollary 9, the map $p \rightarrow \frac{h_{k,p}(u)}{B(p+k, r-p)}$ is log-concave on $(-k+1, r)$. This implies that

$$\frac{d}{dp} \frac{1}{p} \left(\log \frac{h_{k,p}(u)}{B(p+k, r-p)} - \log \frac{h_{k,0}(u)}{B(k, r)} \right) \leq 0.$$

We know from Lemma 11 that, for all $p \in [-\frac{k-1}{2}, \frac{r-1}{2}]$,

$$\frac{d}{dp} \left(\frac{1}{p} \log \frac{B(k+p, r-p)}{B(k, r)} \right) \leq C \left(\frac{1}{k-1} + \frac{1}{r-1} \right).$$

Therefore, for any fixed $u \in SO(n)$,

$$\frac{1}{p^2} \text{Ent}_{\mu_u}(t^p) \leq C h_{k,p}(u) \left(\frac{1}{k-1} + \frac{1}{r-1} \right) - \frac{1}{p^2} h_{k,p}(u) \log h_{k,0}(u).$$

Integrating over $u \in SO(n)$, we deduce that

$$\frac{1}{p^2} \frac{\mathbb{E} \text{Ent}_{\mu_U}(t^p)}{\mathbb{E} h_{k,p}(U)} \leq C \left(\frac{1}{k-1} + \frac{1}{r-1} \right) + \frac{1}{p^2} \frac{\mathbb{E} h_{k,p}(U) \log(h_{k,0}(U)^{-1})}{\mathbb{E} h_{k,p}(U)}. \quad (32)$$

From Jensen and Hölder inequalities,

$$\begin{aligned} \frac{\mathbb{E}(h_{k,p}(U) \log h_{k,0}(U)^{-1})}{\mathbb{E} h_{k,p}(U)} &\leq \log \left(\frac{\mathbb{E}(h_{k,p}(U) h_{k,0}(U)^{-1})}{\mathbb{E} h_{k,p}(U)} \right) \\ &\leq \log \left(\frac{(\mathbb{E} h_{k,p}(U)^2)^{1/2}}{\mathbb{E} h_{k,p}(U)} \right) + \log ((\mathbb{E}(h_{k,0}(U)^{-2}))^{1/2}). \end{aligned}$$

From (3), the first term is upper bounded by $\frac{c}{n} L_{k,p}^2$. For the second term, we first use (3) with $f = h_{k,0}^{-1}$, $q = 2$ and $r = 0$, then we use (3) again with $f = h_{k,0}$, $q = 1$ and $r = 0$. Since $\mathbb{E} h_{k,0}(U) = \mathbb{E}_{\mu_{k,0}}(1) = 1$, we deduce that this term is bounded by $\frac{3c}{n} L_{k,0}^2$. Combining this last inequality with (32), (31) and (30), we conclude that

$$\frac{d}{dp} \log((\mathbb{E} X^{\frac{p}{2}})^{\frac{1}{p}}) \leq \frac{c}{p^2 n} (2L_{k,p}^2 + 3L_{k,0}^2) + \frac{C}{k-1} + \frac{C}{r-1}.$$

□

Acknowledgements

The research of the three authors was partially supported by the ANR project GeMeCoD, ANR 2011 BS01 007 01.

References

- [1] R. Adamczak, O. Guédon, R. Latała, A.E. Litvak, K. Oleszkiewicz, A. Pajor, N. Tomczak-Jaegermann, *Moment estimates for convex measures*, *Electr. J. Probab.* **17** (2012), no 101, 1–19.
- [2] M. Anttila, K. Ball, I. Perissinaki, *The central limit problem for convex bodies*, *Trans. Amer. Math. Soc.* **355** (2003), no. 12, 4723–4735.

- [3] E. Artin, *The gamma function*. Athena Series: Selected Topics in Mathematics, Holt, Rinehart and Winston, New York-Toronto-London 1964 vii+39 pp.
- [4] D. Bakry, M. Émery. *Diffusions hypercontractives*. In Séminaire de probabilités, XIX, 1983/84, volume 1123 of Lecture Notes in Math., pages 177–206. Springer, Berlin, 1985.
- [5] K. Ball, *Logarithmically concave functions and sections of convex sets in \mathbb{R}^n* , *Studia Math.*, **88** (1) (1988), 69–84.
- [6] L. Berwald, *Verallgemeinerung eines Mittelwertsatzes von J. Favard für positive konkave Funktionen*, *Acta Math.*, **79** (1947) 17–37.
- [7] S. G. Bobkov, *On concentration of distributions of random weighted sums*, *Annals of Prob.*, **31**(2003), 195–218.
- [8] S.G. Bobkov, *Convex bodies and norms associated to convex measures*, *Probab. Theory Related Fields*, **147** (2010), 303–332.
- [9] S.G. Bobkov, M. Madiman, *Reverse Brunn-Minkowski and reverse entropy power inequalities for convex measures*, *J. Funct. Anal.*, **262** (2012), no. 7, 3309–3339.
- [10] C. Borell, *Convex measures on locally convex spaces*, *Ark. Math.*, **12** (1974), 239–252.
- [11] C. Borell, *Convex set functions in d -space*, *Period. Math. Hungar.*, **6** (1975), 111–136.
- [12] C. Borell, *Complements of Lyapunov’s inequality*, *Math. Ann.*, **205** (1973), 323–331.
- [13] U. Brehm, J. Voigt, *Asymptotics of cross sections for convex bodies*, *Beiträge Algebra Geom.*, **41** (2000), 437–454.
- [14] U. Brehm, P. Hinow, H. Vogt, J. Voigt, *Moment inequalities and central limit properties of isotropic convex bodies*, *Math. Z.*, **240** (2002) 37–51.
- [15] D. Cordero-Erausquin, M. Fradelizi, G. Paouris, P. Pivovarov, *Volume of the polar of random sets and shadow systems*, preprint available at <http://arxiv.org/abs/1311.3690>.
- [16] B. Fleury, *Concentration in a thin Euclidean shell for log-concave measures*, *J. Funct. Anal.*, **259** (2010) 832–841.

- [17] B. Fleury, O. Guédon, G. Paouris, *A stability result for mean width of L_p -centroid bodies*, Adv. Math., **214** (2007) 865–877.
- [18] B. Grünbaum, *Partitions of mass-distributions and of convex bodies by hyperplanes*, Pacific J. Math. **10** (1960), 1257–1261.
- [19] O. Guédon, E. Milman, *Interpolating thin-shell and sharp large-deviation estimates for isotropic log-concave measures*, Geom. Funct. Anal., **21** (2011), 1043–1068.
- [20] O. Guédon, P. Nayar, T. Tkocz, *Concentration inequalities and geometry of convex bodies*, in IMPAN Lecture Notes 2, Inst. Math., Polish. Acad. Sci., 2014, 9–86.
- [21] A. Guionnet, O. Zeitouni, *An introduction to random matrices*, Cambridge Studies in Advanced Mathematics, **118** Cambridge University Press, Cambridge, 2010.
- [22] B. Klartag, *A central limit theorem for convex sets*, Invent. Math., **168** (2007), 91–131.
- [23] B. Klartag, *Power-law estimates for the central limit theorem for convex sets*, J. Funct. Anal., **245** (2007), 284–310.
- [24] B. Klartag, E. Milman, *Inner regularization of log-concave measures and small-ball estimates*, Geometric aspects of functional analysis, 267–278, Lecture Notes in Math., **2050**, Springer, Heidelberg, 2012.
- [25] V.D. Milman, A. Pajor, *Isotropic position and inertia ellipsoids and zonoids of the unit ball of a normed n -dimensional space*, Lectures Notes in Math. **1376**, Springer, Berlin, 1989, 107–131.
- [26] G. Paouris, *Concentration of mass on convex bodies*, Geom. Funct. Anal., **16** (2006), 1021–1049.
- [27] G. Paouris, *Small ball probability estimates for log-concave measures*, Trans. Amer. Math. Soc., **364** (2012), 287–308.

Matthieu Fradelizi, Olivier Guédon, Alain Pajor
 Université Paris-Est
 Laboratoire d'Analyse et Mathématiques Appliquées (UMR 8050)
 UPEMLV, F-77454, Marne-la-Vallée Cedex 2, France
 olivier.guedon@u-pem.fr, matthieu.fradelizi@u-pem.fr, alain.pajor@u-pem.fr