

Universality versus nonuniversality in asymmetric fluid criticality

M.A. Anisimov

Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742, USA

Received January 14, 2013, in final form March 27, 2013

Critical phenomena in real fluids demonstrate a combination of universal features caused by the divergence of long-range fluctuations of density and nonuniversal (system-dependent) features associated with specific intermolecular interactions. Asymptotically, all fluids belong to the Ising-model class of universality. The asymptotic power laws for the thermodynamic properties are described by two independent universal critical exponents and by two independent nonuniversal critical amplitudes; other critical amplitudes can be obtained by universal relations. The nonuniversal critical parameters (critical temperature, pressure, and density) can be absorbed in the property units. Nonasymptotic critical behavior of fluids can be divided into two parts, symmetric (“Ising-like”) and asymmetric (“fluid-like”). The symmetric nonasymptotic behavior contains a new universal exponent (Wegner exponent) and the system-dependent crossover scale (Ginzburg number) associated with the range of intermolecular interactions, while the asymmetric features are generally described by an additional universal exponent and by three nonasymptotic amplitudes associated with mixing of the physical fields into the scaling fields.

Key words: *fluids, critical point, universality, complete scaling*

PACS: *64.60.F-*

1. Introduction

Universality of critical phenomena is one of the most fascinating concepts in physics of condensed matter [1, 2]. Phase transitions of strikingly different nature, such as para-ferro-magnetism, vaporization, or fluid demixing may be described by the same equation of state near the critical points if a proper (“isomorphic”) set of thermodynamic variables is chosen. There are several classes of universality defined through the dimension of the order parameter. The order parameter may be a scalar, n -component vector or a tensor. For example, the order parameter in fluids is associated with the density or concentration (a scalar) while the order parameter in anisotropic magnetics (magnetization) is a one-component vector. The Ising model of anisotropic ferromagnets is mathematically equivalent to the lattice-gas model which describes the condensation of fluids. In the both cases, the order parameter is one-dimensional ($n = 1$). The isomorphism between the members of a universality class can be established by mapping the thermodynamic variables of one system onto another. In addition, the order parameter can be conserved (such as density) or non-conserved (such as magnetization). This particular nature of the order parameter affects the phase-transition dynamics.

It is well established, primarily through experiments [3, 4], that all fluids and fluid mixtures belong to the Ising-model class of universality in statics and to the conserved-order-parameter class of universality in dynamics. This universality is associated with the universal nature of critical fluctuations. The fluctuations of the order parameter diverge at the critical point. The correlation length of the order-parameter fluctuations becomes much larger than the range of intermolecular interactions, thus making the details of the intermolecular potential unimportant. Landau and Lifshitz stated in an earlier edition of “Statistical Physics” [5], “*Unlike solids and gases, liquids do not allow a general calculation of their thermodynamics quantities or even their temperature dependence. The reason for this is the presence of strong interactions between the molecules of the liquid without having, at the same time, the smallness of vibrations which*

makes the thermal motions of solids so simple.” Undoubtedly, this statement is not applicable to liquids in the vicinity of their critical points. Thermodynamic properties in the critical region of very different substances, such as helium isotopes and other inert gases, organic liquids and water can be theoretically predicted; they are all described by the universal power laws, also known as scaling laws, which are characterized by the universal critical exponents.

However, there are still non-universal features in the critical behavior of fluids. The critical parameters (temperature, pressure, and density) are obviously non-universal, being determined by specific intermolecular potentials. The critical temperature ranges from a few kelvins for helium isotopes to thousands of kelvins for liquid metals. This particular non-universal feature can be eliminated by reducing the properties of a particular substance by its critical parameters. Another non-universal feature is the size of the asymptotic critical region in which the universal scaling laws are valid. This size is controlled by the so-called Ginzburg number which depends on the range of intermolecular interactions. Finally, real fluids, unlike the lattice-gas/Ising model, are asymmetric with respect to the critical isochor. The fluid asymmetry causes additional specific non-asymptotic corrections to the universal critical behavior. In this paper, I present a brief overview of universal and nonuniversal contributions to the equation of state of near-critical fluids.

2. Universal asymptotic criticality

The fluctuation-induced non-analytic critical behavior can be asymptotically described by scaling theory in terms of two independent scaling fields, namely, h_1 (“ordering” field) and h_2 (“thermal” field) and two conjugate scaling densities, namely, the order parameter ϕ_1 (strongly fluctuating) and ϕ_2 (weakly fluctuating). The third field, h_3 , is the critical part of an appropriate field-dependent thermodynamic potential, which is defined as a function exhibiting a minimum at equilibrium with respect to a variation of the order parameter. The differential of the third field is

$$dh_3 = \phi_1 dh_1 + \phi_2 dh_2. \quad (1)$$

In the scaling theory, the field potential h_3 is a homogeneous function of h_1 and h_2 . Asymptotically,

$$h_3 \approx |h_2|^{2-\alpha} f^\pm \left(\frac{h_1}{|h_2|^{2-\alpha-\beta}} \right), \quad (2)$$

where f^\pm is a scaling function and the superscript \pm refers to $h_2 > 0$ and $h_2 < 0$, respectively. Here and below, \approx means asymptotically equal, while \simeq means approximately equal. The critical point is defined by the condition $h_1 = h_2 = h_3 = 0$. The form of the scaling function is universal; however, it contains two thermodynamically independent (but system-specific) amplitudes. All other asymptotic amplitudes are related to the selected ones by universal relations. The critical exponents α and β are universal within a class of critical-point universality. All fluids and fluid mixtures belong to the Ising-model universality class. The Ising values for $\alpha \simeq 0.109$ and $\beta \simeq 0.326$, are well established theoretically and confirmed experimentally [3, 4, 6–16]. Two Ising amplitudes, \hat{A}_0 and \hat{B}_0 are determined by the asymptotic power-law behavior of the two scaling densities in zero ordering field ($h_1 = 0$):

$$\phi_1 = \left(\frac{\partial h_3}{\partial h_1} \right)_{h_2} \approx \pm \hat{B}_0 |h_2|^\beta \quad (h_2 < 0), \quad (3)$$

$$\phi_2 = \left(\frac{\partial h_3}{\partial h_2} \right)_{h_1} \approx \frac{\hat{A}_0^\pm}{1-\alpha} h_2 |h_2|^{-\alpha} + B_{\text{cr}} |h_2|, \quad (4)$$

and of the three scaling susceptibilities, “strong” χ_1 , “weak” χ_2 , and “cross” χ_{12} in zero ordering field:

$$\chi_1 = \left(\frac{\partial \phi_1}{\partial h_1} \right)_{h_2} \approx \hat{\Gamma}_0^\pm |h_2|^{-\gamma}, \quad (5)$$

$$\chi_2 = \left(\frac{\partial \phi_2}{\partial h_2} \right)_{h_1} \approx \hat{A}_0^\pm |h_2|^{-\alpha}, \quad (6)$$

$$\chi_{12} = \left(\frac{\partial \phi_1}{\partial h_2} \right)_{h_1} \approx \beta \hat{B}_0 \frac{|h_2|^\beta}{h_2} \quad (h_2 < 0), \quad (7)$$

where the strong susceptibility critical exponent

$$\gamma = 2 - \alpha - 2\beta \simeq 1.239, \quad (8)$$

and the strong susceptibility critical amplitude $\hat{\Gamma}_0^\pm$ is related to \hat{B}_0 and \hat{A}_0^\pm through universal ratios as [12]:

$$\alpha \hat{\Gamma}_0^+ \hat{A}_0^+ / \hat{B}_0^2 \simeq 0.0581, \quad (9)$$

$$\hat{\Gamma}_0^+ / \hat{\Gamma}_0^- \simeq 4.8, \quad (10)$$

$$\hat{A}_0^+ / \hat{A}_0^- \simeq 0.523. \quad (11)$$

While the superscript \pm refers to the states at $h_2 > 0$ and $h_2 < 0$, the prefactor \pm in equation (3) refers to the two branches of the order parameter corresponding to $h_1 > 0$ and $h_1 < 0$ sides (in the limit $h_1 = 0$), respectively. The field-dependent potential h_3 is symmetric with respect to the sign of the ordering field h_1 and, hence, to the sign of the order parameter ϕ_1 . In these expressions \hat{A}_0^\pm , $\hat{\Gamma}_0^\pm$, and \hat{B}_0 are non-universal critical amplitudes. The term in ϕ_2 proportional to B_{cr} is an analytic fluctuation-induced contribution to the second scaling density [17]. Strictly speaking, this term is not asymptotic since the term $h_2 |h_2|^{-\alpha}$ dominates.

Additional universal relations connect the critical exponent $\nu \simeq 0.63$ of the correlation length (diverging in zero ordering field as $\xi = \xi_0^\pm |h_2|^{-\nu}$) and α ,

$$2 - \alpha = d\nu \quad (12)$$

(where d is the number of space dimensions), and the amplitudes \hat{A}_0^+ and ξ_0^+ ,

$$A_0^+ \rho_c (\xi_0^+)^3 \simeq 0.172. \quad (13)$$

This relation is known as the two-scale factor of universality [1, 12]. The ratio $\xi_0^+ / \xi_0^- \simeq 1.96$ is also universal [12].

In the mean-field approximation, with $\alpha = 0$ and $\beta = 1/2$, equation (2) reduces to the asymptotic Landau expansion [18],

$$-h_3 \approx \frac{1}{2} a_0 h_2 \phi_1^2 + \frac{1}{24} u_0 \phi_1^4 - h_1 \phi_1, \quad (14)$$

where a_0 and u_0 are mean-field system-dependent amplitudes. The amplitude a_0 is unimportant. It can be eliminated by rescaling the fields h_3 and h_1 and the coupling constant $u_0 \rightarrow u = u_0 / a_0$.

In the lattice-gas model, the ordering field h_1 is associated with the chemical potential μ , the thermal field h_2 is associated with the temperature T , and the order parameter is associated with the molecular density ρ , while h_3 is associated with the pressure P . In both, the scaling regime and the mean-field approximation, the thermodynamic properties of the lattice gas are symmetric with respect to the sign of the order parameter. Similar to the lattice-gas model, in real one-component fluids, the thermodynamic fields are the temperature T , the chemical potential μ , and the pressure P , while the conjugate densities are the number density ρ and the entropy density $s = \rho S$ (S is the entropy per molecule). The physical variables are interrelated by the Gibbs-Duhem relation

$$dP = \rho d\mu + s dT. \quad (15)$$

Consequently, the densities, namely, the molecular density and the entropy density are derived from the pressure as

$$\rho = \left(\frac{\partial P}{\partial \mu} \right)_T, \quad s = \left(\frac{\partial P}{\partial T} \right)_\mu. \quad (16)$$

In addition to the reduced density $\Delta\hat{\rho}$ and reduced temperature $\Delta\hat{T}$,

$$\Delta\hat{\rho} = \frac{\rho - \rho_c}{\rho_c}, \quad \Delta\hat{T} = \frac{T - T_c}{T_c}, \quad (17)$$

it is convenient to define

$$\Delta\hat{s} = \frac{s - s_c}{\rho_c k_B}, \quad \Delta\hat{P} = \frac{P - P_c}{\rho_c k_B T_c}, \quad \Delta\hat{\mu} = \frac{\mu - \mu_c}{k_B T_c}, \quad (18)$$

where k_B is Boltzmann's constant. In equations (17)–(18) and below, the subscript “c” denotes the properties at the critical point.

As shown by Anisimov et al. [19], since in classical thermodynamics, the absolute value of entropy is arbitrary, the critical value of entropy can be chosen upon practical convenience. It is clearly seen from the basic thermodynamic relation

$$dP = \rho d\mu + \rho S dT, \quad (19)$$

that

$$\frac{d\hat{\mu}}{d\hat{T}} + \hat{S}_c - \left(\frac{\partial \hat{P}}{\partial \hat{T}} \right)_c = 0. \quad (20)$$

Thus, with adopting $\hat{S}_c = (\partial \hat{P} / \partial \hat{T})_{h_1=0,c}$, we obtain $(\partial \hat{\mu} / \partial \hat{T})_{h_1=0,c} = 0$ meaning that in the linear approximation, the chemical potential along the vapor-liquid coexistence does not depend on temperature. With this choice of the critical entropy, we find for the critical part of pressure (the density of the grand thermodynamic potential $\Omega = -PV$) after subtracting its regular part

$$\Delta \tilde{P} = \Delta \hat{P} - \Delta \hat{\mu} - \hat{S}_c \Delta \hat{T}. \quad (21)$$

Then, asymptotically, for one-component fluids, the scaling fields have the following simple relations to the physical fields:

$$h_1 = \Delta \hat{\mu}, \quad (22)$$

$$h_2 = \Delta \hat{T}, \quad (23)$$

$$h_3 = \Delta \tilde{P}. \quad (24)$$

One can conclude that in the asymptotic regime, in addition to the system-dependent critical parameters, which can be eliminated by rescaling the units of thermodynamic properties, there are only two independent critical amplitudes. The independent critical amplitudes correspond to the relevant nonuniversal coefficients, u and g , in the asymptotic Landau-Ginzburg Hamiltonian, given in terms of the spatially dependent order parameter $\phi_1(\mathbf{x})$ as [18]

$$\mathcal{H} = \frac{1}{2}(\Delta \hat{T})\phi_1^2 + \frac{u}{4!}\phi_1^4 + \frac{1}{2}g\phi_1(\nabla\phi_1)^2 - h_1\phi_1. \quad (25)$$

The universal scaling function f^\pm in equation (2) for practical applications is commonly calculated from a parametric equation of state, such as the linear model [16, 20], which has been shown to be accurate to an order of ϵ^2 in the ϵ -expansion, where $\epsilon = 4 - d$ [21].

3. Size of the critical region and symmetric corrections to asymptotic scaling laws

The universal scaling laws discussed in the previous Section are valid only asymptotically, very close to the critical point. Upon departure from the critical point, corrections to the asymptotic power laws appear. The first correction, also known as the Wegner correction [22] contains a new universal scaling function, $g_1^\pm(z)$, and a new universal critical exponent θ (known as the “Wegner exponent”). In the first-order ϵ -expansion $\theta = \epsilon/2$ [22].

The Wegner correction arises from the difference between the renormalization-group fixed-point coupling constant u^* and the system dependent mean-field value of the coupling constant u . When the Wegner correction is included, the Ising field-dependent potential [equation (2)] reads [23]

$$h_3 = |h_2|^{2-\alpha} f^\pm(z) + |h_2|^\theta f_1^\pm(z) + \dots, \quad (26)$$

where $z = h_1/|h_2|^{2-\alpha-\beta}$ and $f_1^\pm \propto (u^* - u)$. The nonasymptotic (“confluent”) scaling function f_1^\pm contains additional system-dependent parameters, $(u^* - u)$ and the Ginzburg number, $N_G = u^2[\rho_c(\xi_0^+)^3]^{-2}$.

Thus, the scaling power laws are to be complemented by confluent singularities. In terms of physical variables in zero ordering field

$$\Delta\tilde{P} = |\Delta\hat{T}|^{2-\alpha} \left[1 + \left(\frac{|\Delta\hat{T}|}{t_x} \right)^\theta + \left(\frac{|\Delta\hat{T}|}{t_x} \right)^{2\theta} + \dots \right], \quad (27)$$

where the crossover scale $t_x \propto (N_G)(u^* - u)^{-2}$, which can be considered as the effective Ginzburg number, defines the size of the asymptotic critical region, $|\Delta\hat{T}| \ll t_x$. The amplitude B_{cr} in the analytic fluctuation-induced contribution to ϕ_2 given by equation (4) also depends on the crossover scale t_x [17, 24].

4. Nonasymptotic asymmetry corrections

The canonical mapping of the liquid-vapor critical point onto Ising criticality is given by the lattice-gas model [25]. This model can easily be extended to binary fluids and fluid mixtures through a reassignment of variables and the principle of isomorphism [19]. For the remainder of the text, the liquid-vapor one-component system is only discussed. The lattice-gas model preserves the exact symmetry of uniaxial Ising-type ferromagnets and consequently, the liquid-vapor coexistence curve of the lattice gas is symmetric with respect to the density ρ . The order parameter of the lattice gas is the reduced density, $\Delta\hat{\rho} = (\rho - \rho_c)/\rho_c$. If the liquid and vapor branches of the coexistence curve are denoted by “+” and “-” respectively, the asymmetric portion of the density is given by the excess density

$$\Delta\hat{\rho}_d = \frac{\Delta\hat{\rho}^+ + \Delta\hat{\rho}^-}{2}. \quad (28)$$

For the lattice gas, $\Delta\hat{\rho}_d = 0$. However, real fluids do not possess the symmetry of the Ising model, and in general $\Delta\hat{\rho}_d \neq 0$. Even the coexistence curve of ^3He , the most symmetric fluid known, exhibits some small asymmetry [26]. In asymmetric systems, the leading behavior is still determined by the Ising-type behavior, and asymmetric corrections appear as sub-leading terms in the quantities like density. In mean-field models of the liquid-vapor critical point, such as the van der Waals model, the asymmetry of the coexistence curve is described by the “law” of rectilinear diameter [27, 28]

$$\Delta\hat{\rho}_d = D_1|\Delta\hat{T}|, \quad (29)$$

where the reduced temperature is defined by $\Delta\hat{T} = (T - T_c)/T_c$, with T_c being the critical temperature. While some one-component fluids such as xenon [29] seem to asymptotically follow this “law”, others, like SF_6 [30], show strong deviations from rectilinearity in the critical region.

Models such as the one proposed by ... (text is partially obscured and contains some garbled characters)

$$\Delta\hat{\rho}_d \approx D_{1-\alpha}|\Delta\hat{T}|^{1-\alpha} + D_1|\Delta\hat{T}|.$$

At a nontrivial support of a ... (text is partially obscured and contains some garbled characters)

$$\Delta\hat{\rho}_d \approx D_{1-\alpha}|\Delta\hat{T}|^{1-\alpha} + D_1|\Delta\hat{T}| + D_{\beta+\theta_5}|\Delta\hat{T}|^{\beta+\theta_5}.$$

unversa, xpon nt θ_5 was oun to b, $\theta_5 = 1/2 + \epsilon = 3/2$ nt, first or, re, xpans on ω , $\epsilon = 4 - d$ an d st, spat a, ns ona ty - or n to or, $\epsilon^3 \hat{u} Z$ an an Z a. \hat{u} oun t, r r, sut s to b, cons st nt w t t, bou n $\theta_5 \geq 1.0$

or, r, ç nt yúF s, r an co wor, rs ú, av, ar u, or an, xt n, or u at on o sca n ú or na y scuss, by, r an, r n úw c s now nown as “co p, t sca n” st, ory o asy, tr c flu cr t ca ty s an, xt ns on o t, fi, x n n a r, v s, sca n an ncorpora t s t, ypot, s so Gr ffit s an, r t at pr, rab, t, r o yna c var ab, s o not, x st s conc p t p, s t at pr, ssur, P_{uc} , ca pot nt a μ (an t, p, ratur, T s ou a b, tr, at, on, qua, oot n n any or u at on o sca n or t, qu vapor cr t ca, po nt, Is n sca n fi, s s ou ú t, r, or, úb, tr, at, as ana yt c unct ons o a t r, By contrast úa r, v s, sca n ass ns a sp, ca ro, to t, pr, ssur, $P(\mu, T)$ as t, fi, p, n, nt t, r o yna c pot nt a A co p, t sca n pr, cts t at t, xç, ss, ns ty s asy ptot ca y v, n by

$$\Delta \hat{\rho}_d \approx D_{2\beta} |\Delta \hat{T}|^{2\beta} + D_{1-\alpha} |\Delta \hat{T}|^{1-\alpha} + D_1 |\Delta \hat{T}|, \quad (1)$$

w, r, $2\beta \approx 0.65$ s r, sut c, ar y, rs ro t, F pr, ct on ú quat on. In t, an fi, ap prox at on ú, conn ct on b, tw, na co p, t sca n an t, asy, tr c Lan au, xpans on as b, n nv, st at, by An s ov an an ú, úw o, onstrat t at t, two approac, s app ar to b, con s st nt A co p, t sca n as a so b, n, xt n, to n o o, n, ous flu, s by B, rtran an An s ov at t, p, n, trab, sp, r, o, o, s not, x b t co p, t sca n ú as b, n nv, st at, by, n ta úw o oun t at t s s ú to a sp, ca sy, try o t, o,

In a t on to t, a n 2β t r nt, xç, ss, ns ty úa co p, t sca n a so pr, cts a v, r, nc, nt, s, con, r vat v, o t, c, ca pot nt a a on t, co, x st nc, curv,

$$\left(\frac{d^2 \mu}{dT^2}\right)_{cxc} \sim |\Delta \hat{T}|^{-\alpha}, \quad (2)$$

w, r, t, subscr pt cxc, not s t, con t ons o p as, co, x st nc, so ca, Yan Yan ano a y, r v, s ts na, ro t, Yan Yan r, at on

$$\frac{\rho C_V}{T} = \left(\frac{d^2 P}{dT^2}\right)_{cxc} - \rho \left(\frac{d^2 \mu}{dT^2}\right)_{cxc}, \quad (3)$$

w, r, C_V st, soc or c, at capac ty A co p, t sca n p, s t at t, v, r, nc, o t, soc or c, at capac ty s s ar, b, tw, n t, s, con, r vat v, s o t, pr, ssur, an t, c, ca pot nt a By contrast úa r, v s, sca n pr, cts t at $(d^2 \mu/dT^2)_{cxc}$ r, a ns fin t at t, cr t ca po nt co's ana ys s a so pr, cts a non ana yt c b, av or o t, c, ca pot nt a úsp, c fica yú

$$\left(\frac{d^2 \mu}{dT^2}\right)_{cxc} \sim |\Delta \hat{T}|^{-\alpha-\beta+\theta_5}, \quad (4)$$

ow, v, r út, r, at v, y ar, va ú o θ_5 , nsur, s t at t s quant y r, a ns fin t at t, cr t ca po nt F s, r an co wor, rs av, oun support or a co p, t sca n n, at capac ty, asur, nts an co pu t r s u at ons o y asy, tr c flu o, s. An s ov an an av, on strat t at a co p, t sca n s a so support, by t, ata on qu vapor co, x st nc, n y asy, tr c flu s ú, r, s a so at, ast on, o, t, at, x b t s t, typ, o fi, x n c aract r st co co p, t sca n A co p, t sca n r, a ns ú ow, v, r úan, s s nt a yp, no, no o ca t, ory

5. Complete scaling

For t, qu vapor trans t on t, pr nc p, o co p, t sca n ass rts t at t, sca n fi, s can b, xpans, n $\Delta \hat{\mu} \hat{u} \Delta \hat{T} \hat{u} \Delta \hat{P}$ In t, ow, st or, r approx at on t, sca n fi, s ar, v, n by

$$h_1 \approx a_1 \Delta \hat{\mu} + a_2 \Delta \hat{T} + a_3 \Delta \hat{P}, \quad (5)$$

$$h_2 \approx b_1 \Delta \hat{T} + b_2 \Delta \hat{\mu} + b_3 \Delta \hat{P}, \quad (6)$$

$$h_3 \approx c_1 \Delta \hat{P} + c_2 \Delta \hat{\mu} + c_3 \Delta \hat{T} + c_{23} \Delta \hat{\mu} \Delta \hat{T}, \quad (7)$$

K YCúFs, r EúLu t nEú ys v L ttú . 091ú o ys vL tt
 K YCúFs, r Eú ys v Eú . 068ú o ys vE
 K YCú ys v Eú . 071ú o ys vE
 F, r o BúFs, r EúAnn ys ú . 056ú
 F, r o BúFs, r EúAnn ys ú . 056ú
 B, rtran CEúAsy tr cFu Cr t ca tyú D s stú n v rs ty o ary an ú .
 B, rtran CEú ço JFúAn s ov Aú ys v Eú . 085ú o ys vE
 Yu novs Iú as rans tons o t, con r, r Co, ct v, ar ab, s, t o ú or c, nt ficú n a
 por, ú
 Yu novs IúKo ovs ú y y Iú croscop c, or y o as rans tons n r, D, ns ona yst, stú
 Evrosv túLv vú . n -ra n an

Універсальність чи неуніверсальність в асиметричній критичності плинів

М.А. Анісімов

Інститут фізичної науки і технології, Університет Мариленду, Коледж Парк, MD 20742, США

Критичні явища в реальних плинах демонструють комбінацію універсальних рис, спричинених розбіжністю далекосяжних флуктуацій густини, і неуніверсальних (системо залежних) рис, пов'язаних із специфічними міжмолекулярними взаємодіями. Асимптотично всі плинні належать до класу універсальності моделі Ізинга. Асимптотичні степеневі закони для термодинамічних властивостей описуються двома незалежними універсальними критичними показниками і двома незалежними неуніверсальними критичними амплітудами; решту критичних амплітуд можна отримати з універсальних співвідношень. Неуніверсальні критичні параметри (критична температура, тиск і густина) можуть бути включені в одиниці цих властивостей. Неасимптотичну критичну поведінку плиннів можна поділити на дві частини, симетричну ("ізингоподібну") і асиметричну ("плиноподібну"). Симетрична неасимптотична поведінка містить новий універсальний показник (показник Вегнера) і системо залежний масштаб кросоверу (число Гінзбурга), пов'язаний з областю дії міжмолекулярних взаємодій, тоді як асиметричні риси взагалюму описуються додатковим універсальним показником і трьома неасимптотичними амплітудами, пов'язаними зі змішуванням фізичних полів у скейлінгових полях.

Ключові слова: плинні, критична точка, універсальність, повний скейлінг