

Higher symmetries of cotangent coverings for Lax-integrable multi-dimensional partial differential equations and Lagrangian deformations

H Baran¹, I S Krasil'shchik^{1,2}, O I Morozov³ and P Vojčák¹

¹ Mathematical Institute, Silesian University in Opava, Na Rybníčku 1, 746 01 Opava, Czech Republic

² Independent University of Moscow, B. Vlasevsky 11, 119002 Moscow, Russia

³ Institute of Mathematics and Statistics, University of Tromsø, Tromsø 9037, Norway

E-mail: josephkra@gmail.com

Abstract. We present examples of Lax-integrable multi-dimensional systems of partial differential equations with higher local symmetries. We also consider Lagrangian deformations of these equations and construct variational bivectors on them.

1. Introduction

Different approaches to integrability of partial differential equations (PDEs), [11, 22], are based on their diverse but related properties such as existence infinite hierarchies of (local or nonlocal) symmetries and/or conservation laws, zero-curvature representations (ZCR), bi-Hamiltonian structures, recursion operators, etc.

Much progress was achieved in the study of PDEs with two independent variables. In particular, in a big number of examples it was shown that a PDE integrable in the sense of presence of a ZCR with a non-removable parameter (we call such equations *Lax-integrable*) have infinite hierarchies of higher local symmetries, see, e.g., [11]. In the multidimensional case the situation looks different. As far as we know, no nontrivial examples of multi-dimensional equations with local higher symmetries were found, cf. [20, §6].

In this paper, we present five examples of multi-dimensional systems with higher local symmetries. All these systems are defined as cotangent coverings, [8, 9], for Lax-integrable PDEs with three or four independent variables, namely, for the r -th dispersionsless Dym equation, [2, 10, 12, 15, 17],

$$u_{\mathbf{t}\mathbf{y}} = u_{\mathbf{x}} u_{\mathbf{x}\mathbf{y}} - u_{\mathbf{y}} u_{\mathbf{x}\mathbf{x}}, \quad v_{\mathbf{t}\mathbf{y}} = u_{\mathbf{x}} v_{\mathbf{x}\mathbf{y}} - u_{\mathbf{y}} v_{\mathbf{x}\mathbf{x}} + 2(u_{\mathbf{x}\mathbf{x}} v_{\mathbf{y}} - u_{\mathbf{x}\mathbf{y}} v_{\mathbf{x}}), \quad (1)$$

the r -th dispersionsless KP equation, [2, 5, 10, 16, 17, 18],

$$u_{\mathbf{y}\mathbf{y}} = u_{\mathbf{t}\mathbf{x}} + u_{\mathbf{y}} u_{\mathbf{x}\mathbf{x}} - u_{\mathbf{x}} u_{\mathbf{x}\mathbf{y}}, \quad v_{\mathbf{y}\mathbf{y}} = v_{\mathbf{t}\mathbf{x}} + u_{\mathbf{y}} v_{\mathbf{x}\mathbf{x}} - u_{\mathbf{x}} v_{\mathbf{x}\mathbf{y}} + 2(u_{\mathbf{x}\mathbf{y}} v_{\mathbf{x}} - u_{\mathbf{x}\mathbf{x}} v_{\mathbf{y}}), \quad (2)$$

the Veronese web equation, [3, 7, 21],

$$u_{\mathbf{x}\mathbf{y}} = \frac{\alpha u_{\mathbf{x}} u_{\mathbf{t}\mathbf{y}} + (1 - \alpha) u_{\mathbf{y}} u_{\mathbf{t}\mathbf{x}}}{u_{\mathbf{t}}},$$

$$\begin{aligned}
v_{\mathbf{xy}} &= \frac{\alpha u_{\mathbf{x}} v_{\mathbf{ty}} + (1 - \alpha) u_{\mathbf{y}} v_{\mathbf{tx}}}{u_{\mathbf{t}}} \\
&+ \frac{1}{u_{\mathbf{t}}^2} ((\alpha u_{\mathbf{x}} u_{\mathbf{ty}} + (1 - \alpha) u_{\mathbf{y}} u_{\mathbf{tx}}) v_{\mathbf{t}} \\
&+ ((1 - 2\alpha) u_{\mathbf{t}} u_{\mathbf{ty}} - (1 - \alpha) u_{\mathbf{y}} u_{\mathbf{tt}}) v_{\mathbf{x}} - (\alpha u_{\mathbf{x}} u_{\mathbf{tt}} + (1 - 2\alpha) u_{\mathbf{t}} u_{\mathbf{tx}}) v_{\mathbf{y}}) \\
&- \frac{v}{u_{\mathbf{t}}^3} ((1 - \alpha) u_{\mathbf{y}} u_{\mathbf{tt}} u_{\mathbf{tx}} - u_{\mathbf{t}} u_{\mathbf{tx}} u_{\mathbf{ty}} + \alpha u_{\mathbf{x}} u_{\mathbf{tt}} u_{\mathbf{ty}})
\end{aligned} \tag{3}$$

with $\alpha \neq 0, 1$, the universal hierarchy equation, [10, 17],

$$u_{\mathbf{yy}} = u_{\mathbf{y}} u_{\mathbf{tx}} - u_{\mathbf{x}} u_{\mathbf{ty}}, \quad v_{\mathbf{yy}} = u_{\mathbf{y}} v_{\mathbf{tx}} - u_{\mathbf{x}} v_{\mathbf{ty}} + 2(u_{\mathbf{ty}} v_{\mathbf{x}} - u_{\mathbf{tx}} v_{\mathbf{y}}), \tag{4}$$

and for the 4-dimensional analogue of (1), [14],

$$u_{\mathbf{ty}} = u_{\mathbf{z}} u_{\mathbf{xy}} - u_{\mathbf{y}} u_{\mathbf{xz}}, \quad v_{\mathbf{ty}} = u_{\mathbf{z}} v_{\mathbf{xy}} - u_{\mathbf{y}} v_{\mathbf{xz}} + 2(u_{\mathbf{xz}} v_{\mathbf{y}} - u_{\mathbf{xy}} v_{\mathbf{z}}), \tag{5}$$

introduced in [10]. We show that these systems have local symmetries of the third order. Equations (1), (2) and (3) belong to the families of PDEs

$$u_{\mathbf{ty}} = u_{\mathbf{x}} u_{\mathbf{xy}} + \varkappa u_{\mathbf{y}} u_{\mathbf{xx}}, \tag{6}$$

$$u_{\mathbf{yy}} = u_{\mathbf{tx}} + \left(\frac{1}{2}(\varkappa + 1) u_{\mathbf{x}}^2 + u_{\mathbf{y}}\right) u_{\mathbf{xx}} + \varkappa u_{\mathbf{x}} u_{\mathbf{xy}}, \tag{7}$$

$$u_{\mathbf{xy}} = \frac{\alpha u_{\mathbf{x}} u_{\mathbf{ty}} + \beta u_{\mathbf{y}} u_{\mathbf{tx}}}{u_{\mathbf{t}}} \tag{8}$$

and are featured as the only representatives of their families whose ZCRs have a non-removable parameter. We make an observation that all the symmetries of the cotangent coverings of Equations (6) with $\varkappa \neq -1$, (7) with $\varkappa \neq -1$, and (8) with $\beta \neq 1 - \alpha$ up to order 5 are the point symmetries. This leads to a conjecture that the cotangent coverings of Equations (6), (7), (8) have local higher symmetries whenever these equations are Lax-integrable. If this claim turns out to be correct, it will be a valuable refinement of the observations of [20, §6].

In the last section, we consider *Lagrangian deformations* of cotangent coverings to some of the equations considered above and construct variational bivectors, see [8], on these deformations.

2. Basics

We briefly recall here the basic definitions from the geometry of PDEs, including the definition of the cotangent covering, see [8, 9], and its Lagrangian deformations. Let $\pi: \mathbb{R}^{\mathbf{n}} \times \mathbb{R}^{\mathbf{m}} \rightarrow \mathbb{R}^{\mathbf{n}}$ be the trivial bundle with the coordinates $(x^1, \dots, x^{\mathbf{n}})$ in $\mathbb{R}^{\mathbf{n}}$ and $(u^1, \dots, u^{\mathbf{m}})$ in $\mathbb{R}^{\mathbf{m}}$ and $J^\infty(\pi)$ be the space of its infinite jets. Local coordinates on $J^\infty(\pi)$ are $(x^{\mathbf{I}}, u_{\mathbf{I}})$, where $\mathbf{I} = (i_1, \dots, i_{\mathbf{k}})$ is a multi-index, and for every local section f of π and the corresponding infinite jet $j_\infty(f): \mathbb{R}^{\mathbf{n}} \rightarrow J^\infty(\pi)$ the coordinate $u_{\mathbf{I}}$ is defined by $u_{\mathbf{I}}|_{j_\infty(f)} = \frac{\partial^{\#\mathbf{I}} f}{\partial x^{\mathbf{I}}} = \frac{\partial^{i_1 + \dots + i_n} f}{(\partial x^1)^{i_1} \dots (\partial x^n)^{i_n}}$. We put

$$u = u_{(0, \dots, 0)}.$$

The vector fields

$$D_{\mathbf{x}^k} = \frac{\partial}{\partial x^{\mathbf{k}}} + \sum_{\#\mathbf{I} \geq 0} \sum_{i=1}^{\mathbf{m}} u_{\mathbf{I}+1_{\mathbf{k}}} \frac{\partial}{\partial u_{\mathbf{I}}}, \quad k = 1, \dots, n,$$

$(i_1, \dots, i_{\mathbf{k}}, \dots, i_n) + 1_{\mathbf{k}} = (i_1, \dots, i_{\mathbf{k}} + 1, \dots, i_n)$, are called *total derivatives*. They commute everywhere on $J^\infty(\pi)$.

The *evolutionary derivation* associated to an arbitrary smooth function $\varphi: J^\infty(\pi) \rightarrow \mathbb{R}^m$ is the vector field

$$\mathbf{E} = \sum_{\#\mathbf{I} \geq 0} \sum_{\mathbf{I}=1}^m D_{\mathbf{I}}(\varphi) \frac{\partial}{\partial u_{\mathbf{I}}} \quad (9)$$

with $D_{\mathbf{I}} = D_{(i_1, \dots, i_n)} = D_{x^{\mathbf{I}_1}} \circ \dots \circ D_{x^{\mathbf{I}_n}}$.

A system of PDEs $F_a(x^{\mathbf{i}}, u_{\mathbf{I}}) = 0$, $\#\mathbf{I} \leq s$, $a = 1, \dots, r$, of order $s \geq 1$ with $r \geq 1$ defines the submanifold $\mathcal{E} = \{(x^{\mathbf{i}}, u_{\mathbf{I}}) \in J^\infty(\pi) \mid D_{\mathbf{K}}(F_a(x^{\mathbf{i}}, u_{\mathbf{I}})) = 0, \#\mathbf{K} \geq 0\}$ in $J^\infty(\pi)$.

A function $\varphi: \mathcal{E} \rightarrow \mathbb{R}^m$ is called a (*generator of an infinitesimal symmetry*) of \mathcal{E} when $\mathbf{E}(\varphi) = 0$ on \mathcal{E} . The symmetry φ is a solution to the *defining system*

$$\ell_{\mathcal{E}}(\varphi) = 0, \quad (10)$$

where $\ell_{\mathcal{E}} = \ell_{\mathbf{F}}|_{\mathcal{E}}$ with the matrix differential operator

$$\ell_{\mathbf{F}} = \left(\sum_{\#\mathbf{I} \geq 0} \frac{\partial F_a}{\partial u_{\mathbf{I}}} D_{\mathbf{I}} \right).$$

Solutions to (10) constitute the Lie algebra with respect to the *Jacobi bracket* $\{\varphi, \psi\} = \mathbf{E}(\psi) - \mathbf{E}(\varphi)$ denoted by $\mathbf{sym}(\mathcal{E})$. The subalgebra of *contact symmetries* of \mathcal{E} is $\mathbf{sym}_0(\mathcal{E}) = \mathbf{sym}(\mathcal{E}) \cap C^\infty(J^1(\pi), \mathbb{R}^m)$. Symmetries from $\mathbf{sym}(\mathcal{E}) \setminus C^\infty(J^1(\pi), \mathbb{R}^m)$ are said to be *higher ones*.

Dually, a *cosymmetry* is a function $\psi: \mathcal{E} \rightarrow \mathbb{R}^r$ that satisfies the equation

$$\ell_{\mathcal{E}}^*(\psi) = 0,$$

where $\ell_{\mathcal{E}}^*$ denotes the formally adjoint operator

$$\ell_{\mathbf{F}}^* = \left(\sum_{\#\mathbf{I} \geq 0} (-1)^{\#\mathbf{I}} D_{\mathbf{I}} \circ \frac{\partial F_a}{\partial u_{\mathbf{I}}} \right)^T$$

restricted to \mathcal{E} .

A differential operator $\Delta: C^\infty(\mathcal{E}, \mathbb{R}^r) \rightarrow C^\infty(\mathcal{E}, \mathbb{R}^m)$ is called a *variational bivector* if it takes symmetries of the equation \mathcal{E} to its cosymmetries and satisfies $(\ell_{\mathcal{E}} \circ \Delta)^* = \ell_{\mathcal{E}} \circ \Delta$. Any such an operator defines a skew-symmetric bracket $\{\omega_1, \omega_2\}_\Delta = L_{\Delta(\omega_1)}(\omega_2)$ on the space of conservation laws, where $\delta = \delta^{0, \mathbf{n}-1}$ is the differential in the term E_1 of the Vinogradov \mathcal{C} -spectral sequence (see [19]) and L denotes the Lie derivative. If the variational Schouten bracket $[[\Delta, \Delta]]$ vanishes then Δ is a Hamiltonian operator, see [8].

Denote $\mathcal{W} = \mathbb{R}^l$, $l \leq \infty$, with coordinates w^0, \dots, w^s, \dots . Locally, a *differential covering* over \mathcal{E} is a trivial bundle $\tau: \tilde{\mathcal{E}} = \mathcal{E} \times \mathcal{W} \rightarrow \mathcal{E}$ equipped with the *extended total derivatives*

$$\tilde{D}_{\mathbf{x}^k} = D_{\mathbf{x}^k} + \sum_{s=0}^l T_{\mathbf{k}}^s(x^{\mathbf{i}}, u_{\mathbf{I}}, w^{\mathbf{j}}) \frac{\partial}{\partial w^s} \quad (11)$$

such that $[\tilde{D}_{\mathbf{x}^i}, \tilde{D}_{\mathbf{x}^j}] = 0$. This yields the system of the *covering equations*

$$w_{\mathbf{x}^k}^s = T_{\mathbf{k}}^s(x^{\mathbf{i}}, u_{\mathbf{I}}, w^{\mathbf{j}}). \quad (12)$$

This over-determined system of PDEs is compatible whenever $(x^{\mathbf{i}}, u_{\mathbf{I}}) \in \mathcal{E}$.

The *cotangent covering* is defined as follows. Let, as before, $\mathcal{E} \subset J^\infty(\pi)$ be given by the relations $F_1 = \dots = F_r = 0$, where $F_{\mathbf{a}} = F(x^{\mathbf{i}}, u_{\mathbf{j}}^{\mathbf{j}})$. Consider the bundle $\pi': \mathbb{R}^r \times \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ with coordinates v in \mathbb{R}^r . Then $\mathcal{T}^*\mathcal{E} \subset J^\infty(\pi')$ is the equation obtained from \mathcal{E} by adding the relations $\ell_{\mathcal{E}}^*(v) = 0$. The natural projection $\tau^*: \mathcal{T}^*\mathcal{E} \rightarrow \mathcal{E}$ is an infinite-dimensional covering. Note that $\mathcal{T}^*\mathcal{E}$ is always an Euler-Lagrange equation with the Lagrangian density $\mathcal{L} = (F_1 v^1 + \dots + F_r v^r) dx^1 \wedge \dots \wedge dx^n$. Consequently, the spaces of symmetries $\mathbf{sym}(\mathcal{E})$ and cosymmetries $\mathbf{cosym}(\mathcal{E})$ coincide for $\mathcal{T}^*\mathcal{E}$.

3. Higher symmetries of cotangent coverings

In this section we present the results of computation of the third order symmetries for Systems (1)–(5). The computations were held in the JETS software, [1, 6]. We use the notation $u^1 = u$, $u^2 = v$, $x^1 = t$, $x^2 = x$, $x^3 = y$, and $x^4 = z$ in the formulas of the previous section.

Proposition 1. *The cotangent coverings of the Lax-integrable cases of the r -th dispersionsless Dym equation, r -th modified dispersionsless KP equation, the Veronese web equation, the universal hierarchy equation, and the 4-dimensional r -th dispersionsless Dym equation have the following symmetries up to the third order:*

1) System (1):

$$\begin{aligned}
\varphi_0(A_0) &= (A_0 u_t + A_{0,t}(x u_x - u) + \frac{1}{2} A_{0,tt} x^2, A_0 v_t + A_{0,t}(x v_x + 2v)), \\
\varphi_1(A_1) &= (A_1 u_x + A_{1,t} x, A_1 v_x), \\
\varphi_2(A_2) &= (A_2, 0), \\
\varphi_3(A_3) &= (0, A_3(2u_t - 3u_x^2) + A_{3,t}(2x u_x + u) - \frac{1}{2} A_{3,tt} x^2), \\
\varphi_4(A_4) &= (0, 2A_4 u_x - A_{4,t} x), \\
\varphi_5(A_5) &= (0, A_5), \\
\varphi_6(B_0) &= (B_0 u_y, B_0 v_y), \\
\varphi_7(B_1) &= (0, B_1 u_y^{-2}), \\
\varphi_8 &= (x u_x - 2u, x v_x), \\
\varphi_9 &= (0, v), \\
\varphi_{10} &= (0, u_{ttt} - 3u_x(u_{ttx} - u_x u_{txx} - u_{tx} u_{xx}) - u_x^3 u_{xxx} - \frac{3}{2}(u_{tx}^2 + u_x^2 u_{xx})), \\
\varphi_{11} &= (0, u_{ttx} - 2u_x u_{txx} + u_x^2 u_{xxx} - u_{xx}(u_{tx} - u_x u_{xx})), \\
\varphi_{12} &= (0, u_{txx} - u_x u_{xxx} - \frac{1}{2} u_{xx}^2), \\
\varphi_{13} &= (0, u_{xxx}), \\
\varphi_{14} &= (0, (2u_y u_{xxy} - u_{xy}^2) u_y^{-2}), \\
\varphi_{15} &= (0, (u_y u_{xyy} - u_{xy} u_{yy}) u_y^{-3}), \\
\varphi_{16} &= (0, (2u_y u_{yyy} - 3u_{yy}^2) u_y^{-4}),
\end{aligned}$$

where $A_i = A_i(t)$, $B_i = B_i(y)$ are arbitrary functions of their arguments;

2) System (2):

$$\begin{aligned}
\varphi_0(A_0) &= (A_0 u_t + (x u_x + y u_y - u) A_{0,t} + (\frac{1}{2} u_x y^2 - x y) A_{0,tt} - \frac{1}{6} y^3 A_{0,ttt}, \\
&\quad A_0 v_t + (x v_x + y v_y + 2v) A_{0,t} + \frac{1}{2} y^2 v_x A_{0,tt}), \\
\varphi_1(A_1) &= (A_1 u_y + A_{1,t}(y u_x - x) - \frac{1}{2} A_{1,tt} y^2, A_1 v_y + A_{1,t} y v_x), \\
\varphi_2(A_2) &= (A_2 u_x - A_{2,t} y, A_2 v_x), \\
\varphi_3(A_3) &= (A_3, 0),
\end{aligned}$$

$$\begin{aligned}
\varphi_4(A_4) &= (0, 2A_4(u_t + 2u_x^3 + 3u_x u_y) + A_{4,t}(3y u_x^2 + 2x u_x + 2y u_y + u) \\
&\quad + A_{4,tt} y(y u_x + x) + \frac{1}{6} A_{4,ttt} y^3), \\
\varphi_5(A_5) &= (0, A_5(2u_y + 3u_x^2) + A_{5,t}(2y u_x + x) + \frac{1}{2} A_{5,tt} y^2), \\
\varphi_6(A_6) &= (0, 2A_6 u_x + A_{6,t} y), \\
\varphi_7(A_7) &= (0, A_7), \\
\varphi_8 &= (2x u_x + y u_y - 3u, 2x v_x + y v_y + 4v + 2t u_t + u_x(2x + 3y u_x + 4t u_x^2) \\
&\quad + 2u_y(3t u_x + y) + u), \\
\varphi_9 &= (y u_x + 2t u_y, y v_x + 2t v_y), \\
\varphi_{10} &= (0, v), \\
\varphi_{11} &= (0, u_{xxx}), \\
\varphi_{12} &= (0, u_{xxy} + u_x u_{xxx} + \frac{1}{2} u_{xx}^2), \\
\varphi_{13} &= (0, u_{txx} + u_{xxx}(u_x^2 + u_y) + u_x u_{xxy} + u_{xx}(u_{xy} + u_x u_{xx})), \\
\varphi_{14} &= (0, 2u_{txy} + 4u_x u_{txx} + 2u_x u_{xxx}(u_x^2 + 2u_y) + 2u_{xxy}(u_x^2 + u_y) + u_{xx}^2(3u_x^2 + 2u_y) \\
&\quad + u_{xy}^2 + 2u_{xx}(u_{tx} + 2u_x u_{xy})), \\
\varphi_{15} &= (0, u_{ttx} + u_{txx}(3u_x^2 + 2u_y) + 2u_x u_{txy} + u_{xxx}(u_x^4 + 3u_y u_x^2 + u_y^2) + u_x u_{xy}^2 \\
&\quad + u_x u_{xxy}(u_x^2 + 2u_y) + u_{tx}(u_{xy} + 3u_x u_{xx}) + u_x u_{xx}^2(2u_x^2 + 3u_y) \\
&\quad + u_{xx}(u_{ty} + u_{xy}(3u_x^2 + 2u_y))), \\
\varphi_{16} &= (0, 2u_{tty} + 6u_x u_{ttx} + 4u_x u_{txx}(2u_x^2 + 3u_y) + 2u_{txy}(2u_y + 3u_x^2) \\
&\quad + 2u_x u_{xxx}(u_x^4 + 4u_x^2 u_y + 3u_y^2) + 2u_{xxy}(u_x^4 + 3u_x^2 u_y + u_y^2) + 3u_{tx}^2 \\
&\quad + 6u_{tx}(u_x u_{xy} + u_{xx}(u_y + 2u_x^2)) + u_{xx}^2(5u_x^4 + 12u_x^2 u_y + 3u_y^2) \\
&\quad + u_{xy}^2(3u_x^2 + 2u_y) + 2u_{ty}(u_{xy} + 3u_x u_{xx}) + 4u_x u_{xx} u_{xy}(2u_x^2 + 3u_y)), \\
\varphi_{17} &= (0, u_{ttt} + 3(2u_x^2 + u_y) u_{ttx} + 3u_x u_{tty} + (5u_x^4 + 12u_x^2 u_y + 3u_y^2) u_{txx} \\
&\quad + (5u_y u_x^4 + 6u_y^2 u_x^2 + u_x^6 + u_y^3) u_{xxx} + u_x(u_y + u_x^2)(3u_y + u_x^2) u_{xxy} \\
&\quad + 12u_x u_y u_{tx} u_{xx} + 10u_x^3 u_{tx} u_{xx} + 6u_x u_{tx}^2 + 2u_x(2u_x^2 + 3u_y) u_{txy} \\
&\quad + 5u_x^4 u_{xx} u_{xy} + 12u_x^2 u_y u_{xx} u_{xy} + 3u_y u_{tx} u_{xy} + 6u_x^2 u_{tx} u_{xy} + 3u_x u_y u_{xy}^2 \\
&\quad + 3u_y^2 u_{xx} u_{xy} + 3u_x^5 u_{xx}^2 + 6u_x u_y^2 u_{xx}^2 + 10u_y u_x^3 u_{xx}^2 \\
&\quad + 3(u_{tx} + 2u_{xx} u_x^2 + u_y u_{xx} + u_x u_{xy}) u_{ty} + 2u_x^3 u_{xy}^2),
\end{aligned}$$

where $A_i = A_i(t)$ are arbitrary functions of t ;

3) System (3):

$$\begin{aligned}
\varphi_1(P_1) &= (P_1 u_t, P_1 v_t + P_{1,t} v), \\
\varphi_2(Q_1) &= (Q_1 u_x, Q_1 v_x), \\
\varphi_3(R_1) &= (R_1 u_y, R_1 v_y), \\
\varphi_4(S_1) &= (S_1, -S_{1,u} v), \\
\varphi_5 &= (u, 0), \\
\varphi_6(P_2) &= (0, P_2 u_t^{-1}), \\
\varphi_7(Q_2) &= (0, Q_2 u_t u_x^{-2}), \\
\varphi_8(R_2) &= (0, R_2 u_t u_y^{-2}), \\
\varphi_9(S_2) &= (0, S_2 u_t), \\
\varphi_{10} &= (0, (2u_t u_x u_{ttx} - 2u_x u_{tt} u_{tx} - u_t u_{tx}^2) u_t^{-2} u_x^{-2}),
\end{aligned}$$

$$\begin{aligned}
\varphi_{11} &= (0, (2 u_t u_y u_{tty} - 2 u_y u_{tt} u_{ty} - u_t u_{ty}^2) u_t^{-2} u_y^{-2}), \\
\varphi_{12} &= (0, (2 u_t u_x u_{txx} - u_x u_{tx}^2 - 2 u_t u_{tx} u_{xx}) u_t^{-1} u_x^{-3}), \\
\varphi_{13} &= (0, (2 u_t u_y u_{tyy} - u_y u_{ty}^2 - 2 u_t u_{ty} u_{yy}) u_t^{-1} u_y^{-3}), \\
\varphi_{14} &= (0, (2 u_t u_{ttt} - 3 u_{tt}^2) u_t^{-3}), \\
\varphi_{15} &= (0, u_t (2 u_x u_{xxx} - 3 u_{xx}^2) u_x^{-4}), \\
\varphi_{16} &= (0, u_t (2 u_y u_{yyy} - 3 u_{yy}^2) u_y^{-4}),
\end{aligned}$$

where $P_i = P_i(t)$, $Q_i = Q_i(x)$, $R_i = R_i(y)$, $S_i = S_i(u)$ are arbitrary functions of their arguments;

4) System (4):

$$\begin{aligned}
\varphi_0(A_0) &= (A_0 u_t - A_{0,t} u, A_0 v_t + 2 A_{0,t} v), \\
\varphi_1(A_1) &= (A_1, 0), \\
\varphi_2(A_2) &= (0, 2 A_2 u_t + A_{2,t} u), \\
\varphi_3(A_3) &= (0, A_3), \\
\varphi_4(B_0) &= (B_0 u_x + B_{0,x} y u_y, B_0 v_x + B_{0,x} y v_y), \\
\varphi_5(B_1) &= (B_1 u_y, B_1 v_y), \\
\varphi_6(B_2) &= (0, (2 B_2 u_x - B_{2,x} y u_y) u_y^{-3}), \\
\varphi_7(B_3) &= (0, B_3 u_y^{-2}), \\
\varphi_8 &= (y u_y + u, y v_y), \\
\varphi_9 &= (0, v), \\
\varphi_{10} &= (0, u_{ttt}), \\
\varphi_{11} &= (0, (u_y^2 u_{ttx} - u_x u_y u_{tty} + u_x u_{ty}^2 - u_y u_{tx} u_{ty}) u_y^{-3}), \\
\varphi_{12} &= (0, (2 u_y u_{tty} - u_{ty}^2) u_y^{-2}), \\
\varphi_{13} &= (0, (2 u_x^2 u_y^2 (u_x u_{ttx} - 3 u_{txy}) - 2 u_x^4 u_y u_{tty} - 2 u_y^3 (u_{xxy} - 2 u_x u_{txx}) \\
&\quad + 3 u_x u_y^2 (u_x u_{tx}^2 - 2 u_{ty} u_{xx} - 2 u_{tx} u_{xy}) - 4 u_x^2 u_y u_{ty} (2 u_x u_{tx} - 3 u_{xy}) \\
&\quad + u_y^2 (3 u_{xy}^2 + 2 u_y u_{tx} u_{xx}) + 5 u_x^4 u_{ty}^2) u_y^{-6}), \\
\varphi_{14} &= (0, (2 u_x^4 u_{xxx} + 2 u_x^4 u_y (u_y u_{ttx} - 5 u_{tx} u_{ty}) - 4 u_x^3 u_y^2 (2 u_{txy} - u_{tx}^2) \\
&\quad + 6 u_x^2 u_y^2 (u_y u_{txx} - 2 u_{ty} u_{xx} - 2 u_{tx} u_{xy}) - 2 u_x^5 (u_y u_{tty} - 3 u_{ty}^2) \\
&\quad - 6 u_y^3 (u_x u_{xxy} + u_{xx} u_{xy} - u_x u_{tx} u_{xx}) + 4 u_x u_y u_{xy} (3 u_y u_{xy} + 5 u_x^2 u_{ty})) u_y^{-7}), \\
\varphi_{15} &= (0, (u_x^2 (2 u_y u_{tty} - 3 u_{ty}^2) + u_y^2 (2 u_{txy} - 2 u_x u_{ttx} - u_{tx}^2) \\
&\quad - 2 u_y u_{ty} (u_{xy} - 2 u_x u_{tx})) u_y^{-4}), \\
\varphi_{16} &= (0, (u_x^3 (u_y u_{tty} - 2 u_{ty}^2) + u_x u_y^2 (2 u_{txy} - u_{tx}^2) - u_y^2 (u_x^2 u_{ttx} - u_{ty} u_{xx} - u_{tx} u_{xy}) \\
&\quad - u_y^3 u_{txx} + 3 u_x u_y (u_x u_{tx} u_{ty} - u_{ty} u_{xy})) u_y^{-5}),
\end{aligned}$$

where $A_i = A_i(t)$, $B_i = B_i(x)$ are arbitrary functions of their arguments;

5) System (5):

$$\begin{aligned}
\varphi_0(A_0) &= (A_0 u_x - A_{0,x} u + A_{0,t} z, A_0 v_x + 2 A_{0,x} v), \\
\varphi_1(A_1) &= (A_1, 0), \\
\varphi_2(A_2) &= (0, 2 A_2 u_x + A_{2,x} u - A_{2,t} z), \\
\varphi_3(A_3) &= (0, A_3), \\
\varphi_4(B_0) &= (B_0 u_y, B_0 v_y),
\end{aligned}$$

$$\begin{aligned}
\varphi_5(B_1) &= (0, B_1 u_{\mathbf{y}}^{-2}), \\
\varphi_6 &= (t u_{\mathbf{t}} + u, t v_{\mathbf{t}}), \\
\varphi_7 &= (z u_{\mathbf{z}} - u, z v_{\mathbf{z}}), \\
\varphi_8 &= (u_{\mathbf{z}}, v_{\mathbf{z}}), \\
\varphi_9 &= (0, v), \\
\varphi_{10} &= (0, u_{\mathbf{xxx}}), \\
\varphi_{11} &= (0, (2 u_{\mathbf{y}} u_{\mathbf{xxxy}} - u_{\mathbf{xy}}^2) u_{\mathbf{y}}^{-2}), \\
\varphi_{12} &= (0, (u_{\mathbf{y}} u_{\mathbf{xyy}} - u_{\mathbf{xy}} u_{\mathbf{yy}}) u_{\mathbf{y}}^{-3}), \\
\varphi_{13} &= (0, (2 t u_{\mathbf{y}}^2 u_{\mathbf{xxx}} - z (2 u_{\mathbf{y}} u_{\mathbf{xxxy}} - u_{\mathbf{xy}}^2)) u_{\mathbf{y}}^{-2}), \\
\varphi_{14} &= (0, (t u_{\mathbf{y}} (2 u_{\mathbf{y}} u_{\mathbf{xxxy}} - u_{\mathbf{xy}}^2) - 2 z (u_{\mathbf{y}} u_{\mathbf{xyy}} - u_{\mathbf{xy}} u_{\mathbf{yy}})) u_{\mathbf{y}}^{-3}), \\
\varphi_{15} &= (0, (z^2 (u_{\mathbf{y}} u_{\mathbf{xyy}} - u_{\mathbf{xy}} u_{\mathbf{yy}}) - t z u_{\mathbf{y}} (2 u_{\mathbf{y}} u_{\mathbf{xxxy}} - u_{\mathbf{xy}}^2) + t^2 u_{\mathbf{y}}^3 u_{\mathbf{xxx}}) u_{\mathbf{y}}^{-3}), \\
\varphi_{16} &= (0, (2 u_{\mathbf{y}} u_{\mathbf{yyy}} - 3 u_{\mathbf{yy}}^2) u_{\mathbf{y}}^{-4}), \\
\varphi_{17} &= (0, (z (2 u_{\mathbf{y}} u_{\mathbf{yyy}} - 3 u_{\mathbf{yy}}^2) - 2 t u_{\mathbf{y}} (u_{\mathbf{y}} u_{\mathbf{xyy}} - u_{\mathbf{xy}} u_{\mathbf{yy}})) u_{\mathbf{y}}^{-4}), \\
\varphi_{18} &= (0, (z^2 (2 u_{\mathbf{y}} u_{\mathbf{yyy}} - 3 u_{\mathbf{yy}}^2) - 4 t z u_{\mathbf{y}} (u_{\mathbf{y}} u_{\mathbf{xyy}} - u_{\mathbf{xy}} u_{\mathbf{yy}}) \\
&\quad + t^2 u_{\mathbf{y}}^2 (2 u_{\mathbf{y}} u_{\mathbf{xxxy}} - u_{\mathbf{xy}}^2)) u_{\mathbf{y}}^{-4}), \\
\varphi_{19} &= (0, (z^3 (2 u_{\mathbf{y}} u_{\mathbf{yyy}} - 3 u_{\mathbf{yy}}^2) - 6 t z^2 u_{\mathbf{y}} (u_{\mathbf{y}} u_{\mathbf{xyy}} - u_{\mathbf{xy}} u_{\mathbf{yy}}) \\
&\quad + 3 t^2 z u_{\mathbf{y}}^2 (2 u_{\mathbf{y}} u_{\mathbf{xxxy}} - u_{\mathbf{xy}}^2) - 2 t^3 u_{\mathbf{y}}^4 u_{\mathbf{xxx}}) u_{\mathbf{y}}^{-4}),
\end{aligned}$$

where $A_{\mathbf{i}} = A_{\mathbf{i}}(t, x)$, $B_{\mathbf{i}} = B_{\mathbf{i}}(y, z)$ are arbitrary functions of their arguments.

All the functions above are assumed to be smooth and the second subscript (as in, say, $A_{3,\mathbf{t}}$) denotes the corresponding derivative.

Thus, we have the following higher symmetries:

$$\begin{array}{ll}
\text{For Systems (1), (3), (4):} & \varphi_{10}, \dots, \varphi_{16}, \\
\text{For System (2):} & \varphi_{11}, \dots, \varphi_{17}, \\
\text{For System (5):} & \varphi_{10}, \dots, \varphi_{19}.
\end{array}$$

4. Bivectors on the deformed cotangent covering

Let \mathcal{E} be an equation and $\tau^*: \mathcal{T}^*\mathcal{E} \rightarrow \mathcal{E}$ be its tangent covering. As it was mentioned in Section 2, $\mathcal{T}^*\mathcal{E}$ is an Euler-Lagrange equation. Let $\mathcal{T}^*\mathcal{E}$ be given by $F(x, u) = 0$, $\ell_{\mathcal{E}}^*(x, u, v) = 0$. Then we say that the system

$$\tilde{\mathcal{E}}: \quad \ell_{\mathcal{E}}^*(x, u, v) + G(x, u, v) = 0, \quad F(x, u) + H(x, u, v) = 0 \quad (13)$$

is a *Lagrangian deformation* (LD) if $\tilde{\mathcal{E}}$ is also an Euler-Lagrange equation. We know at least two meaningful examples of this construction: (i) The system

$$u_{\mathbf{t}\mathbf{y}} = u_{\mathbf{xy}}u_{\mathbf{x}} - u_{\mathbf{xx}}u_{\mathbf{y}}, \quad v_{\mathbf{t}\mathbf{y}} = 2(u_{\mathbf{xx}}v_{\mathbf{y}} - u_{\mathbf{xy}}v_{\mathbf{x}}) + u_{\mathbf{x}}v_{\mathbf{xy}} - u_{\mathbf{y}}v_{\mathbf{xx}} - 2(u_{\mathbf{xx}}u_{\mathbf{y}} + 2u_{\mathbf{xy}}u_{\mathbf{x}}),$$

see [15], is an LD of (1); (ii) The Dunajski equation, [4],

$$\Theta_{\mathbf{wx}} + \Theta_{\mathbf{zy}} + \Theta_{\mathbf{xx}}\Theta_{\mathbf{yy}} - \Theta_{\mathbf{xy}}^2 = f, \quad f_{\mathbf{xw}} + f_{\mathbf{yz}} + \Theta_{\mathbf{yy}}f_{\mathbf{xx}} + \Theta_{\mathbf{xx}}f_{\mathbf{yy}} - 2\Theta_{\mathbf{xy}}f_{\mathbf{xy}} = 0$$

is an LD of the 2nd Heavenly Equation.

The identical map from $\mathbf{cosym}(\tilde{\mathcal{E}})$ to $\mathbf{sym}(\tilde{\mathcal{E}})$ is always a bivector on any LD $\tilde{\mathcal{E}}$ and we pose the following problem: *Given an equation \mathcal{E} , can $\mathcal{T}^*\mathcal{E}$ be deformed in such a way that the resulting LD $\tilde{\mathcal{E}}$ will admit nontrivial bivectors?*

We studied LDs for Equations (1), (4), and (5) and obtained the following result:

Proposition 2. *Equations (1), (4), and (5) possess two-parameter families of LDs with $H = 0$ and $G = \delta L_{\varkappa_1, \varkappa_2} / \delta u$ and three-dimensional spaces of bivectors $\Delta = \alpha_0 \text{id} + \alpha_1 \Delta^1 + \alpha_2 \Delta^2$, $\Delta^i = (\Delta^i_{\mathbf{jk}})$, $i, j, k = 1, 2$, where*

For Equation (1):

$$\begin{aligned}
L_{\varkappa_1, \varkappa_2} &= (\varkappa_1 u_{\mathbf{y}} + \varkappa_2) u_{\mathbf{x}}^2, \\
\Delta^1_{11} &= u_{\mathbf{y}} D_{\mathbf{x}}^{-1} \circ u_{\mathbf{y}}^{-2} \circ D_{\mathbf{y}}, \\
\Delta^1_{12} &= 0, \\
\Delta^1_{21} &= (v_{\mathbf{y}} - \varkappa_2) D_{\mathbf{x}}^{-1} \circ u_{\mathbf{y}}^{-2} \circ D_{\mathbf{y}} + u_{\mathbf{y}}^{-2} D_{\mathbf{x}}^{-1} \circ (v_{\mathbf{y}} - \varkappa_2) \circ D_{\mathbf{y}}, \\
\Delta^1_{22} &= u_{\mathbf{y}}^{-2} D_{\mathbf{x}}^{-1} \circ u_{\mathbf{y}} \circ D_{\mathbf{y}}, \\
\Delta^2_{11} &= \frac{1}{2} u_{\mathbf{x}} + \frac{1}{2} D_{\mathbf{x}}^{-1} \circ (D_{\mathbf{t}} - 2u_{\mathbf{x}} D_{\mathbf{x}}), \\
\Delta^2_{12} &= 0, \\
\Delta^2_{21} &= v + \varkappa_1 u D_{\mathbf{x}} - \frac{1}{2} D_{\mathbf{x}}^{-1} \circ (2\varkappa_1 u D_{\mathbf{x}} + v_{\mathbf{xx}}) - \varkappa_1 D_{\mathbf{x}}^{-1} \circ (D_{\mathbf{t}} - 2u_{\mathbf{x}} D_{\mathbf{x}}), \\
\Delta^2_{22} &= \frac{1}{2} u D_{\mathbf{x}} - u_{\mathbf{x}} + \frac{1}{2} D_{\mathbf{x}}^{-1} \circ (D_{\mathbf{t}} - u D_{\mathbf{x}}^2).
\end{aligned}$$

For Equation (4):

$$\begin{aligned}
L_{\varkappa_1, \varkappa_2} &= u_{\mathbf{t}} (\varkappa_1 u_{\mathbf{y}} + \varkappa_2 u_{\mathbf{x}}), \\
\Delta^1_{11} &= u D_{\mathbf{t}}^{-1} \circ u_{\mathbf{y}}^{-2} \circ D_{\mathbf{y}}, \\
\Delta^1_{12} &= 0, \\
\Delta^1_{21} &= (v_{\mathbf{y}} + \varkappa_2) D_{\mathbf{t}}^{-1} \circ u_{\mathbf{y}}^{-2} \circ D_{\mathbf{y}} + u_{\mathbf{y}}^{-2} D_{\mathbf{t}}^{-1} \circ (v_{\mathbf{y}} + \varkappa_2) \circ D_{\mathbf{y}}, \\
\Delta^1_{22} &= u_{\mathbf{y}}^{-2} D_{\mathbf{t}}^{-1} \circ u_{\mathbf{y}} \circ D_{\mathbf{y}}, \\
\Delta^2_{11} &= u D_{\mathbf{t}} - D_{\mathbf{y}}^{-1} \circ (u_{\mathbf{y}t} + u D_{\mathbf{y}} D_{\mathbf{t}}), \\
\Delta^2_{12} &= 0, \\
\Delta^2_{21} &= -2v D_{\mathbf{t}} - D_{\mathbf{y}}^{-1} \circ (v_{\mathbf{y}t} + 2\varkappa_2 D_{\mathbf{t}} - 2v D_{\mathbf{y}} D_{\mathbf{t}}), \\
\Delta^2_{22} &= u D_{\mathbf{t}} + D_{\mathbf{y}}^{-1} \circ (u D_{\mathbf{y}} D_{\mathbf{t}} - 2u_{\mathbf{y}t}).
\end{aligned}$$

For Equation (5):

$$\begin{aligned}
L_{\varkappa_1, \varkappa_2} &= u_{\mathbf{x}} (\varkappa_1 u_{\mathbf{z}} + \varkappa_2 u_{\mathbf{y}}), \\
\Delta^1_{11} &= -u D_{\mathbf{x}} + D_{\mathbf{z}}^{-1} \circ (u_{\mathbf{xz}} + D_{\mathbf{t}} + u D_{\mathbf{x}} D_{\mathbf{z}}), \\
\Delta^1_{12} &= 0, \\
\Delta^1_{21} &= -D_{\mathbf{z}}^{-1} \circ (v_{\mathbf{xz}} + 2(v_{\mathbf{z}} + \varkappa_2) D_{\mathbf{x}}), \\
\Delta^1_{22} &= -u D_{\mathbf{x}} + 2u_{\mathbf{x}} + D_{\mathbf{z}}^{-1} \circ (u_{\mathbf{xz}} + D_{\mathbf{t}} + u D_{\mathbf{x}} D_{\mathbf{z}}),
\end{aligned}$$

$$\begin{aligned}\Delta_{11}^2 &= u_{\mathbf{y}} D_{\mathbf{x}}^{-1} \circ u_{\mathbf{y}}^{-2} \circ D_{\mathbf{y}}, \\ \Delta_{12}^2 &= 0, \\ \Delta_{21}^2 &= -u_{\mathbf{y}}^{-2} D_{\mathbf{x}}^{-1} \circ (v_{\mathbf{y}} + \varkappa_1) \circ D_{\mathbf{y}} - (v_{\mathbf{y}} + \varkappa_1) D_{\mathbf{x}}^{-1} \circ u_{\mathbf{y}}^{-2} \circ D_{\mathbf{y}}, \\ \Delta_{22}^2 &= u_{\mathbf{y}}^{-2} D_{\mathbf{x}}^{-1} \circ u_{\mathbf{y}} D_{\mathbf{y}}.\end{aligned}$$

5. Concluding remarks

Remark 1. The identical bivector in Equation (1), after transforming the system to the evolutionary form, corresponds to the first Hamiltonian operator from [15], while the bivector Δ^1 corresponds to the second one.

Remark 2. It can be shown that the LDS for Equations (4) and (5) presented above are in a sense trivial, because the deformed equations are transformed to the initial cotangent space by a point transformation. As for Equation (1), there exists a covering $\tau: \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{E}}$ (that corresponds to a differential substitution) such that $\tilde{\mathcal{E}}$ is equivalent to $\mathcal{J}^*\mathcal{E}$.

Remark 3. As it was mentioned above, all computations were done using the JETS (MAPLE 15), [1, 6], software. We used the IBM Blade cluster consisting of 3 working machines, each of them quad-core Intel Xeon: one E5460, 2.66 GHz, 56 GB RAM two E5430, 2.66 GHz, 16 GB RAM. The classification tree contained 3446 branches, 15 of them were abandoned due to lack of memory. The results presented are due to successful processing of 11 branches only. Other branches are to be processed in future. We also plan to consider LDS of a more general nature.

Remark 4. Actually, presenting results of Proposition 2 in the operator form is not complete and is due to the common tradition only. It is more adequate to deal with them as with Bäcklund transformations between the tangent (see [8]) and cotangent spaces of the equation at hand (cf. [14]), but the operator form seems to be “more visual”.

Acknowledgments

The work of H. Baran and P. Vojčák was supported by RVO funding for IČ47813059. The work of H. Baran was also supported by Czech Science Foundation (GAČR) project P201/11/0356. I. Krasil’shchik was supported by the European Social Fund under the project CZ.1.07/2.3.00/20.0002.

References

- [1] Baran H, Marvan M Jets. A software for differential calculus on jet spaces and diffieties. <http://jets.math.slu.cz>
- [2] Błaszak M 2002 Classical R-matrices on Poisson algebras and related dispersionless systems *Phys. Lett. A* **297** 191–5
- [3] Burovskiy P A, Ferapontov E V, Tsarev S P 2010 Second order quasilinear PDEs and conformal structures on in projective space *Int. J. Math.* **21** 799–841
- [4] Dunajski M 2002 Anti-self-dual four-manifolds with a parallel real spinor *Proc. Roy. Soc. Lond.* **A 458** 1205–1222
- [5] Dunajski M 2004 A class of Einstein-Weil spaces associated to an integrable system of hydrodynamic type *J. Geom. Phys.* **51** 126–37
- [6] Marvan M 2009 Sufficient set of integrability conditions of an orthonomic system *Foundations of Comp. Math.* **9** 651–74
- [7] Marvan M, Sergveyev A 2012 Recursion operators for dispersionless integrable systems in any dimensions *Inverse Problems* **28** 025011
- [8] Krasil’shchik J, Verbovetsky A 2011 Geometry of jet spaces and integrable systems *J. Geom. Phys.* **61** 1633–74
- [9] Krasil’shchik I S, Verbovetsky A M, Vitolo R 2012 A unified approach to computation of integrable structures *Acta Appl. Math.* **120** 199–218

- [10] Martínez Alonso L, Shabat A B 2004 Hydrodynamic reductions and solutions of a universal hierarchy *Theor. Math. Phys.* **140** 1073–85
- [11] Mikhailov A V (Ed) 2009 *Integrability* Lecture Notes in Physics **767** (Berlin: Springer)
- [12] Morozov O I 2009 Contact integrable extensions of symmetry pseudo-groups and coverings of (2+1) dispersionless integrable equations *J. Geom. Phys.* **59** 1461–75
- [13] Morozov O I 2013 A Recursion Operator for the Universal Hierarchy Equation via Cartan’s Method of Equivalence, *Central Euro. J. Math.*, accepted, arXiv:1205.5748 [nlin.SI]
- [14] Morozov O I 2013 A Four-Dimensional Generalization of the Integrable rdDym Equation *Preprint* arXiv:1309.4993 [nlin.SI]
- [15] Ovsienko V 2010 Bi-Hamiltonian nature of the equation $u_{tx} = u_{xy}u_y - u_{yy}u_x$ *Adv. Pure Appl. Math.* **1** 7–17
- [16] Ovsienko V, Roger C 2007 Looped cotangent Virasoro algebra and non-linear integrable systems in dimension $2 + 1$ *Comm. Math. Phys.* **273** 357–88
- [17] Pavlov M V 2003 Integrable hydrodynamic chains *J. Math. Phys.* **44** 4134–56
- [18] Pavlov M V 2006 The Kupershmidt hydrodynamics chains and lattices 2006 *Intern. Math. Research Notes* **2006** 46987
- [19] Vinogradov A M 1984 The \mathcal{C} -spectral sequence, Lagrangian formalism, and conservation laws. I. The linear theory. II. The nonlinear theory *J. Math. Anal. Appl.* **100** 1–129.
- [20] Vinogradov A M 1989 Symmetries and conservation laws of partial differential equations: basic notions and results *Acta Appl. Math.* **15** 3–21
- [21] Zakharevich I 2000 Nonlinear wave equation, nonlinear Riemann problem, and the twistor transform of Veronese webs *Preprint* math-ph/0006001
- [22] Zakharov V E (Ed) 1992 *What is integrability?* Springer series in Nonlinear Dynamics (Berlin: Springer)