

Killing Correspondence in Finsler spaces

Suresh K. Shukla and M. K. Gupta

Department of Pure and Applied Mathematics

Guru Ghasidas Vishwavidyalaya

Bilaspur (C.G.)-495009, INDIA

Email: shuklasureshk@gmail.com; mkgiaps@gmail.com

Abstract

The present paper deals with the Killing correspondence between some Finsler spaces. We consider a Finsler space equipped with a β -change of metric and study the Killing correspondence between the original Finsler space and the Finsler space equipped with β -change of metric. We obtain necessary and sufficient condition for a vector field Killing in the original Finsler space to be Killing in the Finsler space equipped with β -change of metric. Certain consequences of such result are also discussed.

Keywords: Finsler spaces, β -change, Killing vector field

2010 Mathematics Subject Classification: 53B40.

1 Introduction

As a matter of investigation, it is important to observe how properties of a Finsler space change under a change in the metric. Several geometers from different parts of the globe have been working in this direction for the last 2-3 decades. M.S. Knebelman [1], S.Golab [2] and M.Hashiguchi [3] studied conformal change of Finsler metrics. Park and Lee [4] discussed various Randers changes of Finsler spaces with (α, β) -metrics of Douglas type. M. Matsumoto [6] and T. Aikou [5] studied and investigated several properties of projective change and projective Randers change. In 1984, C. Shibata [8] studied β -change of Finsler metrics and discussed certain invariant tensors under such a change.

Killing equations play important role in the study of a Finsler space which undergoes a change in the metric. In fact, they give an equivalent characterization for the transformations to preserve distances. In 1979, Singh, *et. al.* [7] studied a Randers space $F^n \left(M, L(x, y) = (g_{ij}(x) y^i y^j)^{\frac{1}{2}} + b_i(x) y^i \right)$, $n \geq 2$ which undergoes a change $L(x, y) \mapsto L^*(x, y) = L^2(x, y) + (\alpha_i(x) y^i)^2$. They discussed Killing correspondence of the spaces $F^n(M, L)$ and $F^{*n}(M, L^*)$.

In the present paper, we consider a general Finsler space $F^n(M, L)$ which undergoes a β -change, that is $L(x, y) \mapsto \bar{L}(x, y) = f(L, \beta)$, where $\beta(x, y) = b_i(x) y^i$ is a 1-form. We study Killing correspondence of the Finsler spaces $F^n(M, L)$ and $\bar{F}^n(M, \bar{L})$. For the notations and terminology, we refer the reader to the books [9] and [10], and the paper [8] by Shibata.

The paper is organized as follows. In section 2, we give some preliminaries which are used in the discussion of subsequent sections. Section 3 deals with Killing correspondence of $F^n(M, L)$ and $\bar{F}^n(M, \bar{L})$, where $\bar{L}(x, y) = f(L, \beta)$. In section 4, we give conclusion to the results obtained in the paper and discuss future possible work to be done in this direction.

2 Preliminaries

Let $F^n(M, L)$, $n \geq 2$ be an n -dimensional Finsler space. Suppose that the metric function $L(x, y)$ undergoes a change $L(x, y) \mapsto \bar{L}(x, y) = f(L, \beta)$, where $\beta(x, y) = b_i(x) y^i$ is a 1-form and the new space is $\bar{F}^n(M, \bar{L})$. This change of metric is called a β -change (*see* [8] and [9]).

The angular metric tensor \bar{h}_{ij} of the space \bar{F}^n is given by [8]

$$(2.1) \quad \bar{h}_{ij} = p h_{ij} + q_0 m_i m_j,$$

where

$$(2.2) \quad \begin{cases} p = f f_1 / L, & q_0 = f f_{22}, & m_i = b_i - \beta y^i / L^2, \\ f_1 = \partial f / \partial L, & f_2 = \partial f / \partial \beta, \end{cases}$$

h_{ij} being the angular metric tensor of F^n . The fundamental metric tensor \bar{g}_{ij} and its inverse \bar{g}^{ij} of \bar{F}^n are expressed as [8]

$$(2.3) \quad \bar{g}_{ij} = pg_{ij} + p_0b_ib_j + p_{-1}(b_iy_j + b_jy_i) + p_{-2}y_iy_j,$$

$$(2.4) \quad \bar{g}^{ij} = g^{ij}/p - sb^ib^j - s_{-1}(b^iy^j + b^jy^i) - s_{-2}y^iy^j,$$

where

$$(2.5) \quad \left\{ \begin{array}{l} p_0 = q_0 + f_2^2, \\ q_{-1} = f f_{12}/L, \quad p_{-1} = q_{-1} + p f_2/f, \quad q_{-2} = f (f_{11} - f_1/L) /L^2, \\ p_{-2} = q_{-2} + p^2/f^2, \\ b^i = g^{ij}b_j, \quad b^2 = g^{ij}b_ib_j, \quad s_0 = \bar{L}q_0/(\tau pL^2), \\ s_1 = p_{-1}\bar{L}^2/(\tau pL^2), \quad s_{-2} = p_{-1}(\nu pL^2 - b^2\bar{L}^2)/(\tau pL^2\beta), \\ \tau = \bar{L}^2(p + \nu q_0)/L^2, \quad \nu = b^2 - \beta^2/L^2, \end{array} \right.$$

g_{ij} and g^{ij} respectively being the metric tensor and inverse metric tensor of F^n . The Cartan tensor \bar{C}_{ijk} and the associate Cartan tensor \bar{C}_{ij}^h of \bar{F}^n are given by the following expressions:

$$(2.6) \quad \bar{C}_{ijk} = pC_{ijk} + \frac{1}{2}p_{-1}\mathfrak{S}_{(ijk)}\{h_{ij}m_k\} + \frac{1}{2}p_{02}m_im_jm_k,$$

$$(2.7) \quad \bar{C}_{ij}^h = C_{ij}^h - V_{ij}^h,$$

where

$$(2.8) \quad \begin{aligned} V_{ij}^h = & Q^h(pC_{imj}b^m - p_{-1}m_im_j) - \left(\frac{1}{p}m^h - \nu Q^h\right)(p_{02}m_im_j + p_{-2}h_{ij})/2 \\ & - p_{-1}(h_i^hm_j + h_j^hm_i)/(2p), \end{aligned}$$

$$(2.9) \quad Q^h = s_0b^h + s_{-1}y^h, \quad h_i^h = g^{hr}h_{ir}, \quad m^h = g^{hr}m_r, \quad p_{02} = \partial p_0/\partial \beta,$$

$\mathfrak{S}_{(ijk)}$ denote the cyclic sum with respect to the indices i, j and k ; C_{ijk} and C_{ij}^h respectively being the Cartan tensor and associate Cartan tensor of F^n .

The spray coefficients \bar{G}^i of \bar{F}^n in terms of the spray coefficients G^i of F^n are expressed as [8]

$$(2.10) \quad \bar{G}^i = G^i + D^i,$$

where

$$\begin{aligned} D^i &= (q/p) F_0^i + (p E_{00} - 2q F_{r0} b^r)(s_{-1} y^i + s_0 b^i)/2, \\ F_j^i &= g^{ir} F_{rj}, \quad E_{jk} = (1/2)(b_{j|k} + b_{k|j}), \quad F_{jk} = (1/2)(b_{j|k} - b_{k|j}), \end{aligned}$$

the symbol $'|'$ denote the h -covariant derivative with respect to the Cartan connection $C\Gamma$ and the lower index $'_0'$ (except in s_0) denote the contraction by y^i .

The relation between the coefficients \bar{N}_j^i of Cartan nonlinear connection in \bar{F}^n and the coefficients N_j^i of the corresponding Cartan nonlinear connection in F^n is given by [8]

$$(2.11) \quad \bar{N}_j^i = N_j^i + D_j^i,$$

where

$$(2.12) \quad D_j^i = \dot{\partial}_j D^i, \quad \dot{\partial}_j \equiv \partial/\partial y^j.$$

The coefficients \bar{F}_{jk}^i of Cartan connection $C\bar{\Gamma}$ in \bar{F}^n and the coefficients F_{jk}^i of the corresponding Cartan connection $C\Gamma$ in F^n are related as [8]

$$(2.13) \quad \bar{F}_{jk}^i = F_{jk}^i + D_{jk}^i,$$

where

$$\begin{aligned} D_{jk}^i &= \{(1/p)g^{is} - Q^i b^s - y^s(s_{-1} b^i + s_{-2} y^i)\} \\ &\quad (B_{sj} b_{0|k} + B_{sk} b_{0|j} - B_{kj} b_{0|s} + F_{sj} Q_k + F_{sk} Q_j + E_{kj} Q_s + p C_{jkr} D_s^r \\ &\quad + V_{jkr} D_s^r - p C_{skm} D_j^m - V_{sjm} D_k^m - p C_{sjm} D_k^m - V_{skm} D_j^m); \end{aligned}$$

$$B_{kj} = 2\dot{\partial}_j Q_k.$$

The difference tensor D_{jk}^i satisfies the following properties:

$$(i) \quad D_{j0}^i = B_{j0}^i = D_j^i, \quad (ii) \quad D_{00}^i = 2D^i, \quad \text{where } B_{jk}^i = \dot{\partial}_k D_j^i.$$

3 Killing Correspondence of F^n and \bar{F}^n

Let us consider an infinitesimal transformation

$$(3.1) \quad 'x^i = x^i + \epsilon v^i(x),$$

where ϵ is an infinitesimal constant and $v^i(x)$ is a contravariant vector field.

The vector field $v^i(x)$ is said to be a Killing vector field in F^n if the metric tensor of the Finsler space with respect to the infinitesimal transformation (3.1) is Lie invariant, that is

$$(3.2) \quad \mathfrak{L}_v g_{ij} = 0,$$

\mathfrak{L}_v being the operator of Lie differentiation. Equivalently, the vector field $v^i(x)$ is Killing in F^n if

$$(3.3) \quad v_{i|j} + v_{j|i} + 2C_{ij}^h v_{h|0} = 0,$$

where $v_i = g_{il}v^l$.

Now, we prove the following result which gives a necessary and sufficient condition for a Killing vector field in F^n to be Killing in \bar{F}^n :

Theorem 3.1. *A Killing vector field $v^i(x)$ in F^n is Killing in \bar{F}^n if and only if*

$$(3.4) \quad V_{ij}^h v_{h|0} + C_{rjl} v^l D_i^r + C_{ril} v^l D_j^r + v_r D_{ij}^r + \bar{C}_{ij}^h (2C_{rhl} v^l D^r + v_r D_h^r) = 0,$$

where \bar{C}_{ij}^h is the associate Cartan tensor of \bar{F}^n .

Proof. Assume that $v^i(x)$ is Killing in F^n . Then (3.3) is satisfied. By definition, the h -covariant derivatives of v_i with respect to $C\bar{\Gamma}$ and $C\Gamma$ are respectively given as

$$(3.5) \quad (a) \quad v_{i||j} = \partial_j v_i - (\dot{\partial}_r v_i) \bar{G}_j^r - v_r \bar{F}_{ij}^r, \quad (b) \quad v_{i|j} = \partial_j v_i - (\dot{\partial}_r v_i) G_j^r - v_r F_{ij}^r,$$

where $\partial_j \equiv \partial/\partial x^j$ and $'||'$ denote the h -covariant differentiation with respect to $C\bar{\Gamma}$. Equation (3.5)(a), by virtue of (2.10), (2.13) and (3.5)(b), takes the form

$$(3.6) \quad v_{i||j} = v_{i|j} - 2C_{ril} v^l D_j^r - v_r D_{ij}^r.$$

Now, from (3.6), we have

$$(3.7) \quad \begin{aligned} v_{i||j} + v_{j||i} + 2\bar{C}_{ij}^h v_{h||0} = & v_{i|j} + v_{j|i} + 2\bar{C}_{ij}^h v_{h|0} - 2C_{ril} v^l D_j^r - 2C_{rjl} v^l D_i^r \\ & - 2v_r D_{ij}^r - 2\bar{C}_{ij}^h (2C_{rhl} v^l D^r + v_r D_h^r). \end{aligned}$$

Using (2.7) in (3.7) and applying (3.3), we get

$$(3.8) \quad \begin{aligned} v_{i||j} + v_{j||i} + 2\bar{C}_{ij}^h v_{h||0} = & -2V_{ij}^h v_{h|0} - 2C_{ril} v^l D_j^r - 2C_{rjl} v^l D_i^r \\ & - 2v_r D_{ij}^r - 2\bar{C}_{ij}^h (2C_{rhl} v^l D^r + v_r D_h^r). \end{aligned}$$

Proof completes with the observation that $v^i(x)$ is Killing in \bar{F}^n if and only if $v_{i||j} + v_{j||i} + 2\bar{C}_{ij}^h v_{h||0} = 0$, that is, if and only if (3.4) holds. \square

If a vector field $v^i(x)$ is Killing in F^n and \bar{F}^n , then from Theorem 3.1, (3.4) holds, which on transvection by y^i yields

$$(3.9) \quad 2C_{rjl} v^l D^r + v_r D_j^r = 0.$$

Equation (3.4), in view of (3.9), enables us to state the following:

Corollary 3.1. *If a vector field $v^i(x)$ is Killing in F^n and \bar{F}^n , then*

$$(3.10) \quad V_{ij}^h v_{h|0} + C_{rjl} v^l D_i^r + C_{ril} v^l D_j^r + v_r D_{ij}^r = 0.$$

As another important consequence of Theorem 3.1, we have the following:

Corollary 3.2. *If a vector field $v^i(x)$ is Killing in F^n and \bar{F}^n , then the vector $v_i(x, y)$ is orthogonal to the vector $D^i(x, y)$.*

Proof. As $v^i(x)$ is Killing in F^n and \bar{F}^n , (3.4) holds, which on transvection by y^i gives (3.9). Again transvecting (3.9) by y^j , it follows that $v_r D^r = 0$. This proves the result. \square

4 Discussion and Conclusion

We proved Theorem 3.1 as the main result and as its consequences we obtained Corollary 3.1 and Corollary 3.2. Since the Killing equation (3.2) is a necessary and

sufficient condition for the transformation (3.1) to be a motion in F^n (*vide* [10]), the condition (3.4) obtained in Theorem 3.1 may be taken as the necessary and sufficient condition for the vector field $v^i(x)$, generating a motion in F^n , to generate a motion in \bar{F}^n as well. Also, since every motion is an affine motion and every affine motion is a projective motion (*vide* [11]-[13]), it is clear that vector field $v^i(x)$, generating an affine motion (respectively projective motion) in F^n , generates an affine motion (respectively projective motion) in \bar{F}^n if condition (3.4) holds. The main result and its consequences, obtained in the paper, may be further utilized to link various transformations in F^n with corresponding transformations in \bar{F}^n .

References

- [1] M. S. Knebelman, Conformal geometry of generalized metric spaces, *Proc. Nat. Sci., U. S. A.*, **15** (1929), 376-379.
- [2] S. Golab, Einige Bemerkungen uber Winkelmetrik in Finslerschen Raumen, *Verh. Intern. Math. Kongr., Zurich*, **II** (1932), 178-179.
- [3] M. Hashiguchi, On conformal transformations of Finsler metrics, *J. Math. Kyoto Univ.*, **16** (1976), 25-50.
- [4] H. S. Park and I.Y. Lee, The Randers changes of Finsler spaces with (α, β) -metrics of Douglas type, *J. Korean Math. Soc.*, **8(3)** (2001), 503-521.
- [5] T. Aikou, Some remarks on conformal changes of generalized Finsler metrics, *Rep. Fac. Sci., Kagoshima Univ., (Math., Phys. & Chem.)*, **20** (1987), 57-61.
- [6] M. Matsumoto, The Tavakol-van Den Berg conditions in the theories of gravity and projective changes of Finsler metrics, *Publ. Math. Debrecen*, **42** (1993), 155-168.
- [7] U.P. Singh, V. N. John and B. N. Prasad, Finsler spaces preserving Killing vector fields, *J. Math. Phys. Sci. Madras*, **13** (1979), 265-271.

- [8] C. Shibata, On invariant tensors of β -changes of Finsler metrics, *J. Math. Kyoto Univ.*, **24** (1984), 163-188.
- [9] P. L. Antonelli (ed.), *Handbook of Finsler geometry*, Kluwer Acad. Publ., Dordrecht, 2003.
- [10] H. Rund, *The Differential Geometry of Finsler Spaces*, Springer-Verlag, 1959.
- [11] P. N. Pandey, Certain types of affine motion in a Finsler manifold I, II, III. *Colloq. Math.*, **49** (1985), 243-252; **53** (1987), 219-227; **56** (1988), 333-340.
- [12] P. N. Pandey, Certain types of projective motion in a Finsler manifold, *Atti. Accad. Peloritana Pericolanti Cl. Sci. Fis. Mat. Natur.*, **60** (1983) 287-300.
- [13] P. N. Pandey, Certain types of projective motion in a Finsler manifold II, *Atti. Accad. Sci. Torino*, **120** (5-6) (1986), 168-178.