

## STRUCTURE OF FINITE DIHEDRAL GROUP ALGEBRA

F. E. BROCHERO MARTÍNEZ

ABSTRACT. In this article, we show the relation between the irreducible idempotents of the cyclic group algebra  $\mathbb{F}_q C_n$  and the central irreducible idempotents of the group algebras  $\mathbb{F}_q D_{2n}$ , where  $\mathbb{F}_q$  is a finite field with  $q$  elements and  $D_{2n}$  is the dihedral group of order  $2n$ , where  $\gcd(q, n) = 1$ .

In addition, if every divisor of  $n$  divides  $q - 1$ , we show explicitly all central irreducible idempotents of this group algebra and its Wedderburn decomposition.

## 1. INTRODUCTION

Let  $K$  be a field and  $G$  be a group with  $n$  elements. It is known that, if  $\text{char}(K) \nmid n$ , then the group algebra  $KG$  is semisimple and as consequence of Wedderburn Theorem, we have that  $KG$  is isomorphic to a direct sum of matrix algebras over division rings, such that each division algebra is a finite algebra over the field  $K$ , i.e, there exists an isomorphism

$$\rho : KG \rightarrow M_{l_1}(D_1) \oplus M_{l_2}(D_2) \oplus \cdots \oplus M_{l_t}(D_t),$$

where  $D_j$  are division rings such that  $|G| = \sum_{j=1}^t l_j^2 [D_j : K]$ . Observe that  $KG$  has  $t$  central irreducible idempotents, each one of the form

$$e_i = \rho^{-1}(0 \oplus \cdots \oplus 0 \oplus I_i \oplus 0 \cdots \oplus 0),$$

where  $I_i$  is the identity matrix of the component  $M_{l_i}(D_j)$ . Then, the isomorphism  $\rho$  determines explicitly each central irreducible idempotent.

In the case  $K = \mathbb{Q}$ , the calculus of central idempotents and Wedderburn decomposition is widely studied; the classical method to calculate the primitive central idempotents of group algebras depends on computing the character group table. Other method is shown in [8], where Jespers, Leal and Paques describe the central irreducible idempotents when  $G$  is a nilpotent group, using the structure of its subgroups, without employing the characters of the group. Generalizations and improvements of this method can be found in [11], where the authors provide information about the Wedderburn decomposition of  $\mathbb{Q}G$ . This computational method is also used in [2] to compute the Wedderburn decomposition and the primitive central idempotents of a semisimple finite group algebra  $KG$ , where  $G$  is an abelian-by-supersolvable group  $G$  and  $K$  is a finite field.

The structure of  $KG$  when  $G = D_{2n}$  is the dihedral group with  $2n$  elements is well known for  $K = \mathbb{Q}$  (see [7]). In [5], Dutra, Ferraz and Polcino Milies impose conditions over  $q$  and  $n$  in order for  $\mathbb{F}_q D_{2n}$  to have the same number of irreducible components that  $\mathbb{Q}D_{2n}$ . This result is generalized in [6], where Ferraz Goodaire

---

*Date:* September 10, 2018.

*2010 Mathematics Subject Classification.* 20C05 (primary) and 16S34(secondary).

*Key words and phrases.* Central idempotent, Dihedral group.

and Polcino Milies find, for some families of groups, conditions under  $q$  and  $G$  in order for  $\mathbb{F}_q G$  to have the minimum number of simple components.

In this article, assuming that every prime factor of  $n$  divides  $q - 1$ , we show explicitly the central irreducible idempotents of  $\mathbb{F}_q D_{2n}$  and an isomorphism between the group algebra  $\mathbb{F}_q D_{2n}$  and its Wedderburn decomposition. Observe that this isomorphism also shows the structure of  $\mathcal{U}(\mathbb{F}_q D_{2n})$ , the unit group of  $\mathbb{F}_q D_{2n}$ .

## 2. IDEMPOTENTS OF CYCLIC GROUP ALGEBRA

Throughout this article,  $\mathbb{F}_q$  denotes a finite field of order  $q$ , where  $q$  is a power of a prime and  $n$  is a positive integer such that  $\gcd(n, q) = 1$ . For every polynomial  $g(x)$  with  $g(0) \neq 0$ ,  $g^*$  denotes the *reciprocal polynomial* of  $g$ , i.e.,  $g^*(x) = x^{\deg(g)} g(\frac{1}{x})$ . The polynomial  $x^n - 1 \in \mathbb{F}_q[x]$  splits in monic irreducible factors as

$$x^n - 1 = f_1 f_2 \cdots f_r f_{r+1} f_{r+1}^* f_{r+2} f_{r+2}^* \cdots f_{r+s} f_{r+s}^*,$$

where  $f_1 = x - 1$ ,  $f_2 = x + 1$  if  $n$  is even, and  $f_j^* = f_j$  for  $2 \leq j \leq r$ , where  $r$  is the number of auto-reciprocal factors in the factorization and  $2s$  the number of non-auto-reciprocal factors.

We denote by  $C_n$  the cyclic group of order  $n$ . It is well known that  $\mathbb{F}_q C_n \simeq \mathcal{R}_n = \frac{\mathbb{F}_q[x]}{\langle x^n - 1 \rangle}$ , and by the Chinese Remainder Theorem

$$\frac{\mathbb{F}_q[x]}{\langle x^n - 1 \rangle} \simeq \bigoplus_{j=1}^{r+s} \frac{\mathbb{F}_q[x]}{\langle f_j \rangle} \oplus \bigoplus_{j=r+1}^{r+s} \frac{\mathbb{F}_q[x]}{\langle f_j^* \rangle}$$

is exactly the Wedderburn decomposition of the group algebra  $\mathcal{R}_n$ , so every primitive idempotent generates a maximal ideal of  $\mathcal{R}_n$  and also one component of this direct sum.

In addition, since  $\mathcal{R}_n$  is a principal ideal domain, every ideal of  $\mathcal{R}_n$  is generated by a polynomial  $g$  that is a divisor of  $x^n - 1$ . The relation between the generator of the ideal and its principal idempotent is shown in the following lemma.

**Lemma 2.1.** *Let  $\mathcal{I} \subset \mathcal{R}_n$  be an ideal generated by the monic polynomial  $g$ , that is divisor of  $x^n - 1$ , and define  $f = \frac{x^n - 1}{g}$ . Then the principal idempotent of  $\mathcal{I}$  is*

$$e_f = -\frac{((f^*)')^*}{n} \cdot \frac{x^n - 1}{f}.$$

*Proof:* Let  $t$  be an integer such that  $n$  divides  $q^t - 1$ . By Theorem 2.1 in [1] (see also Theorem 3.4 in [3]), every primitive idempotent of  $\frac{\mathbb{F}_t[x]}{\langle x^n - 1 \rangle}$  is given by

$$u_\lambda = \frac{\lambda}{n} \cdot \frac{x^n - 1}{x - \lambda} = \frac{1}{n} \sum_{l=0}^{n-1} \lambda^{-l} x^l$$

where  $\lambda^n = 1$ .

Since  $f$  divides  $x^n - 1$ , then  $f$  splits in  $\mathbb{F}_{q^t}[x]$  as  $(x - \lambda_1) \cdots (x - \lambda_k)$  and

$$(f^*)' = \sum_{i=1}^k (-\lambda_i) \prod_{i \neq j} (1 - \lambda_j x) = f^* \sum_{i=1}^k \frac{-\lambda_i}{x - \lambda_j x},$$

hence

$$e_f = -\frac{((f^*)')^*}{n} \cdot \frac{x^n - 1}{f} = \sum_{i=1}^k \frac{\lambda_i}{n} \cdot \frac{x^n - 1}{x - \lambda} = \sum_{i=1}^k u_{\lambda_i}.$$

Therefore  $e_f$  is an idempotent of  $\mathbb{F}_q[x]$ . In order to prove that  $e_f$  is the principal idempotent of  $\mathcal{I}$ , it is enough to show that  $g \cdot e_f = g$ . Observe that, using partial fraction decomposition we obtain

$$g = \frac{x^n - 1}{f} = \sum_{i=1}^k A_i u_{\lambda_i},$$

where  $A_i = \frac{n}{\lambda_i} \frac{1}{\prod_{j \neq i} (\lambda_i - \lambda_j)}$  and then

$$g \cdot e_f = \sum_{i=1}^k A_i u_{\lambda_i} \cdot \sum_{j=1}^k u_{\lambda_j} = \sum_{1 \leq i, j \leq k} A_i u_{\lambda_i} u_{\lambda_j} = \sum_{i=1}^k A_i u_{\lambda_i} = g$$

as we wanted to prove.  $\square$

**Remark 2.2.** *This lemma is also true for fields with characteristic zero, it suffices to change in the proof the field  $\mathbb{F}_q$  by the splitting field of the polynomial  $f$ .*

**Corollary 2.3.** *The cyclic group ring  $\mathcal{R}_n$  has  $r + 2s$  irreducible idempotents of the form  $e_f$  given by Lemma 2.1, where the polynomials  $f$ 's are the irreducible factors of  $x^n - 1 \in \mathbb{F}_q[x]$ .*

### 3. CENTRAL IDEMPOTENTS OF DIHEDRAL GROUP ALGEBRA

Throughout this section,  $\alpha_j$  denotes a root of the polynomial  $f_j$  and  $D_{2n}$  denotes the dihedral group of order  $2n$ , i.e.

$$D_{2n} = \langle x, y \mid x^n = 1, y^2 = 1, xy = yx^{-1} \rangle.$$

We define integer numbers  $\epsilon$  and  $\delta$  as

$$\epsilon = \begin{cases} 0 & \text{if } \text{char}(q) = 2 \\ 1 & \text{if } \text{char}(q) \neq 2 \text{ and } n \text{ is odd} \\ 2 & \text{if } \text{char}(q) \neq 2 \text{ and } n \text{ is even} \end{cases}$$

and  $\delta = \max\{\epsilon, 1\}$ .

The following theorem shows explicitly the dependence of the Wedderburn decomposition of the Dihedral group algebra over a finite field  $\mathbb{F}_q$  with the factorization of  $x^n - 1 \in \mathbb{F}_q[x]$ .

**Theorem 3.1.** *The group algebra  $\mathbb{F}_q D_{2n}$  has Wedderburn decomposition of the form*

$$\mathbb{F}_q D_{2n} \cong \bigoplus_{j=1}^{r+s} A_j$$

where

$$A_j = \begin{cases} \mathbb{F}_q \oplus \mathbb{F}_q & \text{if } j \leq \delta, \\ M_2(\mathbb{F}_q[\alpha_j + \alpha_j^{-1}]) & \text{if } \delta + 1 \leq j \leq r, \\ M_2(\mathbb{F}_q[\alpha_j]) & \text{if } r + 1 \leq j \leq r + s, \end{cases}$$

*Proof:* For each  $j \in \{1, \dots, s + r\}$ , let  $\tau_j$  be the homomorphism of  $\mathbb{F}_q$ -algebras defined by the generators of the group  $D_{2n}$  as

$$\begin{aligned} \tau_1 : \mathbb{F}_q D_{2n} &\rightarrow \mathbb{F}_q \oplus \mathbb{F}_q \\ x &\mapsto (1, 1) \\ y &\mapsto (1, -1), \end{aligned}$$

in the case  $\epsilon \geq 1$  and

$$\begin{aligned} \tau_2 : \mathbb{F}_q D_{2n} &\rightarrow \mathbb{F}_q \oplus \mathbb{F}_q \\ x &\mapsto (-1, -1) \\ y &\mapsto (1, -1), \end{aligned}$$

in the case  $\epsilon = 2$ , where the sum and product in  $\mathbb{F}_q \oplus \mathbb{F}_q$  is defined by adding and multiplying the corresponding components of the same coordinates. Finally, for every  $j \geq \epsilon + 1$

$$\begin{aligned} \tau_j : \mathbb{F}_q D_{2n} &\rightarrow M_2(\mathbb{F}_q[\alpha_j]) \\ x &\mapsto \begin{pmatrix} \alpha_j & 0 \\ 0 & \alpha_j^{-1} \end{pmatrix} \\ y &\mapsto \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \end{aligned}$$

It is easy to prove that  $(\tau_j(x))^n = I$ ,  $(\tau_j(y))^2 = I$  and  $\tau_j(x)\tau_j(y) = \tau_j(y)\tau_j(x)^{-1}$ .

Observe that in the case of characteristic 2, i.e.  $\epsilon = 0$ , we have that

$$\text{img}(\tau_1) = \left\{ \begin{pmatrix} a & b \\ b & a \end{pmatrix} \mid a, b \in \mathbb{F}_q \right\},$$

that is isomorphic to  $\mathbb{F}_q \oplus \mathbb{F}_q$  by the projection  $\sigma_0$  over the first row of the matrix, where the product is defined by  $(a, b) \cdot (c, d) = (ac + bd, ad + bc)$ . Thus,  $\dim_{\mathbb{F}_q}(\text{img}(\tau_1)) = 2$  in all cases. In addition, if  $n$  is even, then  $\dim_{\mathbb{F}_q}(\text{img}(\tau_2)) = 2$ .

For each  $\delta < j \leq r$ , if we define  $Z_j = \begin{pmatrix} 1 & -\alpha_j \\ 1 & -\alpha_j^{-1} \end{pmatrix}$ , then

$$\begin{aligned} \sigma_j : M_2(\mathbb{F}_q[\alpha_j]) &\rightarrow M_2(\mathbb{F}_q[\alpha_j]) \\ X &\mapsto Z_j^{-1} X Z_j \end{aligned}$$

is an automorphism such that

$$\sigma_j \circ \tau_j(x) = \begin{pmatrix} 0 & 1 \\ -1 & \alpha_j + \alpha_j^{-1} \end{pmatrix} \quad \text{and} \quad \sigma_j \circ \tau_j(y) = \begin{pmatrix} 1 & -(\alpha_j + \alpha_j^{-1}) \\ 0 & -1 \end{pmatrix},$$

so the images of the generators of  $D_n$  are in  $\mathbb{F}_q(\alpha_j + \alpha_j^{-1})$ . It follows that for each  $j$  such that  $\delta < j \leq r$  we have

$$\dim_{\mathbb{F}_q}(\text{img}(\tau_j)) = \dim_{\mathbb{F}_q}(\text{img}(\sigma_j \circ \tau_j)) \leq 4 \dim_{\mathbb{F}_q}(\mathbb{F}_q(\alpha_j + \alpha_j^{-1})) = 2 \deg(f_j)$$

and in the case  $r + 1 \leq j \leq r + s$ , we know that

$$\dim_{\mathbb{F}_q} \text{img}(\tau_j) \leq 4 \dim_{\mathbb{F}_q} \mathbb{F}_q(\alpha_j) = 4 \deg(f_j).$$

Now, let  $\tau$  be the homomorphism of  $\mathbb{F}_q$ -algebras defined by  $\bigoplus_{j=1}^{s+r} \tau_j$ . Observe that this homomorphism is injective. In fact, let  $u$  be an element of  $\mathbb{F}_q D_n$  such that  $\tau(u) = 0$ . If we write  $u = P_1(x) + P_2(x)y$ , where  $P_1$  and  $P_2$  are polynomials of degree less than  $n$ , for each  $j > \epsilon$ , we have

$$\tau_j(u) = \begin{pmatrix} P_1(\alpha_j) & P_2(\alpha_j) \\ P_2(\alpha_j^{-1}) & P_1(\alpha_j^{-1}) \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix},$$

so,  $P_1(\alpha_j) = P_1(\alpha_j^{-1}) = 0$  and  $P_2(\alpha_j) = P_2(\alpha_j^{-1}) = 0$ . In addition, if  $\epsilon \geq 1$ , then

$$\tau_1(u) = (P_1(1) + P_2(1), P_1(1) - P_2(1)) = 0,$$

and if  $\epsilon = 2$  we have

$$\tau_2(u) = (P_1(-1) + P_2(-1), P_1(-1) - P_2(-1)) = 0.$$

It follows that  $P_1$  and  $P_2$  are divisible by the polynomial  $x^n - 1$  and since the degrees of these polynomials are less than  $n$ , we conclude that  $P_1$  and  $P_2$  are null polynomials and therefore  $\tau$  is an injective homomorphism.

Finally, we observe that the homomorphism  $\rho : \mathbb{F}_q D_{2n} \rightarrow \bigoplus_{j=1}^{r+s} A_j$  defined by  $\rho = \bigoplus_{j=1}^{r+s} \rho_j$  where  $\rho_j = \begin{cases} \sigma_j \circ \tau_j & \text{if } \epsilon < j \leq r \\ \tau_j & \text{otherwise} \end{cases}$  is injective. Furthermore,  $\dim_{\mathbb{F}_q}(\mathbb{F}_q D_n) = 2n$  and

$$\begin{aligned} \dim_{\mathbb{F}_q} \left( \bigoplus_{j=1}^{r+s} A_j \right) &= 2\epsilon + 4 \sum_{j=\epsilon+1}^r \dim_{\mathbb{F}_q}(\mathbb{F}_q[\alpha_j + \alpha_j^{-1}]) + 4 \sum_{j=r+1}^{r+s} (\mathbb{F}_q[\alpha_j]) \\ &= 2\epsilon + 2 \sum_{j=\epsilon+1}^r \deg(f_j) + 4 \sum_{j=r+1}^{r+s} \deg(f_j) \\ &= 2 \deg(x^n - 1) = 2n. \end{aligned}$$

Therefore  $\rho$  is an isomorphism.  $\square$

**Remark 3.2.** *In the proof of the theorem we use the following facts: if  $\beta$  is a root of the polynomial  $g \in \mathbb{F}_q[x]$ , then  $\beta^{-1}$  is root of the polynomial  $g^*$ . In addition, when  $g$  is auto-reciprocal and  $\pm 1$  are not roots of  $g$ , there exists a polynomial  $h \in \mathbb{F}_q[x]$  of degree  $\frac{\deg(g)}{2}$ , such that  $\beta$  is a root of  $g$  if and only if  $\beta + \beta^{-1}$  is a root of  $h$ . In fact, since  $g$  is symmetrical, we can write  $g$  as*

$$g(x) = \sum_{j=0}^t a_j(x^{t+j} + x^{t-j}) = x^t \sum_{j=0}^t a_j(x^j + x^{-j}) = x^t \sum_{j=0}^t a_j D_j(z, 1) = x^t h(z)$$

where  $D_j$  is the Dickson polynomial of degree  $j$  and  $z = x + x^{-1}$  (see [9] or [10]).

**Theorem 3.3.** *The dihedral group algebra  $\mathbb{F}_q D_{2n}$  has  $\epsilon + r + s$  central irreducible idempotents:*

- (1)  $2\epsilon$  idempotents of the form  $\frac{1+y}{2}e_{f_j}$  and  $\frac{1-y}{2}e_{f_j}$ , where  $j \leq \epsilon$ .
- (2)  $r - \epsilon$  idempotents  $e_{f_j}$ , where  $j = \epsilon+1, \dots, r$ , generated by the auto-reciprocals factor of  $x^n - 1$ .
- (3)  $s$  idempotents  $e_{f_j} + e_{f_j^*}$ , where  $j = r+1, \dots, r+s$ .

*Proof:* Since the homomorphism  $\tau$  in the proof of Theorem 3.1 is injective, then the image of a central primitive idempotent  $u$  by the homomorphism has to be zero in every component, except for one component where the image is the identity, i.e., for some  $i$  fixed,  $\tau_j(u) = \delta_{i,j} I_j$ , where  $I_j$  is the identity over the component  $A_j$ . Let  $u = P(x) + Q(x)y$  be a representation of  $u$ , where  $P$  and  $Q$  are polynomials in  $\mathbb{F}_q[x]$  of degree less than or equal to  $n-1$ . Observe that  $Q$  is zero when calculated at each root of the polynomial  $x^n - 1 = 0$ , so  $Q$  is the null polynomial. In addition,  $P$  is one when we calculate it at the roots of the polynomials  $f_j$  and  $f_j^*$  and zero when we calculate it at the other roots of the polynomial  $x^n - 1$ . The unique polynomial of degree less or equal to  $n-1$  that satisfies that properties is  $e_{f_j}$ , when  $f_j = f_j^*$  and  $e_{f_j} + e_{f_j^*}$  when  $f_j \neq f_j^*$ . Finally, if  $j \leq \epsilon$  the image  $\tau_j(e_{f_j}) = (1, 1)$  is not a primitive idempotent, and we can decompose this idempotent in two central primitive idempotents,  $(\frac{1+y}{2})e_{f_j}$  and  $(\frac{1-y}{2})e_{f_j}$ , such that  $\tau_j((\frac{1+y}{2})e_{f_j}) = (1, 0)$  and  $\tau_j((\frac{1-y}{2})e_{f_j}) = (0, 1)$ .  $\square$

4. EXPLICIT FORM OF THE IDEMPOTENTS WHEN  $\text{rad}(n)|(q-1)$ 

Throughout this section, we assume that every prime factor of  $n$  divides  $q-1$ ,  $\kappa$  and  $\nu$  denote the numbers  $\text{gcd}(n, q-1)$  and  $\min\{\nu_2(\frac{n}{2}), \nu_2(q+1)\}$  respectively,  $\theta$  and  $\alpha$  are generators of  $\mathbb{F}_q^*$  and  $\mathbb{F}_{q^2}^*$  such that  $\alpha^{q+1} = \theta$ . In the following results, we show the explicit form of the idempotents of the cyclic group algebra  $\mathbb{F}_q C_n$  and the Wedderburn decomposition of the Dihedral group algebra  $\mathbb{F}_q D_{2n}$ . In order to show that representation, we need the following lemma

**Lemma 4.1.** [4, Corollary 3.3 and Corollary 3.6] *The factorization of  $x^n - 1$  in irreducible factors of  $\mathbb{F}_q[x]$  depends on  $n$  and  $q$  in the following form:*

(i) *If  $8 \nmid n$  or  $q \not\equiv 3 \pmod{4}$ , then*

$$x^n - 1 = \prod_{t|m} \prod_{\substack{1 \leq u \leq \text{gcd}(n, q-1) \\ \text{gcd}(u, t)=1}} (x^t - \theta^{ul}),$$

where  $m = \frac{n}{\kappa}$  and  $l = \frac{q-1}{\kappa}$ . In addition, for each  $t$  such that  $t|m$ , the number of irreducible factors of degree  $t$  is  $\frac{\varphi(t)}{t} \cdot \kappa$ , where  $\varphi$  denotes the Euler Totient function.

(ii) *If  $8 | n$  and  $q \equiv 3 \pmod{4}$ , then*

$$x^n - 1 = \prod_{\substack{t|m' \\ t \text{ odd}}} \prod_{\substack{1 \leq w \leq \kappa \\ \text{gcd}(w, t)=1}} (x^t - \theta^{wl}) \cdot \prod_{t|m'} \prod_{u \in \mathcal{S}_t} (x^{2t} - (\alpha^{ul'} + \alpha^{qu l'})x^t + \theta^{ul'}),$$

where  $m' = \frac{n}{2^\nu \kappa}$ ,  $l' = \frac{q^2-1}{2^\nu \kappa}$ , and  $\mathcal{S}_t$  is the set

$$\left\{ u \in \mathbb{N} \mid \begin{array}{l} 1 \leq u \leq 2^\nu \kappa, \text{gcd}(u, t) = 1 \\ 2^\nu \nmid u \text{ and } u < \{qu\}_{2^\nu \kappa} \end{array} \right\},$$

where  $\{a\}_b$  denotes the remainder of the division of  $a$  by  $b$ , i.e. the number  $0 \leq c < b$  such that  $a \equiv c \pmod{b}$ . In addition, for each  $t$  odd such that  $t|m'$ , the number of irreducible binomials of degree  $t$  and  $2t$  is  $\frac{\kappa \cdot \varphi(t)}{t}$  and  $\frac{\kappa \cdot \varphi(t)}{2t}$  respectively, and the number of irreducible trinomials of degree  $2t$  is

$$\begin{cases} \frac{\varphi(t)}{t} \cdot 2^{\nu-1} \kappa, & \text{if } t \text{ is even} \\ \frac{\varphi(t)}{t} \cdot (2^{\nu-1} - 1) \kappa, & \text{if } t \text{ is odd.} \end{cases}$$

The following corollary, direct from Lemmas 2.1 and 4.1, shows the explicit form of each idempotent of the cyclic group algebra  $\mathbb{F}_q C_n$  when  $\text{rad}(n)|(q-1)$ .

**Corollary 4.2.** *Let  $m$ ,  $m'$ ,  $l$  and  $l'$  be as in Lemma 4.1.*

(1) *If  $8 \nmid n$  or  $n \not\equiv 3 \pmod{4}$ , then every irreducible idempotent of the ring  $\mathcal{R}_n$  is of the form*

$$e_{t,ul} = \frac{\theta^{ul} t}{n} \cdot \frac{x^n - 1}{x^t - \theta^{ul}}$$

where  $t$  and  $u$  satisfy the condition of Lemma 4.1 item (i).

(2) *If  $8 | n$  or  $n \equiv 3 \pmod{4}$ , then every irreducible idempotent of the ring  $\mathcal{R}_n$  is of the form shown in (1) and of the form*

$$e_{t,ul'} = \frac{t}{n} \left( (\alpha^{ul'} + \alpha^{ul'q})x^t - 2\theta^{ul'} \right) \frac{x^n - 1}{(x^{2t} - (\alpha^{ul'} + \alpha^{qu l'})x^t + \theta^{ul'})},$$

where  $t$  and  $u$  satisfy the condition of Lemma 4.1 item (ii).

**Remark 4.3.** By Theorem 3.3 If

- $\text{char}(\mathbb{F}_q) = 2$ , or
- $n$  is odd and  $\theta^{ul} \neq 1$ , or
- $n$  is even and  $\theta^{ul} \neq \pm 1$ ,

then every idempotent found in Corollary 4.2 item (1) is a central irreducible idempotent of  $\mathbb{F}_q D_{2n}$ . Otherwise, the idempotent can be reduced to two central primitive idempotents  $\frac{1+y}{2}e_{t,ul}$  and  $\frac{1-y}{2}e_{t,ul}$ .

In addition,  $e_{t,ul'}$  of item (2) is also a central irreducible idempotent of  $\mathbb{F}_q D_{2n}$  if  $\theta^{ul'} = 1$ , otherwise, the central irreducible idempotent is  $e_{t,ul'} + e_{t,-ul'}$ .

**Theorem 4.4.** The Wedderburn decomposition of the group algebra  $\mathbb{F}_q D_{2n}$  depends on  $n$  and  $q$  in the following form:

(1) When  $n$  is odd, the decomposition is

$$2\mathbb{F}_q \oplus \frac{\kappa - 1}{2} M_2(\mathbb{F}_q) \oplus \bigoplus_{\substack{t|m \\ t \neq 1}} \frac{\kappa \cdot \varphi(t)}{2t} M_2(\mathbb{F}_{q^t}).$$

(2) When  $n$  is even,

(2.1) if  $q \equiv 1 \pmod{4}$  or  $8 \nmid n$ , the decomposition is

$$4\mathbb{F}_q \oplus \left(\frac{\kappa}{2} - 1\right) M_2(\mathbb{F}_q) \oplus \bigoplus_{\substack{t|m \\ t \neq 1}} \frac{\kappa \cdot \varphi(t)}{2t} M_2(\mathbb{F}_{q^t}),$$

(2.2) if  $q \equiv 3 \pmod{4}$  and  $8|n$ , the decomposition is

$$\begin{aligned} & 4\mathbb{F}_q \oplus (\kappa + 2^{\nu-1} - 3) M_2(\mathbb{F}_q) \oplus (2^{\nu-2}\kappa - 2^{\nu-1} - \frac{\kappa}{4} + 1) M_2(\mathbb{F}_{q^2}) \oplus \bigoplus_{\substack{t|m' \\ t \text{ odd} \\ t \neq 1}} \frac{\kappa \cdot \varphi(t)}{2t} M_2(\mathbb{F}_{q^t}) \\ & \oplus \bigoplus_{\substack{t|m' \\ t \text{ even}}} \frac{2^{\nu-2}\kappa \cdot \varphi(t)}{t} M_2(\mathbb{F}_{q^{2t}}) \oplus \bigoplus_{\substack{t|m' \\ t \text{ odd} \\ t \neq 1}} \frac{(2^{\nu-1} - 1)\kappa \cdot \varphi(t)}{2t} M_2(\mathbb{F}_{q^{2t}}). \end{aligned}$$

$$\text{where } i = \begin{cases} 0 & \text{if } \nu_2(q+1) > \nu_2(\frac{n}{2}) \\ 1 & \text{if } \nu_2(q+1) \leq \nu_2(\frac{n}{2}). \end{cases}$$

*Proof:* First, we consider the case  $n \not\equiv 3 \pmod{4}$  or  $8 \nmid n$ , so every irreducible factor of  $x^n - 1$  is a binomial, and except for the factors  $x - 1$  and  $x + 1$ , we have that any irreducible factor of the form  $x^t - a$  is not auto-reciprocal. Thus, we have two cases to analyse:

- i) If  $n$  is odd, we have that  $\epsilon = 0$  or  $1$  and  $r = 1$ . By Lemma 4.1 there exist  $\frac{\kappa \varphi(t)}{t}$  irreducible factors of degree  $t$  and by Theorem 3.3 there exist two components isomorphic to  $\mathbb{F}_q$ ,  $\frac{\kappa \varphi(t)}{2t}$  components of the form  $M_2(\mathbb{F}_{q^t})$  if  $t > 1$  and  $\frac{\kappa - 1}{2}$  components of the form  $M_2(\mathbb{F}_q)$  if  $t = 1$ , where  $t$  is a divisor of  $m$ . So we obtain item (1).
- ii) If  $n$  is even, we have that  $\epsilon = 2$  and there exist four components isomorphic to  $\mathbb{F}_q$ . In addition, every factor of  $x^n - 1$  different that  $x \pm 1$  is a non-auto-reciprocal binomial, then  $r = 2$ , and by the same argument of the previous case there exist  $\frac{\kappa \varphi(t)}{2t}$  components of the form  $M_2(\mathbb{F}_{q^t})$  if  $t > 1$  and  $\frac{\kappa - 2}{2}$  components of the form  $M_2(\mathbb{F}_q)$  if  $t = 1$ , where  $t$  is a divisor of  $m$ . So, we obtain item (2.1).

Finally, in the case which  $q \equiv 3 \pmod{4}$  and  $8|n$ , every factor of  $x^n - 1$  is a binomial or a trinomial. The unique auto-reciprocal factor of the form  $x^t - a$  with  $t$  odd is  $f_1 = x - 1$ . Now, suppose that  $x^{2t} - (\alpha^{ul'} + \alpha^{qu'l'})x^t + \theta^{ul'}$  is an irreducible factor of  $x^n - 1$  as in Lemma 4.1 item (b), such that it is an auto-reciprocal polynomial. It follows that  $\theta^{ul'} = 1$  and therefore  $(q-1)|ul'$ . Since

$$l' = \frac{q-1}{\gcd(n, q-1)} \cdot \frac{q+1}{2^\nu},$$

the polynomial is auto-reciprocal when  $\gcd(n, q-1)|u \cdot \frac{q+1}{2^\nu}$  and we have two cases to consider:

- i) If  $\nu_2(q+1) \leq \nu_2(\frac{n}{2})$  then  $\frac{q+1}{2^\nu}$  is odd and  $\gcd(\gcd(n, q-1), \frac{q+1}{2^\nu}) = 1$ , therefore  $\gcd(n, q-1)|u$ . But  $t|m'|n$  and  $\gcd(t, u) = 1$ , then these conditions imply that  $t = 1$  and  $u$  is a multiple of  $\gcd(n, q-1)$  not divisible by  $2^\nu$  and less than  $2^\nu \gcd(n, q-1)$ . So there exist  $2^\nu - 2$  values of  $u$  that generate  $2^{\nu-1} - 1$  auto-reciprocal factors, all of them of degree 2, each one generating a component of the form  $M_2(\mathbb{F}_q)$ . In addition, we have  $\kappa - 2$  irreducible factors of degree 1, each one generating a component of the same type.

Therefore there exist  $(\kappa - 2) + (2^{\nu-1} - 1) = \kappa + 2^{\nu-1} - 3$  components  $M_2(\mathbb{F}_q)$  and

$$\frac{k}{4}(2^\nu - 1) - (2^{\nu-1} - 1) = 2^{\nu-2}\kappa - 2^{\nu-1} - \frac{k}{4} + 1$$

components  $M_2(\mathbb{F}_{q^2})$ .

- ii) If  $\nu_2(q+1) > \nu_2(\frac{n}{2})$  then  $\frac{q+1}{2^\nu}$  is even and  $\gcd(\gcd(n, q-1), \frac{q+1}{2^\nu}) = 2$ , therefore  $\frac{1}{2}\gcd(n, q-1)|u$ . Similarly, we obtain  $t = 1$  and  $u$  is a multiple of  $\frac{1}{2}\gcd(n, q-1)$  non divisible by  $2^\nu$  and less than  $2^\nu \gcd(n, q-1)$ . So there exist  $2^{\nu+1} - 2$  values of  $u$  and then  $2^\nu - 1$  auto-reciprocal factors, all of them of degree 2.

Then there exist  $\kappa + 2^\nu - 3$  components  $M_2(\mathbb{F}_q)$  and  $2^{\nu-2}\kappa - 2^\nu - \frac{k}{4} + 1$  components  $M_2(\mathbb{F}_{q^2})$ .  $\square$

## REFERENCES

- [1] Arora, S. K., Pruthi M. *Minimal Codes of Prime-Power Length*. Finite Fields Appl. **3** (1997) 99-113
- [2] Broche, O., del Río, Á., *Wedderburn decomposition of finite group algebras*, Finite Fields Appl. **13** (2007) 71-79.
- [3] Brochero Martínez, F.E., Giraldo Vergara, C.R., *Explicit Idempotents of Finite Group Algebra* Finite Fields Appl. **28** (2014) 123-131
- [4] Brochero Martínez, F.E., Giraldo Vergara, C.R., Batista de Oliveira, L., *Explicit Factorization of  $x^n - 1 \in \mathbb{F}_q[x]$* , submitted for publication in Designs, Codes and Cryptography. Preprint available on <http://arxiv.org/abs/1404.6281>
- [5] Dutra, F., Ferraz, R., Polcino Milies, C., *Semisimple group codes and dihedral codes*, Algebra Discrete Math. **3** (2009), 28-48.
- [6] Ferraz, R., Goodaire, E., Polcino Milies, C., *Some classes of semisimple group (and loop) algebras over finite fields*, J. of Algebra **324** (2010) 3457-3469
- [7] Giraldo Vergara, C.R., Brochero Martínez, F. E. *Wedderburn decomposition of some special rational group algebras*. Lect. Mat. **23**, no. 2, (2002) 99-106.
- [8] Jespers, E., Leal, G., Paques, A., *Central idempotents in rational group algebras of finite nilpotent groups*, J. Algebra Appl. **2** (1) (2003) 57-62.
- [9] Lidl, R. Mullen, G. L., Turnwald, G., *Dickson polynomials*, Pitman Monographs and Surveys in Pure and Applied Math., Longman, London-Harlow-Essex, 1993.
- [10] Meyn H. *Factorization of the cyclotomic polynomials  $x^{2^n} + 1$  over finite fields*. Finite Fields Appl. **2**, (1996) 439-442



- [11] Olivieri, A., del Río, Á, Simón, J.J., *On monomial characters and central idempotents of rational group algebras*, *Comm. Algebra* **32** (4) (2004) 1531-1550.

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE MINAS GERAIS, UFMG, BELO  
HORIZONTE, MG, 30123-970, BRAZIL,  
*E-mail address:* `fbrocher@mat.ufmg.br`