

BOHR–SOMMERFELD LAGRANGIANS OF MODULI SPACES OF HIGGS BUNDLES

INDRANIL BISWAS, NIELS LETH GAMMELGAARD, AND MARINA LOGARES

ABSTRACT. Let X be a compact connected Riemann surface of genus at least two. Let $M_H(r, d)$ denote the moduli space of semistable Higgs bundles on X of rank r and degree d . We prove that the compact complex Bohr–Sommerfeld Lagrangians of $M_H(r, d)$ are precisely the irreducible components of the nilpotent cone in $M_H(r, d)$. This generalizes to Higgs G –bundles and also to the parabolic Higgs bundles.

1. INTRODUCTION

Let $M_H(r, d)$ be the moduli space of semistable Higgs bundles of rank r and degree d on a compact connected Riemann surface X of genus g at least two. It is an irreducible normal complex projective variety of complex dimension $2(r^2(g-1) + 1)$. This moduli space is equipped with an algebraic symplectic form. In fact, there is a canonical algebraic one-form ω on $M_H(r, d)$ such that $d\omega$ is the symplectic form. Let

$$\mathcal{L} := \mathcal{O}_{M_H(r, d)} = M_H(r, d) \times \mathbb{C}$$

be the trivial holomorphic line bundle on $M_H(r, d)$. Consider the holomorphic connection

$$D := d + \omega$$

on \mathcal{L} , where d denotes the de Rham differential on functions on $M_H(r, d)$. We note that the curvature of D is $d\omega$.

A compact Lagrangian on $M_H(r, d)$ is a reduced irreducible compact complex analytic subset

$$\mathbb{L} \subset M_H(r, d)$$

of dimension $(\dim M_H(r, d))/2 = r^2(g-1) + 1$ such that the restriction of (\mathcal{L}, D) to \mathbb{L} is a flat line bundle. Indeed, this follows immediately from the fact that the curvature of D coincides with the symplectic form on $M_H(r, d)$. A compact Bohr–Sommerfeld Lagrangian on $M_H(r, d)$ is a reduced irreducible compact complex analytic subset $\mathbb{L} \subset M_H(r, d)$ of dimension $r^2(g-1) + 1$ such that \mathcal{L} admits a nonzero flat section over \mathbb{L} . Clearly, a Bohr–Sommerfeld Lagrangian is a Lagrangian.

A flat section of \mathcal{L} over \mathbb{L} gives a holomorphic function on \mathbb{L} because the form ω is holomorphic, and \mathcal{L} coincides with $\mathcal{O}_{M_H(r, d)}$. On the other hand, \mathbb{L} does not admit any nonconstant holomorphic function because it is compact and irreducible. A constant function is a flat section of $\mathcal{L}|_{\mathbb{L}}$ if and only if the pullback of the connection D to $\mathcal{L}|_{\mathbb{L}}$ is

2000 *Mathematics Subject Classification.* 14H60, 14H70, 53D12.

Key words and phrases. Bohr–Sommerfeld Lagrangian, Higgs bundle, \mathbb{C}^* -action, nilpotent cone.

the de Rham differential. Consequently, a reduced irreducible compact complex analytic subset

$$\iota : \mathbb{L} \hookrightarrow M_H(r, d)$$

is a Bohr–Sommerfeld Lagrangian if and only if $\iota^*\omega = 0$.

The Hitchin map

$$\mathcal{H} : M_H(r, d) \longrightarrow \bigoplus_{i=1}^r H^0(X, K_X^{\otimes i})$$

sends any (E, θ) to $\sum_{i=1}^r \text{trace}(\theta^i)$. The fiber of \mathcal{H} over 0 is the nilpotent cone.

We prove the following:

Theorem 1.1. *The compact Bohr–Sommerfeld Lagrangians in $M_H(r, d)$ are precisely the irreducible components of the nilpotent cone.*

Theorem 1.1 generalizes to other contexts; see Section 3.1 and Section 3.2.

Quantization is a procedure which associates to a symplectic manifold M a Hilbert space $Q(M)$. When using geometric quantization, the quantum space is constructed from sections or higher cohomology groups of an appropriate complex line bundle, known as the prequantization bundle, equipped with a connection whose curvature equals the symplectic form. The interesting sections of the prequantization bundle are those provided by a choice of a polarization of M . For real polarizations a theorem by Sniaticky [Sn] determines the dimension of the quantized space in terms of the number of Bohr–Sommerfeld Lagrangians for the symplectic form of M .

In the context of moduli spaces, Bohr–Sommerfeld Lagrangians have previously been studied by Jeffrey and Weitsman [JW] for a certain real polarization, on the moduli space of flat $SU(2)$ -connections on a smooth two-manifold, associated to a pair-of-pants decomposition. Quantizing the moduli space using this polarization results in distributional quantum states supported on its Bohr–Sommerfeld Lagrangians. Andersen proved that the Jeffrey–Weitsman polarization is obtained as the limit of the Kähler polarization on the moduli space of stable rank two bundles with trivial determinant on a Riemann surface by pinching the Riemann surface structure along the curves of the pair-of-pants decomposition [An2]. He used this fact, and the Bohr–Sommerfeld description of the quantum space for the Jeffrey–Weitsman polarization, to construct a mapping class group invariant unitary structure for the Hitchin connection in this setting, he also gives a geometric formula for the Witten–Reshetikhin–Turaev quantum invariants and proves that the colored Jones polynomials detect the unknot [An1, An3].

2. ONE-FORM ON THE MODULI SPACE

Let X be a compact connected Riemann surface of genus g , with $g \geq 2$. The holomorphic cotangent bundle of X will be denoted by K_X . Fix an integer d and also fix a positive integer r . Let $M_H(r, d)$ denote the moduli space of semistable Higgs bundles on

X of rank r and degree d . The space $M_H(r, d)$ is an irreducible normal complex projective variety of complex dimension $2(r^2(g-1) + 1)$.

There is a natural algebraic 1-form on $M_H(r, d)$; we will recall its construction. Take any Higgs bundle (E, θ) . Define

$$C^0 := \text{End}(E) = E \otimes E^\vee \quad \text{and} \quad C^1 := \text{End}(E) \otimes K_X.$$

Let $\text{ad}(\theta) : C^0 \rightarrow C^1$ be the \mathcal{O}_X -linear homomorphism defined by

$$s \mapsto [s, \theta] = s \circ \theta - \theta \circ s,$$

where the composition is the usual composition of endomorphisms. Now consider the two-term complex

$$\mathcal{C}^\bullet : C^0 \xrightarrow{\text{ad}(\theta)} C^1. \quad (2.1)$$

The infinitesimal deformations of the Higgs bundle (E, θ) are parametrized by the first hypercohomology $\mathbb{H}^1(\mathcal{C}^\bullet)$ [Hi2], [BR], [Bot], [Ma].

The natural homomorphism of complexes

$$\begin{array}{ccc} C^0 & \longrightarrow & C^1 \\ \parallel & & \downarrow \\ \text{End}(E) & \longrightarrow & 0 \end{array} \quad (2.2)$$

induces a homomorphism

$$q : \mathbb{H}^1(\mathcal{C}^\bullet) \rightarrow H^1(X, C^0) = H^1(X, \text{End}(E)). \quad (2.3)$$

We note that $H^1(X, \text{End}(E))$ parametrizes the infinitesimal deformations of the holomorphic vector bundle E . The above homomorphism q coincides with the forgetful map that sends an infinitesimal deformation of the Higgs bundle (E, θ) to the infinitesimal deformation of the underlying vector bundle E (forgetting the Higgs field θ). Let

$$\mathcal{SD} : H^1(X, \text{End}(E)) \otimes H^0(X, \text{End}(E) \otimes K_X) \rightarrow H^1(X, K_X) = \mathbb{C}$$

be the pairing given by Serre duality. Define

$$\omega(E, \theta) : \mathbb{H}^1(\mathcal{C}^\bullet) \rightarrow \mathbb{C}, \quad \alpha \mapsto \mathcal{SD}(q(\alpha) \otimes \theta), \quad (2.4)$$

where q is the homomorphism in (2.3). The one-form ω on $M_H(r, d)$ defined by $(E, \theta) \mapsto \omega(E, \theta)$ is algebraic. The algebraic two-form $d\omega$ is the symplectic form on $M_H(r, d)$ [Hi1], [Hi2], [BR], [Bot], [Ma].

Let $N^s(r, d)$ denote the moduli space of stable vector bundles on X of rank r and degree d . The total space $T^*N^s(r, d)$ of the (algebraic) cotangent bundle of $N^s(r, d)$ is a Zariski open subset of $M_H(r, d)$. The restriction of ω to $T^*N^s(r, d)$ coincides with the Liouville one-form on $T^*N^s(r, d)$.

For any nonzero complex number λ , let

$$T_\lambda : M_H(r, d) \rightarrow M_H(r, d), \quad (E, \theta) \mapsto (E, \lambda \cdot \theta) \quad (2.5)$$

be the automorphism. Clearly these automorphisms together define an action of the multiplicative group $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ on $M_H(r, d)$. Let

$$\xi \in H^0(M_H(r, d), TM_H(r, d)) \quad (2.6)$$

be the algebraic vector field associated to this action of \mathbb{C}^* .

Lemma 2.1. *The one-form $i_\xi d\omega$ obtained by contracting the symplectic form $d\omega$ using the vector field ξ satisfies the equation*

$$i_\xi d\omega = \omega.$$

Proof. We first show that $q(\xi) = 0$, where q is the homomorphism in (2.3). This follows from the fact that the action of \mathbb{C}^* does not change the underlying vector bundle (recall that q coincides with the forgetful map that sends the infinitesimal deformations of a Higgs bundle to the infinitesimal deformation of the underlying vector bundle). To prove this directly, first note that the kernel of the homomorphism of complexes in (2.2) is the complex

$$\tilde{\mathcal{C}}^\bullet : \tilde{\mathcal{C}}^0 := 0 \longrightarrow \tilde{\mathcal{C}}^1 := \text{End}(E) \otimes K_X.$$

The tangent vector $\xi((E, \theta)) \in T_{(E, \theta)}M_H(r, d) = \mathbb{H}^1(\mathcal{C}^\bullet)$ lies in the image of the homomorphism

$$H^0(X, \text{End}(E) \otimes K_X) = \mathbb{H}^1(\tilde{\mathcal{C}}^\bullet) \longrightarrow \mathbb{H}^1(\mathcal{C}^\bullet)$$

given by the natural homomorphism $\tilde{\mathcal{C}}^\bullet \longrightarrow \mathcal{C}^\bullet$. In fact, $\xi((E, \theta))$ is the image of

$$\theta \in H^0(X, \text{End}(E) \otimes K_X) = \mathbb{H}^1(\tilde{\mathcal{C}}^\bullet).$$

This immediately implies that $q(\xi) = 0$.

Since $q(\xi) = 0$, from the definitions of ω it follows that

$$\omega(\xi) = 0. \quad (2.7)$$

It also follows from the definitions of ω and T_λ (see (2.5)) that $T_\lambda^* \omega = \lambda \cdot \omega$. Therefore, we have

$$L_\xi \omega = \omega. \quad (2.8)$$

Consider the identity $L_\xi \omega = i_\xi d\omega + d(\omega(\xi))$. In view of (2.7) and (2.8), the lemma follows from this identity. \square

3. BOHR–SOMMERFELD LAGRANGIANS ON THE MODULI SPACE

Consider the symplectic form $d\omega$ on $M_H(r, d)$, where ω is constructed in (2.4). A *compact complex Lagrangian* on $M_H(r, d)$ is a reduced irreducible compact complex analytic subset

$$\iota : \mathbb{L} \hookrightarrow M_H(r, d)$$

of dimension $r^2(g-1) + 1$ such that $\iota^* d\omega = 0$. A *compact Bohr–Sommerfeld Lagrangian* on $M_H(r, d)$ is a reduced irreducible compact complex analytic subset $\iota : \mathbb{L} \hookrightarrow M_H(r, d)$ of dimension $r^2(g-1) + 1$ such that

$$\iota^* \omega = 0.$$

Since $\iota^*d\omega = dt^*\omega$, a Bohr–Sommerfeld Lagrangian is indeed a Lagrangian.

Lemma 3.1. *Let $\iota : \mathbb{L} \hookrightarrow M_H(r, d)$ be a Bohr–Sommerfeld Lagrangian. For any smooth point $z \in \mathbb{L}$, the tangent vector*

$$\xi(z) \in T_z M_H(r, d)$$

lies in the subspace $T_z \mathbb{L} \subset T_z M_H(r, d)$, where ξ is the vector field in (2.6).

Proof. Since \mathbb{L} is a Bohr–Sommerfeld Lagrangian, we have

$$\omega(v) = 0, \quad \forall v \in T_z \mathbb{L}.$$

Hence from Lemma 2.1 it follows that

$$d\omega(\xi(z), v) = 0, \quad \forall v \in T_z \mathbb{L}. \quad (3.1)$$

Since \mathbb{L} is Lagrangian for the symplectic form $d\omega$, from (3.1) it follows immediately that $\xi(z) \in T_z \mathbb{L}$. \square

Let

$$\mathcal{H} : M_H(r, d) \longrightarrow \mathcal{V} := \bigoplus_{i=1}^r H^0(X, K_X^{\otimes i}), \quad (E, \theta) \longmapsto \sum_{i=1}^r \text{trace}(\theta^i) \quad (3.2)$$

be the Hitchin map. Every irreducible component of every fiber of \mathcal{H} is a compact complex Lagrangian [Hi2]. The fiber

$$\mathcal{N} := \mathcal{H}^{-1}(0) \subset M_H(r, d)$$

is known as the nilpotent cone.

Proposition 3.2. *All compact Bohr–Sommerfeld Lagrangians in $M_H(r, d)$ are contained in the nilpotent cone \mathcal{N} .*

Proof. For any nonzero complex number λ , consider the linear automorphism of the vector space \mathcal{V} in (3.2) defined by

$$(c_1, \dots, c_i, \dots, c_r) \longmapsto (\lambda \cdot c_1, \dots, \lambda^i \cdot c_i, \dots, \lambda^r \cdot c_r).$$

These automorphisms together define an action of \mathbb{C}^* on \mathcal{V} . The morphism \mathcal{H} in (3.2) is clearly \mathbb{C}^* -equivariant for this action of \mathbb{C}^* on \mathcal{V} and the action of \mathbb{C}^* on $M_H(r, d)$ in (2.5).

Let

$$\iota : \mathbb{L} \hookrightarrow M_H(r, d)$$

be a compact Bohr–Sommerfeld Lagrangian. So \mathbb{L} does not admit any nonconstant holomorphic map to \mathcal{V} . In particular, $\mathcal{H} \circ \iota$ is a constant map, where \mathcal{H} is the Hitchin map defined in (3.2). So we have

$$\iota(\mathbb{L}) \subset \mathcal{H}^{-1}(t_0) \quad (3.3)$$

for some point $t_0 \in \mathcal{V}$.

From Lemma 3.1 it follows that $\iota(\mathbb{L})$ is preserved by the action of \mathbb{C}^* on $M_H(r, d)$ in (2.5). Since the map \mathcal{H} is \mathbb{C}^* -equivariant, we conclude that t_0 in (3.3) is fixed by the action of \mathbb{C}^* on \mathcal{V} . This implies that $t_0 = 0$. \square

Theorem 3.3. *The compact Bohr–Sommerfeld Lagrangians in $M_H(r, d)$ are precisely the irreducible components of the nilpotent cone \mathcal{N} .*

Proof. In view of Proposition 3.2, it suffices to show each irreducible component of \mathcal{N} is indeed a Bohr–Sommerfeld Lagrangian. Let

$$\mathbb{L} \subset \mathcal{N}$$

be an irreducible component. We note that \mathbb{L} is compact because \mathcal{H} is a proper morphism [Hi1], [Hi2], [Ni].

Take a smooth point

$$z = (E, \theta) \in \mathbb{L}.$$

Take any tangent vector $v \in T_z\mathbb{L}$. We need to show that

$$\omega(z)(v) = 0. \quad (3.4)$$

From Lemma 2.1 we have

$$\omega(z)(v) = d\omega(\xi, v). \quad (3.5)$$

The subvariety $\mathbb{L} \subset M_H(r, d)$ is closed under the action of \mathbb{C}^* on $M_H(r, d)$. Indeed, the Hitchin map \mathcal{H} in (3.2) is \mathbb{C}^* -equivariant, and the point $0 \in \mathcal{V}$ is fixed by the action of \mathbb{C}^* . Therefore, it follows that

$$\xi(z) \in T_z\mathbb{L}.$$

Since \mathbb{L} is a Lagrangian subvariety of $M_H(r, d)$ for the symplectic form $d\omega$ on $M_H(r, d)$ [Hi2], [La], and $\xi(z), v \in T_z\mathbb{L}$, we have

$$d\omega(\xi(z), v) = 0. \quad (3.6)$$

Finally, (3.4) follows from (3.5) and (3.6).

We give an alternative proof which may be of independent interest.

Associated to \mathbb{L} there is an integer $n \geq 2$ and pairs of integers $(r_1 d_1), \dots, (r_n d_n)$ satisfying the following conditions:

- (1) for any $(E, \theta) \in \mathbb{L}$, there is filtration of subbundles

$$0 = E_0 \subset E_1 \subset E_2 \subset \dots \subset E_{n-1} \subset E_n = E \quad (3.7)$$

such that $\text{rank}(E_i/E_{i-1}) = r_i$ and $\text{degree}(E_i/E_{i-1}) = d_i$ for all $1 \leq i \leq n$, and

- (2) $\theta(E_i) \subset E_{i-1} \otimes K_X$ for all $1 \leq i \leq n$.

(See the proof of Theorem 5.3 in [BR, p. 228].)

As before, $z = (E, \theta) \in \mathbb{L}$ is a smooth point. We will describe the tangent space $T_z\mathbb{L}$. Consider the filtration in (3.7). Let

$$\text{End}^p(E) \subset \text{End}(E)$$

be the subbundle defined by the condition that $\text{End}^p(E)(E_i) \subset E_i$ for all $1 \leq i \leq n$. Let

$$\text{End}^n(E) \subset \text{End}^p(E)$$

be the subbundle defined by the condition that $\text{End}^n(E)(E_i) \subset E_{i-1}$ for all $1 \leq i \leq n$. Let \mathcal{D}^\bullet be the two-term complex

$$\mathcal{D}^\bullet : D^0 := \text{End}^p(E) \xrightarrow{\text{ad}(\theta)} D^1 := \text{End}^n(E) \otimes K_X,$$

where $\text{ad}(\theta)$ is the homomorphism in (2.1). Note that the condition $\theta(E_i) \subset E_{i-1} \otimes K_X$, $1 \leq i \leq n$, ensures that $\text{ad}(\theta)(\text{End}^p(E)) \subset \text{End}^n(E) \otimes K_X$. The inclusion of complexes

$$\begin{array}{ccc} D^0 & \xrightarrow{\text{ad}(\theta)} & D^1 \\ \downarrow & & \downarrow \\ \text{End}(E) & \xrightarrow{\text{ad}(\theta)} & \text{End}(E) \otimes K_X \end{array}$$

induces a homomorphism of hypercohomologies

$$\varphi : \mathbb{H}^1(\mathcal{D}^\bullet) \longrightarrow \mathbb{H}^1(\mathcal{C}^\bullet)$$

(see (2.1)). We have

$$T_z \mathbb{L} = \mathbb{H}^1(\mathcal{D}^\bullet)$$

[BR, p. 229, Proposition 5.7] (see also the proof of Proposition (4.9) in [La, p. 662]), and the differential $T_z \mathbb{L} \rightarrow T_z M_H(r, d)$ of the inclusion map $\mathbb{L} \hookrightarrow M_H(r, d)$ coincides with the above homomorphism φ .

We note that

- $[\text{End}^p(E), \text{End}^n(E)] \subset \text{End}^n(E)$, and
- the homomorphism

$$\text{trace} : \text{End}(E) \longrightarrow \mathcal{O}_X$$

vanishes identically on $\text{End}^n(E)$.

Therefore, for the homomorphism $\omega(E, \theta)$ in (2.4), we have

$$\omega(E, \theta) \circ \varphi = 0.$$

Hence $\iota^* \omega = 0$. In other words, \mathbb{L} is a Bohr–Sommerfeld Lagrangian. \square

3.1. Principal Higgs bundles. Theorem 3.3 generalizes to the moduli spaces of G –Higgs bundles of fixed topological type, where G is any reductive complex algebraic group.

Let G be a complex reductive group. The topological type of G –Higgs bundles on X are parametrized by $\pi_1(G)$. For any $\delta \in \pi_1(G)$, let $M_H(G, \delta)$ denote the moduli space of G –Higgs bundles on X of topological type δ .

We have the following:

Theorem 3.4. *The compact Bohr–Sommerfeld Lagrangians in $M_H(G, \delta)$ are precisely the irreducible components of the nilpotent cone of $M_H(G, \delta)$.*

The proof is same as for the case of Higgs vector bundles. The first proof of Theorem 3.3 goes through without any change. The second proof of Theorem 3.3 also works. Indeed, for this the only point to note is that if \mathfrak{p} is the Lie algebra of a parabolic subgroup P of G , and \mathcal{B} is any G -invariant nondegenerate symmetric bilinear form on the Lie algebra of G , then the annihilator of \mathfrak{p} , with respect to \mathcal{B} , is the Lie algebra of the unipotent radical of P .

3.2. Parabolic Higgs bundles. Theorem 3.3 generalizes to the moduli spaces of parabolic Higgs bundles. In this case the assumption that $g \geq 2$ is not needed. It is enough to assume that there are stable parabolic Higgs bundles.

In this case the deformation complex is

$$\mathcal{A}^\bullet : \text{PEnd}(E) \longrightarrow \text{SPEnd}(E) \otimes K(D)$$

where

- $\text{PEnd}(E)$ is the bundle of endomorphism of E which preserves the quasiparabolic filtrations,
- $\text{SPEnd}(E) \subset \text{PEnd}(E)$ is the subsheaf defined by the nilpotent endomorphisms with respect to the quasiparabolic filtrations, and
- $K(D) = K_X \otimes \mathcal{O}_X(D)$ with D being the divisor provided by the parabolic data.

Imitating q in (2.3), consider the homomorphism

$$\tilde{q} : \mathbb{H}^1(\mathcal{A}^\bullet) \longrightarrow H^1(X, \text{PEnd}(E)).$$

Finally in this case Serre duality for parabolic bundles provides the following morphism

$$\widetilde{\mathcal{SD}} : H^1(X, \text{PEnd}(E)) \otimes H^0(X, \text{SPEnd}(E) \otimes K(D)) \longrightarrow \mathbb{C}.$$

Now we define the algebraic one form for parabolic Higgs bundles as

$$\omega(E, \theta) : \mathbb{H}^1(\mathcal{A}^\bullet) \longrightarrow \mathbb{C}, \quad \alpha \longmapsto \widetilde{\mathcal{SD}}(\tilde{q}(\alpha) \otimes \theta).$$

Once the set-up is built, Theorem 3.3 has a straightforward generalization.

Theorem 3.3 also generalizes to parabolic analog of principal Higgs bundles; see [BBN], [Boa], [PR], [He] for parabolic analog of principal Higgs bundles.

ACKNOWLEDGEMENTS

We thank the referee for helpful comments. This work was supported by a Marie Curie International Research Staff Exchange Scheme Fellowship within the 7th European Union Framework Programme (FP7/2007-2013) under grant agreement no. 612534, project MODULI - Indo European Collaboration. The second author was partly supported by the center of excellence grant 'Center for Quantum Geometry of Moduli Spaces' from the Danish National Research Foundation (DNRF95). The first author is supported by the J. C. Bose Fellowship.

REFERENCES

- [An1] J. E. Andersen, Mapping class group invariant unitarity of the Hitchin connection over Teichmüller space, arXiv:1206.2635, 2012.
- [An2] J. E. Andersen, New polarizations on the moduli spaces and the Thurston compactification of Teichmüller space, *Internat. J. Math.* **9** (1998), 1–45.
- [An3] J. E. Andersen, A geometric formula for the Witten-Reshetikhin-Turaev quantum invariants and some applications, arXiv:1206.2785, 2012.
- [BBN] V. Balaji, I. Biswas and D. S. Nagaraj, Principal bundles over projective manifolds with parabolic structure over a divisor, *Tohoku Math. Jour.* **53** (2001), 337–367.
- [BR] I. Biswas and S. Ramanan, An infinitesimal study of the moduli of Hitchin pairs, *Jour. London Math. Soc.* **49** (1994), 219–231.
- [Boa] P. P. Boalch, Riemann-Hilbert for tame complex parahoric connections, *Transform. Groups* **16** (2011), 27–50.
- [Bot] F. Bottacin, Symplectic geometry on moduli spaces of stable pairs, *Ann. Sci. École Norm. Sup.* **28** (1995), 391–433.
- [He] J. Heinloth, Uniformization of \mathcal{G} -bundles, *Math. Ann.* **347** (2010), 499–528.
- [Hi1] N. J. Hitchin, The self-duality equations on a Riemann surface, *Proc. London Math. Soc.* **55** (1987), 59–126.
- [Hi2] N. J. Hitchin, Stable bundles and integrable systems, *Duke Math. Jour.* **54** (1987), 91–114.
- [JW] L. C. Jeffrey and J. Weitsman, Bohr-Sommerfeld orbits in the moduli space of flat connections and the Verlinde dimension formula, *Comm. Math. Phys.* **150** (1992), 593–630.
- [La] G. Laumon, Un analogue global du cône nilpotent, *Duke Math. Jour.* **57** (1988), 647–671.
- [Ma] E. Markman, Spectral curves and integrable systems, *Compos. Math.* **93** (1994), 255–290.
- [Ni] N. Nitsure, Moduli space of semistable pairs on a curve, *Proc. London Math. Soc.* **62** (1991), 275–300.
- [PR] G. Pappas and M. Rapoport, Some questions about \mathcal{G} -bundles on curves. *Algebraic and arithmetic structures of moduli spaces* (Sapporo 2007), 159–171, Adv. Stud. Pure Math., 58, Math. Soc. Japan, Tokyo, 2010.
- [Sn] J. Śniatycki, *On cohomology groups appearing in geometric quantization*, Differential geometrical methods in mathematical physics (Proc. Sympos., Univ. Bonn, Bonn, 1975), 46–66, Lecture Notes in Math., No. 570, Springer, Berlin, 1977.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400005, INDIA

E-mail address: indranil@math.tifr.res.in

CENTER FOR QUANTUM GEOMETRY OF MODULI SPACES, UNIVERSITY OF AARHUS, NY MUNKEGADE 118, DK-8000 AARHUS, DENMARK

E-mail address: nlg@qgm.au.dk

INSTITUTO DE CIENCIAS MATEMÁTICAS (CSIC-UAM-UC3M-UCM), C/ NICOLAS CABRERA 15, 28049 MADRID, SPAIN

E-mail address: marina.logares@icmat.es