

# Supersolutions to degenerated logistic equation type

Marcos Marva.

Departamento de Fısica y Matematicas, Universidad de Alcala,  
28871 Alcala de Henares, Spain  
marcos.marva@uah.es

Mathematics Subject Classification (2010): 35J99 (primary).

Keywords: Supersolution, logistic equation.

## Abstract

In this work we provide a method for building up a strictly positive supersolution for the steady state of a degenerated logistic equation type, i.e., when the weight function vanishes on the boundary of the domain. This degenerated system is related in obtaining the so-called large solutions. Previously, this problem was handled as the limit case of non degenerated approaching problems. Our method can be adapted straightforwardly to degenerated boundary value problems.

## 1 Introduction

In this work we show how to build up a positive strict supersolution to the boundary value problem

$$\begin{cases} -\Delta u = \lambda m(x)u - a(x)f(x, u) & \text{in } \Omega, \\ u = g(x) & \text{on } \partial\Omega, \end{cases} \quad (1)$$

where  $a \in C^\mu(\bar{\Omega}; [0, +\infty))$ ,  $m \in C^\mu(\bar{\Omega}; \mathbb{R})$ ,  $\mu \in (0, 1)$ ,  $a \equiv 0$  on  $\partial\Omega$ ,  $g \in C^{1+\mu}(\partial\Omega; [0, +\infty))$  is such that  $g \geq 0$ ,  $g \not\equiv 0$ ,  $f \in C^{\mu, 1+\mu}(\bar{\Omega} \times [0, +\infty); \mathbb{R})$ ,  $f(x, 0) = 0$ ,  $f(x, u) > 0$ ,  $\partial_u f(x, u) > 0$  and  $\lim_{k \nearrow +\infty} \frac{f(x, ku)}{k} = +\infty$  for each  $u > 0$  uniformly in  $x \in \Omega$ . The set  $\Omega \subset \mathbb{R}^N$  is a bounded domain with  $\partial\Omega$  of class  $C^2$ .

Problem (1) (with  $m \equiv 1$ ) stands for the steady states of the generalized logistic growth law [13]. Typically,  $u$  stands for the distribution in  $\Omega$  of individuals of certain species,  $\lambda$  stands for the net growth rate of  $u$  modulated by  $m(x)$ ,  $a(x)$  simulates demography pressure and, along with  $m(x)$ , environmental heterogeneity. When  $a \equiv 0$  somewhere, model (1) (with  $m \equiv 1$ ) becomes a Malthusian model in the region  $\Omega_0 := \{x \in \Omega; a(x) = 0\}$ , known as *refuge*. We deal with the case  $a \equiv 0$  on  $\partial\Omega$ , but it could be adapted to apply in the general case when  $\Omega_0 \neq \emptyset$  and  $\Omega_0 \subset \Omega$ .

In addition, solving problem (1) is an intermediate step to prove the existence of large solutions related to problem (1), i.e., a function  $u \in C^{2,\nu}(\bar{\Omega})$  such that

$$\begin{cases} -\Delta u = \lambda m(x)u - a(x)f(x, u)u & \text{in } \Omega, \\ \lim_{x \rightarrow \partial\Omega} u(x) = \infty, \end{cases} \quad (2)$$

see [11]. Large solutions come across from studies concerning combustion due to Keller [9] and Osserman [14]. In fact, there are available uniqueness results for problem (2) ([3], [5], [12] and references therein). As long as we know, problem (1) has been handled as the limit case of non degenerated approaching problems.

If we restrict ourselves in problem (1) to  $a(x) \geq \gamma > 0$  on  $\bar{\Omega}$ , it is easy to show that there exists  $\epsilon > 0$  small enough such that  $\epsilon\varphi$  is a positive subsolution to problem (1), being  $\varphi$  the principal eigenfunction of  $-\Delta$  in  $\Omega$  under homogeneous Dirichlet boundary conditions. In this case a positive constant  $K$  such that

$$K > \max \left\{ \max_{x \in \partial\Omega} \{g\}, \frac{\max_{x \in \bar{\Omega}} \{\lambda m(x)\}}{\min_{x \in \bar{\Omega}} \{a(x)\}} \right\}, \quad (3)$$

provides us with a supersolution to problem (1). Now, enlarging  $K$  if necessary, we get an ordered pair  $(\epsilon\varphi, K)$  of sub-supersolution. Thanks to a Theorem by Amann [1], there exists a solution to problem (1) between  $\epsilon\varphi$  and  $K$ . Unfortunately, as soon as we let  $a = 0$  somewhere in  $\bar{\Omega}$ , we lose condition (3), so it is needed something different from a constant to get a supersolution to problem (1). Different strategies have been used to avoid this problem: some authors approximate  $\Omega$  by  $\Omega_n := \{x \in \Omega; \text{dist}(x, \partial\Omega) > 1/n\}$  (for each  $n$   $a(x)$  is uniformly bounded away from 0 in  $\Omega_n$ , that is  $a \geq \gamma > 0$ , see [4] and [11]), and others approach  $a(x)$  by means of  $\frac{1}{n} + a(x)$ , essentially in order to avoid condition  $a \equiv 0$  on  $\partial\Omega$  and to generate a sequence of solutions of approximate problems converging to a solution to problem (1), see [2] and [8].

## 2 Results

The result we present here allows us to obtain via sub and supersolution the existence of a solution to (1). Actually, the construction of the supersolution in  $\Omega$  follows from an extension of the Faber [6] and Krahn [7] inequality due to López-Gómez [10] which provides us with a lower estimate for the main eigenvalue to  $-\Delta$  on the domain  $D$ :

$$\sigma_1[-\Delta, D] \geq \frac{\sigma_1[-\Delta, B_1(0)] \cdot |B_1(0)|^{2/N}}{|D|^{2/N}}, \quad (4)$$

where  $B_1(0) := \{x \in \mathbb{R}^N; |x| \leq 1\}$ ,  $\sigma_1[-\Delta, B_1(0)]$  is the principal eigenvalue of  $-\Delta$  on  $B_1(0)$  and  $|D|$  stands for the Lebesgue's measure of  $D$ .

**Theorem 2.1** *Consider the boundary value problem (1). Then, for each  $\lambda \in \mathbb{R}$ , there exists an ordered pair  $(\underline{u}, \bar{u})$  consisting of a subsolution and a positive strict supersolution, where  $\underline{u}(x) < \bar{u}(x) \forall x \in \Omega$ .*

**Proof.**- Let us define

$$O_\epsilon := \{x \in \mathbb{R}^N; \text{dist}(x, \partial\Omega) < \epsilon\}, \quad \sigma_\epsilon := \sigma[-\Delta, O_\epsilon],$$

where, keeping in mind (4), for a fixed  $\lambda \in \mathbb{R}$  we can choose  $\epsilon > 0$  such that

$$\max_{x \in \overline{\Omega}} \{\lambda m(x)\} < \sigma_\epsilon.$$

Let  $\varphi_o^\epsilon$  be the principal eigenfunction of  $-\Delta$  under homogeneous boundary conditions on  $O_\epsilon$ . We define

$$\Phi := \begin{cases} \varphi_o^\epsilon & \text{in } \overline{\Omega} \cap O_{\epsilon/2}, \\ \Psi & \text{in } \Omega_{\epsilon/2} := \Omega \setminus (\overline{\Omega} \cap O_{\epsilon/2}), \end{cases}$$

where  $\Psi \geq \tau > 0$  is any regular enough function such that  $\Phi \in C^2(\overline{\Omega})$ . We are looking for the existence of a constant  $K > 0$ , large enough, such that function

$$\bar{u} := K\Phi$$

is a strict positive supersolution to problem (1).

We estimate now  $K$  on  $\overline{\Omega} \cap O_{\epsilon/2}$ . Keeping in mind that  $a(x) \geq 0$  and  $K\Phi = K\varphi_o^\epsilon$ , it must be

$$\begin{cases} K(-\Delta\varphi_o^\epsilon) \geq \lambda m K\varphi_o^\epsilon - af(x, K\varphi_o^\epsilon) & \text{in } \overline{\Omega} \cap O_{\epsilon/2}, \\ K\varphi_o^\epsilon \geq g > 0 & \text{on } \partial\overline{\Omega} \cap O_{\epsilon/2} = \partial\Omega. \end{cases}$$

In the interior of the domain it should happen

$$K(-\Delta\varphi_o^\epsilon) \geq \lambda m K\varphi_o^\epsilon - af(x, K\varphi_o^\epsilon),$$

which is equivalent to

$$K\sigma_\epsilon\varphi_o^\epsilon \geq \lambda m K\varphi_o^\epsilon - af(x, K\varphi_o^\epsilon)$$

by the definition of  $\varphi_o^\epsilon$ . Rearranging terms we get

$$af(x, K\varphi_o^\epsilon) \geq K\varphi_o^\epsilon (\lambda m - \sigma_\epsilon), \quad (5)$$

where the left hand side is non negative and the right one is strictly negative because of the selection of  $\epsilon$ . Therefore, (5) is a proper inequality  $\forall K > 0$ . By construction, on the boundary we have  $\varphi_o^\epsilon|_{\partial\Omega}, g > 0$  and therefore, whenever

$$K > \max_{x \in \partial\Omega} \left\{ \frac{g(x)}{\varphi_o^\epsilon(x)} \right\}, \quad (6)$$

the boundary condition is satisfied (with strict inequality).

In the domain  $\Omega_{\epsilon/2}$  it should be verified

$$K(-\Delta\Psi) \geq \lambda m K\Psi - af(x, K\Psi). \quad (7)$$

rearranging terms, expression (7) is equivalent to

$$\frac{f(x, K\Psi)}{K} \geq \max_{x \in \overline{\Omega}_{\epsilon/2}} \left\{ \frac{\Delta\Psi + \lambda m\Psi}{a} \right\}. \quad (8)$$

As  $a(x) > 0 \quad \forall x \in \Omega_{\epsilon/2}$ , function  $\frac{\Delta\Psi + \lambda m\Psi}{a}$  reaches its maximum (provided a regular  $\Psi$ ). By hypothesis on  $f$ , there exists  $K_0$  such that  $\forall K > K_0$  condition (8) holds. Thus, there exists  $K > 0$  such that function  $K\Psi$  is a strict positive supersolution to (1). Function  $\underline{u} := 0$  is a subsolution to problem (1), and  $\underline{u}(x) < \overline{u}(x) \quad \forall x \in \overline{\Omega}$ . ■

**Corollary 2.2** *For each  $\lambda \in \mathbb{R}$ , problem (1) has an unique positive solution.*

**Proof.-** Since we have an ordered pair formed by a subsolution and a supersolution, a Theorem due to Amann [1] guaranties the existence of a solution  $u$  to problem (1) such that  $0 \leq u \leq K\Psi$ . Uniqueness follows, for instance, from [11]. ■

**Acknowledgement.-** The author thanks E. Gómez-Hidalgo for her fruitful comments and support. The author is partially supported by Ministerio de Ciencia e Innovación (Spain), projects MTM2011-24321 and MTM2011-25238.

## References

- [1] H. Amann, *Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces*, SIAM Rev., **18**:219 (1976), 620–709.
- [2] F.C. Cirstea, V. Radulescu, *Asymptotics for the blow-up boundary solution of the logistic equation with absorption*, C. R. Math. Acad. Sci. Paris, **226** (3)336 (2003), 231–236.
- [3] F.C.Cirstea, V. Radulescu, *Nonlinear problems with boundary blow-up: a Karamata regular variation theory approach*, Asymptot. Anal., **46**:3–4 (2006), 275–298.
- [4] Y. Du, Q. Huang, *Blow-up solutions for a class of semilinear elliptic and parabolic problems*, SIAM J. Math. Anal., **31**:1 (1999), 1–18, .
- [5] Y. Du, *Asymptotic behavior and uniqueness results for boundary blow-up solutions*, Differ. Integral Equ. Appl., **17** (2004), 819–834.
- [6] G. Faber *Beweis das unter allen homogenen Membranen von Gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt*, Munch. Ber. (1923), 169–172.
- [7] E. Krahn, *Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises*, Math. Ann., **94** (1925), 97–100.

- [8] J. García-Melián, R. Letelier-Albornoz, J.C. Sabina de Lis, *Uniqueness and asymptotic behaviour for solutions of semilinear problems with boundary blow-up*, Proc. Amer. Math. Soc., **129**:12 (2001), 3593–3602.
- [9] J.B. Keller, *On solutions of  $\Delta u = f(u)$* , Comm. Pure Appl. Math., **10** (1957), 503–510.
- [10] J. López-Gómez, *The maximum principle and the existence of principal eigenvalues for some linear weighted boundary value problems*, J. Diff. Eqns. Conf., **1**, No.1 electronic only (2000), (1996) 263–294.
- [11] J. López-Gómez, *Large solutions, metasolutions, and asymptotic behaviour of the regular positive solutions of sublinear parabolic problems*, J. Diff. Eqns. Conf., **5** (2000), 135–171.
- [12] J. López-Gómez, *Optimal uniqueness theorems and exact blow-up rates of large solutions*, J. Diff. Eqns., **224**:2 (2006), 385–439.
- [13] J.D. Murray, *Mathematical biology*. Berlin: Biomathematics 19 Springer-Verlag, 1993.
- [14] R. Osserman, *On the inequality  $\Delta u \geq f(u)$* , Pacific. J. Math., **7** (1957), 1641–1647.