

Dipolar Dark Matter with Massive Bigravity

Luc Blanchet^a and Lavinia Heisenberg^{b,c}

^a $\mathcal{G}\mathcal{R}\mathcal{E}\mathcal{C}\mathcal{O}$ Institut d'Astrophysique de Paris — UMR 7095 du CNRS, Université Pierre & Marie Curie, 98^{bis} boulevard Arago, 75014 Paris, France

^b Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, 10691 Stockholm, Sweden

^c Department of Physics & The Oskar Klein Centre, AlbaNova University Centre, 10691 Stockholm, Sweden

E-mails: blanchet@iap.fr, laviniah@kth.se

Abstract. Massive gravity theories have been developed as viable IR modifications of gravity motivated by dark energy and the problem of the cosmological constant. On the other hand, modified gravity and modified dark matter theories were developed with the aim of solving the problems of standard cold dark matter at galactic scales. Here we propose to adapt the framework of ghost-free massive bigravity theories to reformulate the problem of dark matter at galactic scales. We investigate a promising alternative to dark matter called dipolar dark matter (DDM) in which two different species of dark matter are separately coupled to the two metrics of bigravity and are linked together by an internal vector field. We show that this model successfully reproduces the phenomenology of dark matter at galactic scales (*i.e.* MOND) as a result of a mechanism of gravitational polarisation. The model is safe in the gravitational sector, but because of the particular couplings of the matter fields and vector field to the metrics, a ghost in the decoupling limit is present in the dark matter sector. However, it might be possible to push the mass of the ghost beyond the strong coupling scale by an appropriate choice of the parameters of the model. Crucial questions to address in future work are the exact mass of the ghost, and the cosmological implications of the model.

Contents

1	Introduction	1
1.1	Massive gravity context	1
1.2	Dark matter context	3
2	The covariant theory	4
3	Linear perturbations	8
4	Polarisation mechanism and MOND	10
5	The decoupling limit	16
6	Conclusions	19

1 Introduction

In the last century, cosmology has progressively developed from a philosophical to an empirical scientific discipline, witnessing high precision cosmological observations, which culminated with the standard model of cosmology, the Λ -CDM model [1]. The standard model is based on General Relativity (GR) and is in great agreement with the abundance of the light elements in the big bang nucleosynthesis, the anisotropies of the cosmic microwave background (CMB), the baryon acoustic oscillations, lensing and the observed large scale structures. It notably relies on the presence of dark energy in form of a cosmological constant Λ , giving rise to the accelerated expansion of the universe.

1.1 Massive gravity context

In the prevailing view, the cosmological constant corresponds to the constant vacuum energy density which receives large quantum corrections. Unfortunately, so far there is no successful mechanism to explain the observed unnatural tiny value of the cosmological constant, see *e.g.* [2, 3]. This difficulty has sparked a whole industry studying modifications of gravity in the infra-red (IR) invoking new dynamical degrees of freedom.

One tempting route is massive gravity, motivated by the possibility that the graviton has a mass. It was a challenge over forty years to construct a covariant non-linear theory for massive gravity. The foregathered remarkable amount of effort finally rose to the challenge [4–11], to construct the potential interactions in a way that got rid of the Boulware–Deser (BD) ghost [12]. This theory has been further extended to more general models by adding additional degrees of freedom. A very important extension is massive bigravity theory [13].

Once the interactions in the gravitational sector were guaranteed to be ghost-free, a natural follow up question was how to couple this theory to the matter fields without spoiling the ghost-freedom. First attempts were already discussed in the original paper of bigravity [13].

If one couples the matter fields to both metrics simultaneously, this reintroduces the BD ghost [14, 15]. Furthermore, the one-loop quantum corrections detune the special potential interactions at an unacceptable low scale and hence this way of coupling is not a consistent one [15]. The safer way is to couple the matter fields to just one metric. In this way, the quantum corrections give rise to contributions in form of cosmological constants. One could also try to couple the matter field to the massless mode, which is unfortunately also not ghost free [16]. Another possible way of coupling the matter fields to both metrics is through a composite effective metric constructed out of both metrics [15, 17–19], which is unique in the sense that it is the only non-minimal matter coupling that maintains ghost-freedom in the decoupling limit [20–22]. Furthermore, the quantum corrections are guaranteed to maintain the nice potential structure. Other important consequence of this new matter coupling is the fact that it helps to evade the no-go result [23] for the flat Friedmann–Lemaître–Robertson–Walker (FLRW) background. A detailed perturbed ADM analysis revealed the existence of a BD ghost originating from an operator involving spatial derivatives [15, 17]. Therefore, the ghost will probably reappear for highly anisotropic solutions. Since the ghost remains absent up to the strong coupling scale, the matter coupling can be considered in an effective field theory sense at the very least till the strong coupling scale. The precise cut-off of the theory or mass of the ghost has still to be established.

The absence of the BD ghost is not only important at the classical level, but also at the quantum level. For the classical ghost-freedom the relative tuning of the potential interactions is the key point. Therefore one has to ensure that the quantum corrections do not detune the potential interactions. Concerning the decoupling limit, it is easy to show that the theory receives no quantum corrections *via* the non-renormalization theorem due to the antisymmetric structure of the interactions [24] (in fact the same antisymmetric structure of the Galileon interactions protect them from quantum corrections [25–27]). Beyond the decoupling limit, the quantum corrections of the matter loops maintain the potential interactions provided that the above criteria are fulfilled. Concerning the graviton loops, they do destroy the relative tuning of the potential interactions. Nevertheless, this detuning is harmless since the mass of the corresponding BD ghost is never below the cut-off scale of the theory [28]. The bimetric version of the theory shares the same property [29].

Massive gravity/biggravity theory has a rich phenomenology. The decoupling limit admits stable self-accelerating solutions [30], where the helicity-0 degree of freedom of the massive graviton plays the role of a condensate whose energy density sources self-acceleration. Unfortunately these solutions suffer from strong coupling issues due to the vanishing kinetic term of the vector modes [30, 31]. With the original massive gravity and the restriction of flat reference metric, one faces the no-go result, namely that there are no flat FLRW solution [23]. One can construct self-accelerating open FLRW solutions [32], which however have three instantaneous modes [33] and suffer from a nonlinear ghost instability [34]. The attempt to promote the reference metric to de Sitter [35, 36] also failed due to the presence of the Higuchi ghost [35]. In order to avoid these difficulties, one either gives up on the FLRW symmetries [23], or invokes new additional degrees of freedom [13, 37, 38].

Thanks to the freedom gained in the inclusion of the second kinetic term in the bimetric extension [13], there now exists many elaborate works concerning the cosmology of the bigrav-

ity theory, see *e.g.* [39–41]. In the case of minimally coupled matter fields and small graviton mass, the theory admits several interesting branches of solutions. Unfortunately, among them the self-accelerating branch is unstable due to the presence of three instantaneous modes, and a second branch of solutions admits an early time gradient instability [42]. Nevertheless, there exists attempts to overcome the gradient instability either by viable though finely tuned solutions in the case of a strongly interacting bimetric theory with $m \gg H_0$ [43, 44], or by demanding that $M_g \gg M_f$ as was proposed in [45]. See also other recent works concerning the phenomenology of bimetric gravity [46–50].

1.2 Dark matter context

On a quite different but equally fascinating topic is the phenomenology of dark matter at galactic scales. The evidence for dark matter is through the measurement of the rotation curves of spiral galaxies which turn out to be approximately flat, contrary to the Newtonian prediction based on ordinary baryonic matter [51, 52]. The standard explanation is that the disk of galaxies is embedded into the quasi-spherical potential of a huge halo of dark matter, and that this dark matter is the same as the cold dark matter (CDM) which is evidenced at large cosmological scales notably with the fit of Λ -CDM to the CMB anisotropies [53, 54]. Unfortunately this explanation faces severe challenges when compared to observations at galactic scales [55, 56]. There are predictions of the Λ -CDM model that are not observed, like the phase-space correlation of galaxy satellites and the generic formation of dark matter cusps in the central regions of galaxies. Even worse, there are observations which are not predicted by Λ -CDM, such as the tight correlation between the mass discrepancy (luminous *vs.* dynamical mass) which measures the presence of dark matter and the involved scale of acceleration, the famous baryonic Tully-Fisher (BTF) relation for spiral galaxies [57–59], and its equivalent for elliptical galaxies, the Faber-Jackson relation [60].

Instead of additional, non-visible mass, Milgrom [61–63] proposed an amendment to the Newtonian laws of motion in order to account for the phenomenology of dark matter in galaxies, dubbed MOND for MODified Newtonian Dynamics. According to the MOND hypothesis the change has a relevant influence on the motion only for very small accelerations, as they occur at astronomical scales, below the critical value $a_0 \simeq 1.2 \times 10^{-10} \text{ m/s}^2$ measured for instance from the BTF relation.¹ A more elaborate version of MOND is the Bekenstein-Milgrom [64] modification of the Poisson equation of Newtonian gravity. All the challenges met by Λ -CDM at galactic scales are then solved — sometimes with incredible success — by the MOND formula [55, 56]. Unfortunately, MOND faces difficulties in explaining the dark matter distribution at the larger scales of galaxy clusters [65–69]. It is also severely constrained by observations in the solar system [70, 71].

Reconciling Λ -CDM at cosmological scales and MOND at galactic scales into a single relativistic theory is a great challenge. There has been extensive works on this. One approach is to modify gravity by invoking new gravitational degrees of freedom without the presence of dark matter [72–81]. Primary examples are the tensor-vector-scalar (TeVeS) theory [73, 74] and generalized Einstein-Æther theories [75, 76]. Another approach is a new form of dark

¹A striking observation is the numerical coincidence that $a_0 \sim \sqrt{\Lambda}$ [55, 56].

matter *à la* MOND, called dipolar dark matter (DDM). It is based on the dielectric analogy of MOND [82] — a remarkable property of MOND with possible far-reaching implications. Indeed the MOND equation represents exactly the gravitational analogue (in the non-relativistic limit) of the Gauss equation of electrostatics modified by polarisation effects in non-linear dielectric media. Some early realizations of this analogy were proposed in [83–85] and shown to also reproduce the cosmological model Λ -CDM. The best way to interpret this property is by a mechanism of gravitational polarisation, involving two different species of particles coupled to different gravitational potentials. This was the motivation for pushing forward a more sophisticated model in the context of a bimetric extension of GR [86, 87]. In this model the two species of dark matter particles that couple to the two metrics are linked through an internal vector field. The model [87] is able to recover MOND at galactic scales and the cosmological model Λ -CDM at large scales. Furthermore the post-Newtonian parameters (PPN) in the solar system are the same as those of GR. Unfortunately, even if this model delivers a very interesting phenomenology, it is plagued by ghost instabilities in the gravitational sector already at the linear order of perturbations and within the decoupling limit [88]. A more promising model was then proposed in [88], with the hope to cure these instabilities in the gravitational sector.

In the present work we will study in detail the theoretical and phenomenological consistency of this recently proposed model for dipolar dark matter in the context of bimetric gravity [88]. We will first work out the covariant field equations in Sec. 2 and analyze the linear field equations in Sec. 3. As next, we will investigate in Sec. 4 the mechanism of gravitational polarisation in the non relativistic approximation, and show how it successfully recovers the MOND phenomenology on galactic scales. Finally we will pay attention in Sec. 5 to the matter sector and investigate the interactions in the decoupling limit. The outcome of Sec. 5 is that a ghost instability is still present in the (dark) matter sector of the model. The paper ends with some short conclusions in Sec. 6.

2 The covariant theory

A new relativistic model for dipolar dark matter has been recently proposed in [88]. Basically this model is defined by combining the specific dark matter sector of a previous model [87] with gravity in the form of ghost-free bimetric extensions of GR [13]. The dark matter in this model consists of two types of particles respectively coupled to the two dynamical metrics $g_{\mu\nu}$ and $f_{\mu\nu}$ of bigravity. In addition, these two sectors are linked together *via* an additional internal field in the form of an Abelian $U(1)$ vector field \mathcal{A}_μ . This vector field is coupled to the effective composite metric g_{eff} of bigravity which is built out of the two metrics g and f , and which has been shown to be allowed in the matter Lagrangian in the effective field theory sense at least up to the strong coupling scale [15, 17]. The total matter-plus-gravity

Lagrangian reads²

$$\begin{aligned} \mathcal{L} = & \sqrt{-g} \left(\frac{M_g^2}{2} R_g - \rho_{\text{bar}} - \rho_g \right) + \sqrt{-f} \left(\frac{M_f^2}{2} R_f - \rho_f \right) \\ & + \sqrt{-g_{\text{eff}}} \left[m^2 M_{\text{eff}}^2 + \mathcal{A}_\mu (j_g^\mu - j_f^\mu) + \lambda M_{\text{eff}}^2 \mathcal{W}(\mathcal{X}) \right]. \end{aligned} \quad (2.1)$$

Here R_g and R_f are the Ricci scalars of the two metrics, and the scalar energy densities of the ordinary matter modelled simply by pressureless baryons,³ and the two species of pressureless dark matter particles are denoted by ρ_{bar} , ρ_g and ρ_f respectively. In addition j_g^μ , j_f^μ stand for the mass currents of the dark matter, defined by

$$j_g^\mu = \frac{\sqrt{-g}}{\sqrt{-g_{\text{eff}}}} J_g^\mu \quad \text{and} \quad j_f^\mu = \frac{\sqrt{-f}}{\sqrt{-g_{\text{eff}}}} J_f^\mu, \quad (2.2)$$

where $J_g^\mu = r_g \rho_g u_g^\mu$ and $J_f^\mu = r_f \rho_f u_f^\mu$ are the corresponding conserved dark matter currents associated with the respective metrics g and f , thus satisfying $\nabla_\mu^g J_g^\mu = 0$ and $\nabla_\mu^f J_f^\mu = 0$. Here ρ_g and ρ_f denote the scalar densities and u_g^μ , u_f^μ are the four velocities normalized to $g_{\mu\nu} u_g^\mu u_g^\nu = -1$ and $f_{\mu\nu} u_f^\mu u_f^\nu = -1$. We also introduced two constants r_g and r_f specifying the ratios between the charge and the inertial mass of the dark matter particles.

The metric independent matter degrees of freedom are the coordinate densities $\rho_g^* = \sqrt{-g} \rho_g u_g^0$ and $\rho_f^* = \sqrt{-f} \rho_f u_f^0$, and the coordinate velocities $v_g^\mu = u_g^\mu / u_g^0$ and $v_f^\mu = u_f^\mu / u_f^0$. The associated metric independent currents are $J_g^{*\mu} = r_g \rho_g^* v_g^\mu$ and $J_f^{*\mu} = r_f \rho_f^* v_f^\mu$. They are conserved in the ordinary sense, $\partial_\mu J_g^{*\mu} = 0$ and $\partial_\mu J_f^{*\mu} = 0$. They relate to the classical currents J_g^μ , J_f^μ or j_g^μ , j_f^μ by

$$J_g^{*\mu} = \sqrt{-g} J_g^\mu = \sqrt{-g_{\text{eff}}} j_g^\mu \quad \text{and} \quad J_f^{*\mu} = \sqrt{-f} J_f^\mu = \sqrt{-g_{\text{eff}}} j_f^\mu.$$

Notice that the coupling term $\sqrt{-g_{\text{eff}}} \mathcal{A}_\mu (j_g^\mu - j_f^\mu)$ in the action (2.1) is actually independent of any metric, neither g nor f nor g_{eff} . We shall study in detail in Sec. 5 the implications of the term $\sqrt{-g_{\text{eff}}} \mathcal{A}_\mu (j_g^\mu - j_f^\mu)$ for the decoupling limit of the theory.

The vector field \mathcal{A}_μ is generated by the dark matter currents and plays the role of a ‘‘graviphoton’’. The presence of this internal field is necessary to stabilize the dipolar medium and will yield the wanted mechanism of gravitational polarisation, as we shall see in Sec. 4. It has a non-canonical kinetic term $\mathcal{W}(\mathcal{X})$, where \mathcal{W} is a function which is left unspecified at this stage, and

$$\mathcal{X} = -\frac{\mathcal{F}^{\mu\nu} \mathcal{F}_{\mu\nu}}{4\lambda}. \quad (2.3)$$

Here λ is a constant and the field strength is constructed with the effective composite metric $g_{\mu\nu}^{\text{eff}}$ given by (2.4) below, *i.e.* $\mathcal{F}^{\mu\nu} = g_{\text{eff}}^{\mu\rho} g_{\text{eff}}^{\nu\sigma} \mathcal{F}_{\rho\sigma}$ with $\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu$ not depending on the metric as usual.

²The metric signature convention is $(-, +, +, +)$. We adopt geometrical units with the gravitational constant, the Planck constant and the speed of light being unity, $G = \hbar = c = 1$, unless specified otherwise.

³We have in mind that the baryons actually represent the full standard model of particle physics.

The model (2.1) is defined by several constant parameters, the coupling constants M_g , M_f and M_{eff} , the mass of the graviton m , the charge to mass ratios r_g and r_f of dark matter, the constant λ associated with the vector field, and the arbitrary constants α and β entering the effective metric (2.4). Parts of these constants, as well as the precise form of the function $\mathcal{W}(\mathcal{X})$, will be determined in Sec. 4 when we demand that the model reproduces the required physics of dark matter at galactic scales. In particular λ will be related to the MOND acceleration scale a_0 , and the model will finally depend on a_0 and the mass of the graviton m . Additionally, there will be still some remaining freedom in the parameters α and β , and in the coupling constants M_g and M_f .

In the model (2.1) the ghost-free potential interactions between the two metrics g and f take the particular form of the square root of the determinant of the effective metric [15, 17, 19]

$$g_{\mu\nu}^{\text{eff}} = \alpha^2 g_{\mu\nu} + 2\alpha\beta \mathcal{G}_{\mu\nu}^{\text{eff}} + \beta^2 f_{\mu\nu}, \quad (2.4)$$

where α and β are arbitrary constants, and we have defined

$$\mathcal{G}_{\mu\nu}^{\text{eff}} = g_{\mu\rho} X_\nu^\rho = f_{\mu\rho} Y_\nu^\rho, \quad (2.5)$$

where the square root matrix is defined by $X = \sqrt{g^{-1}f}$, together with its inverse $Y = \sqrt{f^{-1}g}$. Interchanging the two metrics g and f does not change the form of the metric (2.4)–(2.5) except for a redefinition of the parameters α and β . Notice that $\mathcal{G}_{\mu\nu}^{\text{eff}}$ can be proved to be automatically symmetric [16, 89], and in [88] it was shown, that it corresponds to the composite metric considered in the previous model [87]. The square root of the determinant of $g_{\mu\nu}^{\text{eff}}$ is given in either forms respectively associated with the g or f metrics, by

$$\sqrt{-g_{\text{eff}}} = \sqrt{-g} \det(\alpha\mathbb{1} + \beta X) = \sqrt{-f} \det(\beta\mathbb{1} + \alpha Y). \quad (2.6)$$

More explicitly, introducing the elementary symmetric polynomials $e_n(X)$ and $e_n(Y)$ of the square root matrices X or Y ,⁴

$$e_0(X) = 1, \quad (2.7a)$$

$$e_1(X) = [X], \quad (2.7b)$$

$$e_2(X) = \frac{1}{2}([X]^2 - [X^2]), \quad (2.7c)$$

$$e_3(X) = \frac{1}{6}([X]^3 - 3[X][X^2] + 2[X^3]), \quad (2.7d)$$

$$e_4(X) = \frac{1}{24}([X]^4 - 6[X]^2[X^2] + 3[X^2]^2 + 8[X][X^3] - 6[X^4]), \quad (2.7e)$$

we can write, still in symmetric guise,

$$\sqrt{-g_{\text{eff}}} = \sqrt{-g} \sum_{n=0}^4 \alpha^{4-n} \beta^n e_n(X) = \sqrt{-f} \sum_{n=0}^4 \alpha^n \beta^{4-n} e_n(Y). \quad (2.8)$$

⁴As usual we denote the traces of rank-2 tensors as $X_\mu^\mu = [X]$, $X_\nu^\nu X_\mu^\mu = [X^2]$, and so on. Recall in particular that $\sqrt{-g} e_n(X) = \sqrt{-f} e_{4-n}(Y)$, with $e_4(X) = \det(X)$ and $e_4(Y) = \det(Y)$.

In this form we see that (2.6) corresponds to the right form of the acceptable potential interactions between the metrics g and f .

We can first vary the Lagrangian (2.1) with respect to the two metrics g and f , which yields the following two covariant Einstein field equations [90]

$$\begin{aligned} M_g^2 Y_\rho^{(\mu} G_g^{\nu)\rho} - m^2 M_{\text{eff}}^2 \sum_{n=0}^3 \alpha^{4-n} \beta^n Y_\rho^{(\mu} U_{(n)}^{\nu)\rho} \\ = Y_\rho^{(\mu} \left(T_{\text{bar}}^{\nu)\rho} + T_g^{\nu)\rho} \right) + \alpha \frac{\sqrt{-g_{\text{eff}}}}{\sqrt{-g}} \left(\alpha Y_\rho^{(\mu} T_{g_{\text{eff}}}^{\nu)\rho} + \beta T_{g_{\text{eff}}}^{\mu\nu} \right), \end{aligned} \quad (2.9a)$$

$$\begin{aligned} M_f^2 X_\rho^{(\mu} G_f^{\nu)\rho} - m^2 M_{\text{eff}}^2 \sum_{n=0}^3 \alpha^n \beta^{4-n} X_\rho^{(\mu} V_{(n)}^{\nu)\rho} \\ = X_\rho^{(\mu} T_f^{\nu)\rho} + \beta \frac{\sqrt{-g_{\text{eff}}}}{\sqrt{-f}} \left(\beta X_\rho^{(\mu} T_{g_{\text{eff}}}^{\nu)\rho} + \alpha T_{g_{\text{eff}}}^{\mu\nu} \right), \end{aligned} \quad (2.9b)$$

where $G_g^{\mu\nu}$ and $G_f^{\mu\nu}$ are the Einstein tensors for the g and f metrics. The tensors $U_{(n)}^{\mu\nu}$ and $V_{(n)}^{\mu\nu}$ are defined from the following sums of matrices [8]⁵

$$U_{(n)} = \sum_{p=0}^n (-)^p e_{n-p}(X) X^p, \quad (2.10a)$$

$$V_{(n)} = \sum_{p=0}^n (-)^p e_{n-p}(Y) Y^p, \quad (2.10b)$$

by raising or lowering indices with their respective metrics, thus $U_{(n)}^{\mu\nu} = g^{\mu\rho} U_{(n)\rho}^\nu$ and $V_{(n)}^{\mu\nu} = f^{\mu\rho} V_{(n)\rho}^\nu$. By the same property which makes (2.5) to be symmetric, one can show that $U_{(n)}^{\mu\nu}$ and $V_{(n)}^{\mu\nu}$ are indeed automatically symmetric.

In the right sides of (2.9) the stress-energy tensors $T_{\text{bar}}^{\mu\nu}$ and $T_g^{\mu\nu}$ are defined with respect to the metric g , $T_f^{\mu\nu}$ is defined with respect to f and $T_{g_{\text{eff}}}^{\mu\nu}$ with respect to g_{eff} . Thus, for pressureless baryonic and dark matter fluids we have $T_{\text{bar}}^{\mu\nu} = \rho_{\text{bar}} u_{\text{bar}}^\mu u_{\text{bar}}^\nu$, $T_g^{\mu\nu} = \rho_g u_g^\mu u_g^\nu$ and $T_f^{\mu\nu} = \rho_f u_f^\mu u_f^\nu$ in terms of the corresponding scalar densities and normalized four velocities, while $T_{g_{\text{eff}}}^{\mu\nu}$ represents in fact the stress-energy tensor of the vector field \mathcal{A}_μ , *i.e.*

$$T_{g_{\text{eff}}}^{\mu\nu} = M_{\text{eff}}^2 \left[\mathcal{W}_{\mathcal{X}} \mathcal{F}^{\mu\rho} \mathcal{F}^\nu_\rho + \lambda \mathcal{W} g_{\text{eff}}^{\mu\nu} \right], \quad (2.11)$$

where $\mathcal{W}_{\mathcal{X}}$ is the derivative of \mathcal{W} with respect to its argument \mathcal{X} defined by (2.3).

The equation of motion for the baryons follows a geodesic law $a_\mu^{\text{bar}} \equiv u_{\text{bar}}^\nu \nabla_\nu^g u_\mu^{\text{bar}} = 0$, whereas the equations of motion for the dark matter particles are non-geodesic,

$$a_\mu^g = r_g u_g^\nu \mathcal{F}_{\mu\nu}, \quad (2.12a)$$

$$a_\mu^f = -r_f u_f^\nu \mathcal{F}_{\mu\nu}, \quad (2.12b)$$

⁵Here we adopt a slight change of notation with respect to [8]. Our notation is related to the one of [8] by $U_{(n)} = (-)^n Y_{(n)}(X)$ and $V_{(n)} = (-)^n Y_{(n)}(Y)$.

where the accelerations $a_\mu^g \equiv u_g^\nu \nabla_\nu^g u_\mu^g$ and $a_\mu^f \equiv u_f^\nu \nabla_\nu^f u_\mu^f$ are defined with their respective metrics. On the other hand, the equation for the vector field yields

$$\nabla_\nu^{g_{\text{eff}}} \left[\mathcal{W}_{\mathcal{X}} \mathcal{F}^{\mu\nu} \right] = \frac{1}{M_{\text{eff}}^2} (j_g^\mu - j_f^\mu), \quad (2.13)$$

with $\nabla_\nu^{g_{\text{eff}}}$ denoting the covariant derivative associated with g_{eff} . This equation is obviously compatible with the conservation of the currents, since by (2.2) we have also $\nabla_\mu^{g_{\text{eff}}} j_g^\mu = 0$ and $\nabla_\mu^{g_{\text{eff}}} j_f^\mu = 0$. The equation (2.13) can be also written as

$$\nabla_{g_{\text{eff}}}^\nu T_{\mu\nu}^{g_{\text{eff}}} = -(j_g^\nu - j_f^\nu) \mathcal{F}_{\mu\nu}. \quad (2.14)$$

Combining the equations of motion (2.12) with (2.14) we obtain the conservation law

$$\sqrt{-g} \nabla_g^\nu T_{\mu\nu}^g + \sqrt{-f} \nabla_f^\nu T_{\mu\nu}^f + \sqrt{-g_{\text{eff}}} \nabla_{g_{\text{eff}}}^\nu T_{\mu\nu}^{g_{\text{eff}}} = 0. \quad (2.15)$$

Alternatively, such global conservation law can be obtained from the scalarity of the total matter action under general diffeomorphisms.

3 Linear perturbations

In this section we will be interested in computing the linear perturbations of our model, which will be extensively used in the following section for the study of the polarisation mechanism. Following the ideas of [87] we will assume that the two fluids of dark matter particles are slightly displaced from the equilibrium configuration by displacement vectors y_g^μ and y_f^μ . This makes the dark matter medium to act as an analogue of a relativistic plasma in electromagnetism. The dark matter currents will be slightly displaced from the equilibrium current $j_0^\mu = \rho_0 u_0^\mu$ as well, with j_0^μ satisfying $\nabla_\mu^{g_{\text{eff}}} j_0^\mu = 0$. Furthermore, we will perturb the two metrics around a general background in the following way⁶

$$g_{\mu\nu} = (\bar{g}_{\mu\nu} + h_{\mu\nu})^2, \quad (3.1a)$$

$$f_{\mu\nu} = (\bar{f}_{\mu\nu} + \ell_{\mu\nu})^2, \quad (3.1b)$$

with main interest in background solutions where $\bar{g}_{\mu\nu} = \bar{f}_{\mu\nu}$. In the latter case the potential interactions take the form

$$\sqrt{-g_{\text{eff}}} = \sqrt{-\bar{g}} \sum_{n=0}^4 (\alpha + \beta)^{4-n} e_n(k), \quad (3.2)$$

where $k_{\mu\nu} = \alpha h_{\mu\nu} + \beta \ell_{\mu\nu}$, and traces are defined with the common background metric $\bar{g} = \bar{f}$.

Working at first order in the displacement of the dark matter particles and assuming that their gradients are of the same order as the metric perturbations, the dark matter currents can then be expanded as [87]

$$j_g^\mu = j_0^\mu + 2 \nabla_\nu^{g_{\text{eff}}} (j_0^{[\nu} \perp_\rho^{\mu]} y_g^\rho) + \mathcal{O}(2), \quad (3.3a)$$

$$j_f^\mu = j_0^\mu + 2 \nabla_\nu^{g_{\text{eff}}} (j_0^{[\nu} \perp_\rho^{\mu]} y_f^\rho) + \mathcal{O}(2), \quad (3.3b)$$

⁶By which we mean, for instance, $g_{\mu\nu} = (\bar{g}_{\mu\rho} + h_{\mu\rho}) \bar{g}^{\rho\sigma} (\bar{g}_{\sigma\nu} + h_{\sigma\nu})$.

with the projector operator perpendicular to the four velocity of the equilibrium u_0^μ given as $\perp^{\mu\nu} = g_{\text{eff}}^{\mu\nu} + u_0^\mu u_0^\nu$, and the remainder $\mathcal{O}(2)$ indicating that the expansion is at first order. The desired plasma-like solution is achieved after plugging our ansatz (3.3) into the equations of the vector field (2.13), which results in

$$\mathcal{W}_{\mathcal{X}} \mathcal{F}^{\mu\nu} = -\frac{2}{M_{\text{eff}}^2} j_0^{[\mu} \xi^{\nu]} + \mathcal{O}(2), \quad (3.4)$$

where we have defined the projected relative displacement as $\xi^\mu \equiv \perp^\mu_\nu (y_g^\nu - y_f^\nu)$. Note that ξ^μ is necessarily a space-like vector. In Sec. 4 we shall see that the spatial components ξ^i of this vector, which can be called a dipole moment, define in the non-relativistic limit the polarisation field of the dark matter medium as $P^i = \rho_0^* \xi^i$, where ρ_0^* is the coordinate density associated with ρ_0 . But for the moment we only need to notice that ξ^μ is a first order quantity, therefore the field strength $\mathcal{F}^{\mu\nu}$ is itself a first order quantity, and hence the stress-energy tensor (2.11) with respect to g_{eff} is already of second order,⁷

$$T_{g_{\text{eff}}}^{\mu\nu} = \mathcal{O}(2). \quad (3.5)$$

For this reason, in the case of our desired plasma-like solution the contributions of the self-interactions of the internal vector field will be at least third order in perturbation at the level of the action. For more detail see the comprehensive derivations in [87].

Let us first expand the Lagrangian (2.1) to first order in perturbation around a common background $\bar{g} = \bar{f}$. We assume that there is no matter in the background, so the matter interactions do not contribute at that order. Using (3.2) we obtain (modulo a total divergence)

$$\mathcal{L} = \mathcal{L}_{\bar{g}} + \sqrt{-\bar{g}} \left[- (M_g^2 h^{\mu\nu} + M_f^2 \ell^{\mu\nu}) G_{\mu\nu}^{\bar{g}} + m^2 M_{\text{eff}}^2 (\alpha + \beta)^3 [k] \right] + \mathcal{O}(2), \quad (3.6)$$

where $G_{\mu\nu}^{\bar{g}}$ is the Einstein tensor for the background metric \bar{g} , and $\mathcal{L}_{\bar{g}}$ is the background value of the Lagrangian. In order for the background $\bar{g}_{\mu\nu}$ in our interest to be a solution of the theory, we have to impose that linear perturbations vanish. We find

$$G_{\mu\nu}^{\bar{g}} = m^2 \frac{M_{\text{eff}}^2}{M_g^2} \alpha (\alpha + \beta)^3 \bar{g}_{\mu\nu} = m^2 \frac{M_{\text{eff}}^2}{M_f^2} \beta (\alpha + \beta)^3 \bar{g}_{\mu\nu}. \quad (3.7)$$

This is only possible if

$$\frac{\alpha}{M_g^2} = \frac{\beta}{M_f^2}. \quad (3.8)$$

With this choice we guarantee that we are expanding around the background $\bar{g}_{\mu\nu} = \bar{f}_{\mu\nu}$ that is a solution to the background equations of motion.

As next we shall compute the action to second order in perturbations. Let us emphasize again that thanks to the plasma-like ansatz (3.3) the self-interaction of the internal vector field

⁷We are anticipating the form of \mathcal{W} given by (4.23) and which implies $\mathcal{W} = \mathcal{O}(2)$ and $\mathcal{W}_{\mathcal{X}} = 1 + \mathcal{O}(1)$.

does not contribute to this order, see (3.5). Our quadratic Lagrangian above the background $\bar{g}_{\mu\nu}$ simply becomes

$$\mathcal{L} = \mathcal{L}_{\bar{g}} + \sqrt{-\bar{g}} \left[-M_g^2 h^{\rho\sigma} \bar{\mathcal{E}}_{\rho\sigma}^{\mu\nu} h_{\mu\nu} - M_f^2 \ell^{\rho\sigma} \bar{\mathcal{E}}_{\rho\sigma}^{\mu\nu} \ell_{\mu\nu} + h_{\mu\nu} (T_{\text{bar}}^{\mu\nu} + T_g^{\mu\nu}) + \ell_{\mu\nu} T_f^{\mu\nu} + \frac{1}{2} m^2 M_{\text{eff}}^2 (\alpha + \beta)^2 ([k]^2 - [k^2]) \right] + \mathcal{O}(3), \quad (3.9)$$

where $\bar{\mathcal{E}}_{\mu\nu}^{\rho\sigma}$ is the Lichnerowicz operator on the background \bar{g} given in the general case by

$$\begin{aligned} -2\bar{\mathcal{E}}_{\mu\nu}^{\rho\sigma} h_{\rho\sigma} &= \square_{\bar{g}}(h_{\mu\nu} - \bar{g}_{\mu\nu} h) + \nabla_{\mu}^{\bar{g}} \nabla_{\nu}^{\bar{g}} h - 2\nabla_{(\mu}^{\bar{g}} H_{\nu)} + \bar{g}_{\mu\nu} \nabla_{\rho}^{\bar{g}} H^{\rho} \\ &\quad - 2C_{\mu\rho\sigma\nu}^{\bar{g}} h^{\rho\sigma} - \frac{2}{3} R_{\bar{g}} \left(h^{\mu\nu} - \frac{1}{4} \bar{g}_{\mu\nu} h \right), \end{aligned} \quad (3.10)$$

where $h = [h] = \bar{g}^{\mu\nu} h_{\mu\nu}$, $H_{\mu} = \nabla_{\bar{g}}^{\nu} h_{\mu\nu}$, with $C_{\mu\rho\sigma\nu}^{\bar{g}}$ denoting the Weyl curvature of the background metric. Of course this expression is to be simplified using the background equations (3.7). Also bear in mind that $\alpha M_f^2 = \beta M_g^2$ for having a common background $\bar{g} = \bar{f}$. The field equations obtained by varying with respect to $h_{\mu\nu}$ and $\ell_{\mu\nu}$ yield

$$-2M_g^2 \bar{\mathcal{E}}_{\rho\sigma}^{\mu\nu} h^{\rho\sigma} + T_{\text{bar}}^{\mu\nu} + T_g^{\mu\nu} + \alpha(\alpha + \beta)^2 m^2 M_{\text{eff}}^2 (\bar{g}^{\mu\nu} k - k^{\mu\nu}) = \mathcal{O}(2), \quad (3.11a)$$

$$-2M_f^2 \bar{\mathcal{E}}_{\rho\sigma}^{\mu\nu} \ell^{\rho\sigma} + T_f^{\mu\nu} + \beta(\alpha + \beta)^2 m^2 M_{\text{eff}}^2 (\bar{g}^{\mu\nu} k - k^{\mu\nu}) = \mathcal{O}(2), \quad (3.11b)$$

where we recall that $k^{\mu\nu} = \alpha h^{\mu\nu} + \beta \ell^{\mu\nu}$ and $k = [k] = \bar{g}_{\mu\nu} k^{\mu\nu}$. Of course those equations can also be recovered directly from the general Einstein field equations (2.9).

In the present paper we shall mostly make use of that combination of the two Einstein field equations (3.11) which corresponds to the propagation of a massless spin-2 field. We subtract the two equations from each other so as to cancel the mass term, resulting in

$$-2\bar{\mathcal{E}}_{\rho\sigma}^{\mu\nu} \left(\frac{M_g^2}{\alpha} h^{\rho\sigma} - \frac{M_f^2}{\beta} \ell^{\rho\sigma} \right) + \frac{1}{\alpha} (T_{\text{bar}}^{\mu\nu} + T_g^{\mu\nu}) - \frac{1}{\beta} T_f^{\mu\nu} = \mathcal{O}(2). \quad (3.12)$$

However we shall also use the contribution of the mass term in the linearised Bianchi identities associated with (3.11). Thus, last but not least we can act on (3.11) with $\nabla_{\nu}^{\bar{g}}$, yielding

$$\nabla_{\nu}^{\bar{g}} (k^{\mu\nu} - \bar{g}^{\mu\nu} k) = \mathcal{O}(2). \quad (3.13)$$

In deriving this relation we use the result that, for instance, $\nabla_{\nu}^{\bar{g}} T_g^{\mu\nu} = \rho_g a_g^{\mu} = \mathcal{O}(2)$ which comes from the equations of motion (2.12) and the fact that there is no matter in the background. We shall see in the next section how important is this relation for the present model to recover the looked-for phenomenology of dark matter at galactic scales.

4 Polarisation mechanism and MOND

We next consider the Newtonian or non-relativistic (NR) limit of our model. In the previous section we derived the field equations at linear order around a vacuum background metric \bar{g} ,

which necessarily obeys the equations (3.7). Thus, the background is a de Sitter solution, with cosmological constant given by the graviton's mass, $\Lambda \sim m^2$. When computing the Newtonian limit applied to describe the physics of the local universe, *e.g.* the solar system or a galaxy at low redshift, we shall neglect the cosmological constant and shall approximate the de Sitter metric \bar{g} by a flat Minkowski background η .

The best way to implement (and justify) this approximation is to perform an expansion on small scales, say $r \rightarrow 0$. When we solve for the gravitational field in the solar system or a galaxy embedded in the de Sitter background, we get terms like $1 - 2Mr^{-1} - \Lambda r^2/3$ (de Sitter-Schwarzschild solution), where M is the mass of the Sun or the galaxy. On small scales the term $2Mr^{-1}$ will dominate over $\Lambda r^2/3$, so we can ignore the influence of the cosmological constant if the size of the system is $\ll (6M/\Lambda)^{1/3}$. In the present case we shall require that the size of our system includes the very weak field region far from the system where the MOND formula applies. In that case the relevant scale is $r_0 = \sqrt{M/a_0}$, where $a_0 \simeq 1.2 \times 10^{-10} \text{ m/s}^2$ is the MOND acceleration. So our approximation will make sense provided that $r_0 \ll (6M/\Lambda)^{1/3}$. For a galaxy with mass $M \sim 10^{11} M_\odot$ the MOND transition radius is $r_0 \sim 10 \text{ kpc}$. With the value of the cosmological constant $\Lambda \sim (3 \text{ Gpc})^{-2}$ we find $(6M/\Lambda)^{1/3} \sim 700 \text{ kpc}$, so we are legitimate to neglect the influence of the cosmological constant. For the solar system, $r_0 \sim 0.04 \text{ pc}$ while $(6M_\odot/\Lambda)^{1/3} \sim 150 \text{ pc}$ and the approximation is even better.

Actually we shall make the approximation $\bar{g} \simeq \eta$ only on the particular massless spin-2 combination of the two metrics given by (3.12), namely

$$-2\bar{\mathcal{E}}^{\mu\nu}_{\rho\sigma} \left(\frac{M_g^2}{\alpha} h^{\rho\sigma} - \frac{M_f^2}{\beta} \ell^{\rho\sigma} \right) = -\frac{1}{\alpha} (T_{\text{bar}}^{\mu\nu} + T_g^{\mu\nu}) + \frac{1}{\beta} T_f^{\mu\nu} + \mathcal{O}(2), \quad (4.1)$$

where the Lichnerowicz operator (3.10) is now the flat one, and also in the constraint equation (3.13). We shall not need to consider the other, independent (massive) combination of the two metrics. In the NR limit when $c \rightarrow \infty$ we parametrize the two metric perturbations $h_{\mu\nu}$ and $\ell_{\mu\nu}$ by single Newtonian potentials U_g and U_f such that

$$h_{00} = \frac{U_g}{c^2} + \mathcal{O}\left(\frac{1}{c^4}\right), \quad \ell_{00} = \frac{U_f}{c^2} + \mathcal{O}\left(\frac{1}{c^4}\right), \quad (4.2a)$$

$$h_{0i} = \mathcal{O}\left(\frac{1}{c^3}\right), \quad \ell_{0i} = \mathcal{O}\left(\frac{1}{c^3}\right), \quad (4.2b)$$

$$h_{ij} = \delta_{ij} \frac{U_g}{c^2} + \mathcal{O}\left(\frac{1}{c^4}\right), \quad \ell_{ij} = \delta_{ij} \frac{U_f}{c^2} + \mathcal{O}\left(\frac{1}{c^4}\right). \quad (4.2c)$$

We shall especially pay attention to the potential U_g since it represents the ordinary Newtonian potential felt by ordinary baryonic matter. Similarly we parametrize the internal vector field \mathcal{A}_μ by a single Coulomb scalar potential ϕ such that

$$\mathcal{A}_0 = \frac{\phi}{c^2} + \mathcal{O}\left(\frac{1}{c^4}\right), \quad (4.3a)$$

$$\mathcal{A}_i = \mathcal{O}\left(\frac{1}{c^3}\right). \quad (4.3b)$$

It is now straightforward to show that (4.1) reduces in the NR limit to a single scalar equation for a combination of the Newtonian potentials U_g and U_f ,⁸

$$\Delta \left(\frac{2M_g^2}{\alpha} U_g - \frac{2M_f^2}{\beta} U_f \right) = -\frac{1}{\alpha} (\rho_{\text{bar}}^* + \rho_g^*) + \frac{1}{\beta} \rho_f^*, \quad (4.4)$$

where $\Delta = \nabla^2$ is the ordinary Laplacian, and ρ_{bar}^* , ρ_g^* and ρ_f^* are the ordinary Newtonian (coordinate) densities, satisfying usual continuity equations such as $\partial_t \rho_g^* + \nabla \cdot (\rho_g^* \mathbf{v}_g) = 0$. Notably, all non-linear corrections $\mathcal{O}(2)$ in (4.1) are negligible. In principle, we should *a priori* allow some parametrized post-Newtonian (PPN) coefficients in the spatial components ij of the metrics (4.2), say γ_g and γ_f . However, when plugging (4.2) into (4.1) we determine from the ij components of (4.1) that in fact $\gamma_g = 1$ and $\gamma_f = 1$. Similarly we find that the equations (2.13) governing the internal vector field reduce to a single Coulomb type equation,

$$\nabla \cdot [\mathcal{W}_{\mathcal{X}} \nabla \phi] = \frac{1}{M_{\text{eff}}^2} (r_g \rho_g^* - r_f \rho_f^*). \quad (4.5)$$

Notice that the constants α and β in the effective metric g_{eff} to which is coupled the internal field intervene in the NR limit only in the expression

$$\mathcal{X} = \frac{|\nabla \phi|^2}{2\lambda(\alpha + \beta)^4}. \quad (4.6)$$

Next, the equations of the baryons which are geodesic in the metric g reduce to

$$\frac{d\mathbf{v}_{\text{bar}}}{dt} = \nabla U_g, \quad (4.7)$$

while the equations of motion (2.12) of the dark matter particles become

$$\frac{d\mathbf{v}_g}{dt} = \nabla U_g + r_g \nabla \phi, \quad (4.8a)$$

$$\frac{d\mathbf{v}_f}{dt} = \nabla U_f - r_f \nabla \phi. \quad (4.8b)$$

At this stage the important point is to recall that the two sectors associated with the metrics g and f do not evolve independently but are linked together by the constraint equation (3.13). We now show that this constraint provides a mechanism of gravitational polarisation in the NR limit and yields the MOND equation for the potential U_g felt by baryons. In flat space-time this constraint is

$$\partial_\nu (k^{\mu\nu} - \eta^{\mu\nu} k) = \mathcal{O}(2), \quad (4.9)$$

where $k^{\mu\nu} = \alpha h^{\mu\nu} + \beta \ell^{\mu\nu}$ and $k = \eta^{\rho\sigma} k_{\rho\sigma}$. Plugging (4.2) into (4.9) we readily obtain a constraint on the gravitational forces felt by the DM particles, namely

$$\nabla (\alpha U_g + \beta U_f) = 0. \quad (4.10)$$

⁸We shall adopt the usual boldface notation for ordinary three-dimensional Euclidean vectors. Also, from now on we no longer write the neglected remainders $\mathcal{O}(1/c^2)$.

Note that the constraint (4.10) comes from a combination between the 00 and ij components of the metrics (4.2). Using (4.10) we express the equations of motion (4.8) solely in terms of the potential U_g ruling the baryons,

$$\frac{d\mathbf{v}_g}{dt} = \nabla U_g + r_g \nabla \phi, \quad (4.11a)$$

$$\frac{d\mathbf{v}_f}{dt} = -\frac{\alpha}{\beta} \nabla U_g - r_f \nabla \phi. \quad (4.11b)$$

As we see, the gravitational to inertial mass ratio m_g/m_i of the f particles when measured with respect to the g metric is $m_g/m_i = -\alpha/\beta$.

We look for explicit solutions of (4.11) in the form of plasma-like oscillations around some equilibrium solution. We immediately see from (4.11) the possibility of an equilibrium (*i.e.* for which $d\mathbf{v}_g/dt = d\mathbf{v}_f/dt = 0$) when we have the following relation between constants,

$$\frac{\alpha}{\beta} = \frac{r_f}{r_g}. \quad (4.12)$$

The equilibrium holds when $\nabla U_g + r_g \nabla \phi = 0$, *i.e.* when the Coulomb force annihilates the gravitational force. To describe in the proper way the two DM fluids near or at equilibrium we use the NR limits of the relations (3.3), with an appropriate choice of the equilibrium configuration. From (4.11) we note that $(\alpha + \beta)\mathbf{v}_0 = \alpha\mathbf{v}_g + \beta\mathbf{v}_f$ is constant, hence we define the equilibrium in such a way that the two displacement vectors \mathbf{y}_g and \mathbf{y}_f with respect to that equilibrium obey $\alpha\mathbf{y}_g + \beta\mathbf{y}_f = \mathbf{0}$, and in particular we choose the equilibrium velocity to be $\mathbf{v}_0 = \mathbf{0}$. We shall now define the relative displacement or dipole moment vector by $\boldsymbol{\xi} = \mathbf{y}_g - \mathbf{y}_f$.⁹ Hence (3.3) imply

$$r_g \rho_g^* = \rho_0^* - \frac{\beta}{\alpha + \beta} \nabla \cdot \mathbf{P}, \quad (4.13a)$$

$$r_f \rho_f^* = \rho_0^* + \frac{\alpha}{\alpha + \beta} \nabla \cdot \mathbf{P}, \quad (4.13b)$$

where ρ_0^* is the common density of the two DM fluids at equilibrium in the absence of external perturbations (far from any external mass), and $\mathbf{P} = \rho_0^* \boldsymbol{\xi}$ is the polarisation. The velocity fields of the two fluids are

$$\mathbf{v}_g = \frac{\beta}{\alpha + \beta} \frac{d\boldsymbol{\xi}}{dt}, \quad \mathbf{v}_f = -\frac{\alpha}{\alpha + \beta} \frac{d\boldsymbol{\xi}}{dt}, \quad (4.14)$$

where $d\boldsymbol{\xi}/dt = \partial_i \boldsymbol{\xi} + \mathbf{v}_0 \cdot \nabla \boldsymbol{\xi}$ denotes the convective derivative, which reduces here to the ordinary derivative since $\mathbf{v}_0 = \mathbf{0}$.

By inserting (4.13) into (4.5) we can solve for the polarisation resulting in

$$\mathbf{P} = -M_{\text{eff}}^2 \mathcal{W}_X \nabla \phi. \quad (4.15)$$

⁹Here \mathbf{y}_g and \mathbf{y}_f denote the spatial components of the displacement four vectors $\perp_{\nu}^{\mu} y_g^{\nu}$ and $\perp_{\nu}^{\mu} y_f^{\nu}$ considered in Sec. 3, see (3.3)–(3.4).

Furthermore, by combining (4.14) and (4.11) and making use of the previous solution for the polarisation (4.15), we readily arrive at a simple harmonic oscillator describing plasma like oscillations around equilibrium, namely

$$\frac{d^2 \boldsymbol{\xi}}{dt^2} + \omega_g^2 \boldsymbol{\xi} = \frac{\alpha + \beta}{\beta} \nabla U_g. \quad (4.16)$$

In this case the plasma frequency is given by

$$\omega_g = \sqrt{\frac{\alpha + \beta}{\beta} \frac{r_g \rho_0^*}{M_{\text{eff}}^2 \mathcal{W}_{\mathcal{X}}}}. \quad (4.17)$$

Next, consider the equation for the Newtonian potential U_g in the ordinary sector. Combining (4.10) with (4.4) we readily obtain

$$\Delta U_g = -\frac{1}{2 \left(M_g^2 + \frac{\alpha^2}{\beta^2} M_f^2 \right)} \left[\rho_{\text{bar}}^* + \rho_g^* - \frac{\alpha}{\beta} \rho_f^* \right]. \quad (4.18)$$

To recover the usual Newtonian limit we must impose, in geometrical units,

$$M_g^2 + \frac{\alpha^2}{\beta^2} M_f^2 = \frac{1}{8\pi}. \quad (4.19)$$

This condition being satisfied, the right-hand side of (4.18) can be rewritten with the help of (4.13) and the relation (4.12). We obtain an ordinary Poisson equation but modified by polarisation effects,

$$\nabla \cdot \left[\nabla U_g - 4\pi \frac{\mathbf{P}}{r_g} \right] = -4\pi \rho_{\text{bar}}^*. \quad (4.20)$$

We still have to show that the polarisation will be aligned with the gravitational field ∇U_g , *i.e.* we grasp a mechanism of gravitational polarisation. This is a consequence of the constitutive relation (4.15) taken at the equilibrium point, neglecting plasma like oscillations. At this point we have $\nabla U_g + r_g \nabla \phi = 0$. Thus,

$$\mathbf{P} = \frac{M_{\text{eff}}^2}{r_g} \mathcal{W}_{\mathcal{X}} \nabla U_g. \quad (4.21)$$

Finally (4.20) with (4.21) takes exactly the form of the Bekenstein-Milgrom equation [64]. To recover the correct deep MOND regime we must impose that when $\mathcal{X} \rightarrow 0$

$$1 - 4\pi \frac{M_{\text{eff}}^2}{r_g^2} \mathcal{W}_{\mathcal{X}} = \frac{|\nabla U_g|}{a_0}, \quad (4.22)$$

where a_0 is the MOND acceleration scale. This is easily achieved with the choice

$$\mathcal{W}(\mathcal{X}) = \mathcal{X} - \frac{2}{3}(\alpha + \beta)^2 \mathcal{X}^{3/2} + \mathcal{O}(\mathcal{X}^2), \quad (4.23)$$

together with fixing the constants M_{eff} and λ to the values

$$M_{\text{eff}}^2 = \frac{r_g^2}{4\pi}, \quad \lambda = \frac{a_0^2}{2}. \quad (4.24)$$

We thus recovered gravitational polarisation and the MOND equation (in agreement with the dielectric analogy of MOND [82]) as a natural consequence of bimetric gravity since it is made possible by the constraint equation (4.9) linking together the two metrics of bigravity.¹⁰ The polarisation mechanism is possible only if we can annihilate the gravitational force by some internal force, here chosen to be a vector field, and therefore assume a coupling between the two species of DM particles living in the g and f sectors.

Let us recapitulate the various constraints we have found on the parameters in the original action (2.1). These are given by (4.12) for having a polarisation process, (4.19) for recovering the Poisson equation and (4.24) for having the correct deep MOND regime. In addition, we recall (3.8) which was imposed in order to be able to expand the two metrics around the same background. Strictly speaking the relation (3.8) is not necessary for the present calculation because we neglected the influence of the background, and (3.8) may be relaxed for some applications. Finally, we note that the result can be simplified by absorbing the remaining constant r_g together with the sum $\alpha + \beta$ into the following redefinitions of the vector field and the effective metric: $\mathcal{A}_\mu \rightarrow r_g \mathcal{A}_\mu$ and $g_{\mu\nu}^{\text{eff}} \rightarrow (\alpha + \beta)^{-2} g_{\mu\nu}^{\text{eff}}$. When this is done, and after redefining also the graviton mass m , we see that we can always choose $r_g = 1$ and $\alpha + \beta = 1$ without loss of generality.

Finally the fully reduced form of the action (2.1) reads

$$\begin{aligned} \mathcal{L}_{\text{final}} = & \sqrt{-g} \left(\frac{M_g^2}{2} R_g - \rho_{\text{bar}} - \rho_g \left[1 - \mathcal{A}_\mu u_g^\mu \right] \right) + \sqrt{-f} \left(\frac{M_f^2}{2} R_f - \rho_f \left[1 + \frac{\alpha}{\beta} \mathcal{A}_\mu u_f^\mu \right] \right) \\ & + \sqrt{-g_{\text{eff}}} \left[\frac{m^2}{4\pi} + \frac{a_0^2}{8\pi} \mathcal{W}(\mathcal{X}) \right], \end{aligned} \quad (4.25)$$

where we have moved for convenience the mass currents $J_g^\mu = \rho_g u_g^\mu$ and $J_f^\mu = \frac{\alpha}{\beta} \rho_f u_f^\mu$ to the g and f sectors, where the effective metric $g_{\mu\nu}^{\text{eff}}$ of bigravity is given by (2.4) but with $\alpha + \beta = 1$, and where the kinetic term of the vector field is defined by

$$\mathcal{X} = -\frac{\mathcal{F}^{\mu\nu} \mathcal{F}_{\mu\nu}}{2a_0^2}, \quad (4.26a)$$

$$\mathcal{W}(\mathcal{X}) = \mathcal{X} - \frac{2}{3} \mathcal{X}^{3/2} + \mathcal{O}(\mathcal{X}^2). \quad (4.26b)$$

In addition the coupling constants M_g^2 and M_f^2 are constrained by (4.19). We can still further impose (3.8) if we insist that the two metrics g and f can be expanded around a common

¹⁰In the previous model [87] we had to assume that some coupling constant ε in the action tends to zero. With bimetric gravity and its nice potential interactions we see that no particular assumption is required.

background $\bar{g} = \bar{f}$.¹¹ Finally the theory depends also on the graviton's mass m , hopefully to be related to the observed cosmological constant Λ , and the MOND acceleration scale a_0 . Note that in such unified approach between the cosmological constant and MOND, it is natural to expect that a_0 and Λ should have comparable orders of magnitudes, *i.e.* $a_0 \sim \sqrt{\Lambda}$ which happens to be in very good agreement with observations.

5 The decoupling limit

In this section we would like to focus on the matter sector and address the question whether or not the interactions between the matter fields reintroduces the BD ghost [12]. For this we derive the decoupling limit and pay special attention to the helicity-0 mode. Before doing so, we first restore the broken gauge invariance by introducing the Stueckelberg fields in the f metric (with $a, b = 0, 1, 2, 3$)

$$f_{\mu\nu} \rightarrow \tilde{f}_{\mu\nu} = f_{ab} \partial_\mu \phi^a \partial_\nu \phi^b, \quad (5.1)$$

where the Stueckelberg fields can be further decomposed into their helicity-0 π and helicity-1 A^a counterparts,

$$\phi^a = x^a - \frac{mA^a}{\Lambda_3^3} - \frac{f^{ab} \partial_b \pi}{\Lambda_3^3}. \quad (5.2)$$

As next we can take the decoupling limit by sending $M_g, M_f \rightarrow \infty$, while keeping the scale $\Lambda_3^3 = M_g m^2$ constant. For our purpose, it is enough to keep track of the contributions of the helicity-0 mode π to the matter interactions. The BD ghost is hidden behind the higher derivative terms of the helicity-0 interactions after using all of the covariant equations of motion and constraints. We will therefore neglect the contribution of the helicity-1 field and assume $g_{\mu\nu} = \eta_{\mu\nu}$ and

$$f_{\mu\nu} = \eta_{\mu\nu} \rightarrow \tilde{f}_{\mu\nu} = (\eta_{\mu\nu} - \Pi_{\mu\nu})^2, \quad (5.3)$$

where $\Pi_{\mu\nu}$ stands for $\Pi_{\mu\nu} \equiv \partial_\mu \partial_\nu \pi / \Lambda_3^3$. In the remaining of this section all indices are raised and lowered with respect to the Minkowski metric $\eta_{\mu\nu}$. The effective metric in the decoupling limit corresponds to

$$g_{\mu\nu}^{\text{eff}} \rightarrow \tilde{g}_{\mu\nu}^{\text{eff}} = [(\alpha + \beta)\eta_{\mu\nu} - \beta\Pi_{\mu\nu}]^2. \quad (5.4)$$

In [88] the required criteria for ghost freedom in the kinetic and potential terms were investigated in detail and hence the new model was constructed in such a way that it fulfils these criteria. In the decoupling limit, the contribution to the equation of motion for the helicity-0 field coming from the potential term is at most second order in derivative for the allowed

¹¹A simple choice satisfying all these requirements is $\alpha = \beta = \frac{1}{2}$ and $M_g^2 = M_f^2 = \frac{1}{16\pi}$, *i.e.* each coupling constant takes half the GR value. With this choice the plasma frequency (4.17) becomes

$$\omega_g = \sqrt{\frac{8\pi\rho_0^*}{\mathcal{W}_X}},$$

in agreement with the finding of the previous model [87].

potentials. Therefore, we can concentrate on the problematic terms coming from the matter interactions. Let us for instance consider the coupling between $j_{\tilde{f}}^\mu$ and \mathcal{A}_μ , thus¹²

$$\mathcal{L}_{\text{mat}} = -\sqrt{-\tilde{f}}\rho_{\tilde{f}} - \sqrt{-\tilde{g}_{\text{eff}}}\mathcal{A}_\mu j_{\tilde{f}}^\mu + \frac{1}{4}\sqrt{-\tilde{g}_{\text{eff}}}\mathcal{F}_{\mu\nu}^2, \quad (5.5)$$

which we can also write as

$$\mathcal{L}_{\text{mat}} = -\sqrt{-\tilde{f}}\rho_{\tilde{f}}(1 + \mathcal{A}_\mu u_{\tilde{f}}^\mu) + \frac{1}{4}\sqrt{-\tilde{g}_{\text{eff}}}\mathcal{F}_{\mu\nu}^2. \quad (5.6)$$

Let us first compute the equation of motion with respect to the vector field, which is simply given by

$$\frac{\delta\mathcal{L}_{\text{mat}}}{\delta\mathcal{A}_\mu} = -\sqrt{\tilde{g}_{\text{eff}}}\nabla_\nu^{\tilde{g}_{\text{eff}}}\mathcal{F}^{\mu\nu} - \sqrt{-\tilde{f}}\rho_{\tilde{f}}u_{\tilde{f}}^\mu, \quad (5.7)$$

which can be also expressed as

$$\nabla_\nu^{\tilde{g}_{\text{eff}}}\mathcal{F}^{\mu\nu} = -\frac{\sqrt{-\tilde{f}}}{\sqrt{-\tilde{g}_{\text{eff}}}}\rho_{\tilde{f}}u_{\tilde{f}}^\mu = -j_{\tilde{f}}^\mu. \quad (5.8)$$

As next we can compute the contribution of the matter to the equation of motion with respect to the helicity-0 field,

$$\begin{aligned} \frac{\delta\mathcal{L}_{\text{mat}}}{\delta\pi} &= \frac{\partial_\mu\partial_\nu}{\Lambda_3^3}\left(\frac{\delta\mathcal{L}_{\text{mat}}}{\delta\tilde{f}_{\rho\sigma}}\frac{\delta\tilde{f}_{\rho\sigma}}{\delta\Pi_{\mu\nu}}\right) + \frac{\partial_\mu\partial_\nu}{\Lambda_3^3}\left(\frac{\delta\mathcal{L}_{\text{mat}}}{\delta\tilde{g}_{\rho\sigma}^{\text{eff}}}\frac{\delta\tilde{g}_{\rho\sigma}^{\text{eff}}}{\delta\Pi_{\mu\nu}}\right) \\ &= -\frac{\partial_\mu\partial_\nu}{\Lambda_3^3}\left(\sqrt{-\tilde{f}}T_{\tilde{f}}^{\mu\rho}(\delta_\rho^\nu - \Pi_\rho^\nu) + \beta\sqrt{-\tilde{g}_{\text{eff}}}T_{\tilde{g}_{\text{eff}}}^{\mu\rho}[(\alpha + \beta)\delta_\rho^\nu - \beta\Pi_\rho^\nu]\right), \end{aligned} \quad (5.9)$$

where we made use of

$$T_{\tilde{f}}^{\mu\nu} = \frac{2}{\sqrt{-\tilde{f}}}\frac{\delta\mathcal{L}_{\text{mat}}}{\delta\tilde{f}_{\mu\nu}}, \quad \text{and} \quad T_{\tilde{g}_{\text{eff}}}^{\mu\nu} = \frac{2}{\sqrt{-\tilde{g}_{\text{eff}}}}\frac{\delta\mathcal{L}_{\text{mat}}}{\delta\tilde{g}_{\mu\nu}^{\text{eff}}}. \quad (5.10)$$

Applying one of the derivatives we have

$$\begin{aligned} \frac{\delta\mathcal{L}_{\text{mat}}}{\delta\pi} &= -\frac{\partial_\nu}{\Lambda_3^3}\left\{\partial_\mu(\sqrt{-\tilde{f}}T_{\tilde{f}}^{\mu\rho})(\delta_\rho^\nu - \Pi_\rho^\nu) - \sqrt{-\tilde{f}}T_{\tilde{f}}^{\mu\rho}\partial_\mu\Pi_\rho^\nu\right. \\ &\quad \left.+ \beta\partial_\mu(\sqrt{-\tilde{g}_{\text{eff}}}T_{\tilde{g}_{\text{eff}}}^{\mu\rho})[(\alpha + \beta)\delta_\rho^\nu - \beta\Pi_\rho^\nu] - \beta^2\sqrt{-\tilde{g}_{\text{eff}}}T_{\tilde{g}_{\text{eff}}}^{\mu\rho}\partial_\mu\Pi_\rho^\nu\right\}. \end{aligned} \quad (5.11)$$

The equation of motion for the vector field (5.8) can also be written as

$$\nabla_{\tilde{g}_{\text{eff}}}^\mu T_{\rho\mu}^{\tilde{g}_{\text{eff}}} = j_{\tilde{f}}^\mu \mathcal{F}_{\rho\mu}, \quad (5.12)$$

¹²For simplicity we will drop the constants λ , r_f and M_{eff} and assume that $\mathcal{W}(\mathcal{X}) = \mathcal{X}$ in the following.

which we can use to express

$$\nabla_{\mu}^{\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\rho} = \frac{1}{\sqrt{-\tilde{g}_{\text{eff}}}} \partial_{\mu} \left(\sqrt{-\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\rho} \right) + \Gamma_{\mu\sigma}^{\rho\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\sigma} = j_{\tilde{f}}^{\mu} \tilde{g}_{\text{eff}}^{\rho\kappa} \mathcal{F}_{\kappa\mu}, \quad (5.13)$$

hence we have

$$\partial_{\mu} \left(\sqrt{-\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\rho} \right) = \sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^{\mu} \tilde{g}_{\text{eff}}^{\rho\kappa} \mathcal{F}_{\kappa\mu} - \sqrt{-\tilde{g}_{\text{eff}}} \Gamma_{\mu\sigma}^{\rho\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\sigma}. \quad (5.14)$$

Furthermore, the conservation law [see (2.15)] gives

$$\sqrt{-\tilde{g}_{\text{eff}}} \nabla_{\mu}^{\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\rho} + \sqrt{-\tilde{f}} \nabla_{\mu}^{\tilde{f}} T_{\tilde{f}}^{\mu\rho} = 0. \quad (5.15)$$

Thus, we have further

$$\partial_{\mu} \left(\sqrt{-\tilde{f}} T_{\tilde{f}}^{\mu\rho} \right) = -\sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^{\mu} \tilde{f}^{\rho\kappa} \mathcal{F}_{\kappa\mu} - \sqrt{-\tilde{f}} \Gamma_{\mu\sigma}^{\rho\tilde{f}} T_{\tilde{f}}^{\mu\sigma}. \quad (5.16)$$

Using the equation of motion for the vector field and the conservation equation, the equation for the helicity-0 becomes

$$\begin{aligned} \frac{\delta \mathcal{L}_{\text{mat}}}{\delta \pi} = & -\frac{\partial_{\nu}}{\Lambda_3^3} \left\{ \left(-\sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^{\mu} \tilde{f}^{\rho\kappa} \mathcal{F}_{\kappa\mu} - \sqrt{-\tilde{f}} \Gamma_{\mu\sigma}^{\rho\tilde{f}} T_{\tilde{f}}^{\mu\sigma} \right) (\delta_{\rho}^{\nu} - \Pi_{\rho}^{\nu}) - \sqrt{-\tilde{f}} T_{\tilde{f}}^{\mu\rho} \partial_{\mu} \Pi_{\rho}^{\nu} \right. \\ & \left. + \beta \left(\sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^{\mu} \tilde{g}_{\text{eff}}^{\rho\kappa} \mathcal{F}_{\kappa\mu} - \sqrt{-\tilde{g}_{\text{eff}}} \Gamma_{\mu\sigma}^{\rho\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\sigma} \right) [(\alpha + \beta) \delta_{\rho}^{\nu} - \beta \Pi_{\rho}^{\nu}] - \beta^2 \sqrt{-\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\rho} \partial_{\mu} \Pi_{\rho}^{\nu} \right\}, \end{aligned} \quad (5.17)$$

which we can rewrite as

$$\begin{aligned} \frac{\delta \mathcal{L}_{\text{mat}}}{\delta \pi} = & -\frac{\partial_{\nu}}{\Lambda_3^3} \left\{ -\sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^{\mu} \tilde{f}^{\rho\kappa} \mathcal{F}_{\kappa\mu} (\delta_{\rho}^{\nu} - \Pi_{\rho}^{\nu}) - \sqrt{-\tilde{f}} T_{\tilde{f}}^{\mu\sigma} \mathcal{R}_{\mu\sigma}^{\nu} \right. \\ & \left. + \beta \sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^{\mu} \tilde{g}_{\text{eff}}^{\rho\kappa} \mathcal{F}_{\kappa\mu} [(\alpha + \beta) \delta_{\rho}^{\nu} - \beta \Pi_{\rho}^{\nu}] - \beta^2 \sqrt{-\tilde{g}_{\text{eff}}} T_{\tilde{g}_{\text{eff}}}^{\mu\sigma} \tilde{\mathcal{R}}_{\mu\sigma}^{\nu} \right\}, \end{aligned} \quad (5.18)$$

where we have introduced

$$\mathcal{R}_{\mu\sigma}^{\nu} = \Gamma_{\mu\sigma}^{\rho\tilde{f}} (\delta_{\rho}^{\nu} - \Pi_{\rho}^{\nu}) + \partial^{\nu} \Pi_{\mu\sigma}, \quad (5.19a)$$

$$\tilde{\mathcal{R}}_{\mu\sigma}^{\nu} = \Gamma_{\mu\sigma}^{\rho\tilde{g}_{\text{eff}}} [(\alpha + \beta) \delta_{\rho}^{\nu} - \beta \Pi_{\rho}^{\nu}] + \beta^2 \partial^{\nu} \Pi_{\mu\sigma}. \quad (5.19b)$$

Using the fact that the Christoffel symbol with respect to the \tilde{f} metric is given by

$$\Gamma_{\mu\sigma}^{\rho\tilde{f}} = -\tilde{f}^{\rho\kappa} (\delta_{\kappa}^{\nu} - \Pi_{\kappa}^{\nu}) \partial_{\nu} \Pi_{\mu\sigma}, \quad (5.20)$$

and further taking into account the following relation,

$$(\delta_{\rho}^{\mu} - \Pi_{\rho}^{\mu}) \tilde{f}^{\rho\sigma} (\delta_{\sigma}^{\nu} - \Pi_{\sigma}^{\nu}) = \eta^{\mu\nu}, \quad (5.21)$$

we see immediately that $\mathcal{R}_{\mu\sigma}^\nu = 0$. Similarly, using the definition of the Christoffel symbol with respect to the \tilde{g}_{eff} metric,

$$\Gamma_{\mu\sigma}^{\rho\tilde{g}_{\text{eff}}} = -\tilde{g}_{\text{eff}}^{\rho\kappa} [(\alpha + \beta)\delta_\kappa^\nu - \beta\Pi_\kappa^\nu] \partial_\nu \Pi_{\mu\sigma}, \quad (5.22)$$

and the relation

$$[(\alpha + \beta)\delta_\rho^\mu - \beta\Pi_\rho^\mu] \tilde{g}_{\text{eff}}^{\rho\sigma} [(\alpha + \beta)\delta_\sigma^\nu - \beta\Pi_\sigma^\nu] = \eta^{\mu\nu}, \quad (5.23)$$

we can show that also $\tilde{\mathcal{R}}_{\mu\sigma}^\nu = 0$. Thus, the contribution of the matter part to the equation of motion for the helicity-0 mode simplifies to

$$\frac{\delta\mathcal{L}_{\text{mat}}}{\delta\pi} = -\frac{\partial_\nu}{\Lambda_3^3} \left\{ -\sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^\mu \tilde{f}^{\rho\kappa} \mathcal{F}_{\kappa\mu} (\delta_\rho^\nu - \Pi_\rho^\nu) + \beta \sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^\mu \tilde{g}_{\text{eff}}^{\rho\kappa} \mathcal{F}_{\kappa\mu} [(\alpha + \beta)\delta_\rho^\nu - \beta\Pi_\rho^\nu] \right\}. \quad (5.24)$$

Using (5.21) this can be equally written as

$$\frac{\delta\mathcal{L}_{\text{mat}}}{\delta\pi} = -\frac{\partial_\nu}{\Lambda_3^3} \left\{ -j_*^\mu \mathcal{F}_{\kappa\mu} (\eta^{\nu\kappa} - \Pi^{\nu\kappa})^{-1} + \beta j_*^\mu \mathcal{F}_{\kappa\mu} [(\alpha + \beta)\eta^{\nu\kappa} - \beta\Pi^{\nu\kappa}]^{-1} \right\}, \quad (5.25)$$

where $j_*^\mu = \sqrt{-\tilde{g}_{\text{eff}}} j_{\tilde{f}}^\mu$ is the coordinate current and represents the matter degrees of freedom independent of the metric, while $\mathcal{F}_{\kappa\mu}$ is also independent of the metric. We can now also apply the remaining derivative in front,

$$\begin{aligned} \frac{\delta\mathcal{L}_{\text{mat}}}{\delta\pi} = -\frac{1}{\Lambda_3^3} \left\{ -\partial_\nu (j_*^\mu \tilde{f}^{\rho\kappa} \mathcal{F}_{\kappa\mu}) (\delta_\rho^\nu - \Pi_\rho^\nu) + \frac{1}{\Lambda_3^3} j_*^\mu \tilde{f}^{\rho\kappa} \mathcal{F}_{\kappa\mu} \partial_\rho \square\pi \right. \\ \left. + \beta \partial_\nu (j_*^\mu \tilde{g}_{\text{eff}}^{\rho\kappa} \mathcal{F}_{\kappa\mu}) [(\alpha + \beta)\delta_\rho^\nu - \beta\Pi_\rho^\nu] - \frac{1}{\Lambda_3^3} \beta^2 j_*^\mu \tilde{g}_{\text{eff}}^{\rho\kappa} \mathcal{F}_{\kappa\mu} \partial_\rho \square\pi \right\}. \quad (5.26) \end{aligned}$$

The contribution of the matter interactions to the equation of motion for the π field contains higher derivative terms which we are not able to remove by invoking the covariant equations of motion for the vector field and the dark matter particle. Hence, this reflects the presence of a ghostly degree of freedom in the decoupling limit through the matter interaction term. The scale Λ_3 is not the cut-off scale of this theory anymore, but rather given by the scale of the mass of the introduced ghost m_{BD} . For the precise plasma-like background solution that we considered here, the contributions of the matter interactions and the internal vector field will be at least third order in perturbation at the level of the action. Hence the ghost would enter at non-linear order in perturbations, and does not show up in our linear perturbation analysis of the gravitational polarisation in Sec. 4. This indicates that the mass of the ghost m_{BD} could be large.

6 Conclusions

In this work we followed the philosophy of combining the different approaches used to tackle the cosmological constant and dark matter problems. The origin of this dark sector constitutes one of the most challenging puzzle of contemporary physics. Here we want essentially to

consider them on the same footing. The standard model of cosmology Λ -CDM, despite many observational successes, fails to explain the observed tiny value of the cosmological constant in the presence of large quantum corrections using standard quantum field theory techniques. This has initiated the development of IR modifications of GR, like massive gravity and bigravity. On the other hand, the model Λ -CDM does not account for many observations of dark matter at galactic scales, being unable to explain without fine-tuning the tight correlations between the dark and luminous matter in galaxy halos. Rather, the dark matter phenomenology at galactic scales is in good agreement with MOND [61–63].

In the present work we aimed at addressing these two motivations under the same umbrella using a common framework, namely the one of bigravity. An important clue in this respect is the fact that the MOND acceleration scale is of the order of the cosmological constant, $a_0 \sim \sqrt{\Lambda}$. An additional hope was to be able to promote the MOND formula into a decent relativistic theory in the context of bigravity. For this purpose we followed tightly the same ingredients as in the model proposed in [88], with two species of dark matter particles coupled to the two metrics of bigravity respectively, and linked *via* an internal vector field.

This paper was dedicated to explore the theoretical and phenomenological consistency of this model and verify that it is capable to recover the MOND phenomenology on galactic scales. We first worked out the covariant field equations with a special emphasis on the contributions of the vector field and the dark matter particles. Because of the interaction term between the dark matter particles and the vector field, the stress-energy tensors are not conserved separately, but rather a combination of them, giving rise to a global conservation law. The divergence of the stress-energy tensor with respect to the effective metric g_{eff} to which is coupled the vector field [15, 17], is given by the interaction of the dark matter particles with the vector field, which has important consequences for the polarisation mechanism but also for our decoupling limit analysis.

As next we computed the linear field equations around a de Sitter background where the mass term shall play the role of the cosmological constant on large scales. On small scales where the post-Newtonian limit applies, the de Sitter background can be approximated by a flat Minkowski background. Considering a small perturbation of the Minkowski metric and computing the Newtonian limit, we were able to show that the polarisation mechanism works successfully and recovers the MOND phenomenology on galactic scales. We find that this

Acknowledgments

We would like to thank Jose Beltran Jimenez, Claudia de Rham and Gilles Esposito-Farèse for very useful discussions. L.H. wishes to acknowledge the African Institute for Mathematical Sciences in Muizenberg, South Africa, for hospitality and support at the latest stage of this work.

References

- [1] J. Ostriker and P. Steinhardt, *Cosmic concordance*, *Nature* **377** (1995) 600.
- [2] T. Padmanabhan, *Cosmological constant – the weight of the vacuum*, *Phys. Rept.* **380** (2003) 235, [[arXiv:0212290 \[hep-th\]](#)].
- [3] J. Martin, *Everything you always wanted to know about the cosmological constant problem (but were afraid to ask)*, *Comptes Rendus Physique* **13** (2012) 6, [[arXiv:1205.3365 \[astro-ph\]](#)].
- [4] C. de Rham and G. Gabadadze, *Generalization of the Fierz-Pauli action*, *Phys. Rev.* **D82** (2010) 044020, [[arXiv:1007.0443](#)].
- [5] C. de Rham, G. Gabadadze, and A. J. Tolley, *Resummation of massive gravity*, *Phys. Rev. Lett.* **106** (2011) 231101, [[arXiv:1011.1232](#)].
- [6] C. de Rham, G. Gabadadze, and A. Tolley, *Ghost free Massive Gravity in the Stückelberg language*, *Phys. Lett. B* **711** (2012) 190–195, [[arXiv:1107.3820 \[hep-th\]](#)].
- [7] S. Hassan and R. A. Rosen, *Resolving the ghost problem in non-linear massive gravity*, *Phys. Rev. Lett.* **108** (2012) 041101, [[arXiv:1106.3344](#)].
- [8] S. Hassan and R. A. Rosen, *On Non-Linear Actions for Massive Gravity*, *J. High Energy Phys.* **1107** (2011) 009, [[arXiv:1103.6055](#)], [[doi:10.1007/JHEP07\(2011\)009](#)].
- [9] S. Hassan, R. A. Rosen, and A. Schmidt-May, *Ghost-free Massive Gravity with a General Reference Metric*, *JHEP* **1202** (2012) 026, [[arXiv:1109.3230](#)], [[doi:10.1007/JHEP02\(2012\)026](#)].
- [10] S. Hassan and R. A. Rosen, *Confirmation of the secondary constraint and absence of ghost in massive gravity and bimetric gravity*, *J. High Energy Phys.* **1204** (2012) 123, [[arXiv:1111.2070](#)].
- [11] S. Hassan, A. Schmidt-May, and M. von Strauss, *Proof of Consistency of Nonlinear Massive Gravity in the Stückelberg Formulation*, *Phys.Lett.* **B715** (2012) 335–339, [[arXiv:1203.5283](#)], [[doi:10.1016/j.physletb.2012.07.018](#)].
- [12] D. Boulware and S. Deser, *Can gravitation have a finite range?*, *Phys. Rev.* **D6** (1972) 3368–3382.
- [13] S. Hassan and R. A. Rosen, *Bimetric Gravity from Ghost-free Massive Gravity*, *J. High Energy Phys.* **1202** (2012) 126, [[arXiv:1109.3515](#)], [[doi:10.1007/JHEP02\(2012\)126](#)].
- [14] Y. Yamashita, A. De Felice, and T. Tanaka, *Appearance of Boulware–Deser ghost in bigravity with doubly coupled matter*, *Int. J. Mod. Phys.* **D23** (2014) 1443003, [[arXiv:1408.0487](#)], [[doi:10.1142/S0218271814430032](#)].
- [15] C. de Rham, L. Heisenberg, and R. H. Ribeiro, *On couplings to matter in massive (bi-)gravity*, [arXiv:1408.1678](#).

- [16] S. Hassan, A. Schmidt-May, and M. von Strauss, *On Consistent Theories of Massive Spin-2 Fields Coupled to Gravity*, *JHEP* **1305** (2013) 086, [[arXiv:1208.1515](#)], [[doi:10.1007/JHEP05\(2013\)086](#)].
- [17] C. de Rham, L. Heisenberg, and R. H. Ribeiro, *Ghosts and Matter Couplings in Massive (bi-and multi-)Gravity*, *Phys. Rev.* **D90** (2014) 124042, [[arXiv:1409.3834](#)], [[doi:10.1103/PhysRevD.90.124042](#)].
- [18] J. Noller and S. Melville, *The coupling to matter in Massive, Bi- and Multi-Gravity*, [arXiv:1408.5131](#).
- [19] L. Heisenberg, *Quantum corrections in massive bigravity and new effective composite metrics*, *Class. Quantum Grav.* **32** (2015) 105011, [[arXiv:1410.4239](#)].
- [20] C. de Rham and A. J. Tolley, *Vielbein to the Rescue?*, [arXiv:1505.01450](#).
- [21] Q.-G. Huang, R. H. Ribeiro, Y.-H. Xing, K.-C. Zhang, and S.-Y. Zhou, *On the uniqueness of the non-minimal matter coupling in massive gravity and bigravity*, [arXiv:1505.02616](#).
- [22] L. Heisenberg, *More on effective composite metrics*, [arXiv:1505.02966](#).
- [23] G. D'Amico, C. de Rham, S. Dubovsky, G. Gabadadze, D. Pirtskhalava, and A. J. Tolley, *Massive cosmologies*, *Phys. Rev. D* **84** (Dec, 2011) 124046, [[1108.5231](#)].
- [24] C. de Rham, G. Gabadadze, L. Heisenberg, and D. Pirtskhalava, *Non-renormalization and naturalness in a class of scalar-tensor theories*, *Phys. Rev.* **D87** (2012) [[arXiv:1212.4128](#)].
- [25] M. A. Luty, M. Porrati, and R. Rattazzi, *Strong interactions and stability in the DGP model*, *J. High Energy Phys.* **0309** (2003) 029, [[arXiv:hep-th/0303116](#)].
- [26] A. Nicolis and R. Rattazzi, *Classical and quantum consistency of the DGP model*, *J. High Energy Phys.* **0406** (2004) 059, [[arXiv:hep-th/0404159](#)].
- [27] L. Heisenberg, *Quantum corrections in Galileons from matter loops*, [arXiv:1408.0267](#).
- [28] C. de Rham, L. Heisenberg, and R. H. Ribeiro, *Quantum Corrections in Massive Gravity*, *Phys. Rev.* **D88** (2013) 084058, [[arXiv:1307.7169](#)], [[doi:10.1103/PhysRevD.88.084058](#)].
- [29] L. Heisenberg, *Quantum corrections in massive bigravity and new effective composite metrics*, [arXiv:1410.4239](#).
- [30] C. de Rham, G. Gabadadze, L. Heisenberg, and D. Pirtskhalava, *Cosmic acceleration and the helicity-0 graviton*, *Phys. Rev. D* **83** (2011) 103516, [[arXiv:1010.1780](#)].
- [31] G. Tasinato, K. Koyama, and G. Niz, *Vector instabilities and self-acceleration in the decoupling limit of massive gravity*, *Phys. Rev.* **D87** (2013) 064029, [[arXiv:1210.3627](#)].
- [32] A. E. Gumrukcuoglu, C. Lin, and S. Mukohyama, *Open FRW universes and self-acceleration from nonlinear massive gravity*, *J. Cosm. Astropart.* **1111** (2011) 030, [[arXiv:1109.3845](#)].
- [33] A. E. Gumrukcuoglu, C. Lin, and S. Mukohyama, *Cosmological perturbations of self-accelerating universe in nonlinear massive gravity*, *J. Cosm. Astropart.* **1203** (2012) 006, [[arXiv:1111.4107](#)], [[doi:10.1088/1475-7516/2012/03/006](#)].
- [34] A. De Felice, A. E. Gümrukçüoğlu, and S. Mukohyama, *Massive gravity: Nonlinear instability of a homogeneous and isotropic universe*, *Phys. Rev. Lett.* **109** (Oct, 2012) 171101, [[arXiv:1206.2080](#)], [[doi:10.1103/PhysRevLett.109.171101](#)].

- [35] M. Fasiello and A. J. Tolley, *Cosmological perturbations in Massive Gravity and the Higuchi bound*, *J. Cosm. Astropart.* **1211** (2012) 035, [[arXiv:1206.3852](#)], [[doi:10.1088/1475-7516/2012/11/035](#)].
- [36] D. Langlois and A. Naruko, *Cosmological solutions of massive gravity on de Sitter*, *Class.Quant.Grav.* **29** (2012) 202001, [[arXiv:1206.6810](#)], [[doi:10.1088/0264-9381/29/20/202001](#)].
- [37] Q.-G. Huang, Y.-S. Piao, and S.-Y. Zhou, *Mass-Varying Massive Gravity*, *Phys. Rev.* **D86** (2012) 124014, [[arXiv:1206.5678](#)], [[doi:10.1103/PhysRevD.86.124014](#)].
- [38] S. Mukohyama, *A new quasidilaton theory of massive gravity*, *J. Cosm. Astropart.* **1412** (2014), no. 12 011, [[arXiv:1410.1996](#)], [[doi:10.1088/1475-7516/2014/12/011](#)].
- [39] M. S. Volkov, *Cosmological solutions with massive gravitons in the bigravity theory*, *JHEP* **1201** (2012) 035, [[arXiv:1110.6153](#)], [[doi:10.1007/JHEP01\(2012\)035](#)].
- [40] M. von Strauss, A. Schmidt-May, J. Enander, E. Mortsell, and S. Hassan, *Cosmological Solutions in Bimetric Gravity and their Observational Tests*, *JCAP* **1203** (2012) 042, [[arXiv:1111.1655](#)], [[doi:10.1088/1475-7516/2012/03/042](#)].
- [41] D. Comelli, M. Crisostomi, and L. Pilo, *FRW Cosmological Perturbations in Massive Bigravity*, *Phys. Rev.* **D90** (2014) 084003, [[arXiv:1403.5679](#)], [[doi:10.1103/PhysRevD.90.084003](#)].
- [42] D. Comelli, M. Crisostomi, and L. Pilo, *Perturbations in Massive Gravity Cosmology*, *J. High Energy Phys.* **1206** (2012) 085, [[arXiv:1202.1986](#)], [[doi:10.1007/JHEP06\(2012\)085](#)].
- [43] A. De Felice, T. Nakamura, and T. Tanaka, *Possible existence of viable models of bi-gravity with detectable graviton oscillations by gravitational wave detectors*, *PTEP* **2014** (2014), no. 4 043E01, [[arXiv:1304.3920](#)], [[doi:10.1093/ptep/ptu024](#)].
- [44] A. De Felice, A. E. Gümrükçüoğlu, S. Mukohyama, N. Tanahashi, and T. Tanaka, *Viable cosmology in bimetric theory*, *J. Cosm. Astropart.* **1406** (2014) 037, [[arXiv:1404.0008](#)], [[doi:10.1088/1475-7516/2014/06/037](#)].
- [45] Y. Akrami, S. Hassan, F. Könnig, A. Schmidt-May, and A. R. Solomon, *Bimetric gravity is cosmologically viable*, [arXiv:1503.07521](#), [doi:10.1016/j.physletb.2015.06.062](#).
- [46] J. Enander, Y. Akrami, E. Mortsell, M. Renneby, and A. R. Solomon, *Integrated Sachs-Wolfe effect in massive bigravity*, *Phys.Rev.* **D91** (2015) 084046, [[arXiv:1501.02140](#)], [[doi:10.1103/PhysRevD.91.084046](#)].
- [47] J. Enander and E. Mortsell, *On stars, galaxies and black holes in massive bigravity*, [arXiv:1507.00912](#).
- [48] G. Cusin, R. Durrer, P. Guarato, and M. Motta, *Inflationary perturbations in bimetric gravity*, [arXiv:1505.01091](#).
- [49] F. Könnig, *Higuchi Ghosts and Gradient Instabilities in Bimetric Gravity*, *Phys.Rev.* **D91** (2015) 104019, [[arXiv:1503.07436](#)], [[doi:10.1103/PhysRevD.91.104019](#)].
- [50] M. Fasiello and R. H. Ribeiro, *Mild bounds on bigravity from primordial gravitational waves*, [arXiv:1505.00404](#).
- [51] A. Bosma, *21-cm line studies of spiral galaxies. ii. the distribution and kinematics of neutral hydrogen in spiral galaxies of various morphological types*, *Astron. J.* **86** (1981) 1825.

- [52] V. Rubin, J. Ford, N. Thonnard, and D. Burstein, *Rotational properties of 23 sb galaxies*, *Astrophys. J.* **261** (1982) 439.
- [53] W. Hu and S. Dodelson, *Cosmic microwave background anisotropies*, *Annual Rev. Astron. Astrophys.* **40** (2002) 171.
- [54] G. Bertone, D. Hooper, and J. Silk, *Particle dark matter: Evidence, candidates and constraints*, *Phys. Rept.* **405** (2005) 279, [[hep-ph/0404175](#)].
- [55] R. Sanders and S. McGaugh, *Modified newtonian dynamics as an alternative to dark matter*, *Ann. Rev. Astron. Astrophys.* **40** (2002) 263, [[astro-ph/0204521](#)].
- [56] B. Famaey and S. McGaugh, *Modified Newtonian dynamics (MOND): Observational phenomenology and relativistic extensions*, *Living Rev. Rel.* **15** (2012) 10, [[arXiv:1112.3960](#) [[astro-ph.CO](#)]].
- [57] R. Tully and J. Fisher, *A new method of determining distances to galaxies*, *Astron. Astrophys.* **54** (1977) 661.
- [58] S. McGaugh, J. Schombert, G. Bothun, and W. de Blok, *The baryonic tully-fisher relation*, *Astrophys. J.* **533** (2000), no. 2 L99.
- [59] S. S. McGaugh, *Novel test of modified newtonian dynamics with gas rich galaxies*, *Phys. Rev. Lett.* **106** (2011) 121303.
- [60] R. Sanders, *The universal faber-jackson relation*, *Mon. Not. Roy. Astron. Soc.* **407** (2010) 1128–1134, [[arXiv:1002.2765](#) [[astro-ph.CO](#)]].
- [61] M. Milgrom, *A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis*, *Astrophys. J.* **270** (1983) 365.
- [62] M. Milgrom, *A modification of the Newtonian dynamics: Implications for galaxies*, *Astrophys. J.* **270** (1983) 371.
- [63] M. Milgrom, *A modification of the Newtonian dynamics: Implications for galaxy systems*, *Astrophys. J.* **270** (1983) 384.
- [64] J. Bekenstein and M. Milgrom, *Does the missing mass problem signal the breakdown of newtonian gravity?*, *Astrophys. J.* **286** (1984) 7.
- [65] D. Gerbal, F. Durret, M. Lachièze-Rey, and G. Lima-Neto, *Analysis of x-ray galaxy clusters in the framework of modified newtonian dynamics*, *Astron. Astrophys.* **262** (1992) 395.
- [66] E. Pointecouteau and J. Silk, *New constraints on modified newtonian dynamics from galaxy clusters*, *Mon. Not. R. Astron. Soc.* **364** (2005) 654–658, [[astro-ph/0505017](#)].
- [67] D. Clowe, M. Bradac, A. H. Gonzalez, M. Markevitch, S. W. Randall, C. Jones, and D. Zaritsky, *A direct empirical proof of the existence of dark matter*, *Astrophys. J.* **648** (2006) L109, [[astro-ph/0608407](#)].
- [68] G. W. Angus, B. Famaey, and D. A. Buote, *X-ray group and cluster mass profiles in MOND: Unexplained mass on the group scale*, *Mon. Not. Roy. Astron. Soc.* **387** (2008) 1470, [[arXiv:0709.0108](#) [[astro-ph](#)]].
- [69] G. W. Angus, *Is an 11 ev sterile neutrino consistent with clusters, the cosmic microwave background and modified Newtonian dynamics?*, *Mon. Not. Roy. Astron. Soc.* **394** (2009) 527,

- [arXiv:0805.4014 [astro-ph]].
- [70] M. Milgrom, *Mond effects in the inner solar system*, *Mon. Not. Roy. Astron. Soc.* **399** (2009) 474.
- [71] L. Blanchet and J. Novak, *External field effect of modified newtonian dynamics in the solar system*, *Mon. Not. Roy. Astron. Soc.* **412** (2011) 2530.
- [72] R. Sanders, *A stratified framework for scalar-tensor theories of modified dynamics*, *Astrophys. J.* **480** (1997) 492, [astro-ph/9612099].
- [73] J. Bekenstein, *Relativistic gravitation theory for the modified Newtonian dynamics paradigm*, *Phys. Rev. D* **70** (2004) 083509, [astro-ph/0403694].
- [74] R. Sanders, *A tensor-vector-scalar framework for modified dynamics and cosmic dark matter*, *Mon. Not. Roy. Astron. Soc.* **363** (2005) 459, [astro-ph/0502222].
- [75] T. G. Zlosnik, P. G. Ferreira, and G. D. Starkman, *Modifying gravity with the æther: An alternative to dark matter*, *Phys. Rev. D* **75** (2007) 044017, [arXiv:astro-ph/0607411].
- [76] A. Halle, H. S. Zhao, and B. Li, *A nonuniform dark energy fluid: Perturbation equations*, *Astrophys. J. Suppl.* **177** (2008) 1, [arXiv:0711.0958 [astro-ph]].
- [77] M. Milgrom, *Bimetric mond gravity*, *Phys. Rev. D* **80** (2009) 123536.
- [78] E. Babichev, C. Deffayet, and G. Esposito-Farèse, *Improving relativistic mond with galileon k-mouflage*, *Phys. Rev. D* **84** (2011) 061502(R), [arXiv:1106.2538 [gr-qc]].
- [79] C. Deffayet, G. Esposito-Farèse, and R. Woodard, *Nonlocal metric formulations of mond with sufficient lensing*, *Phys. Rev. D* **84** (2011) 124054, [arXiv:1106.4984 [gr-qc]].
- [80] L. Blanchet and S. Marsat, *Modified gravity approach based on a preferred time foliation*, *Phys. Rev. D* **84** (2011) 044056, [arXiv:1107.5264 [gr-qc]].
- [81] Y. Arraut, *Can a nonlocal model of gravity reproduce dark matter effects in agreement with mond?*, *Int. J. of Modern Phys. D* **23** (2014) 1450008, [arXiv:1310.0675].
- [82] L. Blanchet, *Gravitational polarization and the phenomenology of MOND*, *Class. Quant. Grav.* **24** (2007) 3529, [astro-ph/0605637].
- [83] L. Blanchet and A. Le Tiec, *Model of dark matter and dark energy based on gravitational polarization*, *Phys. Rev. D* **78** (2008) 024031, [astro-ph/0804.3518].
- [84] L. Blanchet and A. Le Tiec, *Dipolar dark matter and dark energy*, *Phys. Rev. D* **80** (2009) 023524, [arXiv:0901.3114 [astro-ph]].
- [85] L. Blanchet, D. Langlois, A. Le Tiec, and S. Marsat, *Non-gaussianity in the cosmic microwave background induced by dipolar dark matter*, *J. Cosm. Astropart.* **22** (2013) 1302, [arXiv:1210.4106 [astro-ph]].
- [86] L. Blanchet and L. Bernard, *Phenomenology of mond and gravitational polarization*, arXiv:1403.5963 [gr-qc]. In the proceedings of the second Workshop on Antimatter and Gravity (WAG 2013).
- [87] L. Bernard and L. Blanchet, *Phenomenology of dark matter via a bimetric extension of general relativity*, arXiv:1410.7708 [astro-ph]. To appear in *Phys. Rev. D*.

- [88] L. Blanchet and L. Heisenberg, *Dark matter via massive (bi-)gravity*, [arXiv:1504.00870](#). To appear in Phys. Rev. D.
- [89] V. Baccetti, P. Martin-Moruno, and M. Visser, *Null Energy Condition violations in bimetric gravity*, *JHEP* **1208** (2012) 148, [[arXiv:1206.3814](#)], [[doi:10.1007/JHEP08\(2012\)148](#)].
- [90] A. Schmidt-May, *Mass eigenstates in bimetric theory with ghost-free matter coupling*, [arXiv:1409.3146](#).