

Evolution of fusion hindrance for asymmetric systems at deep sub barrier energies

A. Shrivastava^{a,b,*}, K. Mahata^{a,b}, S.K. Pandit^{a,b}, V. Nanal^c, T. Ichikawa^d, K. Hagino^e, A. Navin^f, C.S. Palshetkar^a, V.V. Parkar^a, K. Ramachandran^{a,b}, P.C. Rout^{a,b}, Abhinav Kumar^a, A. Chatterjee^a, S. Kailas^a

^aNuclear Physics Division, Bhabha Atomic Research Centre, Mumbai 400085, India

^bHomi Bhabha National Institute, Anushaktinagar, Mumbai 400094, INDIA

^cDNAP, Tata Institute of Fundamental Research, Mumbai 400005, India

^dYukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

^eDepartment of Physics, Tohoku University, Sendai 980-8578, Japan

^fGANIL, CEA/DRF - CNRS/IN2P3, Bd Henri Becquerel, BP 55027, F-14076 Caen Cedex 5, France

Abstract

Measurements of fusion cross-sections of ${}^7\text{Li}$ and ${}^{12}\text{C}$ with ${}^{198}\text{Pt}$ at deep sub-barrier energies are reported to unravel the role of the entrance channel in the occurrence of fusion hindrance. The onset of fusion hindrance has been clearly observed in ${}^{12}\text{C} + {}^{198}\text{Pt}$ system but not in ${}^7\text{Li} + {}^{198}\text{Pt}$ system, within the measured energy range. Emergence of the hindrance, moving from lighter (${}^6,{}^7\text{Li}$) to heavier (${}^{12}\text{C}, {}^{16}\text{O}$) projectiles is explained employing a model that considers a gradual transition from a sudden to adiabatic regime at low energies. The model calculation reveals a weak effect of the damping of coupling to collective motion for the present systems as compared to that obtained for systems with heavier projectiles.

Keywords: fusion cross sections, deep sub barrier energies, coupled channels calculations, adiabatic model

1. Introduction

Fusion reactions in the vicinity of the Coulomb barrier have been investigated in the past to explore the mechanism of tunneling through multidimensional barriers, thereby giving an insight into the role of different intrinsic properties of the entrance channel. Recent efforts towards developing new methods to precisely measure very low fusion cross-sections have stimulated new activities, distinct to energies deep below the barrier. Fusion data at these low energies can be uniquely used to interpret the reaction dynamics from the touching point to the region of complete overlap of the density distribution of the colliding nuclei, not accessible through any other reaction [1, 2]. This opens up the possibility to study effects of dissipative quantum tunneling, which has relevance in many fields of physics and chemistry [3]. The data in this energy range was shown to have strong implications on the fusion with light nuclei of astrophysical interest [2].


At deep sub-barrier energies, a change of slope of the fusion excitation function compared to coupled-channels (CC) calculations was observed initially in symmetric systems involving medium-heavy nuclei and was referred to as the phenomenon of fusion hindrance [4, 5]. The models suggested to explain this behavior have different physical basis. The model proposed by Mişicu and Esbensen is based on a sudden approximation [6], where a repulsive core is included to take into account the nuclear compressibility arising due to Pauli exclusion principle when the two nuclei overlap. On the other hand at low energies, the nucleus-nucleus interaction potentials extracted from

the microscopic time-dependent Hartree-Fock theory indicate that after overlap of two nuclei, internal degrees of freedom reorganise adiabatically [7]. The model proposed by Ichikawa *et al.* [8] to explain the deep sub-barrier fusion data is based on such an adiabatic picture. Here a damping factor imposed on the coupling strength as a function of the inter-nuclear distance, takes into account a gradual change from the sudden to the adiabatic formalism [9, 10]. A recent work, applying the random-phase-approximation (RPA) demonstrates that the fusion hindrance originates from damping of quantum vibrations when the two nuclei adiabatically approach each other [11, 12]. The role of quantum de-coherence that effectively cause a reduction in coupling effects has also been investigated [13, 14].

In all the above models, fusion hindrance is a generic property of heavy-ion collision below certain threshold energy. Due to challenges involved with measurement of low cross-section ($\sim\text{nb}$), there are only a limited number of studies involving fusion hindrance. As discussed in a recent review article [2], these studies have mainly concentrated around medium-heavy ($A\sim 100$), medium ($A\sim 50$) and light ($A\sim 10$) symmetric systems [4, 5, 15, 16, 17, 18, 19, 20, 21], covering a wide range of reduced masses, Q -values and nuclear structure properties. Most of the measurements employed recoil mass analyzers and hence are restricted to symmetric or nearly symmetric systems. In such cases the evaporation residues have sufficient recoil velocities for being detected at the focal plane of the spectrometer. The data corresponding to asymmetric systems, presently scarce, are vital to establish the generic nature of the fusion hindrance and for the improvement of current theoretical models. The only exception being the two systems ${}^{16}\text{O} + {}^{208}\text{Pb}$ [14] and ${}^6\text{Li} + {}^{198}\text{Pt}$ [22] that used different methods for fusion cross-

*Corresponding author

Email address: aradhana@barc.gov.in (A. Shrivastava)






Figure 2: (Color online) Results from the adiabatic model calculation for $^{12}\text{C} + ^{198}\text{Pt}$ system compared with the experimental (a) fusion cross-sections, (b) logarithmic derivative along with S-factor (inset) (c) average angular momentum and (d) fusion barrier distribution. Calculations using with and without a damping factor for the coupling strength are shown as solid and dashed-dot curves respectively.

3. Calculations

Coupled-channels calculations using the code CCFULL [29] were performed for both the systems. In the case of $^7\text{Li} + ^{198}\text{Pt}$ system, a standard Woods-Saxon potential (WS) was used with $V_0=110$ MeV, $r_0=1.1$ fm and $a=0.63$ fm. These calculations included two phonon quadrupole excitation of ^{198}Pt in the vibrational and the first excited state of ^7Li in the rotational mode. The CC calculations reproduce the data well for energies around and well below the barrier as seen in Fig. 1(a). Fusion hindrance has not been observed in this system in the measured energy range with the cross-section as low as ≈ 180 nb. The threshold energy for observing fusion hindrance obtained from the systematics of Ref. [30] and the adiabatic model [8] is 20.4 MeV and 21.1 MeV, respectively. However, from an extrapolation of the experimental data, this energy is found to be ≈ 19 MeV (Fig. 1(a) inset). Hence it will be interesting to extend the measurement of fusion cross-sections below the lowest energy of the present measurement (20 MeV).

The corresponding calculations for $^{12}\text{C} + ^{198}\text{Pt}$ were performed using a WS potential with $V_0=95$ MeV, $r_0=1.13$ fm and $a=0.66$ fm. The coupling to the quadrupole phonon excitation for ^{198}Pt and the first two excited states of ^{12}C belonging to the ground state rotational band were included. The quadrupole and hexadecapole deformation parameters used were taken from Ref. [31]. The effect of coupling to the ^{12}C rotational states is not as strong as in the well deformed heavy nuclei. Coupling to one neutron, two neutron and one proton transfer reaction were not included in the present scheme as their effect was found to be negligible for this system [25]. The result from the CC calculations are compared with the experimental data in Fig. 1(b). A change of slope as compared to CC calculations is clearly observed, both in the measured fusion excitation function as well as in the $L(E)$ plot, confirming the onset of fusion hindrance. The energy at which the deviation in the slope occurs

was estimated to be 50 ± 1 MeV using the method described in Ref. [32]. The calculated threshold energy according to the adiabatic model [8] is 49 MeV while that from the systematics (43.7 MeV) [30] is much lower than the observed value.

In order to explain the fusion data at energies deep below the barrier in case of $^{12}\text{C} + ^{198}\text{Pt}$ system, calculations were performed using the adiabatic model of Ref. [9, 10]. This model employs a gradual diminishing of the coupling strength while going from the two body sudden to one body adiabatic potential as the two nuclei begin to overlap. The calculations adopted a Yukawa-plus-exponential (YPE) potential as a basic ion-ion potential with radius, $r_0=1.20$ fm and diffuseness, $a=0.68$ fm. The coupling scheme was the same as that described earlier for this system. The calculated fusion cross-sections without damping, shown as the dot-dashed curves in Fig. 2(a), already provide a good fit, although the calculation underestimates the data for $L(E)$ at the lowest energies (see Fig. 2(b)). The calculations shown here differ slightly from those in Fig 1(b). This is due to the use of different potentials (YPE and Woods-Saxon) in these two calculations, and the fact that the YPE potential is thicker than the Woods-Saxon potential (due to the saturation condition at the touching point in the YPE potential). Further discussion about the choice of potentials used in the present work can be found in section IIC of Ref. [10]. Fig. 2 also shows the results of the calculation with the inclusion of a damping factor ($r_{damp}=1.18$ fm and $a_{damp}=0.5$ fm), which are in excellent agreement with both the fusion and the $L(E)$ data. As can be seen from the figure, the effect of the damping is observed to be small in the present case when compared to that observed in studies involving heavier projectiles [10]. A systematic investigation of various systems showed that the radius parameter, related to the density distribution of the colliding nuclei, is almost constant [10]. On the other hand a_{damp} , associated with the damping strength of quantum vibrations that depends on the structure of interacting nuclei, was found to vary between 0.5 and 1.2 fm. The values of r_{damp} and a_{damp} obtained in the present work are within the range of the values reported in Ref. [10].

The adiabatic calculations were compared with other observables derived from the fusion data. The astrophysical S-factor representation ($S(E)$) of the experimental data is shown in the inset of Fig. 2(b). The observed S-factor maximum is not as pronounced as found for the case of the symmetric systems involving medium mass nuclei, but similar to that for $^{16}\text{O} + ^{208}\text{Pb}$ system [1, 2, 33]. The calculated $S(E)$ match well with the data over the entire energy range. The average angular momenta ($\langle l \rangle$) computed from the fusion excitation function as suggested in Ref. [34] and the fusion barrier distribution (DB) are also well described by the adiabatic calculation (Fig. 2(c) and (d)).

Similar calculations were performed for $^7\text{Li} + ^{198}\text{Pt}$ using the YPE potential ($r_0=1.195$ fm and $a=0.68$ fm) with the coupling scheme being the same as that described above for this system. The calculations explain the data well up to the measured energy. The values of r_{damp} and a_{damp} can not be determined uniquely with the present data as no change of slope of the fusion excitation function was observed. For example, values of the damping factor parameters $r_{damp}=1.16$ fm and $a_{damp}=0.5$

able.

5. Conclusion

In summary, the occurrence of fusion hindrance is clearly observed in case of $^{12}\text{C} + ^{198}\text{Pt}$. The adiabatic model calculation indicates a weak effect of the damping for the present system as compared to that obtained for systems with heavier projectiles. On the other hand fusion hindrance has not been observed in case of $^7\text{Li} + ^{198}\text{Pt}$, within the measured energy range. The corresponding threshold energy estimated from the present measurement is found to be lower than the predicted values [8, 30]. The fusion hindrance at energies deep below the barrier becomes progressively significant in going from the light ($^{6,7}\text{Li}$) to heavier (^{12}C , ^{16}O) projectiles. A strong correlation has been obtained between the degree of hindrance and the charge product over a wide range of target-projectile combinations. The observed trend reveals a weaker influence of hindrance on fusion involving lighter nuclei. This result together with a nearly flat energy dependence of $S(E)$ in the adiabatic model at very low energies, implies that the effect of fusion hindrance will be less substantial on astrophysical reaction rates for the production of light elements in stellar environments. New measurements of fusion cross-sections involving low Z elements including those of astrophysical relevance, and extension of existing theoretical models that explain fusion hindrance to the energies close to the Gamow peak would be of interest.

6. Acknowledgments

We are thankful to the accelerator staff for a smooth operation. AN acknowledges the support through the LIA France-India agreement.

References

- [1] K. Hagino and N. Takigawa, *Prog. Theor. Phys.* 128 (2012) 1001.
- [2] B.B. Back, H. Esbensen, C.L. Jiang and K.E. Rehm, *Rev. Mod. Phys.* 86 (2014) 317, and references therein.
- [3] A.O. Caldeira and A.J. Leggett, *Phys. Rev. Lett.* 46 (1981) 211; *Ann. of Phys.* 149 (1983) 374.
- [4] C.L. Jiang *et al.*, *Phys. Rev. Lett.* 89 (2002) 052701.
- [5] C.L. Jiang *et al.*, *Phys. Rev. Lett.* 93 (2004) 012701.
- [6] S. Mišicu and H. Esbensen, *Phys. Rev. Lett.* 96 (2006) 112701; *ibid.* *Phys. Rev. C* 75 (2007) 034606.
- [7] K. Washiyama and D. Lacroix, *Phys. Rev. C* 78 (2008) 024610; A.S. Umar and V.E. Oberacker, *Phys. Rev. C* 74 (2006) 021601(R).
- [8] T. Ichikawa, K. Hagino and A. Iwamoto, *Phys. Rev. C* 75 (2007) 057603; *ibid.* *Phys. Rev. C* 75 (2007) 064612.
- [9] T. Ichikawa, K. Hagino and A. Iwamoto, *Phys. Rev. Lett.* 103 (2009) 202701; *ibid.* *Prog. theor. Phys. suppl.* 196 (2012) 269.
- [10] Takatoshi Ichikawa, *Phys. Rev. C* 92 (2015) 064604.
- [11] Takatoshi Ichikawa and Kenichi Matsuyanagi, *Phys. Rev. C* 88 (2013) 011602(R).
- [12] Takatoshi Ichikawa and Kenichi Matsuyanagi, *Phys. Rev. C* 92 (2015) 021602(R).
- [13] A. Diaz-Torres *et al.*, *Phys. Rev. C* 78 (2008) 064604; *Phys. Rev. C* 81 (2010) 041603(R).
- [14] M. Dasgupta *et al.*, *Phys. Rev. Lett.* 99 (2007) 192701.
- [15] C.L. Jiang *et al.*, *Phys. Rev. C* 71 (2005) 044613.
- [16] A. M. Stefanini *et al.* *Phys. Rev. C* 82 (2010) 014614.
- [17] C.L. Jiang *et al.*, *Phys. Rev. C* 82 (2010) 041601.
- [18] G. Montagnoli *et al.*, *Phys. Rev. C* 85 (2012) 024607; *ibid.* 82 (2010) 064609.
- [19] G. Montagnoli *et al.*, *Phys. Rev. C* 87 (2013) 014611.
- [20] A. M. Stefanini *et al.*, *Phys. Lett. B* 728 (2014) 639.
- [21] C.L. Jiang *et al.*, *Phys. Rev. Lett.* 113 (2014) 022701.
- [22] A. Shrivastava *et al.*, *Phys. Rev. Lett.* 103 (2009) 232702.
- [23] A. Lemasson *et al.*, *Nucl. Instr. Meth. A* 598 (2009) 445.
- [24] A. Gavron, *Phys. Rev. C* 21 (1980) 230.
- [25] A. Shrivastava *et al.* *Phys. Rev. Lett.* 82 (1999) 699, *ibid.* *Phys. Rev. C* 63 (2001) 054602.
- [26] W.-M. Yao *et al.*, *J. Phys. G* 33 (2006) 1.
- [27] A. Shrivastava *et al.*, *Phys. Lett. B* 718 (2013) 931.
- [28] C.L. Jiang *et al.*, *Phys. Rev. C* 73 (2006) 014613.
- [29] K. Hagino *et al.*, *Comp. Phys. Comm.* 123 (1999) 143.
- [30] C.L. Jiang *et al.*, *Phys. Rev. C* 79 (2009) 044601.
- [31] M. Yasue *et al.*, *Nucl. Phys. A* 394 (1983) 29.
- [32] Ei Shwe Zin Thein, N. W. Lwin, and K. Hagino, *Phys. Rev. C* 85 (2012) 057602.
- [33] C.L. Jiang *et al.*, *Phys. Rev. C* 75 (2007) 057604.
- [34] A.B. Balantekin and P.E. Reimer, *Phys. Rev. C* 33 (1986) 379; C.V.K. Baba, *Nucl. Phys. A* 553 (1993) 719c.
- [35] C.R. Morton *et al.*, *Phys. Rev. C* 60 (1999) 044608.
- [36] L.F. Canto, P.R.S. Gomes, R. Donangelo, J. Lubian, M.S. Hussein, *Phys. Rep.* 596 (2015) 1.
- [37] X.P. Yang, G. L. Zhang and H. Q. Zhang *et al.*, *Phys. Rev. C* 87 (2013) 014603.
- [38] D. H. Luong *et al.*, *Phys. Lett. B* 695 (2011) 105.
- [39] A. Shrivastava *et al.*, *Phys. Lett. B* 633 (2006) 433.
- [40] C.L. Jiang *et al.*, *Phys. Rev. C* 75 (2007) 015803.
- [41] L.R. Gasques *et al.*, *Phys. Rev. C* 76 (2007) 035802.