



Studying the molecular gas towards the R Coronae Australis dark cloud

S. Paron^{1,2}, M. Celis Peña¹, M. E. Ortega¹, M. Cunningham³, P. A. Jones³ & M. Rubio⁴

¹ *Instituto de Astronomía y Física del Espacio (CONICET-UBA)*

² *FADU y CBC, Universidad de Buenos Aires*

³ *School of Physics, University of New South Wales, Sydney, NSW 2052, Australia*

⁴ *Departamento de Astronomía, Universidad de Chile, Casilla 36-D, Santiago, Chile*

Contact / sparon@iafe.uba.ar

Resumen / La nube oscura R Coronae Australis es una de las regiones de formación estelar más cercanas al Sol. Ciertamente se sabe que dicha nube es muy activa en formación estelar conteniendo una gran cantidad de objetos Herbig-Haro (HHs) y objetos de emisión de línea del hidrógeno molecular (MHOs). En este trabajo presentamos los resultados del análisis de observaciones moleculares (un mapa de $5.5' \times 5.5'$ de ^{12}CO J=3-2 y HCO^+ J=4-3, y un espectro único de N_2H^+ J=4-3) obtenidas con el telescopio Atacama Submillimeter Telescope Experiment (ASTE) hacia la nube oscura R CrA con una resolución angular y espectral de $22''$ y 0.11 km s^{-1} , respectivamente. Analizando los espectros de la emisión del ^{12}CO J=3-2 encontramos evidencia cinemática que sugiere la presencia de chorros moleculares hacia una región poblada por varios objetos HHs y MHOs. La mayoría de dichos objetos se encuentran localizados sobre el máximo del HCO^+ , sugiriendo que su emisión proviene del incremento de su abundancia debido a la química generada por los chorros moleculares. De manera adicional, presentamos un espectro de N_2H^+ , siendo la primer detección reportada de esta molécula en la línea J=4-3 hacia la nube oscura R CrA.

Abstract / The R Coronae Australis dark cloud is one of the closest star-forming regions to the Sun. The cloud is known to be very active in star formation, harboring many Herbig-Haro objects (HHs) and Molecular Hydrogen emission-line Objects (MHOs). In this work we present results from molecular observations (a $5.5' \times 5.5'$ map of ^{12}CO J=3-2 and HCO^+ J=4-3, and a single spectrum of N_2H^+ J=4-3) obtained with the Atacama Submillimeter Telescope Experiment (ASTE) towards the R CrA dark cloud with an angular and spectral resolution of $22''$ and 0.11 km s^{-1} , respectively. From the ^{12}CO J=3-2 line we found kinematical spectral features strongly suggesting the presence of outflows towards a region populated by several HHs and MHOs. Moreover, most of these objects lie within an HCO^+ maximum, suggesting that its emission arises from an increase of its abundance due to the chemistry triggered by the outflow activity. Additionally, we are presenting the first reported detection of N_2H^+ in the J=4-3 line towards the R CrA dark cloud.

Keywords / ISM: clouds – ISM: jets and outflows – ISM: molecules – stars: formation

1. Introduction

The R Coronae Australis dark cloud (R CrA dark cloud), located at about 130 pc (Neuhäuser & Forbrich, 2008), is one of the closest star-forming regions to the Sun. The cloud is centered on the Herbig Ae/Be stars R CrA and T CrA, which are the exciting sources of the reflection nebula NGC 6729. It is known that this dark cloud is very active in star formation (e.g. Lindberg & Jørgensen 2012), harboring many Herbig-Haro (HHs) objects and Molecular Hydrogen emission-line Objects (MHOs). Due to its proximity, this region is a very interesting target to study the molecular gas related to the star-forming processes.

2. Observations

A region of $5.5' \times 5.5'$ in size centered at 19:01:51, $-36:58:20$ (J2000) in the R CrA dark cloud was mapped in the ^{12}CO J=3-2 and HCO^+ J=4-3 lines using the

Atacama Submillimeter Telescope Experiment (ASTE). The observations were carried out in August 2010 and were performed in on-the-fly mapping mode achieving an angular sampling of $6''$. In addition, the N_2H^+ J=4-3 line was observed as a single point at 19:01:52.7, $-36:57:49$ (J2000). The angular and spectral resolution are $22''$ and 0.11 km s^{-1} , respectively. The data were reduced with NEWSTAR* and the spectra processed using the XSpec software package**. Polynomials between first and third order were used for baseline fitting.

* Reduction software based on AIPS developed at NRAO, extended to treat single dish data with a graphical user interface (GUI).

** XSpec is a spectral line reduction package for astronomy which has been developed by Per Bergman at Onsala Space Observatory.

3. Results

Figure 1 shows the ^{12}CO J=3-2 emission integrated between -15 and $+25$ km s^{-1} towards the R CrA dark cloud. In addition to the molecular study, we search for HHs objects and MHOs lying in the region from the catalogues of Reipurth (2000) and Davis et al. (2010). This kind of objects are strong evidences of active star formation. The crosses in Fig. 1 show the position of HHs and MHOs, which lie mainly within the molecular gas emission, strongly suggesting that they are embedded in the molecular cloud.




Fig. 1: ^{12}CO J=3-2 emission integrated between -15 and $+25$ km s^{-1} . The contour levels are 30, 50, 70, and 130 K km s^{-1} . The green and cyan crosses represent the positions of MHO and HH objects, respectively. The red and blue squares are the positions from which we extracted the spectra presented in Fig. 2. The beam is presented in the bottom-left corner.

From the ^{12}CO data we look for kinematical spectral evidences of perturbed gas due to the star-forming processes. Analyzing the whole data cube we found that in a region of about $1'$ centered at the position indicated with a blue square in Fig. 1 the spectra present a blue-shifted wing, while in a region with the same size centered at the red square (Fig. 1) the spectra show a red-shifted wing. Two representative spectra of both regions are presented in Fig. 2. These spectral features are strong evidences of molecular outflows driven by the young stellar objects (YSOs) embedded in the cloud. A dynamical study of the outflows will be presented in a further work. Besides, the spectra present a dip at about 5 km s^{-1} , which shows that the emission is self-absorbed, suggesting the presence of high-density gas.

Figure 3 (up) displays the HCO^+ J=4-3 emission integrated between $+2$ and $+10$ km s^{-1} , the range in which this line presents emission, towards the R CrA




Fig. 2: Up: ^{12}CO J=3-2 spectrum with a blue-shifted wing obtained towards the position represented by a blue square in Fig. 1. Bottom: ^{12}CO J=3-2 spectrum with a red-shifted wing obtained towards the position indicated with a red square in Fig. 1. The dashed box in both spectra remarks the shifted emission components.

dark cloud. For comparison, Fig. 3 (bottom) presents the integrated ^{12}CO emission and the stellar sources (as displayed in Fig. 1) with the HCO^+ contours superimposed. It can be appreciated that the HCO^+ peak coincides with the ^{12}CO maximum. Most of the HHs and MHOs lie in the region delimited by the HCO^+ emission, suggesting that it arises due to the outflowing phenomenon. It is known that such molecular species enhances in molecular outflows (Rawlings et al., 2004). Indeed an enhancement of the HCO^+ abundance is expected to occur due to the liberation and photoprocessing by shocks of the molecular material stored in the icy mantles of the dust.

Figure 4 shows the spectrum of the N_2H^+ J=4-3 line obtained towards the position indicated with a yellow cross in Fig. 3. It is important to note that this is the first reported detection of N_2H^+ in the J=4-3 line towards CrA dark cloud. Its detection, centered at $v \sim 6.12$ km s^{-1} , indicates the presence of high-




Fig. 3: Up: HCO^+ $J=4-3$ emission integrated between $+2$ and $+10 \text{ km s}^{-1}$. The contour levels are 5, 10, and 15 K km s^{-1} . The beam is presented in the bottom-left corner. The yellow cross is the position at which was observed the N_2H^+ $J=4-3$ line. Bottom: same as Fig.1 with the HCO^+ contours (in red) of the upper panel.

density gas and it is an interesting line to study the ionization and chemistry related to the star-forming processes. For instance, it is believed that as star formation progresses, the temperature rises and thus the CO is released from the dust, which then interacts with N_2H^+ through $\text{N}_2\text{H}^+ + \text{CO} \rightarrow \text{HCO}^+ + \text{N}_2$ (Busquet et al., 2011).

Finally, we obtained a line integrated intensity ratio of $I(\text{N}_2\text{H}^+)/I(\text{HCO}^+) \sim 0.33$ at the position in which the N_2H^+ was observed. This value is close to the lower cutoff value assigned to the sources named as “ N_2H^+




Fig. 4: N_2H^+ $J=4-3$ emission obtained at 19:01:52.7, $-36:57:49$ (J2000) in the R CrA dark cloud (yellow cross in Fig. 3-up).

poor” in Hoq et al. (2013). Even though this study was performed for high-mass star forming regions and R CrA dark cloud hosts low and intermediate mass YSOs, we can suggest a similar chemistry for both types of YSOs. In this way, the YSOs related to the molecular gas analyzed here may be in the later stages of evolution (Hoq et al., 2013).

Acknowledgements: The ASTE project is driven by Nobeyama Radio Observatory (NRO), a branch of National Astronomical Observatory of Japan (NAOJ), in collaboration with University of Chile, and Japanese institutes including University of Tokyo, Nagoya University, Osaka Prefecture University, Ibaraki University, Hokkaido University and Joetsu University of Education. S.P. and M.O. are members of the *Carrera del investigador científico* of CONICET, Argentina. M.C.P. is a doctoral fellow of CONICET, Argentina. This work was partially supported by Argentina grants awarded by UBA (UBACyT), CONICET and ANPCYT. M.R. wishes to acknowledge support from CONICYT through FONDECYT grant N° 1140839.

References

- Busquet G., et al., 2011, *A&A*, 525, A141
- Davis C. J., et al., 2010, *A&A*, 511, A24
- Hoq S., et al., 2013, *ApJ*, 777, 157
- Lindberg J. E., Jørgensen J. K., 2012, *A&A*, 548, A24
- Neuhäuser R., Forbrich J., 2008, *The Corona Australis Star Forming Region in Handbook of Star Forming Regions, Volume II*, Ed. B. Reipurth. p. 735
- Rawlings J. M. C., et al., 2004, *MNRAS*, 351, 1054
- Reipurth B., 2000, *VizieR Online Data Catalog*, 5104, 0