

# SYMMETRIC GROUP REPRESENTATIONS AND $\mathbb{Z}$

ANSHUL ADVE AND ALEXANDER YONG

Let  $\mathfrak{S}_n$  be the symmetric group of permutations of  $\{1, 2, \dots, n\}$ . A *representation* is a homomorphism  $\rho : \mathfrak{S}_n \rightarrow \mathrm{GL}(V)$  where  $V$  is a vector space over  $\mathbb{C}$ . Equivalently,  $V$  is an  $\mathfrak{S}_n$ -module under the action defined by  $\sigma \cdot v = \rho(\sigma)v$ , for  $\sigma \in \mathfrak{S}_n$  and  $v \in V$ . Then  $\rho$  is *irreducible* if there is no proper  $\mathfrak{S}_n$ -submodule of  $V$ . Conjugacy classes and hence irreducible representations of  $\mathfrak{S}_n$  biject with  $\mathrm{Par}(n)$ , the partitions of size  $n$ .

Consider three families of numbers from the theory:

(I) The *character* of  $\rho$  is

$$\chi^\rho : \mathfrak{S}_n \rightarrow \mathbb{C}; \quad \sigma \mapsto \mathrm{tr}(\rho(\sigma)).$$

Textbooks focus on the case  $V = V_\lambda$  is irreducible (because of Maschke's theorem). Since characters are constant on each conjugacy class  $\mu$ , one needs only  $\chi^\lambda(\mu)$ . These are computed by the Murnaghan-Nakayama rule (see below). More recent results include bounds on (normalized) character evaluations [Ro96, FePi11].

(II) If  $V_\lambda$  and  $V_\mu$  are irreducible  $\mathfrak{S}_m$  and  $\mathfrak{S}_n$ -modules, respectively, then  $V_\lambda \otimes V_\mu$  is an irreducible  $\mathfrak{S}_m \times \mathfrak{S}_n$ -module. If  $V_\nu$  is an irreducible  $\mathfrak{S}_{m+n}$ -representation, it restricts to a  $\mathfrak{S}_m \times \mathfrak{S}_n$ -representation  $V_\nu \downarrow_{\mathfrak{S}_m \times \mathfrak{S}_n}^{\mathfrak{S}_{m+n}}$ . The *Littlewood-Richardson coefficient* is

$$c_{\lambda, \mu}^\nu = \text{multiplicity of } V_\lambda \otimes V_\mu \text{ in } V_\nu \downarrow_{\mathfrak{S}_m \times \mathfrak{S}_n}^{\mathfrak{S}_{m+n}}.$$

Many *Littlewood-Richardson rules* are available to count  $c_{\lambda, \mu}^\nu$  [St99].

(III) If  $V_\lambda, V_\mu$  are  $\mathfrak{S}_n$ -modules then so is  $V_\lambda \otimes V_\mu$ . Hence we may write

$$V_\lambda \otimes V_\mu \cong \bigoplus_{\nu \in \mathrm{Par}(n)} V_\nu^{\oplus g_{\lambda, \mu, \nu}}.$$

Here,  $g_{\lambda, \mu, \nu}$  is the *Kronecker coefficient*. One has an  $\mathfrak{S}_3$ -symmetric but cancellative formula  $g_{\lambda, \mu, \nu} = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \chi^\lambda(\sigma) \chi^\mu(\sigma) \chi^\nu(\sigma)$ ; it is an old open problem to give a manifestly nonnegative combinatorial rule. The study of Kronecker coefficients has been given new impetus from *Geometric Complexity Theory*, an approach to the P vs NP problem; see [BlMuSo15] and the references therein.

This note visits a rudimentary point. While for finite groups, character evaluations are algebraic integers, for  $\mathfrak{S}_n$ , in fact  $\chi^\lambda(\mu) \in \mathbb{Z}$ . Moreover, by definition,  $c_{\lambda, \mu}^\nu, g_{\lambda, \mu, \nu} \in \mathbb{Z}_{\geq 0}$ . We remark the three converses hold.<sup>1</sup> The proof uses standard facts, but we are unaware of any specific reference in the textbooks [Ja78, FuHa99, St99, Sa01], or elsewhere.

**Theorem.** *Every integer is infinitely often an irreducible  $\mathfrak{S}_n$ -character evaluation. Every non-negative integer is infinitely often a Littlewood-Richardson coefficient, and a Kronecker coefficient.*

**Corollary A.** *There exists a value-preserving multiset bijection between the Littlewood-Richardson and Kronecker coefficients.*

*Date:* June 30, 2017.

<sup>1</sup>inspired by P. Polo [Po99]: every  $f \in 1 + q\mathbb{Z}_{\geq 0}[q]$  is a Kazhdan-Lusztig polynomial for some  $\mathfrak{S}_n$

*Proof.* Clearly, the Theorem implies that for each  $k \in \mathbb{Z}_{\geq 0}$ , the sets

$$\text{LR}_k = \{(\lambda, \mu, \nu) : c'_{\lambda, \mu} = k\} \text{ and } \text{Kron}_k = \{(\lambda, \mu, \nu) : g_{\lambda, \mu, \nu} = k\}$$

are countably infinite and thus in bijection.  $\square$

Desirable would be a construction of an injection  $\text{Kron}_k \hookrightarrow \text{LR}_k$  for each  $k \in \mathbb{Z}_{\geq 0}$  (avoiding the countable axiom of choice). That should solve the Kronecker problem in (III), by reduction to (II). This we cannot do. However, there has been success in this vein [KnMiSh04] on another counting problem. See the Remark at the end of this paper.<sup>2</sup>

*Proof of the Theorem:* The Murnaghan-Nakayama rule states  $\chi^\lambda(\mu) = \sum_T (-1)^{\text{ht}(T)}$ , where  $T$  is a tableaux of shape  $\lambda$  with  $\mu_i$  many labels  $i$ , the entries are weakly increasing along rows and columns, and the labels  $i$  form a connected skew shape  $T_i$  with no  $2 \times 2$  sub-square;  $\text{ht}(T)$  is the sum of the heights of each  $T_i$ , i.e., one less than the number of rows of  $T_i$ .

We sharpen the assertion about  $\chi^\lambda(\mu)$ . In particular, for a given  $n$ , we consider the intervals of consecutive integers achievable as character evaluations for  $\mathfrak{S}_n$ . From the rule, the character of the *defining representation* satisfies  $\chi^{(n-1,1)}(\mu) = \#(1\text{'s in } \mu) - 1$  (see also [Ja78, Lemma 6.9]). Hence,  $\chi^{(n-1,1)}$  takes the values  $[0, n-2]$ . Similarly,  $\chi^{(2,1^{n-2})}$  achieves an interval of negative integers: Take  $k \in [1, n-5] \cup \{n-3\}$ . If  $k \not\equiv n \pmod 2$ , let  $\mu = (n-k-1, 1^{k+1})$ . Otherwise, if  $k \equiv n \pmod 2$ , let  $\mu = (n-k-4, 3, 1^{k+1})$ . Note that if  $k = n-6$ , let  $\mu$  be these parts in decreasing order. In either case, the rule shows  $\chi^{(2,1^{n-2})}(\mu) = -k$ . Thus, for  $n \geq 5$ ,  $[-(n-5), n-2] \subseteq \{\chi^\lambda(\mu) : \lambda, \mu \in \text{Par}(n)\}$ . Taking  $n \rightarrow \infty$  implies the statement regarding character evaluations.

The *Kostka coefficient*  $K_{\lambda, \mu}$  is the number of *semistandard Young tableaux* of shape  $\lambda$  with content  $\mu$ , i.e., fillings of  $\lambda$  with  $\mu_i$  many  $i$ 's such that rows are weakly increasing and columns are strictly increasing.

**Lemma.** *Every nonnegative integer is infinitely often a Kostka coefficient.*

*Proof.* Clearly,  $K_{(1+j, 1^{k-1}), (j, 1^k)} = k$  for  $j \geq 1$ . The lemma then follows.  $\square$

The Littlewood-Richardson coefficient claim holds since it is long known that Kostka coefficients are a special case. To be specific,  $K_{\lambda, \mu} = c_{\sigma, \lambda}^\tau$  where

$$\tau_i = \mu_i + \mu_{i+1} + \cdots, \quad i = 1, 2, \dots, \ell(\mu), \text{ and}$$

$$\sigma_i = \mu_{i+1} + \mu_{i+2} + \cdots, \quad i = 1, 2, \dots, \ell(\mu) - 1.$$
<sup>3</sup>

For  $\lambda = (\lambda_1, \lambda_2, \dots)$ , let  $\lambda[N] := (N - |\lambda|, \lambda_1, \lambda_2, \dots)$ . F. D. Murnaghan [Mu38] proved that for an integer  $N \gg 0$ ,  $\chi^{\lambda[N]} \otimes \chi^{\mu[N]} = \sum_\nu \overline{g_{\lambda, \mu, \nu}} \chi^{\nu[N]}$ . The  $\overline{g_{\lambda, \mu, \nu}}$  are called *stable Kronecker coefficients* and are evidently a special case of Kronecker coefficients. When  $|\lambda| + |\mu| = |\nu|$  one has  $\overline{g_{\lambda, \mu, \nu}} = c'_{\lambda, \mu}$ . Hence one infers the Kronecker coefficient assertion.  $\square$

<sup>2</sup>There is debate about the idiomatic meaning of *counting rule* or *manifestly nonnegative combinatorial rule* etc. Consider the (adjusted) Fibonacci numbers  $(1, 1, 2, 3, 5, 8, 13, \dots)$ . A counting rule is that  $F_n$  counts the number of  $(1, 2)$ -lists whose sum is  $n$ . The recursive (and computationally efficient) description is  $F_n = F_{n-1} + F_{n-2}$  ( $n \geq 2$ ) where  $F_0 = F_1 = 1$ . Construct a binary tree  $\mathcal{T}_n$  with root labelled  $F_n$ ; each node of label  $F_i$  has a left child  $F_{i-1}$  and right child  $F_{i-2}$ . Leaves of  $\mathcal{T}_n$  are labelled  $F_1$  or  $F_0$ .  $F_n$  counts the number of leaves of  $\mathcal{T}_n$ . The latter description restates the recurrence and is not, *per se*, a counting rule.

<sup>3</sup>This reduction is used by H. Narayanan [Na06] to show computing  $c'_{\lambda, \mu}$  is a  $\#\text{P}$  problem.

When, e.g.,  $n = 25$ , all of  $[-853, 949]$  appear as some  $\chi^\lambda(\mu)$ , but the proof merely guarantees  $[-20, 23]$ . Let  $\ell_n$  be the maximum size of an interval of consecutive character evaluations for  $\mathfrak{S}_n$ . Trivially, the results of [Ro96, FePi11] imply upper bounds for  $\ell_n$ . Can one prove better upper or lower bounds for  $\ell_n$ ?

Let  $A_n$  be the *alternating group* of even permutations in  $\mathfrak{S}_n$ . Sources about the representation theory of  $A_n$  include [JaKe09, Section 2.5] and [FuHa99, Section 5.1]. Character evaluations of  $A_n$  are not always integral, however:

**Corollary B.** *Every integer appears infinitely often as an  $A_n$ -irreducible character evaluation.*

*Proof.* Let  $\psi^\lambda = \chi^\lambda \downarrow_{A_n}^{\mathfrak{S}_n}$  be the character of the restriction of the  $\mathfrak{S}_n$ -irreducible  $V_\lambda$ . If  $\mu$  is not a partition with distinct odd parts then the conjugacy class in  $\mathfrak{S}_n$  of cycle type  $\mu$  is also a conjugacy class of  $A_n$ . If  $\lambda$  is not a self-conjugate partition, the restriction is an  $A_n$ -irreducible and also  $\psi^\lambda(\mu) = \chi^\lambda(\mu)$ . Repeat the Theorem's character argument, since for  $n \geq 4$  neither  $\lambda$  used is self-conjugate, and since for  $k \geq 1$ ,  $\mu$  has equal parts.  $\square$

**Definition.** For a countable indexing set  $A$ , a family of nonnegative integers  $(a_\alpha)_{\alpha \in A}$  is *entire* if every  $k \in \mathbb{Z}_{\geq 0}$  appears infinitely often.

Many of the nonnegative integers arising in algebraic combinatorics are entire. For example, this is true for the theory of *Schubert polynomials* (we refer to [Ma01] for references). If  $w_0 \in \mathfrak{S}_n$  is the longest permutation then  $\mathbb{S}_{w_0}(x_1, \dots, x_n) = x_1^{n-1} x_2^{n-2} \cdots x_{n-1}$ . If  $w \neq w_0$ ,  $w(i) < w(i+1)$  for some  $i$ . Then  $\mathbb{S}_w(x_1, \dots, x_n) = \partial_i \mathbb{S}_{ws_i}(x_1, \dots, x_n)$  where  $\partial_i = \frac{f - s_i(f)}{x_i - x_{i+1}}$  and  $s_i$  is the simple transposition interchanging  $i, i+1$ . Nontrivially, each  $\mathbb{S}_w \in \mathbb{Z}_{\geq 0}[x_1, x_2, \dots]$ . Moreover,  $\mathbb{S}_w = \mathbb{S}_{w \times 1}$  where  $w \times 1 \in \mathfrak{S}_{n+1}$  is the usual image of  $w \in \mathfrak{S}_n$ . Thus we can discuss  $\mathbb{S}_w$  for  $w \in \mathfrak{S}_\infty$ ; these form a  $\mathbb{Z}$ -linear basis of  $\mathbb{Z}[x_1, x_2, \dots]$ . The *Schubert structure constants*  $C_{u,v}^w := [\mathbb{S}_w] \mathbb{S}_u \mathbb{S}_v \in \mathbb{Z}_{\geq 0}$  for geometric reasons. The *Stanley symmetric function* is defined by  $F_w = \lim_{m \rightarrow \infty} \mathbb{S}_{1^m \times w} \in \mathbb{Z}[[x_1, x_2, \dots]]$ ; here  $1^m \times w \in \mathfrak{S}_{m+n}$  sets  $1^m \times w(i)$  equal to  $i$  if  $1 \leq i \leq m$  and equal to  $w(i - m + 1) + m$  otherwise.  $F_w$  is Schur-nonnegative.

**Corollary C.** *These families of nonnegative integers are entire:*

- (a) *The coefficients of monomials in Schubert polynomials.*
- (b) *The Schubert structure constants.*
- (c) *The coefficients of Schur functions in Stanley symmetric functions.*

*Proof.* (a) is true by the Lemma since when  $w$  is *Grassmannian* (has at most one descent),  $\mathbb{S}_w(x_1, \dots, x_n)$  is a Schur polynomial  $s_\lambda$ . When  $u, v$  and  $w$  are Grassmannian with descent position  $d$ , then  $C_{u,v}^w$  is a Littlewood-Richardson coefficient so the Theorem implies (b). Finally, when  $w$  is 321-avoiding (i.e., there does not exist indices  $i < j < k$  such that  $w(i) > w(j) > w(k)$ ),  $F_w = s_{\nu/\lambda} = \sum_{\mu} c_{\lambda, \mu}^{\nu} s_\mu$  is a skew Schur function. Hence, here the coefficient (c) is  $c_{\lambda, \mu}^{\nu}$  and we apply the Theorem.  $\square$

Abstractly, all entire families are mutually in value-preserving bijection. However, for Corollary C one can say more: (a) and (c) are a special cases of (b) (see [BeSo98] and [BuSoYo05]). Can one construct a “wrong way map” (as in  $\mathbb{Q} \hookrightarrow \mathbb{N}$ ) for either (b) $\hookrightarrow$ (a) or (b) $\hookrightarrow$ (c) (thereby finding a rule for  $C_{u,v}^w$ )? A special case indicating the difficulty is:

**Problem.** *Construct an explicit value-preserving injection between Littlewood-Richardson and Kostka coefficients.*

**Remark.** Finding a wrong way map has solved a significant counting rule problem concerning A. Buch-W. Fulton’s *quiver coefficients*. These arise in the study of degeneracy loci of vector bundles over a smooth projective algebraic variety. It was conjectured by those two authors that these integers are nonnegative, with a conjectural counting rule. Also, A. Buch showed that special cases of the quiver coefficients are the numbers from (c) above. The resolution of this problem, due to A. Knutson-E. Miller-M. Shimozono, came by establishing the *opposite*: quiver coefficients are special cases of the well-understood numbers (c). We refer to the solution [KnMiSh04] for background and references.

**Acknowledgements.** We thank Ezra Miller and Bernd Sturmfels for stimulating remarks (over a dozen years ago) concerning [KnMiSh04, BuSoYo05]. We are grateful to Alexander Miller suggesting the character evaluation argument, for data, as well as many other helpful comments. We thank Nantel Bergeron, Colleen Robichaux, Hugh Thomas and Mike Zabrocki for informative discussions. AY was partially supported by an NSF grant.

## REFERENCES

- [BeSo98] N. Bergeron and F. Sottile, *Schubert polynomials, the Bruhat order, and the geometry of flag manifolds*, Duke Math. J. **95**(1998), no. 2, 373–423.
- [BlMuSo15] J. Blasiak, K. D. Mulmuley and M. Sohoni, *Geometric Complexity Theory IV: Nonstandard Quantum Group for the Kronecker Problem*, Mem. Amer. Math. Soc., Vol. 235, No. 1109, 2015.
- [BuSoYo05] A. S. Buch, F. Sottile and A. Yong, *Quiver coefficients are Schubert structure constants*, Math. Res. Lett. **12**(2005), no. 4, 567–574.
- [FePi11] V. Féray and P. Sniady, *Asymptotics of characters of symmetric groups related to Stanley character formula*, Ann. Math., Vol 173(2011), Issue 2, 887–906.
- [FuHa99] W. Fulton and J. Harris, *Representation theory, a first course*, Springer-Verlag, 1999.
- [Ja78] G. D. James, *The Representation Theory of the Symmetric Groups*, Lecture Notes in Mathematics, Volume 682, Springer, 1978.
- [JaKe09] G. D. James and A. Kerber, *The Representation Theory of the Symmetric Group*, Cambridge University Press, Cambridge, 2009.
- [KnMiSh04] A. Knutson, E. Miller and M. Shimozono, *Four positive formulae for type A quiver polynomials*, Invent. Math. **166**(2006), no. 2, 229–325.
- [Ma01] L. Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*. Translated from the 1998 French original by John R. Swallow. SMF / AMS Texts and Monographs, American Mathematical Society, Providence, 2001.
- [Mu38] F. D. Murnaghan, *The analysis of the Kronecker product of irreducible representations of the symmetric group*, Amer. J. Math. **60**(1938), no. 3, 761–784.
- [Na06] H. Narayanan, *On the complexity of computing Kostka numbers and Littlewood-Richardson coefficients*, J. Alg. Comb., Vol. 24, N. 3, 2006, 347–354.
- [Po99] P. Polo, *Construction of arbitrary Kazhdan-Lusztig polynomials in symmetric groups*, Represent. Theory. **3**(1999), 90–104.
- [Ro96] Y. Roichman, *Upper bound on the characters of the symmetric groups*, Invent. Math., Vol 125 (1996), Issue 3, 451–485.
- [Sa01] B. Sagan, *The symmetric group*, Second edition, Graduate Texts in Mathematics, 203. Springer-Verlag, New York, 2001.
- [St99] R. P. Stanley, *Enumerative Combinatorics, Volume 2*, Cambridge University Press, 1999.

UNIVERSITY LABORATORY HIGH SCHOOL, URBANA, IL 61801

*E-mail address:* anshul.adve@gmail.com

DEPT. OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801

*E-mail address:* ayong@uiuc.edu