

Spectroscopy of the All-Charm Tetraquark

V. R. Debastiani*

Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC Institutos de Investigación de Paterna, Aptdo. 22085, 46071 Valencia, Spain *E-mail:* vinicius.rodrigues@ific.uv.es

F. S. Navarra

Instituto de Física, Universidade de São Paulo, C.P. 66318, 05389-970 São Paulo, SP, Brazil E-mail: navarra@if.usp.br

We use a non-relativistic model to study the mass spectroscopy of a tetraquark composed by $c \bar{c} c \bar{c}$ quarks in the diquark-antidiquark picture. By numerically solving the Schrödinger equation with a Cornell-inspired potential, we separate the four-body problem into three two-body problems. Spin-dependent terms (spin-spin, spin-orbit and tensor) are used to describe the splitting structure of the $c\bar{c}$ spectrum and are also extended to the interaction between diquarks. Recent experimental data on charmonium states are used to fix the parameters of the model and a satisfactory description of the spectrum is obtained. We find that the spin-dependent interaction is sizable in the diquark-antidiquark system, despite of the heavy diquark mass, and that the diquark has a finite size if treated in analogy to the $c\bar{c}$ systems. We find that the lowest *S*-wave T_{4c} tetraquarks might be below their thresholds of spontaneous dissociation into low-lying charmonium pairs, while orbital and radial excitations would be mostly above the corresponding charmonium pair threshold. These states could be investigated in the forthcoming experiments at LHCb and Belle II.

XVII International Conference on Hadron Spectroscopy and Structure - Hadron2017 25-29 September, 2017 University of Salamanca, Salamanca, Spain

arXiv:1711.11495v1 [hep-ph] 30 Nov 2017

^{*}Speaker.

[©] Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

1. Introduction

The possible $cc\bar{c}\bar{c}$ states have gained attention recently due to the increasing fauna of exotics and the possibility of producing charmonium pairs in high energy facilities like LHC and KEK. Due to the absence of light quarks, it is unlikely that these states are meson molecules and the tetraquark picture is favored. Assuming a diquark-antidiquark structure, we study the $[cc][\bar{c}\bar{c}]$ tetraquark with a non-relativistic model [1].

The T_{4c} was first proposed by Iwasaki in 1975 [2], just after the discovery of the J/ψ . A few years later, Chao studied this tetraquark in the diquark picture including orbital momentum and spin-dependent forces [3]. After that, only quite few works have investigated the role of orbital momentum or radial excitations. We have thus chosen to focus on these features, extending a successful charmonium model [4] in order to test the diquark assumption and the contribution of spin-dependent terms.



Figure 1: T_{4c} in diquark-antidiquark picture.

2. Formalism

We adopt a quarkonium model [4], solving the Schödinger equation [5] with the potential $V^{(0)}(r)$, which has a Coulomb term from one gluon exchange, linear confinement and a smeared spin-spin interaction:

$$V^{(0)}(r) = \kappa_s \frac{\alpha_s}{r} + br - \frac{8\pi\kappa_s \alpha_s}{3m_c^2} \left(\frac{\sigma}{\sqrt{\pi}}\right)^3 e^{-\sigma^2 r^2} \mathbf{S_1} \cdot \mathbf{S_2}$$
(2.1)

where the color factor κ_s is -4/3 for color singlet and -2/3 for vector diquark in antitriplet color state, while the QCD strong coupling constant α_s , string tension *b* and gaussian parameter σ , together with the charm quark mass m_c , are fitted to the $c\bar{c}$ spectrum (Fig. 2). The spin-orbit and tensor interactions are included as first order perturbation:

$$V^{(1)}(r) = \left(-\frac{3\kappa_s\alpha_s}{2m^2}\frac{1}{r^3} - \frac{b}{2m^2}\frac{1}{r}\right)\mathbf{L}\cdot\mathbf{S} - \frac{\kappa_s\alpha_s}{4m^2}\frac{1}{r^3}\mathbf{S}_{12}$$
(2.2)

where the operator

$$\mathbf{S}_{12} = 4[3(\mathbf{S}_1 \cdot \hat{\mathbf{r}})(\mathbf{S}_2 \cdot \hat{\mathbf{r}}) - \mathbf{S}_1 \cdot \mathbf{S}_2]$$
(2.3)

can be written in terms of spherical harmonics Y_2^m and spin operators S_+ , S_- , S_z to be applied directly in the tetraquark wavefunction [1]. First we calculate the diquark mass using $V_{cc} = V_{c\bar{c}}/2$, where the attractive antitriplet color structure implies the diquark has total spin 1. Next, the diquark mass is used as input to obtain the tetraquark spectrum as a two-body system, allowing radial and orbital excitations between diquark and antidiquark.

3. Results and Conclusions

In Fig. 2 we compare the model results with recent experimental data of $c\bar{c}$ states.



Figure 2: A good reproduction of $c\bar{c}$ mass spectrum is obtained with $m_c = 1.4622$ GeV, $\alpha_s = 0.5202$, b = 0.1463 GeV², $\sigma = 1.0831$ GeV.

We find that the non-relativistic approximation is reasonable, with $\langle v^2/c^2 \rangle < 0.2$. The lowest 1S states of the T_{4c} are found to be very compact, with $\langle r^2 \rangle^{1/2} < 0.3$ fm, where the binding is mostly due to the Coulomb term from one gluon exchange, which also creates a big overlap of the wavefunction at the origin and a strong contribution of the spin-spin interaction.

On the other hand, the lowest diquark is found to have $\langle r^2 \rangle^{1/2} \sim 0.6$ fm, warning against the point-like approximation, which is very common in the literature, even though the two-body approximation should still be reasonable due to the color structure behind it.

In Fig. 3 we show the results for the T_{4c} mass spectrum in analogy to the $c\bar{c}$ one, considering one radial and one orbital excitation and the splitting structure from the spin-dependent terms, in a total of 20 states. We can see that the three lowest 1*S* states with $J^{PC} = 0^{++}$, 1^{+-} , 2^{++} and masses 5969.4, 6020.9, 6115.4 MeV seem to be below¹ the two-meson thresholds $\eta_c \eta_c$, $\eta_c J/\psi$ and $J/\psi J/\psi$, respectively. The inclusion of orbital momentum increases the possibilities in quantum numbers and decay channels. One interesting case is the 1^3P_1 state with exotic quantum numbers $J^{PC} = 1^{-+}$ and mass 6577.4 MeV, about 80 MeV above the corresponding $\eta_c(1S) \chi_{c1}(1P)$ threshold. Detailed results and discussions are presented in Ref. [1].

¹For self-consistence we use the $c\bar{c}$ meson masses calculated with the model. If experimental values were used, the 0⁺⁺ state would actually be a few MeV above the η_c pair threshold.

Table 1: Results for [cc] diquark and T_{4c} tetraquark masses in MeV.

$N^{2S+1}\ell_J$	M_{T4c}	J^{PC}
Diquark		
$1^{3}S_{1}$	3133.4	1+
Tetraquark		
$1^{1}S_{0}$	5969.4	0^{++}
$1^{3}S_{1}$	6020.9	1^{+-}
1^5S_2	6115.4	2^{++}
$1^{1}P_{1}$	6577.1	1
$1^{3}P_{0}$	6480.4	0^{-+}
$1^{3}P_{1}$	6577.4	1^{-+}
$1^{3}P_{2}$	6609.9	2^{-+}
$1^5 P_1$	6495.4	1
$1^5 P_2$	6600.2	$2^{}$
1^5P_3	6641.2	3
$2^{1}S_{0}$	6663.3	0^{++}
$2^{3}S_{1}$	6674.5	1^{+-}
$2^{5}S_{2}$	6698.1	2^{++}
$2^{1}P_{1}$	6944.1	1
$2^{3}P_{0}$	6866.5	0^{-+}
$2^{3}P_{1}$	6943.9	1^{-+}
$2^{3}P_{2}$	6970.4	2^{-+}
$2^5 P_1$	6875.6	1
$2^5 P_2$	6962.1	$2^{}$
$2^{5}P_{2}$	6996.7	3



Figure 3: Tetraquark spectrum including radial and orbital excitations relative to the diquark-antidiquark system and spin-dependent splitting. Ground state ($N^{2S+1}\ell_J = 1^3S_1$) diquarks are used, with $m_{cc} = 3133.4$ MeV.

4. Acknowledgments

The authors acknowledge the support received from the brazilian funding agencies FAPESP (contract 12/50984-4), CNPq and CAPES. We also thank Profs. W. Lucha, F. F. Schöberl for the Mathematica notebook; and M. R. Robilotta, E. Oset, A. Valcarce, J. Vijande for the fruitful discussions. V. R. Debastiani also acknowledges the support from Generalitat Valenciana in the Program Santiago Grisolia (Exp. GRISOLIA/2015/005) and wishes to thank the organizers of the Hadron 2017 for the award this work received in the poster section and S. Neubert, I. Belyaev for the interest and invitation to present this work for LHCb Collaboration in the Hadron Spectroscopy Phenomenology Workshop.

References

- [1] V. R. Debastiani and F. S. Navarra, "A non-relativistic model for the $[cc][\bar{c}\bar{c}]$ tetraquark," arXiv:1706.07553 [hep-ph].
- Y. Iwasaki, Prog. Theor. Phys. 54, 492 (1975); UTHEP-2 (1975); Phys. Rev. Lett. 36, 1266 (1976);
 Phys. Rev. D 16, 220 (1977).
- [3] K. T. Chao, Z. Phys. C 7, 317 (1981).
- [4] T. Barnes, S. Godfrey, E. S. Swanson, Phys. Rev. D 72, 054026 (2005).
- [5] W. Lucha, F. F. Schöberl, Int. J. Mod. Phys. C 10, 607 (1999).