

Effective-energy universality approach describing total multiplicity centrality dependence in heavy-ion collisions

Edward K. Sarkisyan-Grinbaum,^{1,2,*} Aditya Nath Mishra,^{3,†} Raghunath Sahoo,^{4,‡} and Alexander S. Sakharov^{1,5,6,§}

¹*Experimental Physics Department, CERN, 1211 Geneva 23, Switzerland*

²*Department of Physics, The University of Texas at Arlington, Arlington, TX 76019, USA*

³*Instituto de Ciencias Nucleares, Universidad Nacional Autonoma de Mexico, Mexico City, 04510, Mexico*

⁴*Discipline of Physics, School of Basic Science, Indian Institute of Technology, Indore 452020, India*

⁵*Department of Physics, New York University, New York, NY 10003, USA*

⁶*Physics Department, Manhattan College, Riverdale, NY 10471, USA*

The recently proposed participant dissipating effective-energy approach is applied to describe the dependence on centrality of the multiplicity of charged particles measured in heavy-ion collisions at the collision energies up to the LHC energy of 5 TeV. The effective-energy approach relates multihadron production in different types of collisions, by combining, under the proper collision energy scaling, the constituent quark picture with Landau relativistic hydrodynamics. The measurements are shown to be well described in terms of the centrality-dependent effective energy of participants and an explanation of the differences in the measurements at RHIC and LHC are given by means of the recently introduced hypothesis of the energy-balanced limiting fragmentation scaling. A similarity between the centrality data and the data from most central collisions is proposed pointing to the central character of participant interactions independent of centrality. The findings complement our earlier studies of the similar midrapidity pseudorapidity density measurements extending the description to the full pseudorapidity range in view of the similarity of multihadron production in nucleon interactions and heavy-ion collisions.

PACS numbers: 25.75.Dw, 25.75.Ag, 24.85.+p, 13.85.Ni

Recently, the measurements of the centrality dependence of the (mean) total multiplicity of charged particles in PbPb collisions at the center-of-mass (c.m.) energy $\sqrt{s_{NN}} = 5.02$ TeV have been reported by ALICE [1]. Here, we describe these measurements in the framework of the dissipating energy of constituent quark participants, or, for brevity, the effective-energy approach, proposed by two of us in [2, 3]. Exploiting a concept of centrality-dependent effective energy on nucleon participants [4], one demonstrates that the 5.02 TeV ALICE measurements can be well accommodated by the above approach and thus complements the results of our findings [6] of a very good description of the data on the measurements of the midrapidity pseudorapidity densities in nucleus-nucleus collisions within the energy range up to 5.02 TeV, provided that the total multiplicity is the critical variable for obtaining the information on the multihadron production dynamics [7–9].

Let us give a brief description of the effective-energy approach. Within this approach, which interrelates different types of collisions [2], the particle production process is quantified in terms of the amount of *effective* energy deposited into the small Lorentz-contracted volume which is formed at the early stage of a collision. Then, the whole process of the particle production is considered as the expansion of an initial state and the sub-

sequent break-up into particles. This picture resembles the Landau relativistic hydrodynamic model of multiparticle production [10]. In the meantime, the effective-energy approach considers the Landau hydrodynamics being treated in the framework of constituent (or dressed) quarks, in accordance with the additive quark model [11, 12]. This makes the secondary particle production to be basically driven by the amount of the initial energy of constituent quarks pumped into the Lorentz-contracted overlap region of colliding objects. Then, in $pp/\bar{p}p$ collisions, a single constituent quark from each nucleon is assumed to contribute in a collision. The remaining quarks are treated as spectators. The spectator quarks do not participate in the particle production, while result into formation of leading particles and carrying away a significant part of the collision energy. Thus, the effective energy for multiparticle production in $pp/\bar{p}p$ collisions is the energy of a single quark pair interaction, i.e. represents 1/3 of the entire nucleon energy. In collisions of nuclei, however, due to the large size of the nucleus and the long travel path inside the nucleus, more than one quark per nucleon can interact. In the most central (head-on) heavy-ion collisions, where the colliding nuclei are almost fully overlapped, all three constituent quarks from each of the participating nucleons may interact and deposit their energy into the collision zone. Then the whole energy of the nucleons becomes available for the particle production. Within this picture, the bulk measurements in head-on heavy-ion collisions at $\sqrt{s_{NN}}$ are expected to be similar to those from $pp/\bar{p}p$ collisions but at a three times larger c.m. energy $\sqrt{s_{pp}}$, i.e. at $\sqrt{s_{pp}} \simeq 3\sqrt{s_{NN}}$. Let us stress here that the effective-energy approach is

*Electronic address: Edward.Sarkisyan-Grinbaum@cern.ch

†Electronic address: Aditya.Nath.Mishra@cern.ch

‡Electronic address: Raghunath.Sahoo@cern.ch

§Electronic address: Alexandre.Sakharov@cern.ch

considered being applied to the bulk variables, while it is well understood that collective effects such as elliptic flow, correlations would provide further important details but to be addressed in separate studies.

Combining within the above consideration the two contributing ingredients, namely the constituent quark picture and the pseudorapidity density from the Landau hydrodynamics, one obtains the relationship between the charged particle rapidity density per participant pair, $\rho(\eta) = (2/N_{\text{part}})dN_{\text{ch}}/d\eta$, in heavy-ion collisions and in $pp/\bar{p}p$ interactions [5]:

$$\frac{\rho(\eta)}{\rho_{pp}(\eta)} = \frac{2N_{\text{ch}}}{N_{\text{part}} N_{\text{ch}}^{pp}} \sqrt{\frac{L_{pp}}{L_{NN}}} \exp\left[\frac{\eta^2}{2} \left(\frac{1}{L_{pp}} - \frac{1}{L_{NN}}\right)\right],$$

$$\sqrt{s_{pp}} = 3\sqrt{s_{NN}}. \quad (1)$$

Here, N_{ch} and N_{ch}^{pp} are the (total) mean multiplicities in nucleus-nucleus and nucleon-nucleon collisions, respectively, and N_{part} is the number of nucleon participants. The relation of the pseudorapidity density and the mean multiplicity is applied in its Gaussian form as obtained in Landau hydrodynamics. The factor L , which is related to the Lorentz contraction of the system, is defined as $L = \ln(\sqrt{s}/2m)$. According to the approach considered, m is the proton mass, m_p , in nucleus-nucleus collisions and represents the constituent quark mass (m_q) in $pp/\bar{p}p$ collisions set to $\frac{1}{3}m_p$.

At the midrapidity, $\eta \approx 0$, Eq. (1) simplifies to:

$$\frac{\rho(0)}{\rho_{pp}(0)} = \frac{2N_{\text{ch}}}{N_{\text{part}} N_{\text{ch}}^{pp}} \sqrt{\frac{L_{pp}}{L_{NN}}},$$

$$\sqrt{s_{NN}} = \sqrt{s_{pp}}/3. \quad (2)$$

Taking into account that $L_{pp} = \ln(\sqrt{s_{pp}}/2m_q)$ and $L_{NN} = \ln(\sqrt{s_{NN}}/2m_p)$ and setting $m_q = m_p/3$ and $\sqrt{s_{pp}} = 3\sqrt{s_{NN}}$, one gets for Eq. (2):

$$\frac{\rho(0)}{\rho_{pp}(0)} = \frac{2N_{\text{ch}}}{N_{\text{part}} N_{\text{ch}}^{pp}} \sqrt{1 - \frac{4 \ln 3}{\ln(4m_p^2/s_{NN})}},$$

$$\sqrt{s_{NN}} = \sqrt{s_{pp}}/3. \quad (3)$$

Solving Eq. (3), for the multiplicity N_{ch} at a given midrapidity density pseudorapidity $\rho(\eta \approx 0)$ at $\sqrt{s_{NN}}$, and for the rapidity density $\rho_{pp}(0)$ and the multiplicity N_{ch}^{pp} at $3\sqrt{s_{NN}}$, one finds:

$$\frac{2N_{\text{ch}}}{N_{\text{part}}} = N_{\text{ch}}^{pp} \frac{\rho(0)}{\rho_{pp}(0)} \sqrt{1 - \frac{2 \ln 3}{\ln(4.5\sqrt{s_{NN}}/m_p)}},$$

$$\sqrt{s_{NN}} = \sqrt{s_{pp}}/3. \quad (4)$$

In the further development [4], one considers this dependence in terms of centrality. The centrality is closely related to the number of nucleon participants determined

using Monte Carlo Glauber calculations, so that the largest number of participants contribute to the most central heavy-ion collisions. In the meantime, the centrality is regarded as the degree of the overlap of the volumes of the two colliding nuclei, characterized by the impact parameter. The smaller the impact parameter (more central collisions), the larger is the overlap zone. Considering the volumes of the colliding nuclei projected onto the overlapped area as being populated by the number of participants depositing their energy into the Lorentz-contracted volume of the early stage collision zone, the centrality, in the participant effective-energy approach, can be related to the amount of the energy released for particle production, i.e. to the effective energy, ε_{NN} , of the participants. Then this effective energy can be defined as a fraction of the c.m. energy available in a collision according to the centrality, α :

$$\varepsilon_{NN} = \sqrt{s_{NN}}(1 - \alpha). \quad (5)$$

One has to note that this relation is satisfied for all but most peripheral collisions, where no overlap zone can be actually defined and the collisions rather resemble diffractive interactions of nucleons, so no scale factor 1/3 to be applied anymore.

Conventionally, the data are divided into classes of centrality, or centrality intervals, so that α is the average centrality per centrality interval, e.g. $\alpha = 0.25$ for the centrality interval of 20–30% centrality.

Then, for non-central collisions, Eq. (4) reads

$$\frac{2N_{\text{ch}}}{N_{\text{part}}} = N_{\text{ch}}^{pp} \frac{\rho(0)}{\rho_{pp}(0)} \sqrt{1 - \frac{2 \ln 3}{\ln(4.5 \varepsilon_{NN}/m_p)}},$$

$$\varepsilon_{NN} = \sqrt{s_{pp}}/3. \quad (6)$$

considering central collisions of nuclei at the effective c.m. energy ε_{NN} . Here $\rho_{pp}(0)$ and N_{ch}^{pp} are taken from $pp/\bar{p}p$ data at $\sqrt{s_{pp}} = 3\varepsilon_{NN}$.

Let us note that each of the scalings introduced by Eqs. (4) and (5) regulates a particular physics ingredient of the modelling within the participant dissipative effective-energy approach. The scaling embedded in Eq. (4) reflects the constituent quark picture and then reveals a similarity of multihadron production in hadronic and nuclear collisions. The scaling driven by Eq. (5) addresses the energy budget effectively retained in the most central collisions while defining the energy availability for the global variables in non-central collisions.

Figure 1 shows the effective-energy calculations by Eq. (6) compared to the charged particle multiplicity, $N_{\text{ch}}/(N_{\text{part}}/2)$, as a function of the number of participants N_{part} as measured in heavy-ion collisions at $\sqrt{s_{NN}}$ of TeV energies by the ALICE experiment at the LHC [1, 13] and, at GeV energies, by the PHOBOS experiment at RHIC [14]. In the calculations, the midrapidity density $\rho_{pp}(0)$ and the multiplicity N_{ch}^{pp} are taken from the existing $pp/\bar{p}p$ data [9, 15], and the $\rho(0)$ values are




FIG. 1: The charged particle mean multiplicity per participant pair as a function of the number of participants, N_{part} . The solid stars show the dependence measured in PbPb collisions at the LHC by the ALICE experiment at $\sqrt{s_{\text{NN}}} = 2.76$ TeV [13] and 5.02 TeV [1], and the solid circles show the measurements from AuAu collisions at RHIC by the PHOBOS experiment at $\sqrt{s_{\text{NN}}} = 19.6, 62.4, 130$ and 200 GeV [14] (the symbols indicate $\sqrt{s_{\text{NN}}}$ increasing bottom to top). The triangles show the calculations by Eq. (6) using $pp/\bar{p}p$ data within the participant dissipating effective-energy approach. The open squares show the effective-energy calculations which include the energy-balanced limiting fragmentation scaling (see text); the solid lines connect the calculations to guide the eye. The dashed lines represent the calculations using the ALICE fit [1] to the c.m. energy dependence of the mean multiplicity in the most central heavy-ion collisions. The open stars show the ALICE measurements at $\sqrt{s_{\text{NN}}} = 2.76$ TeV multiplied by 1.25, and the open circles show the PHOBOS measurements at $\sqrt{s_{\text{NN}}} = 200$ GeV multiplied by 2.87.

taken from the heavy-ion collision data [14, 16, 17] where available. Where no data exist, the corresponding experimental c.m. energy fits are used. The linear-log [9] and power-law [18] s_{pp} -fits for $\rho_{pp}(0)$ at $\sqrt{s_{pp}} \leq 53$ GeV and at $\sqrt{s_{pp}} > 53$ GeV, respectively, along with the power law c.m. energy fits for N_{ch}^{pp} [5] and $\rho(0)$ [16] are used.

One can see from Fig. 1 that within the dissipating participants effective-energy approach, where the collisions

are driven by the centrality-defined effective c.m. energy ε_{NN} , the calculations well reproduce the centrality dependence obtained in the TeV-energy region from LHC, slightly underestimating a couple of the most peripheral measurements. However, for the RHIC data, the deviation between the measurements and the calculations is seen already for middle N_{part} values. The difference in the behaviour of the data obtained at RHIC and at LHC

becomes more clearer as soon as one multiplies the RHIC 200 GeV data by a factor 2.87 in order to match the ALICE 2.76 TeV data from highly central collisions. In the meantime, one can observe that there is almost no difference between the 2.76 TeV and 5.02 TeV LHC data, where the lower-energy measurements are multiplied by 1.25 to match the higher-energy ones.

The differences observed have been discussed in [5] and an explanation has been given by introducing the energy-balanced limiting fragmentation scaling for the pseudorapidity spectra in non-central collisions. By means of this scaling, the pseudorapidity distributions in heavy-ion collisions at RHIC energies are shown to be reproduced resulting into the centrality independence of the multiplicity, see Fig. 1.

In what follows, the energy-balanced limiting fragmentation hypothesis is applied as it is elaborated in [5].

First, let us notice that, as it is outlined above, in the picture of the effective energy approach, the global observables are defined by the energy of the participating constituent quarks pumped into the overlapped zone of the colliding nuclei. Hence, the bulk production is driven by the initial energy deposited at zero time at rapidity $\eta = 0$, similar to the Landau hydrodynamics. Then, as is expected and indeed found in [2–4, 6], pseudorapidity density (and pseudorapidity transverse energy density) at *midrapidity* is well reproduced for *all* centralities and all available energies.

Meantime, the data shown in Fig. 1, represent the *total* multiplicities, i.e. addresses Eq. (1) after its integration over the *full- η* spectrum. Then, the fragmentation regions contribute, in addition to the midrapidity region, and this contribution has to be taken into account. This point has been addressed in [5], where the rapidity spectra were studied.

As it was found in [5], the calculations well reproduce the data on the full pseudorapidity spectra at all c.m. energies as soon as the effective-energy approach is applied to the most central collisions. As a consequence, the total multiplicity values for central collisions are well reproduced, as one can indeed see in Fig. 1, the points at the large N_{part} . This is understood as soon as in the calculations, the energy is considered to be deposited into the overlapped zone and then, $\sqrt{s_{NN}}$ is one driving the particle-production process in the most central collisions. Consequently, in Eq. (1) the values of the contributed variables are taken from the most central collisions.

For non-central collisions, as discussed above, not the full c.m. energy is considered to contribute to the particle production but the effective energy ε_{NN} to be assumed instead. As soon as this has been applied to the calculations, the calculated pseudorapidity spectra were found [5] to be narrower than the measured ones for the pre-LHC data, so that the fragmentation region were not well reproduced. Consequently, as shown in Fig. 1, the non-central values, measured at RHIC, are seen to be higher and to follow surprisingly constant values in disagreement with the lower, monotonically increasing calcula-

tions and, interestingly, with the LHC data, to which the calculations are found well agree. The deviation between the calculations and the pre-LHC measurements is not surprising and can be explained due to a smaller value of ε_{NN} , used in the calculations, compared to the value of the actual collision energy $\sqrt{s_{NN}}$, as well as due to the fact that the calculations, similar to the Landau approach, are undertaken in the assumption of the head-on character of nuclei collisions, which is clearly not the case for non-central collisions. Therefore, the pseudorapidity distributions have to be corrected in order to balance the energy used in the calculations vs. that in the measurements, to account for the fragmentation region. These, as discussed below and in [5], seem also to explain the difference between the LHC and pre-LHC total multiplicity measurements.

To address the point of the fragmentation region, let us first recall the hypothesis of the limiting fragmentation scaling [19], which states that at high enough energies, the (pseudo)rapidity density spectra for given interacting particle types become independent of the c.m. energy in the fragmentation region when shifted by the beam rapidity $y_{\text{beam}} = \ln(\sqrt{s_{NN}}/m_p)$: $\eta \rightarrow \eta - y_{\text{beam}}$.

As soon as, in the effective-energy approach, the particle production is considered to be driven by the energy of the participants involved in the overlapped zone, one would naturally expect that the behaviour similar to that of the limiting fragmentation to hold for the calculated pseudorapidity distribution but when the latter is shifted by the rapidity defined by the effective energy ε_{NN} , namely $y_{\text{eff}} = \ln(\varepsilon_{NN}/m_p)$. Indeed, in [5], it was found that for a non-central collision, the effective-energy calculation of the pseudorapidity distribution matches immediately the measured spectra as soon as the former is shifted by y_{eff} while the latter by y_{beam} . This effect is observed to hold independently of centrality, as soon as the corresponding y_{eff} shift was applied.

As so, this unambiguously points out that, in order to describe the unshifted η -distribution, one needs to take into account the difference between the c.m. energy in the measurements and the effective energy of the participants in the calculations, i.e. in the other words to *balance* the energy. To this end, the calculated distribution, Eq. (1), is shifted by the energy difference, $y_{\text{eff}} - y_{\text{beam}} = \ln(\varepsilon_{NN}/\sqrt{s_{NN}})$, in the fragmentation region, or, due to Eq. (5), by $\ln(1 - \alpha)$. This immediately put in agreement the calculated distribution and the measurements for non-central collisions with no deficit in the fragmentation region [5]. Due to this, the hypothesis was named the energy-balanced limiting fragmentation scaling. Using this scaling, the calculated pseudorapidity density spectrum gets corrected outside the central- η region for all centralities. It is clear that in head-on a collisions, α tends to zero what makes the shift negligible.

To add is that where no pseudorapidity density distributions are available in $pp/\bar{p}p$ data at $\sqrt{s_{pp}} = 3\varepsilon_{NN}$, and therefore no integration is possible using Eq. (1), the

energy-balanced limiting fragmentation scaling is applied to reproduce the calculated $\rho(\eta)$. The measured distribution from a non-central heavy-ion collision is shifted by $(y_{\text{beam}} - y_{\text{eff}})$, i.e. $\eta \rightarrow \eta + \ln(1 - \alpha)$. Then N_{ch} is calculated by adding the difference between the integral from the obtained shifted distribution and the measured multiplicity to the midrapidity calculation of Eq. (6). Note that for central (midrapidity) region, there is no need to apply the energy-balanced limiting fragmentation, as soon as particle production in this region is well described by the effective energy approach of centrally colliding participants, as it is discussed above.

Using this ansatz, the values of N_{ch} , calculated for each centrality for the RHIC measurements, found to reproduce well the pre-LHC data, as shown in Fig. 1.

In the meantime, one can also conclude from Fig. 1 that, in contrast to the pre-LHC energies, almost no energy-balanced contribution is needed for the calculations to describe the LHC mean multiplicity measurements at TeV c.m. energies. Given this observation and the fact that the calculations imply they are made in an assumption of central nucleus-nucleus collisions at the c.m. energy of the value of ε_{NN} , one can conclude that in TeV-energy heavy-ion collisions, the multihadron production obeys a head-on collision regime, for all the centralities measured. This conclusion is supported by the one made in [5] where the ε_{NN} dependence of the centrality multiplicity data are shown to well follow and so complementing the $\sqrt{s_{NN}}$ dependence of the head-on multiplicity data without taking into account the energy balance. This feature is also clearly seen here, as shown in Fig. 1, where the ALICE heavy-ion *head-on* $\sqrt{s_{NN}}$ fit [1] is demonstrated to well follow the LHC *centrality* data expressed in terms of ε_{NN} .

The above findings may be treated pointing to apparently different regime of hadroproduction occurring in heavy-ion collisions as $\sqrt{s_{NN}}$ moves from GeV to TeV energies. This conclusion finds its support also in our results [6] of describing the midrapidity density $\sqrt{s_{NN}}$ -dependence as soon as the LHC data are included.

The energy-balanced limiting fragmentation provides also an explanation to the unexpected difference of the midrapidity pseudorapidity density increase vs. the total multiplicity independence with the increase of the number of nucleon participants as measured at RHIC (RHIC ‘‘puzzle’’ [5]). At GeV energies, the multiplicity gets an additional fraction of the energy as the difference between the collision c.m. energy shared by all nucleons and the effective energy of the participants driving the particle production process. No such difference appears at the LHC where both variables are found to increase with the number of participants, in agreement with the central character of all-centrality collisions as discussed

just above.

Let us note in conclusion that the picture of the effective-energy approach is shown as well to explain [2, 3] the similarity of the measurements in other collisions, such the scaling between the charged particle mean multiplicity in e^+e^- and $pp/\bar{p}p$ collisions [20] and the universality of both the multiplicity and the midrapidity density measured in the most central nuclear collisions and in e^+e^- annihilation [21]; see [8, 9] for discussion. In the latter case, colliding leptons are considered to be structureless and deposit their total energy into the Lorentz-contracted volume. This is shown [9] to be supported by the observation that the multiplicity and η -distributions in $pp/\bar{p}p$ interactions are well reproduced by e^+e^- data as soon as the inelasticity is set to ≈ 0.35 , i.e. effectively 1/3 of the hadronic interaction energy. For recent discussion on the universality of hadroproduction up to LHC energies, see [15].

Summarizing, the effective-energy dissipation approach based on the picture combining the constituent quark model together with Landau relativistic hydrodynamics in sense of the universality of the multihadron production in hadronic and nuclear collisions is shown to well describe the total (mean) multiplicity data in GeV to TeV c.m. energy heavy-ion collisions. In particular, the centrality dependence of the multiplicity of charged particles measured up to 5 TeV are shown to be well reproduced. In addition, the centrality dependence of the data is shown to be in good agreement with the fit to the head-on c.m. energy dependence confirming the central character of the collisions independent of the centrality measured. The observation of differences between the RHIC and the LHC measurements suggests the change of the particle production regime as the c.m. energy of heavy-ion collisions moves from GeV to TeV energy region. This study complements and supports the results from our earlier investigations [6] of the midrapidity pseudorapidity density centrality dependence measurements in the same energy range while now made for the full-rapidity interval. Foreseen measurements at LHC and future colliders at higher energies and with different types of colliding objects are of high interest in clarifying the features of the participant effective-energy approach and in view of better understanding of the mechanism of the hadroproduction process.

Raghunath Sahoo acknowledges the financial support from Project No. SR/MF/PS-01/2014-IITI(G) of Department of Science & Technology, Government of India. The work of Alexander Sakharov is partially supported by the US National Science Foundation under Grants No. PHY-1505463 and No. PHY-1402964.

[1] ALICE Collab., J. Adam *et al.*, Phys. Lett. B **772**, 567 (2017).

[2] E.K.G. Sarkisyan, A.S. Sakharov, AIP Conf. Proc. **828**,

- 35 (2006).
- [3] E.K.G. Sarkisyan, A.S. Sakharov, *Eur. Phys. J. C* **70**, 533 (2010).
- [4] A.N. Mishra, R. Sahoo, E.K.G. Sarkisyan, A.S. Sakharov, *Eur. Phys. J. C* **74**, 3147 (2014).
- [5] E.K.G. Sarkisyan, A.N. Mishra, R. Sahoo, A.S. Sakharov, *Phys. Rev. D* **93**, 054046 (2016).
- [6] E.K.G. Sarkisyan, A.N. Mishra, R. Sahoo, A.S. Sakharov, *Phys. Rev. D* **94**, 011501 (2016).
- [7] I.M. Dremin, J.W. Gary, *Phys. Rep.* **349**, 301 (2001).
- [8] W. Kittel, E.A. De Wolf, *Soft Multihadron Dynamics* (World Scientific, Singapore, 2005)
- [9] J.F. Grosse-Oetringhaus, K. Reygers, *J. Phys. G* **37**, 083001 (2010)
- [10] L.D. Landau, *Izv. Akad. Nauk SSSR: Ser. Fiz.* **17**, 51 (1953). English translation: *Collected Papers of L. D. Landau*, ed. by D. Ter-Haarp (Pergamon, Oxford, 1965), p. 569. Reprinted in: *Quark-Gluon Plasma: Theoretical Foundations*, ed. by J. Kapusta, B. Müller, J. Rafelski (Elsevier, Amsterdam, 2003), p. 283.
- [11] For review and a collection of reprints on original papers on quarks and composite models, see: J.J.J. Kokkedee, *The Quark Model* (W.A. Benjamin, Inc., New York, 1969).
- [12] For recent comprehensive review on soft hadron interactions in the additive quark model, see: V.V. Anisovich, N.M. Kobrinsky, J. Nyiri, Yu.M. Shabelsky, *Quark Model and High Energy Collisions* (World Scientific, Singapore, 2004).
- [13] ALICE Collab., J. Adam *et al.*, *Phys. Lett. B* **754**, 373 (2016).
- [14] B. Alver *et al.*, *Phys. Rev. C* **83**, 024913 (2011).
- [15] The Review of Particle Physics, M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018).
- [16] ALICE Collab., J. Adam *et al.*, *Phys. Rev. Lett.* **116**, 222302 (2016).
- [17] ALICE Collab., K. Aamodt *et al.*, *Phys. Rev. Lett.* **106**, 032301 (2011).
- [18] ALICE Collab., J. Adam *et al.*, *Eur. Phys. J. C* **77**, 33 (2017).
- [19] J. Benecke, T.T. Chou, C.N. Yang, E. Yen, *Phys. Rev.* **188**, 2159 (1969).
- [20] P.V. Chliapnikov, V.A. Uvarov, *Phys. Lett. B* **251**, 192 (1990)
- [21] PHOBOS Collab., B.B. Back *et al.*, nucl-ex/0301017, *Phys. Rev. C* **74**, 021902 (2006).