

HIGH-DIMENSIONAL INSTRUMENTAL VARIABLES REGRESSION AND CONFIDENCE SETS - V2/2012

ERIC GAUTIER AND ALEXANDRE B. TSYBAKOV

Date: First version December 2009, this version: November 2012.

Background facts: This was a revision of the arXiv preprint arXiv:1105.2454v1 which was online here. A link is given in footnote 2 of arXiv:1105.2454v5. (3.5) is a different *STIV* estimator from the one in arXiv:1105.2454v1 where, instead of one conic constraint, there are as many conic constraints as moments (instruments) allowing to use more directly moderate deviations for self-normalized sums. The idea first appeared in formula (6.5) in arXiv:1105.2454v1 when some instruments can be endogenous. This was called the *STIV* until the 2014 version. Eventually, the original *STIV* was put back in the v2 on arXiv because we did not agree on the usefulness of the additional complication. For reference and to avoid confusion with the *STIV* estimator, this estimator should be called *C-STIV*.

Keywords: Instrumental variables, sparsity, *STIV* estimator, endogeneity, high-dimensional regression, conic programming, heteroscedasticity, confidence regions, non-Gaussian errors, variable selection, unknown variance, sign consistency.

We thank James Stock and three anonymous referees for comments that greatly improved this paper. We also thank Don Andrews, Ivan Canay, Victor Chernozhukov, Azeem Shaikh and the seminar participants at Brown, Bocconi, CEMFI, CREST, Compiègne, Harvard-MIT, Institut Henri Poincaré, LSE, Paris 6 and 7, Princeton, Queen Mary, Toulouse, Valparaíso, Wharton, Yale, as well as participants of SPA, Saint-Flour, ERCIM 2011, the 2012 CIREQ conference on High Dimensional Problems in Econometrics and 4th French Econometrics Conference for helpful comments.

ABSTRACT. We propose an instrumental variables method for inference in structural linear models with endogenous regressors in the high-dimensional setting where the number of possible regressors K can be much larger than n . Our new procedure, called *STIV* (Self Tuning Instrumental Variables) estimator, is realized as a solution of a conic optimization program. We allow for partial identification and very weak distributional assumptions including heteroscedasticity. We do not need prior knowledge of the variances of the errors. A key ingredient is sparsity, *i.e.*, the vector of coefficients has many zeros. The main result of the paper is nested confidence sets, sometimes non-asymptotic, around the identified region under various level of sparsity. In the presence of very many instruments, a number exponential in the sample size, too many non-zero coefficients, or when there is an endogenous regressor which is only instrumented by instruments which are too weak, our confidence sets have infinite volume. We show that a variation on the *STIV* estimator is a new robust method to estimate low dimensional models without sparsity and possibly many weak instruments. We obtain rates of estimation and show that, under appropriate assumptions, a thresholded *STIV* estimator correctly selects the non-zero coefficients with probability close to 1. In a non sparse setting, we obtain a sparse oracle inequality that shows how well the regression coefficients are estimated when the model can be well approximated by a sparse model. In our *IV* regression setting, the standard tools from the literature on sparsity, such as the restricted eigenvalue assumption are inapplicable. Therefore, for our analysis we develop a new approach based on data-driven sensitivity characteristics. We also obtain confidence sets, when the endogenous regressors in the high-dimensional structural equation have a sparse reduced form and we use a two-stage procedure, akin to two-stage least squares. We finally study the properties of a two-stage procedure to deal with the detection of non-valid (endogenous) instruments.

1. INTRODUCTION

In this article we consider a structural model of the form

$$(1.1) \quad y_i = x_i^T \beta + u_i, \quad i = 1, \dots, n,$$

where x_i are vectors of explanatory variables of dimension $K \times 1$, u_i is a zero-mean random error possibly correlated with some or all regressors. We consider the problem of inference on the structural parameter β from n independent, not necessarily identically distributed, realizations (y_i, x_i^T, z_i^T) , $i = 1, \dots, n$. This allows for heteroscedasticity. We denote by x_{ki} , $k = 1, \dots, K$, the components of x_i . The regressors x_{ki} are called endogenous if they are correlated with u_i and they are called exogenous otherwise. It is well known that endogeneity occurs when a regressor correlated both with y_i and regressors in the model is unobserved; in the errors-in-variables model when the measurement error

is independent of the underlying variable; when a regressor is determined simultaneously¹ with the response variable y_i . A random vector z_i of dimension $L \times 1$ is called a vector of instrumental variables (or instruments) if it satisfies

$$(1.2) \quad \forall i = 1, \dots, n, \quad \mathbb{E}[z_i u_i] = 0,$$

where $\mathbb{E}[\cdot]$ denotes the expectation. Having access to instrumental variables makes it possible to identify the vector β when $\mathbb{E}[z_i x_i^T]$ has full column rank. The case where $K \geq L$ is typically a case where identification fails (see Example 6 below).

Though the method of this paper can be applied in standard setups², we are mainly interested in the more challenging high-dimensional setting where K can be much larger than n and one of the following two assumptions is satisfied:

- (i) only few coefficients β_k are non-zero (β is *sparse*),
- (ii) β can be well approximated by a sparse vector (β is *approximately sparse*).

The first assumption means that K is not the actual number of regressors but the number of potential regressors. All of the regressors do not necessarily appear in the true model underlying (1.1). The second assumption is likely to be satisfied in a wage equation because in some data sets there are many variables that have a nonzero effect on wage but their marginal effect is too small to matter. The large K relative to n problem allows to deal with various important situations encountered in empirical applications. Here are some examples of such applications. In examples 3, 4 and 5 the most likely assumption is the approximate sparsity assumption (ii).

Example 1. Economic theory is not explicit enough about which variables belong to the true model. Sala-i-Martin (1997) and Belloni and Chernozhukov (2001b) give examples from development economics where model selection is important. In macroeconomics, development economics or international finance, it is common to consider cross-country regressions. Because n is of the order of a few dozens, it is important to allow for K to be much larger than n .

Example 2. Rich heterogeneity. When there is a rich heterogeneity one usually wants to control for many variables and possibly interactions, or to carry out a stratified analysis where models are estimated in small population sub-groups (*e.g.*, groups defined by the value of an exogenous discrete variable). In both cases K can be large relative to n .

¹This is the case where β is actually structural.

²This is also very attractive because it is a feasible, non combinatoric, method, unlike a variable selection method like BIC.

Example 3. Many endogenous regressors due to a type of non-parametric specification.

This occurs when one considers a structural equation of the form

$$(1.3) \quad \begin{aligned} y_i &= f(x_{\text{end},i}) + u_i \\ &= \sum_{k=1}^K \alpha_k f_k(x_{\text{end},i}) + u_i \end{aligned}$$

$$(1.4) \quad = x_i^T \beta + u_i$$

and the stronger notion of exogeneity $\mathbb{E}[u_i|z_i] = 0$ (zero conditional mean assumption), $x_{\text{end},i}$ is a low dimensional vector of endogenous regressor, $x_i = (f_1(x_{\text{end},i}), \dots, f_K(x_{\text{end},i}))^T$, $\beta = (\alpha_1, \dots, \alpha_K)^T$ and $(f_k)_{k=1}^K$ are functions from a dictionary³. Exogenous regressors could also be included in the right hand side of (1.3) similar to a partial linear model (see next example). Belloni and Chernozhukov (2011b) gives the example of a wage equation with many transformation of education to properly account for nonlinearities. Another typical application is the estimation of Engle curves where it is important to include nonlinearities in the total budget (see, *e.g.*, Blundell, Chen and Kristensen (2007)). When one estimates Engle curves using aggregate data, n is again usually of the order of a few dozens. It is well known that education in a wage equation and total budget in Engle curves are endogenous variables.

Example 4. Many exogenous regressors due to a type of semi-parametric specification.

Consider a partially linear model of the form

$$(1.5) \quad y_i = x_{\text{end},i}^T \beta_{\text{end}} + f(x_{\text{exo},i}) + u_i \quad \mathbb{E}[u_i|x_{\text{exo},i}] = 0.$$

$x_{\text{exo},i}$ is a low dimensional vector of exogenous regressors. If f can be properly decomposed as a linear combination of functions from a large enough dictionary $(f_k)_{k=1}^{K_c}$, with coefficients α_k , we obtain

$$(1.6) \quad y_i = x_{\text{end},i}^T \beta_{\text{end}} + \sum_{k=1}^{K_c} \alpha_k f_k(x_{\text{exo},i}) + u_i$$

$$(1.7) \quad = x_i^T \beta + u_i$$

³We use the word dictionary, common in the machine learning community, because high-dimensional methods allow to consider very many series terms and to mix bases. This is not exactly the nonparametric *IV* setup because in a truly nonparametric model there would be an extra approximation error term in (1.4), of order at most $n^{-1/2}$ when K is very large and we make minimal smoothness assumptions. High dimensional methods allow in several cases to obtain an adaptive estimation method (that does not need to know the smoothness of the unknown function) but we will not discuss this aspect in this article.

where $\beta = (\beta_{\text{end}}^T, \alpha_1, \dots, \alpha_{K_c})^T$ and $x_i = (x_{\text{end},i}^T, f_1(x_{\text{exo},i}), \dots, f_{K_c}(x_{\text{exo},i}))^T$. In that case, one is usually interested in the marginal effects β_{end} of $x_{\text{end},i}$ holding fixed $x_{\text{exo},i}$, rather than the whole vector β .

Example 5. Many control variables to justify the use of an instrument. Suppose that we are interested in the parameter $\bar{\beta}$ in

$$(1.8) \quad y_i = \bar{x}_i^T \bar{\beta} + v_i,$$

where some of the variables in \bar{x}_i are endogenous, but that we have at our disposal a variable z_i that we want to use as an instrument but which does not satisfy $\mathbb{E}[z_i v_i] = 0$. Suppose that we also have observations of vectors of controls w_i such that $\mathbb{E}[v_i | w_i, z_i] = \mathbb{E}[v_i | w_i]$ (conditional mean independence). Then we can rewrite (1.8) as

$$(1.9) \quad y_i = \bar{x}_i^T \bar{\beta} + f(w_i) + u_i$$

where $f(w_i) = \mathbb{E}[v_i | w_i]$ and $u_i = v_i - \mathbb{E}[v_i | w_i, z_i]$ is such that $\mathbb{E}[z_i u_i] = 0^4$. It yields

$$(1.10) \quad y_i = \bar{x}_i^T \bar{\beta} + \sum_{k=1}^{K_c} \alpha_k f_k(w_i) + u_i$$

when f can be decomposed on $(f_k)_{k=1}^{K_c}$. This model can be rewritten in the form (1.1) with $\beta = (\bar{\beta}^T, \alpha_1, \dots, \alpha_{K_c})^T$ and $x_i = (\bar{x}_i^T, f_1(w_i), \dots, f_{K_c}(w_i))^T$. Again one is usually interested in the sub-vector $\bar{\beta}$.

Example 6. The instruments can have a direct effect on the outcome. Kolesár, Chetty, Friedman, et al. (2011) considers the case where one wants to allow the instruments to have a direct effect on the outcome, *i.e.*, to be potentially on the right hand side of (1.1). We are thus in a setting where $K > L$ which implies that β is not identified from the moment conditions (1.2). One typically needs exclusion restrictions (*i.e.*, that some exogenous variables are instruments and do not appear on the right hand side of (1.1)) which corresponds to some coefficients being equal to zero in the specification where all instruments can potentially appear on the right hand side of (1.1). Exclusion restrictions is therefore a reason for sparsity. This specification is more flexible as we assume that there exists some exclusion restrictions without telling which one in advance. In a setup where there

⁴By the law of iterated conditional expectations.

are many (weak) instruments⁵, allowing the instruments to potentially have a direct effect, implies that K is large.

Statistical inference under the sparsity scenario when the dimension is larger than the sample size is now an active and challenging field. The most studied techniques are the Lasso, the Dantzig selector (see, *e.g.*, Candès and Tao (2007), Bickel, Ritov and Tsybakov (2009); more references can be found in the recent book by Bühlmann and van de Geer (2011), as well as in the lecture notes by Koltchinskii (2011), Belloni and Chernozhukov (2011b)), and agregation methods (see Dalalyan and Tsybakov (2008), Rigollet and Tsybakov (2011) and the papers cited therein). A central concern in this literature is to propose methods that are computationally feasible. The Lasso for example is a convex relaxation of l^0 penalized least squares methods like BIC. The last are NP -hard and it is impossible in practice to consider the case where K is larger than a few dozens. The Dantzig selector is solution of a simple linear program. In recent years, these techniques became a reference in several areas, such as biostatistics and imaging. Some important extensions to model from econometrics have been obtained by Belloni and Chernozhukov (2011a) who study the ℓ_1 -penalized quantile regression and give an application to cross-country growth analysis and by Belloni, Chernozhukov and Hansen (2010) who use the Lasso to estimate the optimal instruments with an application to the impact of eminent domain on economic outcomes. Städler, Buhlmann and van de Geer (2010) studies the estimation of mixtures of high-dimensional regressions, this is important in econometrics to handle group heterogeneity. Caner (2009) studies a Lasso-type GMM estimator. Rosenbaum and Tsybakov (2010) deal with the high-dimensional errors-in-variables problem and discuss an application to hedge fund portfolio replication. Belloni and Chernozhukov (2011b) also presents several applications of high-dimensional methods to economics. The high-dimensional setting in a structural model with endogenous regressors that we are considering here has not yet been analyzed. Note that the direct implementation of the Lasso or Dantzig selector fails in the presence of a single endogenous regressor simply because the zero coefficients in the structural equation (1.1) do not correspond to the zero coefficients in a linear projection type model. We also obtain confidence sets in a high-dimensional framework.

The main message of this paper is that, in model (1.1) containing endogenous regressors, under point identification, the high-dimensional vector of coefficients or a subset of important coefficients can be estimated together with proper confidence sets using instrumental variables. In partially identified settings, we obtain confidence sets around the identified region under a sequence of sparsity scenarios

⁵As exemplified in Angrist and Krueger (1991), under the stronger notion of exogeneity based on the zero conditional mean assumption, considering interactions or functionals of instruments can lead to a large number of instruments.

varying the upper bound s on the number of non-zero coefficients of β (that we call *sparsity certificate*). Thus there is no restriction on the size of K . When K is too large, one loses identification. Our confidence sets can have infinite volume when s and/or K is too large. There is no restriction either on the strength of the instruments. If a very large number of instruments is used (exponential in the sample size), or when all the instruments are weak, the method yields infinite volume confidence sets. The price to pay for including an irrelevant instrument in a preliminary set of L instruments is just a factor of $\sqrt{\log(L+1)}/\sqrt{\log(L)}$ in the size of the confidence sets. The *STIV* estimator is thus a method that is robust to weak instruments and can handle very many (*i.e.* exponential in the sample size⁶) instruments. Also it is not required in principle to know in advance which regressor is endogenous and which is not, one needs a set of valid (exogenous) instruments. This is achieved by the *STIV* estimator (Self Tuning Instrumental Variables estimator) that we introduce below. Based on it, we can also perform variable selection. Under various assumptions on the data generating process we obtain non-asymptotic or asymptotic results. We also provide meaningful bounds when either (i) or (ii) above holds and $\log(L)$ is small compared to n . We believe that a non-asymptotic framework is the most natural setting to consider in high dimensions. We also restrict mostly our attention to a one stage method. This is even more justified in *IV* estimation of structural equations with endogenous regressors and weak instruments where inference usually relies on non-standard asymptotics (see, *e.g.*, Stock, Wright and Yogo (2002) and Andrews and Stock (2007) for a review and the references cited therein). Nelson and Startz (1990a, b) contains a simulation study where, due to weak instruments, the finite sample distribution of the two-stage least squares estimator can be non normal and even bimodal.

The *STIV* estimator is an extension of the Dantzig selector of Candès and Tao (2007). The results of this paper extend those on the Dantzig selector (see Candès and Tao (2007), Bickel, Ritov and Tsybakov (2009) and further references in Bühlmann and van de Geer (2011)) in several ways: By allowing for endogenous regressors when instruments are available, by working under weaker sensitivity assumptions than the restricted eigenvalue assumption of Bickel, Ritov and Tsybakov (2009), which in turns yields tighter bounds, by imposing weak distributional assumptions, by introducing a procedure independent of the noise level and by providing confidence sets.

The *STIV* estimator is also very much inspired by the Square-root Lasso of Belloni, Chernozhukov and Wang (2010) which proposes a pivotal method independent of the variance of the errors

⁶The terminology very many is borrowed from Belloni, Chen, Chernozhukov et al. (2010).

in the Gaussian linear model with fixed regressors. The Square-root Lasso is a very important contribution to the literature on the Lasso where almost all articles require to know the variance of the errors which is related to the degree of penalization required. The most common practice is to adjust the degree of penalization is to use cross validation or BIC. The *STIV* estimator adds extra linear constraints coming from the restrictions (1.2) to the Square-root Lasso which allows one to deal with endogenous regressors. The implementation of the *STIV* estimator also correspond to solving a simple conic optimization program. Confidence sets require as well solving at most $2K^2$ linear programs for lower bounding the sensitivities that we introduce below. So our method is easy and fast to implement in practice. Our confidence sets rely on moderate deviations for self-normalized sums and either on the sparsity certificates or on perfect model selection. Perfect model selection requires a separation from zero of the non-zero coefficients⁷, while the sparsity certificate approach does not. We are indebted to Belloni, Chernozhukov and Wang (2010) who are the first to use moderate deviations for self-normalized sums results for high-dimensional regression (see also Belloni, Chen, Chernozhukov and Hansen (2011)). We will see that under some distributional assumptions these can lead to confidence sets that have finite sample coverage properties.

2. BASIC DEFINITIONS AND NOTATION

We set $\mathbf{Y} = (y_1, \dots, y_n)^T$, $\mathbf{U} = (u_1, \dots, u_n)^T$, and we denote by \mathbf{X} and \mathbf{Z} the matrices of dimension $n \times K$ and $n \times L$ respectively with rows x_i^T and z_i^T , $i = 1, \dots, n$.

The sample mean is denoted by $\mathbb{E}_n[\cdot]$. We use the notation

$$\mathbb{E}_n[X_k^a U^b] \triangleq \frac{1}{n} \sum_{i=1}^n x_{ki}^a u_i^b, \quad \mathbb{E}_n[Z_l^a U^b] \triangleq \frac{1}{n} \sum_{i=1}^n z_{li}^a u_i^b,$$

where x_{ki} is the k th component of vector x_i , and z_{li} is the l th component of z_i for some $k \in \{1, \dots, K\}$, $l \in \{1, \dots, L\}$, $a \geq 0, b \geq 0$. Similarly, we define the sample mean for vectors; for example, $\mathbb{E}_n[UX]$ is a row vector with components $\mathbb{E}_n[UX_k]$. We also define the corresponding population means:

$$\mathbb{E}[X_k^a U^b] \triangleq \frac{1}{n} \sum_{i=1}^n \mathbb{E}[x_{ki}^a u_i^b], \quad \mathbb{E}[Z_l^a U^b] \triangleq \frac{1}{n} \sum_{i=1}^n \mathbb{E}[z_{li}^a u_i^b],$$

⁷We believe that this is unavoidable for model selection but presumably strong for some econometrics applications. However this result is a useful addition to the current state of the art in the theory of high-dimensional regression even with exogenous regressors.

and set, for $l = 1, \dots, L$,

$$z_{l*} \triangleq \max_i |z_{li}|, \quad (x \bullet z)_l = \left\{ \max_{k=1, \dots, K} \mathbb{E}_n \left[\left(\frac{X_k}{\mathbb{E}_n[X_k^2]^{1/2}} Z_l \right)^2 \right] \right\}^{1/2}.$$

We denote by \mathbf{D}_X the diagonal $K \times K$ matrices with diagonal entries $\mathbb{E}_n[X_k^2]^{-1/2}$ for $k = 1, \dots, K$. We use the notation $\mathbf{D}_Z^{(I)}$ for the $L \times L$ matrix of entries $(x \bullet z)_l^{-1}$ for $l \in I$ and z_{l*}^{-1} otherwise. The set I is a subset of $\{1, \dots, L\}$. It always contains the index of the instrument which is identically equal to 1. The corresponding coefficient β_k is the usual constant in (1.1). The matrices \mathbf{D}_X and $\mathbf{D}_Z^{(I)}$ are normalization matrices. The user specifies the set I based on the observed distribution. For fast implementation of our algorithm, z_{l*}^{-1} is the preferred normalization. For heavy tail distributions, though, one may want to use the second normalization. The more heavy tail instruments we have, the more computationally intensive our method will be. Indeed, we will add a conic constraint for each heavy tail instrument. For a vector $\beta \in \mathbb{R}^K$, let $J(\beta) = \{k \in \{1, \dots, K\} : \beta_k \neq 0\}$ be its support, *i.e.*, the set of indices corresponding to its non-zero components β_k . We denote by $|J|$ the cardinality of a set $J \subseteq \{1, \dots, K\}$ and by J^c its complement: $J^c = \{1, \dots, K\} \setminus J$. The subset of indices $\{1, \dots, K\}$ corresponding to variables in (1.1) that are known in advance to be exogenous and serve as their own instruments is denoted by J_{exo} . There might be more true exogenous variables in (1.1), they are excluded from this list of indices if they have not been included as instruments. The ℓ_p norm of a vector Δ is denoted by $|\Delta|_p$, $1 \leq p \leq \infty$. For $\Delta = (\Delta_1, \dots, \Delta_K)^T \in \mathbb{R}^K$ and a set of indices $J \subseteq \{1, \dots, K\}$, we consider $\Delta_J \triangleq (\Delta_1 \mathbb{1}_{\{1 \in J\}}, \dots, \Delta_K \mathbb{1}_{\{K \in J\}})^T$, where $\mathbb{1}_{\{\cdot\}}$ is the indicator function. For a vector $\beta \in \mathbb{R}^K$, we set $\overrightarrow{\text{sign}(\beta)} \triangleq (\text{sign}(\beta_1), \dots, \text{sign}(\beta_K))$ where

$$\text{sign}(t) \triangleq \begin{cases} 1 & \text{if } t > 0 \\ 0 & \text{if } t = 0 \\ -1 & \text{if } t < 0 \end{cases}$$

For $a \in \mathbb{R}$, we set $a_+ \triangleq \max(0, a)$, $a_+^{-1} \triangleq (a_+)^{-1}$, and $a/0 \triangleq \infty$ for $a > 0$. We adopt the convention $0/0 \triangleq 0$ and $1/\infty \triangleq 0$.

We denote by

$$\mathcal{I}_{\text{ident}} = \{\beta : \mathbb{E}[z_i(y_i - x_i^T \beta)] = 0\}$$

the identified region. It is an affine space which is reduced to a point when the model (1.1)-(1.2) is point identified. It is possible to impose some restrictions like a known sign or a prior upper bound on the size of the coefficients. However, for simplicity, we will just consider the case when we know

an a priori upper bound s on the sparsity of β , *i.e.*, we know that $|J(\beta)| \leq s$ for some integer s . We call this a *sparsity certificate*. We introduce

$$\mathcal{B}_s = \mathcal{I}dent \bigcap \{\beta : |J(\beta)| \leq s\}.$$

Note that $\mathcal{B}_K = \mathcal{I}dent$. Thus, considering confidence sets around \mathcal{B}_s allows to deal with both the case when we rely on sparsity certificates and the case when we do not.

3. THE *STIV* ESTIMATOR

The sample counterpart of the moment conditions (1.2) can be written in the form

$$(3.1) \quad \frac{1}{n} \mathbf{Z}^T (\mathbf{Y} - \mathbf{X}\beta) = 0.$$

This is a system of L equations with K unknown parameters. If $L > K$, it is overdetermined. In general $\text{rank}(\mathbf{Z}^T \mathbf{X}) \leq \min(K, L, n)$, thus when $L \leq K$ or when $n < K$ the matrix does not have full column rank. Furthermore, replacing the population equations (1.2) by (3.1) induces a huge error when L, K or both are larger than n . So, looking for the exact solution of (3.1) in the high-dimensional setting makes no sense. However, we can stabilize the problem by restricting our attention to a suitable “small” candidate set of vectors β , for example, to those satisfying the constraint

$$(3.2) \quad \left| \frac{1}{n} \mathbf{Z}^T (\mathbf{Y} - \mathbf{X}\beta) \right|_{\infty} \leq \tau,$$

where $\tau > 0$ is chosen such that (3.2) holds for β in \mathcal{B}_s with high probability. We can then refine the search of the estimator in this “small” random set of vectors β by minimizing an appropriate criterion. It is possible to consider different small sets in (3.2), however the use of the sup-norm makes the inference robust in the presence of weak instruments. This will be clarified later.

In what follows, we use this idea with suitable modifications. First, notice that it makes sense to normalize the matrix \mathbf{Z} . This is quite intuitive because, otherwise, the larger the instrumental variable, the more influential it is on the estimation of the vector of coefficients. For technical reasons, we choose normalization where we multiply \mathbf{Z} by $\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})}$. The constraint (3.2) is modified as follows:

$$(3.3) \quad \left| \frac{1}{n} \mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})} \mathbf{Z}^T (\mathbf{Y} - \mathbf{X}\beta) \right|_{\infty} \leq \tau.$$

Along with the constraint of the form (3.3), we include more constraints to account for the unknown (average in the case of heteroscedasticity) level σ of the “effective noise” $z_{il}u_i$. Specifically,

we say that a pair $(\beta, \sigma) \in \mathbb{R}^K \times \mathbb{R}^+$ satisfies the *IV-constraint* if it belongs to the set

$$(3.4) \quad \widehat{\mathcal{I}}^{(I)} \triangleq \left\{ (\beta, \sigma) : \beta \in \mathbb{R}^K, \sigma > 0, \left| \frac{1}{n} \mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})} \mathbf{Z}^T (\mathbf{Y} - \mathbf{X}\beta) \right|_{\infty} \leq \sigma r, (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll}^2 \widehat{Q}_l(\beta) \leq \sigma^2, \forall l \in I \right\}$$

for some $r > 0$ (specified below), and

$$\widehat{Q}_l(\beta) \triangleq \frac{1}{n} \sum_{i=1}^n z_{li}^2 (y_i - x_i^T \beta)^2.$$

Note that the instrument $z_{li} = 1$ for all $i = 1, \dots, n$ belongs to I and the corresponding value of $\widehat{Q}_l(\beta)$ is

$$\widehat{Q}_l(\beta) = \frac{1}{n} \sum_{i=1}^n (y_i - x_i^T \beta)^2.$$

Definition 3.1. We call the *STIV estimator* any solution $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ of the following minimization problem:

$$(3.5) \quad \min_{(\beta, \sigma) \in \widehat{\mathcal{I}}^{(I)}} (|\mathbf{D}_{\mathbf{X}}^{-1} \beta|_1 + c\sigma),$$

where $0 < c < 1$.

In this formulation the instruments show up both in the *IV-constraint* and in the penalization through σ . As discussed above, the *IV-constraint* includes the constraints coming from the moment conditions (1.2) accounting for instrument exogeneity. We use $\widehat{\beta}^{(c,I)}$ as an estimator of β in \mathcal{B}_g . Finding the *STIV estimator* is a conic program; it can be efficiently solved, see Section 10.1. Note that the *STIV estimator* is not necessarily unique. Minimizing the ℓ_1 criterion $|\mathbf{D}_{\mathbf{X}}^{-1} \beta|_1$ is a convex relaxation of minimizing the ℓ_0 norm, *i.e.*, the number of non-zero coordinates of β . This usually ensures that the resulting solution is sparse. The term $c\sigma$ is included in the criterion to prevent from choosing σ arbitrarily large; indeed, the *IV-constraint* does not prevent from this. The matrix $\mathbf{D}_{\mathbf{X}}^{-1}$ arises from re-scaling of \mathbf{X} . It is natural to normalize the regressors by their size in a procedure that can do variable selection. This way changing units does not change which coefficients are found to be zeros or non-zeros. For the particular case where $\mathbf{Z} = \mathbf{X}$ and $\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})} = \mathbf{D}_{\mathbf{X}}$ is the diagonal matrix with entries $(\mathbb{E}_n[X_k^2]^{-1/2})_{k=1}^K$, the *STIV estimator* provides an extension of the Dantzig selector to the setting with unknown variance of the noise.

In this particular case, the *STIV estimator* can also be related to the Square-root Lasso of Belloni, Chernozhukov and Wang (2010), which solves the problem of unknown variance in high-dimensional regression with deterministic regressors and i.i.d. errors. The definition of *STIV estimator*

contains the additional constraint (3.3), which is not present in the conic program for the Square-root Lasso. This is due to the fact that we have to handle the endogeneity.

4. SUMMARY OF THE MAIN RESULTS

The main result of this paper are nested confidence sets around the regions $(\mathcal{B}_s)_{s \leq K}$. We allow for set-identification, arbitrarily weak instruments, for a situation where $K > L$ (for example when all the instruments can have a direct effect on the outcome or for various combinations of discrete regressors and instruments). The dimension K could be as large as we want, even larger than an exponential in the sample size, in that case the model (1.1) together with (1.2) becomes set-identified. Also, in finite samples and when L and/or K are much larger than n , we can obtain finite volume confidence sets for low values of s and infinite volume confidence sets for larger values of s , even in point-identified settings. These results are presented in Section 6. We first present various assumptions that one can make on the data generating process. These sometimes limit the number of instruments. We give the corresponding way to adjust the constant r in the definition of the set $\widehat{\mathcal{I}}^{(I)}$ appearing in the constrained optimization for the *STIV* estimator.

In all but the first scenario on the data generating process, our confidence sets are of the form: For every β in \mathcal{B}_s , with probability at least $1 - \alpha$, for any c in $(0, 1)$ and any set of indices I containing the index of the instrument which is unity, any solution $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ of the minimization problem (3.5) satisfies

$$\left| \left(\mathbf{D}_{\mathbf{X}}^{-1} (\widehat{\beta}^{(c,I)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\widehat{\sigma}^{(c,I)} r}{\overline{\kappa}_{p, J_0}^{(c,I)}(s)} \left(1 - \frac{r}{\kappa_1^{(c,I)}(s)} \right)_+^{-1} \quad \forall p \in [1, \infty], \quad \forall J_0 \subset \{1, \dots, K\},$$

where the set J_0 is specified by the econometrician, depending on which subset of the regressor he is interested in, and for all $k = 1, \dots, K$,

$$|\widehat{\beta}_k^{(c,I)} - \beta_k| \leq \frac{2\widehat{\sigma}^{(c,I)} r}{\mathbb{E}_n[X_k^2]^{1/2} \kappa_k^{(c,I)*}(s)} \left(1 - \frac{r}{\kappa_1^{(c,I)}(s)} \right)_+^{-1}.$$

Under the first scenario on the data generating process the $1 - \alpha$ probability event depends on I so that the statement is not uniform in I but for fixed I . The constants $\overline{\kappa}_{p, J_0}^{(c,I)}(s)$, $\kappa_1^{(c,I)}(s)$ or $\kappa_k^{(c,I)*}(s)$ are easy to calculate, data-dependent, lower bounds on the sensitivities that we introduce in Section 5. The sensitivities provide a generalization of the restricted eigenvalues to non-symmetric and non-square matrices (see Section 10.1 in the appendix for a comparison). The most difficult constant to calculate is $\kappa_1^{(c,I)}(s)$, it requires to solve $2K^2$ linear programs. The confidence sets could be infinite when $\kappa_1^{(c,I)}(s) \leq r$. We will see that this occurs either when we have very many instruments or when

there exists a regressor for which all instruments are too weak instruments or s is too large. We will also discuss how the constants $\kappa_k^{(c,I)*}(s)$ and $\kappa_1^{(c,I)}(s)$ compare to the usual concentration parameter defined for low-dimensional structural equations with homoscedastic Gaussian errors. Recall that we relax both heteroscedasticity and Gaussian errors. When we are only interested in a subset of the coefficients, our confidence sets are smaller than those that one would obtain by projection of joint confidence sets. They account for the a priori upper bound on the sparsity s . The confidence sets do not require that the non-zero coefficients are large enough but require to make a stand and give a prior upper bound s . As we will see in Section 7, the *STIV* estimator usually estimates too many non-zero coefficients when the underlying vector is sparse so that $|J(\widehat{\beta}^{(c,I)})|$ can be considered as a conservative upper bound on the true sparsity. For increasing values of s one obtain nested confidence sets. These become infinite when s approaches K and K is larger than n . If one is interested in a few coefficients, it is possible to choose the value of c in $(0, 1)$ that yields the smaller confidence sets. This is easily obtained by taking grid values for c . Similarly, for all but the first scenario on the data generating process, it is possible to try several sets I and choose according to the size of the confidence sets around the vector of the coefficients of interest.

In Section 6, we present a sparse oracle inequality in the case (ii) above where the underlying model is not sparse. It shows that the *STIV* estimator estimates the coefficients as well as a method that would know in advance which is the best sparse approximation in terms of bias/variance trade-off.

In Section 7, we present rates of convergence and model selection results. Indeed, the right hand-side in the above two upper bounds are random and these inequalities do not tell, for example, how well is the vector of coefficient estimated depending on L and n . We also obtain that, if the absolute values of the coefficients $|\beta_k|$ are large enough on the support of β , then with probability close to one $J(\beta) \subseteq J(\widehat{\beta}^{(c,I)})$. This is used to obtain a second type of confidence sets where we plug-in an estimator of $J(\beta)$ in the definition of the sensitivities to obtain a lower bound, and thus, a proper data-driven upper bound on the estimation error. Recall that the first type of confidence sets uses sparsity certificates and does not rely on such a separation from zero of the non-zero coefficients assumption. The fact that, with probability close to one $J(\beta) \subseteq J(\widehat{\beta}^{(c,I)})$ is confirmed in the simulation study where the *STIV* estimator usually selects too many regressors. This result allows to recover exactly, with probability close to 1, the true support of the vector of coefficients as well as the sign of the coefficients by a thresholding rule.

Section 8 discusses some special cases and extensions. Because the *STIV* estimator is a one stage method that is robust to weak-instruments and allows for very many instruments, we present

a variation on the *STIV* estimator to estimate low-dimensional ($K < n$) and non sparse structural models. This is a new method to construct confidence sets that is robust to weak instruments, heteroscedasticity and non-Gaussian errors, and allows for very-many-instruments. We also present the refined properties of our estimator when none of the regressors and instruments have heavy tail distribution. This simplifies its calculation and gives finer results, especially in the very many or weak-instruments case. Finally, we present the properties of a two-stage *STIV* procedure with estimated sparse linear projection type instruments, akin to two-stage least squares. Here the two stages are high-dimensional regressions and the second stage has endogenous regressors. This is related to the literature on selection of instruments and optimal instruments. Here we do not touch on optimality because it is an open question to define optimality with a high-dimensional structural equation with endogenous regressors. This method only works if the endogenous regressors have a sparse reduced form. In the case of an approximately sparse reduced form (case (ii) above) we would obtain rates of convergence but our method do not yield confidence sets.

In Section 9, we consider the following model

$$\begin{aligned} y_i &= x_i^T \beta + u_i, \\ \mathbb{E}[z_i u_i] &= 0, \\ \mathbb{E}[\bar{z}_i u_i] &= \theta. \end{aligned}$$

Here, z_{li} for $i = 1, \dots, n$ and $l = 1, \dots, L$, is a set of instruments that are known in advance to be valid (exogenous) and \bar{z}_i for $i = 1, \dots, n$ and $l = 1, \dots, L$, is a second set of instruments. One wants to decide which instrument from this second list is valid and which is invalid (endogenous). In this setup, we again allow for the dimensions of x_i , z_i and \bar{z}_i to be much larger than n .

Section 10 discusses way to implement our algorithm and to calculate the constants that drive the size of the confidence sets. This is based either on conic or linear programs. These are readily available in many classical softwares. In this section we also present a simulation study. All the proofs are given in the appendix.

5. SENSITIVITY CHARACTERISTICS

In the usual linear regression in low dimension, when $\mathbf{Z} = \mathbf{X}$ and the Gram matrix $\mathbf{X}^T \mathbf{X}/n$ is positive definite, the sensitivity is given by the minimal eigenvalue of this matrix. In high-dimensional regression, the theory of the Lasso and the Dantzig selector comes up with a more sophisticated sensitivity analysis; there the Gram matrix cannot be positive definite and the eigenvalue conditions

are imposed on its sufficiently small submatrices. This is typically expressed via the restricted isometry property of Candès and Tao (2007) or the more general restricted eigenvalue condition of Bickel, Ritov and Tsybakov (2009). In our structural model with endogenous regressors, these sensitivity characteristics cannot be used, since instead of a symmetric Gram matrix we have a rectangular matrix $\mathbf{Z}^T \mathbf{X}/n$ involving the instruments. More precisely, we will deal with its normalized version

$$\Psi_n \triangleq \frac{1}{n} \mathbf{D}_Z^{(I)} \mathbf{Z}^T \mathbf{X} \mathbf{D}_X.$$

In general, $\Psi_n^{(I)}$ is not a square matrix. For $L = K$, it is a square matrix but, in the presence of at least one endogenous regressor, $\Psi_n^{(I)}$ is not symmetric.

We now introduce some scalar sensitivity characteristics related to the action of the matrix $\Psi_n^{(I)}$ on vectors in the cone

$$C_J^{(c)} \triangleq \left\{ \Delta \in \mathbb{R}^K : |\Delta_{J^c}|_1 \leq \frac{1+c}{1-c} |\Delta_J|_1 \right\},$$

where $0 < c < 1$ is the constant in the definition of the *STIV* estimator, J is any subset of $\{1, \dots, K\}$. When the cardinality of J is small, the vectors Δ in the cone $C_J^{(c)}$ have a substantial part of their mass concentrated on a set of small cardinality. We call $C_J^{(c)}$ the *cone of dominant coordinates*. The set J that will be used later is the set $J(\beta)$ which is small if β is sparse. The use of similar cones to define sensitivity characteristics is standard in the literature on the Lasso and the Dantzig selector (see, Bickel, Ritov and Tsybakov (2009)); the particular choice of the constant $\frac{1+c}{1-c}$ will become clear from the proofs. It follows from the definition of $C_J^{(c)}$ that

$$(5.1) \quad |\Delta|_1 \leq \frac{2}{1-c} |\Delta_J|_1 \leq \frac{2}{1-c} |J|^{1-1/p} |\Delta_J|_p, \quad \forall \Delta \in C_J^{(c)}, \quad 1 \leq p \leq \infty.$$

For $p \in [1, \infty]$, we define the ℓ_p *sensitivity* as the following random variable:

$$\kappa_{p,J}^{(c,I)} \triangleq \inf_{\Delta \in C_J^{(c)}: |\Delta|_p=1} \left| \Psi_n^{(I)} \Delta \right|_\infty.$$

Similar, but different, quantities named cone invertibility factors have been introduced in Ye and Zhang (2010).

Given a subset $J_0 \subset \{1, \dots, K\}$ and $p \in [1, \infty]$, we define the l_p - J_0 -*block sensitivity* as

$$(5.2) \quad \kappa_{p,J_0,J}^{(c,I)} \triangleq \inf_{\Delta \in C_J^{(c)}: |\Delta_{J_0}|_p=1} \left| \Psi_n^{(I)} \Delta \right|_\infty.$$

By convention, we set $\kappa_{p,\emptyset,J(\beta^*)}^{(c,I)} = \infty$. We use the notation $\kappa_{k,J}^{(c,I)*}$ for *coordinate-wise sensitivities*, *i.e.*, for block sensitivities when $J_0 = \{k\}$ is a singleton⁸:

$$(5.3) \quad \kappa_{k,J}^{(c,I)*} \triangleq \inf_{\Delta \in C_J^{(c)}: \Delta_k=1} \left| \Psi_n^{(I)} \Delta \right|_{\infty}.$$

Note that here we restrict the minimization to vectors Δ with positive k th coordinate, $\Delta_k = 1$, since replacing Δ by $-\Delta$ yields the same value of $|\Psi_n^{(I)} \Delta|_{\infty}$.

The heuristic behind the sensitivity characteristics is the following. As we will see in the Appendix, for a fixed value of β in \mathcal{B}_s , we adjust r in the definition of $\widehat{\mathcal{I}}^{(I)}$ such that $\left(\beta, \max_{l \in I} (\mathbf{D}_Z^{(I)})_{ll} \sqrt{\widehat{Q}_l(\beta)} \right)$ belongs to $\widehat{\mathcal{I}}^{(I)}$ on an event E_{α} of probability $1 - \alpha$. This yields that for a proper τ , on E_{α} ,

$$(5.4) \quad |\Psi_n^{(I)} \Delta|_{\infty} \leq \tau$$

where $\Delta = \mathbf{D}_X^{-1}(\widehat{\beta}^{(c,I)} - \beta)$. When τ can be calculated directly from the data, this yields a confidence region for β in \mathcal{B}_s . Moreover, because we minimize the objective function (3.5), we will see that, on the same event E_{α} , Δ is constrained to belong to the subset $C_{J(\beta)}^{(c,I)}$ of \mathbb{R}^K . The sensitivities allow to deduce from (5.4) and the cone condition, confidence regions for various losses. Suppose, for example, that one is interested in a confidence region for the subvector of β in \mathcal{B}_s , which corresponds to the coefficients of indices in $J_0 \subset \{1, \dots, K\}$, and considers the l_p -loss, then one gets

$$|\Delta_{J_0}|_p \leq \frac{\tau}{\kappa_{p,J_0,J(\beta)}^{(c,I)}}.$$

Indeed, this is trivial when $|\Delta_{J_0}|_p = 0$ and, when it is not, it immediately follows from

$$\frac{|\Psi_n^{(I)} \Delta|_{\infty}}{|\Delta_{J_0}|_p} \geq \inf_{\tilde{\Delta}: \tilde{\Delta} \neq 0, \tilde{\Delta} \in C_{J(\beta)}^{(c,I)}} \frac{|\Psi_n^{(I)} \tilde{\Delta}|_{\infty}}{|\tilde{\Delta}_{J_0}|_p}.$$

Remark that, working with the coordinate-wise sensitivities $\kappa_{k,J(\beta)}^{(c,I)*}$ is much better than projecting the region (5.4) onto the axes because it also takes into account the cone condition and, thus, the sparsity of the underlying β . We see that the sensitivities are quantities that are intrinsic to the estimation properties of the *STIV* estimator⁹. We show in Section 11.1 that the assumption that the sensitivities $\kappa_{p,J}^{(c,I)}$ are positive is weaker and more flexible than the restricted eigenvalue (RE) assumption of Bickel, Ritov and Tsybakov (2009). Unlike the RE assumption, it is applicable to non-square non-symmetric matrices. Another nice feature of the sensitivities that we introduce is that lower bounds on these

⁸They coincide for all values of $p \in [1, \infty]$.

⁹They are also intrinsic to the Dantzig selector and the Lasso.

sensitivities can be efficiently calculated (see below), this opens the path to confidence statements in high-dimensional regression.

The coordinate-wise sensitivities are measures of the strength of the instruments. The coordinate-wise sensitivity for index k could be interpreted as *restricted maximal partial empirical correlation between the instruments and the regressor* $(x_{ki})_{i=1}^n$. Indeed, (5.3) can be written as

$$(5.5) \quad \kappa_{k,J}^{(c,I)*} = \inf_{\lambda \in \mathbb{R}^{K-1}: \sum_{l \in J^c} |\mathbb{E}_n[X_l^2]^{1/2} \lambda_l| \leq \frac{1+c}{1-c} (1 + \sum_{l \in J \setminus \{k\}} |\mathbb{E}_n[X_l^2]^{1/2} \lambda_l|)} \max_{l=1, \dots, L} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \left| \frac{1}{n} \sum_{i=1}^n z_{li} (x_{ki} - x_{\{k\}^c i}^T \lambda) \right|.$$

It is easy to check that (5.5) yields that when $L < |J|$, $\kappa_{k,J}^{(c,I)*} = 0$. For confidence sets or rates of estimation statements, the set J will correspond to $J(\beta)$ for β in \mathcal{B}_s so that the restriction in the infimum corresponds to vectors having most of their mass on the support of the underlying vector of coefficients β from the set \mathcal{B}_s (which includes the constant). When an exogenous variable serves as its own instrument, the coordinate-wise sensitivity for that regressor should be large. This is because there exists, among the list of instruments, $(x_{ki})_{i=1}^n$ themselves. Otherwise, the coordinate-wise sensitivity of a regressor is small if all instruments have small restricted partial empirical correlation with that regressor. Because of the maximum, one good instrument is enough to have a large coordinate-wise sensitivity. It is small only if all instruments are weak.

Example 7. For comparison, consider a structural equation with only one endogenous regressor $(x_{k_{\text{end}}i})_{i=1}^n$. Assume that $(z_{li})_{i=1}^n$ for $l = 1, \dots, L$ are fixed and write the reduced form equation

$$(5.6) \quad x_{k_{\text{end}}i} = \sum_{l \in J^c} \tilde{z}_{li} \zeta_l + \sum_{l \in J} \tilde{z}_{li} \zeta_l + v_i, \quad i = 1, \dots, n,$$

where $(\zeta_l)_{l=1}^L$ are unknown coefficients, J is the set of indices of the exogenous regressors that have non-zero coefficients in (1.1) (this is possible when $|J| \leq n$), $\tilde{z}_{li} = z_{li}$ for $l \in J$ and $i = 1, \dots, n$, while $(\tilde{z}_{li})_{i=1}^n$ are residuals from the regression of the original instruments $(z_{li})_{i=1}^n$ for $l \in J^c$ on $(z_{li})_{i=1}^n$ for $l \in J$. Assume finally that (u_i, v_i) are i.i.d. and have a mean zero bivariate normal distribution. Denote by σ_v^2 the variance of v_i . It is easy to check that λ in \mathbb{R}^{K-1} such that $x_{\{k_{\text{end}}\}^c i}^T \lambda = \sum_{l \in J} \tilde{z}_{li} \zeta_l$ satisfies the constraint in (5.5) which yields using (5.6)

$$(5.7) \quad \kappa_{k_{\text{end}},J}^{(c,I)*} \geq \max \left(\max_{l \in J} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \left| \frac{1}{n} \sum_{i=1}^n z_{li} v_i \right|, \max_{l \in J^c} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \left| \frac{1}{n} \sum_{i=1}^n z_{li} v_i + \frac{1}{n} e_l^T \tilde{\mathbf{Z}}_J^T \tilde{\mathbf{Z}}_{J^c} \zeta_{J^c} \right| \right)$$

where $(e_l)_{l=1}^L$ is the canonical basis of \mathbb{R}^L . Note that $|\frac{1}{n} \sum_{i=1}^n z_{li} v_i|$ is small for large n because the instruments are fixed and v_i is mean zero. This shares similarity with the *concentration parameter*

(see, *e.g.*, Andrews and Stock (2007)), defined as

$$\mu^2 = \frac{\zeta_{J^c}^T \tilde{\mathbf{Z}}_J^T \tilde{\mathbf{Z}}_J \zeta_{J^c}}{\sigma_v^2},$$

which is a classical measure of the strength of the set of instruments for low dimensional structural equations, under homoscedasticity. The quantity in (5.7) isolates the strength of each instrument. Thus we see that, unlike the concentration parameter which considers the set of all instruments, the coordinate-wise sensitivities depend on the strength of the best instrument. Due to our rescaling of the instruments, which allows to relax a lot the usual distributional assumptions and to handle heteroscedasticity, the coordinate-wise sensitivities and concentration parameter are not comparable in terms of the dependence on σ_v^2 .

A first reason to come up with lower bounds on the sensitivities is that they depend on $J(\beta)$ and thus on the unknown β . A second reason is to obtain an easy to calculate lower bound. For every sensitivity, we propose an easy to calculate lower bound. The following proposition will be useful.

Proposition 5.1. (i) Let J, \hat{J} be two subsets of $\{1, \dots, K\}$ such that $J \subseteq \hat{J}$. Then, for all $J_0 \subset$

$\{1, \dots, K\}$, all $p \in [1, \infty]$, all c in $(0, 1)$ and all set of indices I containing the index of unity,

$$\kappa_{p, J_0, J}^{(c, I)} \geq \kappa_{p, J_0, \hat{J}}^{(c, I)}$$

(ii) For all $J_0 \subset \{1, \dots, K\}$ all $p \in [1, \infty]$, all c in $(0, 1)$ and all set of indices I containing the index of unity, $\kappa_{p, J_0, J}^{(c, I)} \geq \kappa_{p, J}^{(c, I)}$.

(iii) For all $p \in [1, \infty]$, all c in $(0, 1)$ and all set of indices I containing the index of unity,

$$(5.8) \quad \left(\frac{2|J|}{1-c} \right)^{-1/p} \kappa_{\infty, J}^{(c, I)} \leq \kappa_{p, J}^{(c, I)} \leq \frac{2}{1-c} |J|^{1-1/p} \kappa_{1, J}^{(c, I)},$$

and for all $J_0 \subset \{1, \dots, K\}$, all $p \in [1, \infty]$, all c in $(0, 1)$ and all set of indices I containing the index of unity,

$$(5.9) \quad |J_0|^{-1/p} \kappa_{\infty, J_0, J}^{(c, I)} \leq \kappa_{p, J_0, J}^{(c, I)} \leq |J_0|^{1-1/p} \kappa_{1, J_0, J}^{(c, I)};$$

(iv) For all $J_0 \subset \{1, \dots, K\}$, $\kappa_{\infty, J_0, J}^{(c, I)} = \min_{k \in J_0} \kappa_{k, J}^{(c, I)*}$.

Results for the l_p -sensitivities are easily deduced from (i) and (iv) by taking $J_0 = \{1, \dots, K\}$. (ii) and (5.9) allow to minorize the bloc sensitivities when J_0 can be large and a direct calculation would be too difficult from a numerical point of view.

We can control $\kappa_{p, J_0, J(\beta)}^{(c, I)}$ without knowing $J(\beta)$ by means of *sparsity certificate*. Assume that we have an upper bound s on the sparsity of β , *i.e.*, we know that $|J(\beta)| \leq s$ for some integer s and that we use it as well to define \mathcal{B}_s . This does not require that non-zero coefficients are large enough

but just that a maximum of s are non-zero. In view of (5.1), if $|J| \leq s$, then for any Δ in the cone $C_J^{(c)}$ we have $|\Delta|_1 \leq \frac{2s}{1-c} |\Delta|_\infty$. Thus, for all J such that $|J| \leq s$, we can bound the coordinate-wise sensitivities as follows:

$$(5.10) \quad \begin{aligned} \kappa_{k,J}^{(c,I)*} &\geq \inf_{\Delta_k=1, |\Delta|_1 \leq a |\Delta|_\infty} \left| \Psi_n^{(I)} \Delta \right|_\infty \\ &\geq \min_{j=1, \dots, K} \left\{ \min_{\Delta_k=1, |\Delta|_1 \leq a |\Delta_j|} \left| \Psi_n^{(I)} \Delta \right|_\infty \right\} \triangleq \kappa_k^{(c,I)*}(s), \end{aligned}$$

where $a = \frac{2s}{1-c}$. For given s , this bound is data-driven since the minimum in curly brackets can be computed by solving $2K$ linear programs (see Section 10.1). Then, using (iv), we can deduce a lower bound on $\kappa_{\infty, J_0, J}^{(c,I)}$

$$(5.11) \quad \kappa_{\infty, J_0, J}^{(c,I)} \geq \min_{k \in J_0} \kappa_k^{(c,I)*}(s).$$

Using (5.8) and (5.11) we get computable lower bounds for all $\kappa_{p,J}^{(c,I)}$, $p \in [1, \infty]$, which depend only on s and on the data. In particular, for $|J| \leq s$,

$$(5.12) \quad \kappa_{1,J}^{(c,I)} \geq \frac{1-c}{2s} \min_{k=1, \dots, K} \kappa_k^{(c,I)*}(s) \triangleq \kappa_1^{(c,I)}(s).$$

This can thus be obtained by solving $2K^2$ linear programs. Analogously to (5.10), the sparsity certificate approach yields a bound for block sensitivities

$$(5.13) \quad \begin{aligned} \kappa_{1, J_0, J}^{(c,I)} &\geq \inf_{|\Delta_{J_0}|_1=1, |\Delta|_1 \leq a |\Delta|_\infty} \left| \Psi_n^{(I)} \Delta \right|_\infty \\ &\geq \min_{j=1, \dots, K} \left\{ \min_{|\Delta_{J_0}|_1=1, |\Delta|_1 \leq a |\Delta_j|} \left| \Psi_n^{(I)} \Delta \right|_\infty \right\} \triangleq \kappa_{1, J_0}^{(c,I)}(s). \end{aligned}$$

In Section 10.1 we show that the expression in curly brackets in (5.13) can be computed by solving $2^{|J_0|}$ linear programs. Thus, the values $\kappa_{1, J_0}^{(c,I)}(s)$ can be readily obtained for sets J_0 of small cardinality. Otherwise we do not advertise this lower bound but simply to use the following lower bound for $p = 1$. For $|J| \leq s$, (ii), (5.8) and (5.11) yield

$$(5.14) \quad \kappa_{p, J_0, J}^{(c,I)} \geq \max \left(|J_0|^{-1/p} \min_{k \in J_0} \kappa_k^{(c,I)*}(s), \left(\frac{1-c}{2s} \min_{k=1, \dots, K} \kappa_k^{(c,I)*}(s) \right) \right).$$

This bound can be calculated by solving $2K^2$ linear programs even for large sets J_0 .

For the sake of completeness, we present an alternative to the sparsity certificate approach. It corresponds to computing $\kappa_{1,J}^{(c,I)}$ and $\kappa_{k,J}^{(c,I)*}$ directly. This is numerically feasible only for J of small cardinality. Indeed, we show in Section 10.1 that obtaining the coordinate-wise sensitivities corresponds to solving $2^{|J|}$ linear programs. Using (5.8) and (5.11), we obtain computable lower

bounds for all $\kappa_{p,J}^{(c,I)}$, $p \in [1, \infty]$. The lower bounds are valid for any given index set J . However, we will need to compute the characteristics for the inaccessible set $J = J(\beta)$, where β is the unknown parameter from \mathcal{B}_s on which we are making inference. To circumvent this problem, we can plug in an estimator \hat{J} of $J(\beta)$. For example, we can take $\hat{J} = J(\hat{\beta}^{(c,I)})$. The confidence bounds remain valid whenever $J(\beta) \subseteq \hat{J}$, since then $\kappa_{p,J(\beta)}^{(c,I)} \geq \kappa_{p,\hat{J}}^{(c,I)}$, by Proposition 5.1 (i). Theoretical guarantees for the inclusion $J(\beta) \subseteq J(\hat{\beta}^{(c,I)})$ to hold with probability close to 1 require a separation from zero: $|\beta_k|$ is not too small on the support of β (see Theorem 7.4 (iv)). On the other hand, one typically observes in simulations that the relevant set $J(\beta)$ is either estimated exactly or overestimated by its empirical counterpart $\hat{J} = J(\hat{\beta}^{(c,I)})$, so that the required inclusion is satisfied for such a simple choice of \hat{J} . Belloni and Chernozhukov (2010) study the property of post model selection least squares estimation in the case of prediction loss, they obtain theoretical bounds on the error made when we do not have $J(\beta) \subset J(\hat{\beta}^{(c,I)})$. We do not touch upon the very difficult question of obtaining confidence sets with coverage at most $1 - \alpha$ that allow to deal with the possibility that $J(\beta) \subset J(\hat{\beta}^{(c,I)})$ is not satisfied. We state that this type of confidence sets has coverage $1 - \gamma$ where $\gamma > \alpha$. They are approximate $1 - \alpha$ confidence sets. In our simulation study we obtain that the confidence regions obtained via the sparsity certificate using for $s = |J(\hat{\beta}^{(c,I)})|$ are almost identical to the second ones which also require the separation from zero. Because the first method does not require such an assumption and is feasible (non combinatoric) even when $|J(\beta)|$ is large, we strongly advertise the first approach based on the sparsity certificate and to use a conservative sparsity certificate s . We also suggest to draw nested confidence sets varying the degree of sparsity that we assume.

In the next proposition, we present a simple lower bound on $\kappa_{p,J}^{(c,I)}$ for general $L \times K$ rectangular matrices $\Psi_n^{(I)}$. Its proof, as well as other lower bounds on $\kappa_{1,J}^{(c,I)}$, can be found in Section 11. It is important to note that adding rows to matrix $\Psi_n^{(I)}$ (*i.e.*, adding instruments) increases the sup-norm $|\Psi_n^{(I)} \Delta|_\infty$, and thus potentially increases the sensitivities $\kappa_{p,J(\beta)}^{(c,I)}$ and their computable lower bounds. This has a positive effect since the inverse of the computable lower bounds of the sensitivities drive the width of the confidence set for β in \mathcal{B}_s , see Theorem 6.5. Thus, adding instruments potentially improves the confidence set, which is quite intuitive. On the other hand, the price for adding instruments in terms of the rate of convergence is only logarithmic in the number of instruments, as we will see in the next section.

Proposition 5.2. Fix $J \subseteq \{1, \dots, K\}$. Assume that there exist $\eta_1 > 0$ and $0 < \eta_2 < 1$ such that

$$(5.15) \quad \forall k \in J, \exists l(k) : \begin{cases} |(\Psi_n^{(I)})_{l(k)k}| \geq \frac{\eta_1}{1-c}, \\ \frac{\max_{k' \neq k} |(\Psi_n^{(I)})_{l(k)k'}|}{|(\Psi_n^{(I)})_{l(k)k}|} \leq \frac{(1-\eta_2)(1-c)}{2|J|}. \end{cases}$$

Then

$$\kappa_{p,J}^{(c,I)} \geq (2|J|)^{-1/p} (1-c)^{-1+1/p} \eta_1 \eta_2.$$

conditional on z_i for $i = 1, \dots, n$ can be obtained numerically, for example using a Monte-Carlo method, and r is adjusted accordingly. This is an approach proposed by Belloni and Chernozhukov (2010) for the linear regression model without endogeneity. This approach is worth mentioning because it has the advantage to yield a smaller r , and thus smaller confidence sets as we will see later. Indeed, it does not rely on a crude union bound but account for the correlation between the regressors.

Scenario 2. Under the assumption

Assumption 6.1. *For every $i = 1, \dots, n$ and $l = 1, \dots, L$, $z_{li}u_i$ are symmetric and neither of $z_{li}u_i$ is almost surely equal to 0.*

we choose

$$(6.1) \quad r = \sqrt{\frac{2 \log(L/2\alpha)}{n}}.$$

Symmetry is a very likely assumption if (1.1) is a first difference between two time periods in a linear panel data model.

Scenario 3. Under the assumption

Assumption 6.2. *For every $l = 1, \dots, L$, $z_{li}u_i$ are i.i.d. and symmetric, neither of $z_{li}u_i$ is almost surely equal to 0 and L is such that*

$$L < \frac{9\alpha}{4e^3 \Phi(-\sqrt{n})}.$$

we choose

$$(6.2) \quad r = -\frac{1}{\sqrt{n}} \Phi^{-1} \left(\frac{9\alpha}{4Le^3} \right).$$

This is a slightly tighter constant than the one given for Scenario 2 but Assumption (6.2) does not allow for heteroscedastic errors. The upper bound on the number of instruments is of the order of an exponential in n .

Scenario 4. We denote by $d_{n,\delta} = \min_{l=1,\dots,L} d_{n,\delta,l}$ where

$$d_{n,\delta,l} \triangleq \frac{(\sum_{i=1}^n \mathbb{E}[z_{li}^2 u_i^2])^{1/2}}{(\sum_{i=1}^n \mathbb{E}[|z_{li} u_i|^{2+\delta}])^{1/(2+\delta)}}$$

when the numerator and denominator are well defined. Under the assumption

Assumption 6.3. *There exists δ positive such that, for all $i = 1, \dots, n$, $l = 1, \dots, L$, $\mathbb{E}[|z_{li} u_i|^{2+\delta}] < \infty$, neither of $z_{li} u_i$ is almost surely equal to 0 and L is such that*

$$L \leq \frac{\alpha}{2\Phi(-d_{n,\delta}) \left(1 + A_0 \left(1 + d_{n,\delta}^{-1} \right)^{2+\delta} \right)}$$

where $A_0 > 0$ is the absolute constant of Theorem 11.6 in Section 11.2.

we choose

$$(6.3) \quad r = -\frac{1}{\sqrt{n}} \Phi^{-1} \left(\frac{\alpha}{2L} \right).$$

Note that if, for any fixed l in $\{1, \dots, L\}$, the variables $z_{li}u_i$ are i.i.d., then for $l = 1, \dots, L$, $d_{n,\delta,l} = n^{\frac{\delta}{4+2\delta}} \frac{(\mathbb{E}[z_{l1}^2 u_1^2])^{1/2}}{(\mathbb{E}[|z_{l1} u_1|^{2+\delta}])^{1/(2+\delta)}}$ which tends to infinity with n . With this choice of r the confidence sets will only be asymptotically valid¹⁰, in the asymptotic where L and n increase in such a way that $\Phi^{-1} \left(\frac{\alpha}{2L} \right) d_{n,\delta}^{-1} \rightarrow 0$. This scenario allows for heteroscedasticity and non-symmetric errors. This is very desirable in an *IV* setup.

Scenario 5. We denote by $\gamma_4 \triangleq \max_{l=1,\dots,L} \gamma_{4l}$ where $\gamma_{4l} = \mathbb{E}[(z_{li}u_i)^4]/(\mathbb{E}[(z_{li}u_i)^2])^2$. Under the assumption

Assumption 6.4. For every $l = 1, \dots, L$, $z_{li}u_i$ are i.i.d., $\mathbb{E}[(z_{li}u_i)^4] < \infty$ and $z_{li}u_i$ is not 0 almost surely, for some c_4 positive $\gamma_4 \leq c_4$ and

$$(6.4) \quad L < \frac{\alpha}{2e+1} \exp \left(\frac{n}{c_4} \right).$$

we choose

$$(6.5) \quad r = \sqrt{\frac{2 \log(L(2e+1)/\alpha)}{n - c_4 \log(L(2e+1)/\alpha)}}.$$

It is reasonable to assume that $n - c_4 \log(L(2e+1)/\alpha) \geq n/2$ as soon as n is relatively large relative compared to $\log(L/\alpha)$. In that case, we can take

$$(6.6) \quad r = 2\sqrt{\frac{\log(L(2e+1)/\alpha)}{n}}$$

and have confidence sets with valid finite sample properties. It is also possible to proceed in two stages. We first start by choosing r as in (6.5) with a very rough upper bound on γ_4 , for example using (6.6). As we will see, it yields a point estimate and a confidence region for β , which in turns yields a point estimate and a confidence set for γ_4 . In a second stage, it is possible to plug-in either the point estimate for γ_4 or the upper bond from the confidence set for γ_4 . This two-stage approach yields approximately valid confidence sets. A similar approach has been used in the simulations in

¹⁰We only consider asymptotically valid confidence sets because the moderate deviations result for the self-normalized sums that we use depends on the constant A_0 which is universal but not explicit.

Bertail, Gauth rat and Harari-Kermadec (2005) to obtain finite sample confidence sets for inference based on empirical φ^* -discrepancies and quasi-empirical likelihood methods¹¹.

Scenarios 2-5 rely on moderate deviations for self-normalized sums that are recalled in Section 11.2 and are respectively from Efron (1969), Pinelis (1994), Jing, Shao and Wang (2003) and Bertail, Gauth rat and Harari-Kermadec (2009)¹². Each moderate deviations result relies on a different set of assumptions and our confidence sets with finite sample validity will rely on a restricted class of distributions for the data generating process. This is related to the Bahadur and Savage (1956) impossibility result, see also Romano and Wolf (2000)¹³.

6.2. Confidence Sets.

Theorem 6.5. *For every β in \mathcal{B}_s , under one of the scenarios 2-5 of Section 6.1, together with its respective choice of r , with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5), for any c in $(0, 1)$ and any set of indices I containing the index of the instrument which is unity, for any solution $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ of the minimization problem (3.5) we have*

$$(6.7) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1} (\widehat{\beta}^{(c,I)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\widehat{\sigma}^{(c,I)} r}{\kappa_{p, J_0, J(\beta)}^{(c,I)}} \left(1 - \frac{r}{\kappa_{1, J(\beta)}^{(c,I)}} \right)_+^{-1} \quad \forall p \in [1, \infty], \quad \forall J_0 \subset \{1, \dots, K\},$$

$$(6.8) \quad |\widehat{\beta}_k^{(c,I)} - \beta_k| \leq \frac{2\widehat{\sigma}^{(c,I)} r}{\mathbb{E}_n[X_k^2]^{1/2} \kappa_{k, J(\beta)}^{(c,I)*}} \left(1 - \frac{r}{\kappa_{1, J(\beta)}^{(c,I)}} \right)_+^{-1} \quad \forall k = 1, \dots, K,$$

and

$$(6.9) \quad \widehat{\sigma}^{(c,I)} \leq \max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(l)})_{ll} \sqrt{\widehat{Q}_l(\beta)} \left(1 + \frac{r}{c\kappa_{1, J(\beta), J(\beta)}^{(c,I)}} \right) \left(1 - \frac{r}{c\kappa_{1, J(\beta), J(\beta)}^{(c,I)}} \right)_+^{-1}.$$

¹¹Self-normalized sums also appear naturally in empirical likelihood contexts.

¹²Bertail, Gauth rat and Harari-Kermadec (2009) provides an upper bound of tail probabilities of self-normalized sums with an explicit constant unlike Jing, Shao and Wang (2003). The result of Jing, Shao and Wang (2003) is mostly useful to study large deviations, that is why under Scenario 4 we only consider asymptotically valid confidence sets.

¹³As explained in Romano and Wolf (2000), the Bahadur and Savage result implies that we need to make some restriction on the set of distributions \mathcal{P} on the line in order to construct conservative confidence intervals for the mean of \mathbb{P} in \mathcal{P} that are bounded. A conservative interval of coverage level $1 - \alpha$ is such that the parameter is contained in the interval with probability at least $1 - \alpha$ for all \mathbb{P} in \mathcal{P} and sample size n .

Under scenario 1 of Section 6.1 with the corresponding choice of r , for any set of indices I containing the index of the instrument which is unity, for every β in \mathcal{B}_s , with probability at least $1 - \alpha$, for any c in $(0, 1)$, any solution $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ of the minimization problem (3.5) satisfies (6.7), (6.8) and (6.9).

Consider the most likely case where $L \geq K$ and the parameter β is point identified. One can then consider Theorem 6.5 with $s = K$ and $\mathcal{B}_s = \mathcal{Ident} = \{\beta^*\}$. Still, in the high-dimensional framework, we have that $\text{rank } \Psi_n^{(I)} \leq \min(n, K) = n < K$, thus $\dim(\text{Ker } \Psi_n^{(I)}) \geq K - n > 0$. Hence, in the high-dimensional framework, $\Psi_n^{(I)} \Delta = 0$ can have non-zero solutions. It is because of the cone constraint that the sensitivities $\kappa_{p, J_0, J(\beta^*)}^{(c,I)}$ and $\kappa_{k, J(\beta^*)}^{(c,I)*}$ on the right hand-side of (6.7) and (6.8) can be different from 0 and thus yield non-trivial upper bounds.

Consider now the case of Example 6 where the instruments can have a direct effect on the outcome. In that case $L < K$ and identification fails without further assumptions. This is simply because $\dim \text{Ker}(\mathbb{E}[z_i x_i^T]) \geq K - L > 0$. If we knew the number s of non-zero coefficients, the underlying vector β is the solution of one system

$$(6.10) \quad \begin{cases} \beta_{J^c} = 0 \\ \mathbb{E}[z_i(y_i - x_{J_i}^T \beta_J)] = 0 \end{cases}$$

for each subset J of $\{1, \dots, K\}$ of size s (including the index of the constant regressor). For every set J such that $\mathbb{E}[z_i x_{J_i}^T]$ has full column rank, (6.10) has at most one solution. When $s < L$, $\mathbb{E}[z_i x_{J_i}^T] \beta_J = \mathbb{E}[z_i y_i]$ does not necessarily have a solution because the system is overdetermined. Note as well that, because s is the number of non-zero coefficients, in addition to (6.10), we know that for every k in $\{1, \dots, K\}$, $\beta_k \neq 0$. This again restricts the set of solutions. Note that there are also cases where the set of solutions of (6.10) is an affine space. For example, if $s > L$, $\mathbb{E}[z_i x_{J_i}^T]$ does not have full column rank. Geometrically \mathcal{B}_s is the union of the intersections of \mathcal{Ident} with the spaces $\{\forall k \in J, \beta_k = 0\}$ if for every k in $\{1, \dots, K\}$, $\beta_k \neq 0$. This is a union of affine spaces. We have observed that, adding the prior information that there are at most $s < L$ non-zero coefficients, is very likely to yield a set \mathcal{B}_s which is a union of points. Recall that, in the context of Example 6, $s < L$ means that there are in reality instruments that do not have a direct effect on the outcome. There are thus unknown exclusion restrictions which have an identification power. Again, because of the cone constraints, the sensitivities $\kappa_{p, J_0, J(\beta^*)}^{(c,I)}$ and $\kappa_{k, J(\beta^*)}^{(c,I)*}$ on the right hand-side of (6.7) and (6.8) can be different from 0 even in the high-dimensional framework where n is small relative K .

Consider now the term $\left(1 - r/\kappa_{1, J(\beta)}^{(c,I)}\right)_+^{-1}$ in the upper bounds of Theorem 6.5. Observe that the confidence sets of level at least $1 - \alpha$ for β in \mathcal{B}_s can have infinite volume on the random event

$\frac{r}{\kappa_{1,J(\beta)}^{(c,I)}} > 1$. Indeed, the upper bounds (6.7) and (6.8) are infinite. This occurs either when r is large or when $\kappa_{1,J(\beta)}^{(c,I)}$ is too small. Recall that r is of the order of $\sqrt{\log(L)/n}$ in all scenarios. Because increasing L increases the sensitivities, once $\kappa_{1,J(\beta)}^{(c,I)}$ is sufficiently bounded away from zero, the first condition corresponds to a situation with very many instruments relative to the sample size. This occurs when L is as large as an exponential in n . To interpret the second condition, note that Proposition 5.2 (ii) yields that for any k in $J(\beta)$,

$$\kappa_{1,J(\beta)}^{(c,I)} \leq \kappa_{k,J(\beta)}^{(c,I)*}.$$

This can occur on the event where a regressor $(x_{ki})_{i=1}^n$ has a restricted maximal partial empirical correlation with the instruments less than r . In the setting of Example 7, this becomes increasingly likely when the parameter ζ_{J^c} in the data generating approaches zero (see (5.7)). Confidence sets of infinite volume with positive probability is a desired feature of a procedure that is robust to weak instruments (see Dufour (1997) building on Gleser and Hwang (1987)). We will show in Section 8.1 that for low dimensional models (K is small and there is no sparsity) a slight modification of the *STIV* estimator is a new procedure that is robust to weak instruments (see Andrews and Stock (2007) for a review of existing methods). Note also that the term $\left(1 - r/\kappa_{1,J(\beta)}^{(c,I)}\right)_+^{-1}$ appears because we have a procedure that relaxes distributional assumptions and, in the homoscedastic case, does not require to know the variance of the errors or an upper bound. When the regressors are deterministic and the errors are Gaussian, an upper bound on the variance is sufficient to adjust the penalization and obtain a consistent Lasso or Dantzig estimator. In the linear regression with exogenous regressors, $\mathbb{E}[y_i^2]$ provides such an upper bound on the variance. This no longer works in the presence of endogenous regressors.

When $\tau_1 \triangleq 1 - \frac{r}{\kappa_{1,J(\beta)}^{(c,I)}}$ is close to 1 and the sensitivities $\kappa_{p,J_0,J(\beta)}^{(c,I)}$ and $\kappa_{k,J(\beta)}^{(c,I)*}$ are bounded away from zero, the upper bounds in (6.8) is of the order $O(r) = O(\sqrt{\log(L)/n})$. Thus, we have an extra $\sqrt{\log(L)}$ factor as compared to the usual root- n rate. It is a modest price for using a large number L of instruments. Under the premise of Proposition 5.2, for $\tau_1 \approx 1$ it is sufficient to have $|J(\beta)| \leq Cr^{-1} = O(\sqrt{n/\log(L)})$ where $C > 0$ is a proper constant. This is quite a reasonable condition on the sparsity $|J(\beta)|$.

The only unknown ingredient of the inequalities (6.7) and (6.8) is the set $J(\beta)$ that determines the sensitivities. To turn these inequalities into valid confidence bounds, it suffices to provide data-driven lower estimates on the sensitivities. As discussed in Section 5, there are two ways to do it. The

first one is based on the sparsity certificate, *i.e.*, assuming some known upper bound s on $|J(\beta)|$; then we get bounds depending only on s and on the data.

Corollary 6.6. *For every β in \mathcal{B}_s , under scenarios 2-5 of Section 6.1 together with its respective assumption and choice of r , with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5), for any c in $(0, 1)$ and any set of indices I containing the index of the instrument which is unity, for any solution $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ of the minimization problem (3.5) we have*

$$(6.11) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1}(\widehat{\beta}^{(c,I)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\widehat{\sigma}^{(c,I)}r}{\overline{\kappa}_{p,J_0}^{(c,I)}(s)} \left(1 - \frac{r}{\kappa_1^{(c,I)}(s)} \right)_+^{-1} \quad \forall p \in [1, \infty], \quad \forall J_0 \subset \{1, \dots, K\},$$

where $\overline{\kappa}_{p,J_0}^{(c,I)}(s)$ is any lower bound on $\kappa_{p,J_0,J(\beta)}^{(c,I)}$ based on the sparsity certificates that is convenient to calculate (see, *e.g.*, (5.13) and (5.14))

$$(6.12) \quad |\widehat{\beta}_k^{(c,I)} - \beta_k| \leq \frac{2\widehat{\sigma}^{(c,I)}r}{\mathbb{E}_n[X_k^2]^{1/2} \kappa_k^{(c,I)*}(s)} \left(1 - \frac{r}{\kappa_1^{(c,I)}(s)} \right)_+^{-1} \quad \forall k = 1, \dots, K, .$$

Under scenario 1 of Section 6.1 with the corresponding choice of r , for any set of indices I containing the index of the instrument which is unity, for every β in \mathcal{B}_s , with probability at least $1 - \alpha$, for any c in $(0, 1)$, any solution $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ of the minimization problem (3.5) satisfies (6.11) and (6.12).

In simulations the *STIV* estimator has always more non-zeros than the truth so it is reasonable to take $s = |J(\widehat{\beta}^{(c,I)})|$ or some larger value. We also advertise the possibility of drawing nested confidence sets for increasing values of s .

The second way is to plug in, instead of $J(\beta)$, some data-driven upper estimate \widehat{J} , *i.e.*, a set satisfying $J(\beta) \subseteq \widehat{J}$ on the intersection of the event of probability at least $1 - \alpha$ (or approximately $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5) of Theorem 6.5 and an event of probability close to 1. Theorem 7.4 provides examples of such estimators \widehat{J} . In particular, assertion (iv) of Theorem 7.4 guarantees that, under some assumptions, the estimator $\widehat{J} = J(\widehat{\beta}^{(c,I)})$ has the required property. The statement of the corresponding result is postponed to Section 7.

We do not touch upon optimality of the confidence regions in this article. Because of the popularity of two-stage least squares, we present the properties of a high dimensional two-stage method akin to two-stage least squares in Section 8.3. The *STIV* estimator depends on the tuning parameter c . A smaller c implies a smaller cone $C_{J(\beta)}^{(c,I)}$ and thus a larger sensitivity. On the other hand, because we penalize less σ in (3.5), $\widehat{\sigma}^{(c,I)}$ is larger. Because the dependence is nonlinear and the sensitivities

are random and depend on the distribution of the data generating process, there does not exist a universally good value for c ¹⁴. It is important to note that our results hold on the $1 - \alpha$ probability event, for any value of c and any set of indices I containing the index of the instrument which is unity (for scenarios 2-5 only for the set I). It is therefore possible to consider values of c or I which are random and depend on the data. If one is interested in a specific coefficient, it is possible to pick the values of c and I that yields the smaller confidence interval for that specific regressor. Because the procedure is fast to implement, varying c on a grid on $(0, 1)$ and picking the value that yields the smaller confidence interval is an easy thing to do. Varying I is too complicated but, if one hesitates on a few specifications of I , it is possible to build confidence sets for these various possibilities and choose the set I that yields the smaller confidence sets.

6.3. Sparse Oracle Inequality. We now consider the approximately sparse setting. The sparsity assumption is quite natural in empirical economics since usually only a moderate number of covariates is included in the model. However, one might be also interested in the case when β is only approximately sparse. This means that most of the coefficients β are not exactly zero but too small to matter, whereas the remaining ones are relatively large. This setting received some attention in the statistical literature. For example, the performance of Dantzig selector and MU -selector under such assumptions is studied by Candès and Tao (2007) and Rosenbaum and Tsybakov (2010) respectively. We will derive a similar result for the $STIV$ estimator.

Consider the enlarged cone

$$\tilde{C}_J^{(c)} \triangleq \left\{ \Delta \in \mathbb{R}^K : |\Delta_{J^c}|_1 \leq \frac{2+c}{1-c} |\Delta_J|_1 \right\}$$

and define, for $p \in [1, \infty]$ and $J_0 \subset \{1, \dots, K\}$

$$\tilde{\kappa}_{p, J_0, J}^{(c, I)} \triangleq \inf_{\Delta \in \mathbb{R}^K : |\Delta_{J_0}|_p = 1, \Delta \in \tilde{C}_J^{(c)}} \left| \Psi_n^{(I)} \Delta \right|_\infty$$

and $\tilde{\kappa}_{1, J}^{(c, I)}$ corresponds to $\tilde{\kappa}_{p, J_0, J}^{(c, I)}$ with $p = 1$ and $J_0 = \{1, \dots, K\}$.

The following theorem is an analog of the above results for the approximately sparse case.

Theorem 6.7. *For every β in \mathcal{B}_K , under scenarios 2-5 of Section 6.1 together with its respective assumption and choice of r , with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5), for any c in $(0, 1)$ and any set of indices I containing*

¹⁴The same is true for the Square-root Lasso.

the index of the instrument which is unity, for any solution $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ of the minimization problem (3.5) we have, for every $J_0 \subset \{1, \dots, K\}$,

$$(6.13) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1} \left(\widehat{\beta}^{(c,I)} - \beta \right) \right)_{J_0} \right|_p \leq \min_{J \subset \{1, \dots, K\}} \left\{ \max \left(\frac{2\widehat{\sigma}^{(c,I)} r}{\widetilde{\kappa}_{p, J_0, J}^{(c,I)}} \left(1 - \frac{r}{\widetilde{\kappa}_{1, J}^{(c,I)}} \right)^{-1}, \frac{6 |(\mathbf{D}_{\mathbf{X}}^{-1} \beta)_{J^c}|_1}{1 - c} \right) \right\}.$$

Under scenario 1 of Section 6.1 with the corresponding choice of r , for any set of indices I containing the index of the instrument which is unity, for every β in \mathcal{B}_s , with probability at least $1 - \alpha$, for any c in $(0, 1)$, any solution $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ of the minimization problem (3.5) satisfies (6.13) for every $J_0 \subset \{1, \dots, K\}$.

We can interpret Theorem 6.7 as the fact that the *STIV* estimator automatically realizes a “bias/variance” trade-off related to a non-linear approximation. Inequality (6.13) means that this estimator performs as well as if the optimal subset J were known.

7. RATES OF CONVERGENCE AND SELECTION OF THE VARIABLES

Let us consider rates of convergence of our estimator. We need to replace the random right hand-side of (6.7), (6.8) and (6.9) by deterministic upper bounds. In this section we consider that c in $(0, 1)$ and the set I are fixed.

Assumption 7.1. *For every $\beta \in \mathcal{B}_s$ and $\gamma_1 \in (0, 1)$, there exists a constant $\sigma_* > 0$ such that*

$$\mathbb{P} \left(\max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(I)})_{ll}^2 \mathbb{E}_n [Z_l^2 (Y - X^T \beta)^2] \leq \sigma_*^2 \right) \geq 1 - \gamma_1.$$

The second assumption concerns the population counterparts of the sensitivities.

Assumption 7.2. *For every $\beta \in \mathcal{B}_s$ and $\gamma_2 \in (0, 1)$, there exists constants $c_p^{(c,I)} > 0$, $c_{1, J_0}^{(c,I)} > 0$ for $J_0 = J(\beta)$ and $J_0 = \{k\}$ for $k = 1, \dots, K$, such that, with probability at least $1 - \gamma_2$,*

$$(7.1) \quad \kappa_{p, J(\beta)}^{(c,I)} \geq c_p^{(c,I)} |J(\beta)|^{-1/p},$$

$$(7.2) \quad \kappa_{1, J_0, J(\beta)}^{(c,I)} \geq c_{1, J_0}^{(c,I)}.$$

If $J_0 = \{k\}$ is a singleton we write for brevity $c_{1, J_0}^{(c,I)} = c_k^{(c,I)*}$.

The dependence on $|J(\beta)|$ of the right hand-side of (7.1) is motivated by Proposition 5.2. In (7.2), we do not indicate the dependence of the bounds on $|J(\beta)|$ explicitly because it can be different for different sets J_0 . For general J_0 , combining Proposition 5.1 (ii) and Proposition 5.2 suggests that

the value $c_{1,J_0}^{(c,I)}$ can be bounded from below by a quantity of the order $|J(\beta)|^{-1}$. Note, however, that this is a coarse bound valid for any set J_0 .

The last assumption defines a population counterpart of $\mathbb{E}_n[X_k^2]^{1/2}$.

Assumption 7.3. *For every $\gamma_3 \in (0, 1)$ and $k \in \{1, \dots, K\}$, there exist constants $v_k > 0$ such that*

$$\mathbb{P}\left(\mathbb{E}_n[X_k^2]^{1/2} \geq v_k, \forall k \in \{1, \dots, K\}\right) \geq 1 - \gamma_3.$$

Assumptions 7.1, 7.2 and 7.3 are very weak. They are required to obtain rates of convergence, *i.e.*, a deterministic bound on the estimation error on a high probability event (see, *e.g.*, (7.4) below). We set $\gamma = \alpha + \sum_{j=1}^3 \gamma_j$, and

$$\tau^{(c,I)*} \triangleq \left(1 + \frac{r}{cc_{1,J(\beta)}^{(c,I)}}\right) \left(1 - \frac{r}{cc_{1,J(\beta)}^{(c,I)}}\right)_+^{-1} \left(1 - \frac{r|J(\beta)|}{c_1^{(c,I)}}\right)_+^{-1}.$$

Theorem 7.4. *For every β in \mathcal{B}_s , under the assumptions of Theorem 6.5 and Assumption 7.1, the following holds.*

(i) *Let part (7.2) of Assumption 7.2 be satisfied. Then, with probability at least $1 - \alpha - \gamma_1 - \gamma_2$, for any solution $\hat{\sigma}^{(c,I)}$ of (3.5) we have*

$$\hat{\sigma}^{(c,I)} \leq \sigma_* \left(1 + \frac{r}{cc_{1,J(\beta)}^{(c,I)}}\right) \left(1 - \frac{r}{cc_{1,J(\beta)}^{(c,I)}}\right)_+^{-1}.$$

(ii) *Fix $p \in [1, \infty]$. Let Assumption 7.2 be satisfied. Then, with probability at least $1 - \alpha - \gamma_1 - \gamma_2$, for any solution $\hat{\beta}^{(c,I)}$ of (3.5) we have*

$$(7.3) \quad \left|\mathbf{D}_X^{-1} \left(\hat{\beta}^{(c,I)} - \beta\right)\right|_p \leq \frac{2\sigma_* r |J(\beta)|^{1/p} \tau^{(c,I)*}}{c_p^{(c,I)}}.$$

(iii) *Let Assumptions 7.2 and 7.3 be satisfied. Then with probability at least $1 - \gamma$, for any solution $\hat{\beta}^{(c,I)}$ of (3.5) we have*

$$(7.4) \quad |\hat{\beta}_k^{(c,I)} - \beta_k| \leq \frac{2\sigma_* r \tau^{(c,I)*}}{c_k^{(c,I)*} v_k}, \quad k = 1, \dots, K.$$

(iv) *Let the assumptions of (iii) hold, and $|\beta_k| > \frac{2\sigma_* r \tau^{(c,I)*}}{c_k^{(c,I)*} v_k}$ for all $k \in J(\beta)$. Then, with probability at least $1 - \gamma$, for any solution $\hat{\beta}^{(c,I)}$ of (3.5) we have*

$$J(\beta) \subseteq J(\hat{\beta}^{(c,I)}).$$

For reasonably large sample size ($n \gg \log(L)$), the value r is small, and $\tau^{(c,I)*}$ is approaching 1 as $r \rightarrow 0$. Thus, the bounds (7.3) and (7.4) are of the order of magnitude $O(r|J(\beta)|^{1/p})$ and $O(r)$ respectively. These are the same rates, in terms of the sparsity $|J(\beta)|$, the dimension L , and the sample size n , that were proved for the Lasso and Dantzig selector in high-dimensional regression with Gaussian errors and without endogenous variables (here $L = K$) in Candès and Tao (2007), Bickel, Ritov and Tsybakov (2009), Lounici (2008) (see also Bühlmann and van de Geer (2011) for references to more recent work).

The lower bound for estimation of a high dimensional linear regression model, without endogeneity, under fixed and Gaussian design, with K potential regressor and at most s non-zero coefficients, is known to be $\sqrt{\frac{|J(\beta)| \log(K/|J(\beta)|)}{n}}$ (see Verzelen (2010), Ye and Zhang (2010) and Raskutti, Wainwright and Yu (2011), rates for prediction are also obtained in Rigollet and Tsybakov (2011)). Proving minimax lower bounds for the model (1.1)-(1.2) with $L \neq K$ is subject of future investigation.

Note that Theorem 7.4 assumes that we work on some event such that the lower bounds $c_p^{(c,I)}$, $c_k^{(c,I)*}$ and $c_{1,J(\beta)}^{(c,I)}$ are positive. When for a particular value of β there are 0 on a large probability event then on that event we do not have convergence rates or model selection results. Also the extra condition in (iv) is not satisfied for every β in \mathcal{B}_s . Therefore when \mathcal{B}_s is not a point then the conclusion of (iv) only holds for certain values of the parameter in the set \mathcal{B}_s .

From (6.8), Theorem 7.4 (iv) and Proposition 5.1 (i), we obtain the following confidence sets of level $1 - \gamma$ for β_k in \mathcal{B}_s .

Corollary 7.5. *For every β in \mathcal{B}_s , under one of the 5 scenarios of Section 6.1, together with its respective assumption and choice of r , under the assumptions of Theorem 7.4 (iv), with probability at least $1 - \gamma$ (approximately at least $1 - \gamma$ for scenario 4 and the two-stage procedure with scenario 5) for any solution $(\hat{\beta}, \hat{\sigma})$ of the minimization problem (3.5) we have*

$$(7.5) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1}(\hat{\beta}^{(c,I)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\hat{\sigma}^{(c,I)}r}{\bar{\kappa}_{p,J_0,J(\hat{\beta}^{(c,I)})}^{(c,I)}} \left(1 - \frac{r}{\kappa_{1,J(\hat{\beta}^{(c,I)})}^{(c,I)}} \right)_+^{-1}, \quad \forall p \in [1, \infty], \quad \forall J_0 \subset \{1, \dots, K\},$$

where $\bar{\kappa}_{p,J_0,J(\hat{\beta}^{(c,I)})}^{(c,I)}$ is any lower bound on $\kappa_{p,J_0,J(\hat{\beta}^{(c,I)})}^{(c,I)}$ that is convenient to calculate (see, e.g., (5.13) and (5.14)), and for all $k = 1, \dots, K$,

$$(7.6) \quad |\hat{\beta}_k^{(c,I)} - \beta_k| \leq \frac{2\hat{\sigma}^{(c,I)}r}{\mathbb{E}_n[X_k^2]^{1/2} \kappa_{k,J(\hat{\beta}^{(c,I)})}^{(c,I)*}} \left(1 - \frac{r}{\kappa_{1,J(\hat{\beta}^{(c,I)})}^{(c,I)}} \right)_+^{-1}.$$

Here we do not control exactly the coverage probability because its probability is now $1 - \gamma$. Having γ_j very small for $j = 1, 2, 3$ and thus coverage probability of approximately at least $1 - \alpha$ requires taking a large σ^* and lower bounds on the sensitivities. Such a choice is compatible with (iv) from Theorem 7.4 only when β_k are large enough for $k \in J(\beta)$.

Theorem 7.4 (iv) provides an upper estimate on the set of non-zero components of β . We now consider the problem of the exact selection of variables. For this purpose, we use the thresholded *STIV* estimator whose coordinates are defined by

$$(7.7) \quad \tilde{\beta}_k^{(c,I)}(\omega_k^{(c,I)}) \triangleq \begin{cases} \widehat{\beta}_k^{(c,I)} & \text{if } |\widehat{\beta}_k^{(c,I)}| > \omega_k^{(c,I)}, \\ 0 & \text{otherwise,} \end{cases}$$

where $\widehat{\beta}_k^{(c,I)}$ are the coordinates of the *STIV* estimator $\widehat{\beta}^{(c,I)}$, and $\omega_k^{(c,I)} > 0$, $k = 1, \dots, K$, are thresholds that will be specified below. We will use the sparsity certificate approach, so that the thresholds will depend on the upper bound s on the number of non-zero components of β . Because, for practical use, we want to a data driven thresholding rule we need to strengthen Assumption 7.2 as follows.

Assumption 7.6. *Fix an integer $s \geq 1$. For every $\gamma_2 \in (0, 1)$, for β in \mathcal{B}_s , there exist constants $c_{1,J(\beta)}^{(c,I)} > 0$, $c_{1,J_0}^{(c,I)}(s) > 0$ for $J_0 = \{k\}$, $\forall k$ and $J_0 = \{1, \dots, K\}$, such that, with probability at least $1 - \gamma_2$,*

$$(7.8) \quad \kappa_{1,J(\beta),J(\beta)}^{(c,I)} \geq c_{1,J(\beta)}^{(c,I)} \quad \text{and} \quad \kappa_{1,J_0}^{(c,I)}(s) \geq c_{1,J_0}^{(c,I)}(s)$$

If $J_0 = \{k\}$ is a singleton we write for brevity $c_{1,J_0}^{(c,I)}(s) = c_k^{(c,I)*}(s)$ and $c_{1,\{1,\dots,K\}}^{(c,I)}(s) = c_1^{(c,I)}(s)$.

Set

$$\tau^{(c,I)*}(s) \triangleq \left(1 + \frac{r}{c c_{1,J(\beta)}^{(c,I)}}\right) \left(1 - \frac{r}{c c_{1,J(\beta)}^{(c,I)}}\right)_+^{-1} \left(1 - \frac{r}{c_1^{(c,I)}(s)}\right)_+^{-1}.$$

The following theorem shows that, based on thresholding of the *STIV* estimator, we can reconstruct exactly the set of non-zero coefficients $J(\beta)$ with probability close to 1. Even more, we achieve the sign consistency, *i.e.*, we reconstruct exactly the vector of signs of the coefficients of β with probability close to 1.

Theorem 7.7. *For every β in \mathcal{B}_s , let the assumptions of Theorem 6.5 and Assumptions 7.1, 7.3, 7.6 be satisfied. Assume that $|J(\beta)| \leq s$, and $|\beta_k| > \frac{4\sigma_* r \tau^{(c,I)*}(s)}{c_k^{(c,I)*}(s)v_k}$ for all $k \in J(\beta)$. Take the thresholds*

$$\omega_k^{(c,I)}(s) \triangleq \frac{2\widehat{\sigma}^{(c,I)} r}{\kappa_k^{(c,I)*}(s) \mathbb{E}_n[X_k^2]^{1/2}} \left(1 - \frac{r}{\kappa_1^{(c,I)}(s)}\right)_+^{-1},$$

and consider the estimator $\tilde{\beta}^{(c,I)}$ with coordinates $\tilde{\beta}_k^{(c,I)}(\omega_k^{(c,I)}(s)), k = 1, \dots, K$. Then, with probability at least $1 - \gamma$, we have

$$(7.9) \quad \overrightarrow{\text{sign}(\tilde{\beta}^{(c,I)})} = \overrightarrow{\text{sign}(\beta)}.$$

As a consequence, $J(\tilde{\beta}^{(c,I)}) = J(\beta)$.

Sign consistency implies perfect model selection but it is also of independent interest. Indeed, the above result gives conditions under which one can make valid statements on the sign of the coefficients. Unfortunately this is only a theoretical result because it is not possible to verify the condition that $|\beta_k| > \frac{4\sigma_* r \tau^{(c,I)*}(s)}{c_k^{(c,I)*}(s) v_k}$ for all $k \in J(\beta)$.

8. FURTHER RESULTS ON THE STIV ESTIMATOR

8.1. Low Dimensional Models and Very-Many-Weak-Instruments. In this section we consider a slight modification of the *STIV* estimator to deal with the case where K is small relative to n and we know that all K coefficients are nonzero. It is a new procedure that is robust to a situation where each individual instrument is weak. It can handle very-many instruments, *i.e.* L can be much larger than n . It does not rely on normal errors and the confidence sets under Scenarios 2 and 4 (asymptotic) allow for heteroscedasticity. Again, we do not rely on non-standard asymptotics.

Because this is a very active and challenging area of research in econometrics we cannot make an exhaustive literature review and refer to Andrews and Stock (2007) and the references therein.

Because K is small and L is large (1.1)-(1.2) is usually point identified, but in the presence of weak instruments one is at the verge of identification. Thus for the sake of completeness we do not exclude partial identification from our result.

Definition 8.1. We call the *STIV-R estimator* any solution $(\hat{\beta}^{(I)}, \hat{\sigma}^{(I)})$ of the following minimization problem:

$$(8.1) \quad \min_{(\beta, \sigma) \in \mathcal{I}^{(I)}} \sigma.$$

In that case there are no cone constraints in the definition of the sensitivities and we simply drop the index $J(\beta)$ in the notation of the sensitivities. We also drop the exponent c everywhere because the minimization problem (8.1) no longer involves the constant c because the model is low dimensional. Unlike the high-dimensional setup of the previous sections, the sensitivities can be directly calculated from the data and we do not have to rely on lower bounds. Obtaining confidence

sets in a non high-dimensional structural equation is much more direct and easy. We obtain the following result.

Theorem 8.2. *For every β in $\mathcal{I}dent$, under one of the 5 scenarios of Section 6.1 together with its respective assumption and choice of r , with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5) for any solution $(\widehat{\beta}^{(I)}, \widehat{\sigma}^{(I)})$ of the minimization problem (8.1)*

$$(8.2) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1}(\widehat{\beta}^{(I)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\widehat{\sigma}^{(I)}r}{\kappa_{p,J_0}^{(I)}} \left(1 - \frac{r}{\kappa_1^{(I)}} \right)_+^{-1}, \quad \forall p \in [1, \infty], \quad \forall J_0 \subset \{1, \dots, K\},$$

for all $k = 1, \dots, K$,

$$(8.3) \quad |\widehat{\beta}_k^{(I)} - \beta_k| \leq \frac{2\widehat{\sigma}^{(I)}r}{\mathbb{E}_n[X_k^2]^{1/2}\kappa_k^{(I)*}} \left(1 - \frac{r}{\kappa_1^{(I)}} \right)_+^{-1}$$

and

$$(8.4) \quad \widehat{\sigma}^{(I)} \leq \max_{l=1, \dots, L} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\beta)}.$$

The same discussion that we had in the case of a high-dimensional structural equation with endogenous regressors holds. The term

$$\left(1 - \frac{r}{\kappa_1^{(I)}} \right)_+^{-1}$$

in the upper bounds of Theorem 8.2 can yield infinite volume confidence regions. This occurs either when r , of the order of $\sqrt{\log(L)/n}$ in all scenarios, is large or when $\kappa_1^{(I)}$ is too small. The first condition occurs when L is as large as an exponential in n . To interpret the second condition, again Proposition 5.2 (ii) yields that for any k in $\{1, \dots, K\}$,

$$\kappa_1^{(I)} \leq \kappa_k^{(I)*}.$$

Therefore a necessary condition for $\kappa_1^{(I)}$ to be small is when there exists in model (1.1) a regressor $(x_{ki})_{i=1}^n$ with low coordinate-wise sensitivity. Without the cone constraint the coordinate wise sensitivities are defined through

$$\kappa_k^{(I)*} = \inf_{\lambda \in \mathbb{R}^{K-1}} \max_{l=1, \dots, L} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \left| \frac{1}{n} \sum_{i=1}^n z_{li} \left(x_{ki} - x_{\{k\}^c_i}^T \lambda \right) \right|$$

and can be interpreted as *maximal partial empirical correlations with the instruments*. This means that there is a regressor for which all¹⁵ instruments are weak in the sense that it is such that

$$\inf_{\lambda \in \mathbb{R}^{K-1}} \max_{l=1, \dots, L} (\mathbf{D}_{\mathbf{Z}}^{(l)})_{ll} \left| \frac{1}{n} \sum_{i=1}^n z_{li} (x_{ki} - x_{\{k\}^c_i}^T \lambda) \right| \leq r.$$

Again, in the setting of Example 7, this becomes increasingly likely as ζ_{J^c} approaches zero (see (5.7)). Rates of convergence could be obtained as well as in Section 7.

8.2. Potentially Sharper Results When None of the Instruments Have Heavy Tail. In the subsequent sections we restrict the subset of indices I in (3.4) to a singleton (the index of the regressor which is constant and equal to 1). When I is a singleton, the *STIV* estimator is straightforward to obtain because (3.4) only involves one conic constraint. This procedure should be used when all regressors and instruments are bounded or have relatively thin tails. In this section we consider the properties of the *STIV* estimator with a different matrix $\mathbf{D}_{\mathbf{X}}$ in (3.5). We use for the matrix $\mathbf{D}_{\mathbf{X}}$ the diagonal $K \times K$ matrix with diagonal entries x_{k*}^{-1} , $k = 1, \dots, K$ where $x_{k*} \triangleq \max_i |x_{ki}|$.

The following theorem which is in the same spirit as Theorem 6.5.

Theorem 8.3. *For every β in \mathcal{B}_s , under one of the 5 scenarios of Section 6.1 together with its respective assumption and choice of r , with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5), for any c in $(0, 1)$, for any solution $(\widehat{\beta}^{(c)}, \widehat{\sigma}^{(c)})$ of the minimization problem (3.5) we have for all $p \in [1, \infty]$ and $J_0 \subset \{1, \dots, K\}$,*

$$(8.5) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1} (\widehat{\beta}^{(c)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\widehat{\sigma}^{(c)} r}{\kappa_{p, J_0, J(\beta)}^{(c)}} \left(1 - \frac{r}{\kappa_{1, J_{\text{exo}}, J(\beta)}^{(c)}} - \frac{r^2}{\kappa_{1, J_{\text{exo}}, J(\beta)}^{(c)}} \right)_+^{-1}$$

for all $k = 1, \dots, K$,

$$(8.6) \quad |\widehat{\beta}_k^{(c)} - \beta_k| \leq \frac{2\widehat{\sigma}^{(c)} r}{x_{k*} \kappa_{k, J(\beta)}^{(c)*}} \left(1 - \frac{r}{\kappa_{1, J_{\text{exo}}, J(\beta)}^{(c)}} - \frac{r^2}{\kappa_{1, J_{\text{exo}}, J(\beta)}^{(c)}} \right)_+^{-1}$$

and

$$(8.7) \quad \widehat{\sigma}^{(c)} \leq \sqrt{\widehat{Q}(\beta)} \left(1 + \frac{r}{c\kappa_{1, J(\beta), J(\beta)}^{(c)}} \right) \left(1 - \frac{r}{c\kappa_{1, J(\beta), J(\beta)}^{(c)}} \right)_+^{-1}.$$

¹⁵Not as a set, see the discussion in Section 5 about the comparison with the usual concentration parameter.

Note that the upper bound (6.7) involves the term

$$\tau_1 = 1 - \frac{r}{\kappa_{1,J(\beta)}^{(c)}}$$

while (8.5) in Theorem 8.3 involves

$$\tau_2 \triangleq 1 - \frac{r}{\kappa_{1,J_{\text{exo}}^c,J(\beta)}^{(c)}} - \frac{r^2}{\kappa_{1,J_{\text{exo}},J(\beta)}^{(c)}}.$$

Though the sensitivities in the two cases are not entirely comparable because we use different matrices $\mathbf{D}_{\mathbf{X}}$, we know, from Proposition 5.1 (ii) that restricting a sensitivity to a sub-block increases the sensitivity. When the set J_{exo}^c of regressors that have not been identified as exogenous and therefore used as instruments is small, the sensitivity $\kappa_{1,J_{\text{exo}}^c,J(\beta)}^{(c)}$ is large, whereas the small sensitivity of its complement $\kappa_{1,J_{\text{exo}},J(\beta)}^{(c)}$ is compensated by the small value r^2 in the numerator. In the extreme case where $J_{\text{exo}}^c = \emptyset$, we have $\frac{r}{\kappa_{1,J_{\text{exo}}^c,J(\beta)}^{(c)}} = 0$, so that $\tau_2 \leq 1 - \frac{r^2}{\kappa_{1,J_{\text{exo}},J(\beta)}^{(c)}}$. Under the premise of Proposition 5.2, for it to be positive it is sufficient to have $|J(\beta)| \leq Cr^{-2} = O(n/\log(L))$ where $C > 0$ is a proper constant. Recall that in the case of Theorem 6.5 we need that $|J(\beta)| \leq Cr^{-1} = O(\sqrt{n/\log(L)})$.

The following results gives proper confidence sets.

Corollary 8.4. *For every β in \mathcal{B}_s , under one of the 5 scenarios of Section 6.1 together with its respective assumption and choice of r , with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5), for any c in $(0, 1)$, for any solution $(\widehat{\beta}^{(c)}, \widehat{\sigma}^{(c)})$ of the minimization problem (3.5) we have*

(8.8)

$$\left| \left(\mathbf{D}_{\mathbf{X}}^{-1} (\widehat{\beta}^{(c)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\widehat{\sigma}^{(c)}r}{\overline{\kappa}_{p,J_0}^{(c)}(s)} \left(1 - \frac{r}{\overline{\kappa}_{1,J_{\text{exo}}^c}^{(c)}(s)} - \frac{r^2}{\overline{\kappa}_{1,J_{\text{exo}}}^{(c)}(s)} \right)_+^{-1}, \quad \forall p \in [1, \infty], \forall J_0 \subset \{1, \dots, K\},$$

where $\overline{\kappa}_{p,J_0}^{(c)}(s)$, $\overline{\kappa}_{1,J_{\text{exo}}^c}^{(c)}(s)$, $\overline{\kappa}_{1,J_{\text{exo}}}^{(c)}(s)$ are any lower bounds on $\kappa_{p,J_0,J(\beta)}^{(c)}$, $\kappa_{1,J_{\text{exo}}^c,J(\beta)}^{(c)}$ and $\kappa_{1,J_{\text{exo}},J(\beta)}^{(c)}$ based on the sparsity certificates that is convenient to calculate (see, e.g., (5.13) and (5.14)), and for all $k = 1, \dots, K$,

$$(8.9) \quad |\widehat{\beta}_k^{(c)} - \beta_k| \leq \frac{2\widehat{\sigma}^{(c)}r}{x_{k*} \kappa_k^{(c)*}(s)} \left(1 - \frac{r}{\overline{\kappa}_{1,J_{\text{exo}}^c}^{(c)}(s)} - \frac{r^2}{\overline{\kappa}_{1,J_{\text{exo}}}^{(c)}(s)} \right)_+^{-1}.$$

As we have seen in Section 5, equation (5.13) yields a lower bound for the block sensitivities. It is only computationally feasible for small blocks. So this lower bound could only be used at most for one of the two block sensitivities appearing in τ_2 , the other one (possibly the two) should be lower

bounded using (5.14). When J_{exo}^c is small there are cases when the method of this section yields finite volume confidence regions while due to τ_1 we can only deduce from Theorem 6.5 infinite volume confidence regions.

It is easy to check that Theorem 7.4 and Corollary 7.5 also hold replacing Assumption 7.3 by

Assumption 8.5. *For every $0 < \gamma_3 < 1$, there exist constants $v_k > 0$ such that*

$$\mathbb{P}(x_{k*} \geq v_k, \forall k \in J(\beta)) \geq 1 - \gamma_3.$$

by assuming Assumption 7.6 allowing for J_0 to also be J_{exo} and J_{exo}^c and using the following expression for $\tau^{(c)*}$.

$$\tau^{(c)*} \triangleq \left(1 + \frac{r}{cc_{1,J(\beta)}^{(c)}}\right) \left(1 - \frac{r}{cc_{1,J(\beta)}^{(c)}}\right)_+^{-1} \left(1 - \frac{r}{c_{1,J_{\text{exo}}}^{(c)}} - \frac{r^2}{c_{1,J_{\text{exo}}}^{(c)}}\right)_+^{-1}.$$

Theorem 7.7 also holds using

$$\omega_k^{(c)}(s) \triangleq \frac{2\widehat{\sigma}^{(c)}r}{\kappa_k^{(c)*}(s)x_{k*}} \left(1 - \frac{r}{\kappa_{1,J_{\text{exo}}}^{(c)}(s)} - \frac{r^2}{\kappa_{1,J_{\text{exo}}}^{(c)}(s)}\right)_+^{-1},$$

with

$$\tau^{(c)*}(s) \triangleq \left(1 + \frac{r}{cc_{1,J(\beta)}^{(c)}}\right) \left(1 - \frac{r}{cc_{1,J(\beta)}^{(c)}}\right)_+^{-1} \left(1 - \frac{r}{c_{1,J_{\text{exo}}}^{(c)}(s)} - \frac{r^2}{c_{1,J_{\text{exo}}}^{(c)}(s)}\right)_+^{-1}.$$

Note that due to the new scaling, the constant v_k can be much larger than in Section 7. This does not change the rates when the regressors are bounded, but it does if the regressors are unbounded. If a regressor has a normal distribution, this implies an increased $\log(n)$ in the rate. When $\tau^{(c,I)*}$ would be infinite under the scaling of Section 3 this is still a mild loss. We present in Section 10.2 the implementation of this method in the difficult setting where the regressors are Gaussian. This method still performs well in this case.

8.3. The *STIV* Estimator with Linear Projection Instruments. The results of the previous sections show that the *STIV* estimator can handle a very large number of instruments, up to an exponential in the sample size. Moreover, adding instruments always improves the sensitivities. In this section, we consider the case where $L > K$ but we look for a smaller set of instruments, namely, of size K .

This is a classical problem when the structural equation is low-dimensional. The two-stage least square is a leading and very popular example. Under the stronger zero conditional mean assumption,

optimal instruments attain the semi-parametric efficiency bound (see Amemiya (1974), Chamberlain (1987), Newey (1990) and Hahn (2002)). In the homoscedastic case, the optimal instruments correspond to the projection of the endogenous variables on the space of variables measurable with respect to all the instruments¹⁶. These optimal instruments are expressed in terms of conditional expectations that are not available in practice and should be estimated. Related work includes, for example, Donald and Newey (2001), Chamberlain and Imbens (2004), Bai and Ng (2009), Okui (2010), Carrasco and Florens (2000) and Carrasco (2012). Belloni, Chen, Chernozhukov et al. (2010) is very related to our approach because it proposes to use the Lasso or post-Lasso to estimate the optimal instrument. This allows to approximate the conditional expectations with a number of functionals of a low dimensional vector of instrument that is exponential in the sample size. They consider a second stage which is the heteroscedastic robust *IV* estimator and obtain confidence sets for the low dimensional structural equation.

In this section we investigate the properties of a high-dimensional version of the two-stage least squares for high-dimensional structural equations with endogenous regressors. In that case, we have $L > K$ and K is potentially much larger than n . For simplicity, we will not consider estimating the optimal instruments in the first stage. One would also have to properly define optimality in high-dimension. This is a difficult question that should be addressed in the future. Moreover, we are aiming at robust confidence sets that have a good approximate finite sample validity. For simplicity of exposure we assume that there is only one endogenous regressor $(x_{k_{\text{end}}i})_{i=1}^n$ in (1.1). We write the reduced form equation in the form

$$(8.10) \quad x_{k_{\text{end}}i} = \sum_{l=1}^L z_{li}\zeta_l + v_i, \quad i = 1, \dots, n,$$

where ζ_l are unknown coefficients of the linear combination of instruments and

$$(8.11) \quad \mathbb{E}[z_{li}v_i] = 0$$

for $i = 1, \dots, n$, $l = 1, \dots, L$. We call $\sum_{l=1}^L z_{li}\zeta_l$ the linear projection instrument.

For simplicity of the proofs we use the normalization of Section 8.2. What we present in this section can easily be extended to a situation with more than one endogenous regressor and with the normalization of Section 3.

¹⁶This is extremely hard in the presence of many instruments due to the curse of dimensionality but it can be done when the instruments are functionals of a low dimensional original instrument.

The first stage consists in estimating the unknown coefficients ζ_l . If $L \geq K > n$ and if the reduced form model (8.10) has some sparsity (or approximate sparsity), it is natural to use a high-dimensional procedure, such as the Lasso, the Dantzig selector, the Square-root Lasso or the post-Lasso, to produce estimators $\widehat{\zeta}_l$ of the coefficients. Because we are after confidence sets for the coefficients of the structural equation, we will assume that (8.10) is sparse¹⁷. When the reduced form equation is not sparse or approximately sparse, for example when all the instruments are equally weak but relevant, this method will fail. This is what we observe in the simulations in Section 10.2. In that situation we recommend to use the one stage *STIV* estimator.

It is easy to check, from the first order condition that the *STIV* estimator is, up to the normalization, equivalent to the Square-root Lasso when all the regressors are exogenous. We can deduce from Theorem 8.3, that under one of the 5 scenarios, on an event E_α of probability at least $1 - \alpha$ (or approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5), for any c_{RF} in $(0, 1)$, any solution $(\widehat{\zeta}^{(c_{RF})}, \widehat{\sigma}_1^{(c_{RF})})$ to the *STIV* estimator applied to the reduced form equation is such that

$$\left| \left(\mathbf{D}_{\mathbf{X}}^{-1} (\widehat{\zeta}^{(c_{RF})} - \zeta) \right) \right|_1 \leq \frac{2\widehat{\sigma}_1^{(c_{RF})} r}{\kappa_{1,J(\zeta)}^{(c_{RF})}} \left(1 - \frac{r^2}{\kappa_{1,J(\zeta)}^{(c_{RF})}} \right)_+^{-1}.$$

This yields, if one is given a maximum sparsity s_{RF} , the computable upper bound

$$(8.12) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1} (\widehat{\zeta} - \zeta) \right) \right|_1 \leq \frac{2\widehat{\sigma}_{RF} r}{\kappa_1^{(c_{RF})}(s_{RF})} \left(1 - \frac{r^2}{\kappa_1^{(c_{RF})}(s_{RF})} \right)_+^{-1} \triangleq C_1(r, c_{RF}, s_{RF}).$$

The second stage now consists in using the *STIV* estimator from Section 8.2 with the enlarged *IV*-constraint

$$(8.13) \quad \widehat{\mathcal{I}}_{2S} \triangleq \left\{ (\beta, \sigma) : \beta \in \mathbb{R}^K, \sigma > 0, \left| \frac{1}{n} \mathbf{D}_{\mathbf{Z},2S} \mathbf{Z}_{2S}^T (\mathbf{Y} - \mathbf{X}\beta) \right|_\infty \leq \sigma r, \widehat{Q}(\beta) \leq \sigma^2 \right\}$$

where $\mathbf{D}_{\mathbf{Z},2S}$ is a $K \times K$ diagonal matrix such that $(\mathbf{D}_{\mathbf{Z},2S})_{k_{\text{end}} k_{\text{end}}} = (\max_{i=1, \dots, n} |z_i^T \widehat{\zeta}| + 2C_1(r, c_{RF}, s_{RF}))^{-1}$ and for every k in $\{1, \dots, K\} \setminus \{k_{\text{end}}\}$, $(\mathbf{D}_{\mathbf{Z},2S})_{kk} = x_{k^*}^{-1}$, and the matrix \mathbf{Z}_{2S} is the stacked matrix of the endogenous regressors and the estimated linear projection instrument $(z_i^T \widehat{\lambda})_{i=1}^n$. We enlarge the *IV*-constraint to account for the error made in the first stage estimation of the linear projection instrument. In this section, for simplicity, we restrict our attention to the case where I is a singleton corresponding to the index of the regressor being unity in $\widehat{\mathcal{I}}_{2S}$. For this reason we drop the index

¹⁷Belloni, Chen, Chernozhukov et al. (2010) also study the setting where the reduced form has an approximately sparse vectors of coefficients. We could obtain rates of convergence in that case or asymptotically valid confidence sets. Because we are aiming at confidence sets that have some finite sample validity we do not study this case in this article.

exponent I everywhere. We obtain the following result where we denote by $\kappa^{(c)2S}$ the sensitivities where we replace in the definition of $\Psi_n^{(I)}$, \mathbf{Z} by \mathbf{Z}_{2S} and $\mathbf{D}_{\mathbf{Z}}^{(I)}$ by $\mathbf{D}_{\mathbf{Z},2S}$.

Theorem 8.6. *For every β in \mathcal{B}_s , under one of the 5 scenarios of Section 6.1 together with its respective assumption and choice of r , with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5), for any c in $(0, 1)$, for any solution $(\widehat{\beta}^{(c)}, \widehat{\sigma}^{(c)})$ of the minimization problem (3.5), replacing $\widehat{\mathcal{I}}$ by $\widehat{\mathcal{I}}_{2S}$, we have for all $p \in [1, \infty]$ and $J_0 \subset \{1, \dots, K\}$,*

$$(8.14) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1}(\widehat{\beta}^{(c)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\widehat{\sigma}^{(c)}r}{\kappa_{p,J_0,J(\beta)}^{(c)2S}} \left(1 - \frac{r}{\kappa_{k_{\text{end}},J(\beta)}^{(c)2S^*}} - \frac{r^2}{\kappa_{1,\{1,\dots,K\}\setminus\{k_{\text{end}}\},J(\beta)}^{(c)2S}} \right)_+^{-1}$$

for all $k = 1, \dots, K$,

$$(8.15) \quad |\widehat{\beta}_k^{(c)} - \beta_k| \leq \frac{2\widehat{\sigma}^{(c)}r}{x_{k^*} \kappa_{k,J(\beta)}^{(c)2S^*}} \left(1 - \frac{r}{\kappa_{k_{\text{end}},J(\beta)}^{(c)2S^*}} - \frac{r^2}{\kappa_{1,\{1,\dots,K\}\setminus\{k_{\text{end}}\},J(\beta)}^{(c)2S}} \right)_+^{-1}$$

and

$$(8.16) \quad \widehat{\sigma}^{(c)} \leq \sqrt{\widehat{Q}(\beta)} \left(1 + \frac{r}{c\kappa_{1,J(\beta),J(\beta)}^{(c)2S}} \right) \left(1 - \frac{r}{c\kappa_{1,J(\beta),J(\beta)}^{(c)2S}} \right)_+^{-1}.$$

In turn, this yields the following *bona-fide* confidence sets.

Corollary 8.7. *For every β in \mathcal{B}_s , under one of the 5 scenarios of Section 6.1 together with its respective assumption and choice of r , with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5), for any c in $(0, 1)$, for any solution $(\widehat{\beta}^{(c)}, \widehat{\sigma}^{(c)})$ of the minimization problem (3.5), replacing $\widehat{\mathcal{I}}$ by $\widehat{\mathcal{I}}_{2S}$, we have*

$$(8.17) \quad \left| \left(\mathbf{D}_{\mathbf{X}}^{-1}(\widehat{\beta}^{(c)} - \beta) \right)_{J_0} \right|_p \leq \frac{2\widehat{\sigma}^{(c)}r}{\overline{\kappa}_{p,J_0}^{(c)2S}(s)} \left(1 - \frac{r}{\kappa_{k_{\text{end}}}^{(c)2S^*}(s)} - \frac{r^2}{\kappa_1^{(c)2S}(s)} \right)_+^{-1}, \quad \forall p \in [1, \infty], \forall J_0 \subset \{1, \dots, K\},$$

where $\overline{\kappa}_{p,J_0}^{(c)2S}(s)$, is any lower bounds on $\kappa_{p,J_0,J(\beta)}^{(c)2S}$ based on the sparsity certificates that is convenient to calculate (see, e.g., (5.13) and (5.14)), and for all $k = 1, \dots, K$,

$$(8.18) \quad |\widehat{\beta}_k^{(c)} - \beta_k| \leq \frac{2\widehat{\sigma}^{(c)}r}{x_{k^*} \kappa_k^{(c)*}(s)} \left(1 - \frac{r}{\kappa_{k_{\text{end}}}^{(c)2S^*}(s)} - \frac{r^2}{\kappa_1^{(c)2S}(s)} \right)_+^{-1}.$$

All lower bounds on the sensitivities appearing in (8.17) and (8.18) are easy to calculate. Indeed, we have minorized $\kappa_{1,\{1,\dots,K\}\setminus\{k_{\text{end}}\}}^{(c)2S}(s)$ by $\kappa_1^{(c)2S}(s)$ using Proposition 5.1 (ii). It is also possible

to proceed using the plug-in strategy where we replace $J(\beta)$ by an estimate \hat{J} and obtain confidence sets of level at least $1 - \gamma$ where $\gamma > \alpha$ but γ is very close to α under the assumption that the non-zero coefficients are sufficiently separated from zero.

Rates of convergence and model selection results, similar to those of Section 7, could also be obtained.

9. MODELS WITH POSSIBLY NON-VALID INSTRUMENTS

In this section, we study the problem of checking for instrument exogeneity when there is overidentification. This is a classic problem in econometrics (see, *e.g.*, Sargan (1958) and Basmann (1960) for the linear IV model, and Hansen (1982) for GMM). See also Andrews (1999) and Liao (2010) where one can find more references. We propose a two-stage method based on the *STIV* estimator. The main purpose of the suggested method is to construct confidence sets for non-validity indicators, and to detect non-valid (*i.e.*, endogenous) instruments in the high-dimensional framework. We restrict our attention to the case where (1.1) is point identified, *i.e.* $\mathcal{I}dent = \{\beta^*\}$. The model can be written in the form:

$$(9.1) \quad y_i = x_i^T \beta^* + u_i,$$

$$(9.2) \quad \mathbb{E}[z_i u_i] = 0,$$

$$(9.3) \quad \mathbb{E}[\bar{z}_i u_i] = \theta^*,$$

where x_i , z_i , and \bar{z}_i are vectors of dimensions K , L and L_1 , respectively. For simplicity of the discussion, we assume in this section that (9.1)-(9.2) is point identified. for this reason we use the notation β^* instead of β . The instruments are decomposed in two parts, z_i and \bar{z}_i , where $\bar{z}_i^T = (\bar{z}_{1i}, \dots, \bar{z}_{L_1 i})$ is a vector of possibly non-valid instruments. A component of the unknown vector $\theta^* \in \mathbb{R}^{L_1}$ is equal to zero when the corresponding instrument is indeed valid. The component θ_l^* of θ^* will be called the *non-validity indicator* of the instrument \bar{z}_{li} . Our study covers the models with dimensions K , L and L_1 that can be much larger than the sample size.

As above, we assume independence and allow for heteroscedasticity. The difference from the previous sections is only in introducing equation (9.3). We observe realizations of independent random vectors $(x_i^T, y_i, z_i^T, \bar{z}_i^T)$, $i = 1, \dots, n$. The components \bar{z}_{li} satisfy $\mathbb{E}[\bar{z}_{li} u_i] = \theta_l^*$ for all $l = 1, \dots, L_1$, $i = 1, \dots, n$. We denote by $\bar{\mathbf{Z}}$ the matrix of dimension $n \times L_1$ with rows \bar{z}_i^T , $i = 1, \dots, n$. Set

$$\bar{z}_* = \max_{l=1, \dots, L_1} \left(\frac{1}{n} \sum_{i=1}^n \bar{z}_{li}^2 \right)^{1/2}.$$

In this section, we assume that we have a pilot estimator $\widehat{\beta}$ and a statistic \widehat{b} such that, with probability close to 1,

$$(9.4) \quad \left| \mathbf{D}_{\mathbf{X}}^{-1}(\widehat{\beta} - \beta^*) \right|_1 \leq \widehat{b}.$$

For example, $\widehat{\beta}$ can be the *STIV* estimator based only on the vectors of valid instruments z_1, \dots, z_n . In this case, an explicit expression for \widehat{b} can be obtained from (6.11)¹⁸, (7.5), (8.2) or (8.9), depending on the situation.

We define the *STIV-NV* estimator $(\widehat{\theta}^{(c)}, \widehat{\sigma}_1^{(c)})$ as any solution of the problem

$$(9.5) \quad \min_{(\theta, \sigma_1) \in \widehat{\mathcal{I}}_1} (|\theta|_1 + c\sigma_1),$$

where $0 < c < 1$,

$$\widehat{\mathcal{I}}_1 \triangleq \left\{ (\theta, \sigma_1) : \theta \in \mathbb{R}^{L_1}, \sigma_1 > 0, \left| \frac{1}{n} \overline{\mathbf{Z}}^T (\mathbf{Y} - \mathbf{X}\widehat{\beta}) - \theta \right|_{\infty} \leq \sigma_1 r_1 + \widehat{b} \overline{z}_*, F(\theta, \widehat{\beta}) \leq \sigma_1 + \widehat{b} \overline{z}_* \right\}$$

for some $r_1 > 0$ (to be specified below), where for all $\theta = (\theta_1, \dots, \theta_{L_1}) \in \mathbb{R}^{L_1}$, $\beta \in \mathbb{R}^K$,

$$F(\theta, \beta) \triangleq \max_{l=1, \dots, L_1} \sqrt{\widehat{Q}_l(\theta_l, \beta)}$$

with

$$\widehat{Q}_l(\theta_l, \beta) \triangleq \frac{1}{n} \sum_{i=1}^n (\overline{z}_{li}(y_i - x_i^T \beta) - \theta_l)^2.$$

It is not hard to see that the optimization problem (9.5) can be re-written as a conic program.

The following theorem provides a basis for constructing confidence sets for the non-validity indicators.

Theorem 9.1. *Under one of the 5 scenarios of Section 6.1, replacing z_{li} by \overline{z}_{li} and L by L_1 , together with its respective assumption and choice of r with α replaced by α_1 , when $\widehat{\beta}$ is an estimator satisfying (9.4) with probability at least $1 - \alpha_2$ for some $0 < \alpha_2 < 1$, then, with probability at least $1 - \alpha_1 - \alpha_2$ (approximately at least $1 - \alpha_1 - \alpha_2$ for scenario 4 and the two-stage procedure with scenario 5), for any c in $(0, 1)$, for any solution $(\widehat{\theta}^{(c)}, \widehat{\sigma}_1^{(c)})$ of the minimization problem (9.5), we have*

$$(9.6) \quad |\widehat{\theta}^{(c)} - \theta^*|_{\infty} \leq \frac{2 \left[\widehat{\sigma}_1^{(c)} r_1 + (1 + r_1(1 - c)^{-1}) \widehat{b} \overline{z}_* \right]}{(1 - 2r_1(1 - c)^{-1} |J(\theta^*)|)_+} \triangleq V(\widehat{\sigma}_1, c, \widehat{b}, |J(\theta^*)|),$$

and

$$(9.7) \quad |\widehat{\theta}^{(c)} - \theta^*|_1 \leq \frac{2 \left[2|J(\theta^*)| \left(\widehat{\sigma}_1^{(c)} r_1 + (1 + r_1) \widehat{b} \overline{z}_* \right) + c \widehat{b} \overline{z}_* \right]}{(1 - c - 2r_1 |J(\theta^*)|)_+}.$$

¹⁸(6.13) is not explicit.

This theorem should be naturally applied when r_1 is small, *i.e.*, $n \gg \log(L_1)$. In addition, we need a small \widehat{b} , which is guaranteed by the results of Section 6 under the condition $n \gg \log(L)$ if the pilot estimator $\widehat{\beta}$ is the *STIV* estimator. Note also that the bounds (9.6) and (9.7) are meaningful if their denominators are positive, which is roughly equivalent to the following bound on the sparsity of θ^* : $|J(\theta^*)| = O(1/r_1) = O(\sqrt{n/\log(L_1)})$.

Bounds for all the norms $|\widehat{\theta}^{(c)} - \theta^*|_p$, $\forall 1 < p < \infty$, follow immediately from (9.6) and (9.7) by the standard interpolation argument. We note that, in Theorem 9.1, $\widehat{\beta}^{(c)}$ can be any estimator satisfying (9.4), not necessarily the *STIV* estimator.

We now consider that c in $(0, 1)$ is fixed. To turn (9.6) and (9.7) into valid confidence bounds, we can replace there $|J(\theta^*)|$ by $|J(\widehat{\theta}^{(c)})|$, as follows from Theorem 9.3 (ii) below. In addition, Theorem 9.3 establishes the rate of convergence of the *STIV-NV* estimator and justifies the selection of non-valid instruments by thresholding. To state the theorem, we will need an extra assumption that the random variable $F(\theta^*, \beta^*)$ is bounded in probability by a constant $\sigma_{1*} > 0$:

Assumption 9.2. *There exist constants $\sigma_{1*} > 0$ and $0 < \varepsilon < 1$ such that, with probability at least $1 - \varepsilon$,*

$$(9.8) \quad \max_{l=1, \dots, L_1} \frac{1}{n} \sum_{i=1}^n (\bar{z}_{li} u_i - \theta_l^*)^2 \leq \sigma_{1*}^2.$$

As in (7.7) we define a thresholded estimator

$$(9.9) \quad \widetilde{\theta}_l^{(c)} \triangleq \begin{cases} \widehat{\theta}_l^{(c)} & \text{if } |\widehat{\theta}_l^{(c)}| > \omega^{(c)}, \\ 0 & \text{otherwise,} \end{cases}$$

where $\omega^{(c)} > 0$ is some threshold. For $b_* > 0$, $s_1 > 0$, define

$$\bar{\sigma}_* = \left(1 - \frac{4r_1 s_1}{c(1-c-2r_1 s_1)_+} \right)_+^{-1} \left[\sigma_{1*} + \frac{2b_* \bar{z}_* (1 + 2(1+r_1)s_1/c)}{(1-c-2r_1 s_1)_+} \right].$$

Theorem 9.3. *Let the assumptions of Theorem 9.1 and Assumption 9.2 be satisfied. Then the following holds.*

(i) *Let $\widehat{\beta}$ be an estimator satisfying*

$$(9.10) \quad \left| \mathbf{D}_{\mathbf{X}}^{-1}(\widehat{\beta} - \beta^*) \right|_1 \leq b_*$$

with probability at least $1 - \alpha_2$ for some $0 < \alpha_2 < 1$ and some constant b_ . Assume that $|J(\theta^*)| \leq s_1$. Then, with probability at least $1 - \alpha_1 - \alpha_2 - \varepsilon$, for any solution $\widehat{\theta}^{(c)}$ of the*

minimization problem (9.5) we have

$$(9.11) \quad |\widehat{\theta}^{(c)} - \theta^*|_\infty \leq V(\bar{\sigma}_*, c, b_*, s_1).$$

- (ii) Let $(\widehat{\beta}, \widehat{\sigma})$ be the STIV estimator, and let the assumptions of all the items of Theorem 7.4 be satisfied (with $p = 1$ in item (ii)). Assume that $|J(\theta^*)| \leq s_1$, $|J(\beta^*)| \leq s$, and $|\theta_l^*| > V(\bar{\sigma}_*, c, b_*, s_1)$ for all $l \in J(\theta^*)$, where

$$(9.12) \quad b_* = \frac{2\sigma_* r s \tau^{(c, J)^*}(s)}{c_1}.$$

Then, with probability at least $1 - \alpha_1 - \varepsilon - \gamma$, for any solution $\widehat{\theta}^{(c)}$ of the minimization problem (9.5) we have

$$(9.13) \quad J(\theta^*) \subseteq J(\widehat{\theta}^{(c)}).$$

- (iii) Let the assumptions of item (ii) and Assumption 7.6 hold. Assume that $|\theta_l^*| > 2V(\bar{\sigma}_*, c, b_*, s_1)$ for all $l \in J(\theta^*)$. Let $\tilde{\theta}$ be the thresholded estimator defined in (9.9) where $\widehat{\theta}^{(c)}$ is any solution of the minimization problem (9.5), and the threshold is defined by $\omega^{(c)} = V(\widehat{\sigma}_1, c, \widehat{b}, s_1)$ with

$$\widehat{b} = \frac{2\widehat{\sigma} r s}{\kappa_1(s)} \left(1 - \frac{r}{\kappa_1(s)} \right)_+^{-1}.$$

Then, with probability at least $1 - \alpha_1 - \varepsilon - \gamma$, we have

$$(9.14) \quad \overrightarrow{\text{sign}(\widehat{\theta}^{(c)})} = \overrightarrow{\text{sign}(\theta^*)}.$$

As a consequence, $J(\widehat{\theta}^{(c)}) = J(\theta^*)$.

In practice, the parameter s may not be known and it can be replaced by $|J(\widehat{\theta})|$; this is a reasonable upper bound on $|J(\theta^*)|$ as suggested by Theorem 9.3 (ii). It is interesting to analyze the dependence of the rate of convergence in (9.11) on r, r_1, s , and s_1 . As discussed above, a meaningful framework is to consider small r, r_1 and the sparsities s, s_1 such that $rs, r_1 s_1$ are comfortably smaller than 1. In this case, the value b_* given in (9.14) is of the order $O(rs)$ and the rate of convergence in (9.11) is of the order $O(r_1) + O(rs)$. We see that the rate does not depend on the sparsity s_1 of θ^* but it does depend on the sparsity s of β^* . It is interesting to explore whether this rate is optimal, *i.e.*, whether it can be improved by estimators different from the STIV-NV estimator.

10. PRACTICAL IMPLEMENTATION

10.1. Computational Aspects. For simplicity of exposition, in this section we only consider the case where $I = \{i_0\}$ and i_0 is the index of unity. Extension to several cones is easy. Finding a solution $(\widehat{\beta}^{(c, \{i_0\})}, \widehat{\sigma}^{(c, \{i_0\})})$ of the minimization problem (3.5) reduces to the following conic program: find $\beta \in \mathbb{R}^K$ and $t > 0$ ($\sigma = t/\sqrt{n}$), which achieve the minimum

$$(10.1) \quad \min_{(\beta, t, v, w) \in \mathcal{V}} \left(\sum_{k=1}^K w_k + c \frac{t}{\sqrt{n}} \right)$$

where \mathcal{V} is the set of (β, t, v, w) , with satisfying:

$$\begin{aligned} v &= \mathbf{Y} - \mathbf{X}\beta, & -rt\mathbf{1} &\leq \frac{1}{\sqrt{n}}\mathbf{D}_z\mathbf{Z}^T(\mathbf{Y} - \mathbf{X}\beta) \leq rt\mathbf{1}, \\ -w &\leq \mathbf{D}_x^{-1}\beta \leq w, & w &\geq \mathbf{0}, & (t, v) &\in C. \end{aligned}$$

Here and below $\mathbf{0}$ and $\mathbf{1}$ are vectors of zeros and ones respectively, the inequality between vectors is understood in the componentwise sense, and C is a cone: $C \triangleq \{(t, v) \in \mathbb{R} \times \mathbb{R}^n : t \geq |v|_2\}$. Conic programming is a standard tool in optimization and many open source toolboxes are available to implement it (see, *e.g.*, Sturm (1999)).

The expression in curly brackets in the lower bound (5.10) is equal to the value of the following optimization program:

$$(10.2) \quad \min_{\epsilon = \pm 1} \min_{(w, \Delta, v) \in \mathcal{V}_{k,j}} v$$

where $\mathcal{V}_{k,j}$ is the set of (w, Δ, v) with $w \in \mathbb{R}^K$, $\Delta \in \mathbb{R}^K$, $v \in \mathbb{R}$ satisfying:

$$\begin{aligned} v &\geq 0, & -v\mathbf{1} &\leq \Psi_n^{(I)}\Delta \leq v\mathbf{1}, & w &\geq \mathbf{0}, & -w_{I^c} &\leq \Delta_{I^c} \leq w_{I^c} \text{ for } I = \{j, k\}, \\ w_I &= \mathbf{0}, & \Delta_k &= 1, & \epsilon\Delta_j &\geq 0, & \sum_{i=1}^K w_i + 1 &\leq \epsilon(a + g)\Delta_j \end{aligned}$$

where g is the constant such that

$$g = \begin{cases} 0 & \text{if } k = j \\ -1 & \text{otherwise.} \end{cases}$$

Note that, here, ϵ is the sign of Δ_j , and (10.2) is the minimum of two terms, each of which is the value of a linear program. Analogously, the expression in curly brackets in (5.13) can be computed by solving $2^{|J_0|}$ linear programs. The reduction is done in the same way as in (10.2) with the only difference that instead of ϵ we introduce a vector $(\epsilon_k)_{k \in J_0}$ of signs of the coordinates Δ_k for indices $k \in J_0$.

The coordinate-wise sensitivities

$$\kappa_{k,J}^{(c,\{i_0\})^*} = \inf_{\Delta_k=1, |\Delta_{J^c}|_1 \leq \frac{1+c}{1-c} |\Delta_J|_1} \left| \Psi_n^{(I)} \Delta \right|_\infty$$

can be efficiently computed for given J when the cardinality $|J|$ is small. Indeed, it is enough to find the minimum of the values of $2^{|J|}$ linear programs:

$$(10.3) \quad \min_{(\epsilon_j)_{j \in J} \in \{-1,1\}^{|J|}} \min_{(w, \Delta, v) \in \mathcal{U}_{k,J}} v$$

where $\mathcal{U}_{k,J}$ is the set of (w, Δ, v) with $w \in \mathbb{R}^K$, $\Delta \in \mathbb{R}^K$, $v \in \mathbb{R}$ satisfying:

$$\begin{aligned} v \geq 0, \quad -v \mathbf{1} \leq \Psi_n^{(I)} \Delta \leq v \mathbf{1}, \quad w \geq \mathbf{0}, \quad -w_{I^c} \leq \Delta_{I^c} \leq w_{I^c} \quad \text{for } I = J \cup \{k\}, \\ w_I = \mathbf{0}, \quad \Delta_k = 1, \quad \epsilon_j \Delta_j \geq 0, \quad \text{for } j \in J, \quad \sum_{i=1}^K w_i \leq \frac{1+c}{1-c} \sum_{j \in J} \epsilon_j \Delta_j + g. \end{aligned}$$

Here $(\epsilon_j)_{j \in J}$ is the vector of signs of the coordinates Δ_j with $j \in J$ and g is the constant defined by

$$g = \begin{cases} 0 & \text{if } k \in J, \\ -1 & \text{otherwise.} \end{cases}$$

10.2. Simulations. In this section, we consider the performance of the *STIV* estimator of Section 8.2 on simulated data. The model is as follows:

$$\begin{aligned} y_i &= \sum_{k=1}^K x_{ki} \beta_k^* + u_i, \\ x_{1i} &= \sum_{l=1}^{L-K+1} z_{li} \zeta_l + v_i, \\ x_{l'i} &= z_{li} \quad \text{for } l' = l - L + K \quad \text{and } l \in \{L - K + 2, \dots, L\}, \end{aligned}$$

where $(y_i, x_i^T, z_i^T, u_i, v_i)$ are i.i.d., (u_i, v_i) have the joint normal distribution

$$\mathcal{N} \left(0, \begin{pmatrix} \sigma_{\text{struct}}^2 & \rho \sigma_{\text{struct}} \sigma_{\text{end}} \\ \rho \sigma_{\text{struct}} \sigma_{\text{end}} & \sigma_{\text{end}}^2 \end{pmatrix} \right),$$

z_i^T is a vector of independent standard normal random variables, and z_i^T is independent of (u_i, v_i) . Clearly, in this model $\mathbb{E}[z_i u_i] = 0$. We take $n = 49$, $L = 50$, $K = 25$, $\sigma_{\text{struct}} = \sigma_{\text{end}} = \rho = 0.3$, $\beta^* = (1, 1, 1, 1, 1, 0, \dots, 0)^T$ and $\zeta_l = 0.15$ for $l = 1, \dots, L - K + 1$. We have 50 instruments and only 49 observations, so we are in a framework of application of high-dimensional techniques. We set $c = 0.1$ and set r according to (6.3) with $\alpha = 0.05$. The three columns on the left of Table 1 present simulation results for the *STIV* estimator. It is straightforward to see that only the first five variables

(the true support of β^*) are eligible to be considered as relevant. This set will be denoted by \widehat{J} . The second and third columns in Table 1 present the true coordinate-wise sensitivities $\kappa_{k,\widehat{J}}^*$ as well as their lower bounds $\kappa_k^*(5)$ obtained via the sparsity certificate with $s = 5$. These lower bounds are easy to compute, and we see that they yield reasonable approximations from below of the true sensitivities. The estimate $\widehat{\sigma}$ is 0.247 which is quite close to σ_{struct} . Next, based on (6.8), the fact that $J_{\text{exo}}^c = \{1\}$, and the bounds on the sensitivities in Proposition 5.1 and in (5.10) – (5.13), we have the following formulas for the confidence intervals

$$(10.4) \quad |\widehat{\beta}_k - \beta_k^*| \leq \frac{2\widehat{\sigma}r}{x_{k^*} \kappa_{k,\widehat{J}}^*} \left(1 - \right.$$

TABLE 1. Results for the *STIV* estimator without and with estimated instruments, $n = 49$

	$\hat{\beta}$ (1)	$\kappa_{k,\hat{\mathcal{J}}}^*$ (1)	$\kappa_k^*(5)$ (1)	$\hat{\beta}$ (2)	$\kappa_{k,\hat{\mathcal{J}}}^*$ (2)	$\kappa_k^*(5)$ (2)
β_1^*	1.03	0.107	0.103	1.03	0.085	0.068
β_2^*	0.98	0.308	0.157	0.98	0.367	0.075
β_3^*	0.96	0.129	0.103	0.96	0.126	0.071
β_4^*	0.95	0.150	0.109	0.95	0.115	0.057
β_5^*	0.90	0.253	0.175	0.90	0.177	0.086
β_6^*	0.00	0.166	0.095	0.00	0.126	0.065
β_7^*	0.00	0.155	0.080	0.00	0.148	0.060
β_8^*	0.00	0.154	0.110	0.00	0.122	0.056
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
β_{23}^*	0.02	0.287	0.170	0.02	0.231	0.128
β_{24}^*	0.00	0.243	0.137	0.00	0.195	0.105
β_{25}^*	0.00	0.141	0.109	0.00	0.106	0.067

We use dots because the values that do not appear are similar.

(1): With all the 50 instruments,

(2): With 25 instruments including an estimate of the linear projection instrument.

TABLE 2. Estimates of the coefficients of the linear projection instrument

$\hat{\zeta}_1$	$\hat{\zeta}_2$	$\hat{\zeta}_3$	$\hat{\zeta}_4$	$\hat{\zeta}_6$	$\hat{\zeta}_8$	$\hat{\zeta}_9$	$\hat{\zeta}_{10}$	$\hat{\zeta}_{14}$	$\hat{\zeta}_{15}$	$\hat{\zeta}_{16}$	$\hat{\zeta}_{17}$	$\hat{\zeta}_{18}$	$\hat{\zeta}_{20}$
0.084	0.130	0.190	0.142	0.115	0.083	0.104	0.126	0.176	0.030	0.023	0.157	0.135	0.082
$\hat{\zeta}_{21}$	$\hat{\zeta}_{23}$	$\hat{\zeta}_{24}$	$\hat{\zeta}_{25}$	$\hat{\zeta}_{26}$	$\hat{\zeta}_{27}$	$\hat{\zeta}_{32}$	$\hat{\zeta}_{33}$	$\hat{\zeta}_{34}$	$\hat{\zeta}_{44}$	$\hat{\zeta}_{47}$	$\hat{\zeta}_{49}$	$\hat{\zeta}_{50}$	
0.100	0.125	0.038	0.025	0.026	-0.058	0.108	0.005	-0.053	-0.006	-0.009	-0.063	0.033	

We only show the non-zero coefficients.

set to 0 and several superfluous variables are included, sometimes with significant coefficients, such as $\hat{\zeta}_{32}$. We get $\kappa_{1,\hat{\mathcal{J}}}^* = 0.0076$ and $\kappa_1^*(5) = 0.0040$. Again, $r/\kappa_{1,\hat{\mathcal{J}}}^* > 1$, so that we cannot use (10.4) and (10.5) to get the confidence intervals. Note that this approach based on the estimated linear projection instrument gives sensitivities, which are lower than with the full set of instruments. This is mainly due to the fact that the estimation of the linear projection instrument is quite imprecise. Indeed, we add an instrument \hat{x}_{i1} , which is not so good, and at the same time we drop a large number of other instruments, which may be not so bad. The overall effect on the sensitivities turns out to be negative. Recall that since the sensitivities involve the maximum of the scalar products of the rows of $\Psi_n^{(I)}$ with Δ , the more we have rows (*i.e.*, instruments) the higher is the sensitivity. The same deterioration of the sensitivities occurred in other simulated data sets. In conclusion, the approach based on estimation of the linear projection instrument was not helpful to realize the above confidence

TABLE 3. Monte-Carlo study, 1000 replications

	5 th percentile	Median	95 th percentile		5 th percentile	Median	95 th percentile
β_1^*	0.872	0.986	1.093	β_8^*	-0.057	0.000	0.055
β_2^*	0.877	0.970	1.048	β_9^*	-0.052	0.000	0.059
β_3^*	0.879	0.970	1.049	\vdots	\vdots	\vdots	\vdots
β_4^*	0.886	0.971	1.051	β_{23}^*	-0.051	0.000	0.051
β_5^*	0.877	0.968	1.049	β_{24}^*	-0.057	0.000	0.051
β_6^*	-0.048	0.000	0.055	β_{25}^*	-0.053	0.000	0.049
β_7^*	-0.059	0.000	0.063	$\hat{\sigma}$	0.181	0.233	0.291

intervals in this small sample situation. However, we will see that it achieves the task when the sample size gets large.

Although in this numerical example we were not able to use (10.4) and (10.5) for the confidence intervals, we got evidence that the performance of the *STIV* estimator is quite satisfactory. Table 3 shows a Monte-Carlo study where we keep the same values of the parameters of the model, of the sample size $n = 49$, and of the parameter A defining the set $\hat{\mathcal{L}}$, simulate 1000 data sets, and compute 1000 estimates. The empirical performance of the *STIV* estimator is extremely good, even for the endogenous variable. The Monte-Carlo estimation of the variability of $\hat{\beta}_1$ is very similar to that of the exogenous variables. With $c = 0.1$ the estimate $\hat{\sigma}$ is larger than σ_{struct} in 95% of the simulations. This suggests that there remains some margin to penalize less for the “variance” in (3.5), *i.e.*, to decrease c and thus to obtain higher sensitivities.

Next, we study the empirical behavior of the non-pivotal *STIV* estimator. We consider the same model and the same values of all the parameters, and we choose $\sigma_* = 2 \cdot 0.233$ where 0.233 is the median of $\hat{\sigma}$ from Table 3. Indeed $\mathbb{P}(\mathbb{E}_n[U^2] \leq \sigma_*^2)$ should be close to 1 (see Assumption 7.1). The results are given in Table 4. The non-pivotal procedure seems to better estimate as zeros the zero coefficients. This is because we minimize the ℓ_1 norm of the coefficients without an additional $c\sigma$ term. On the other hand, the non-zero coefficients are better estimated using the pivotal estimator. The non-pivotal procedure yields some shrinkage to zero (especially for large σ_*). Using the pivotal procedure in the first place allows us to have a good initial guess of σ_* .

Let us now increase n to see whether we can obtain interval estimates and take advantage of thresholding for variable selection. We consider the same model as above and the same values of the parameters of the method but we replace $n = 49$ by $n = 8000$. Then we are no longer in a situation where we must use specific high-dimensional techniques. However, it is still a challenging

TABLE 4. Monte-Carlo study of the non-pivotal estimator, 1000 replications

	5 th percentile	Median	95 th percentile		5 th percentile	Median	95 th percentile
β_1^*	0.714	0.914	1.110	β_8^*	-0.003	0.000	0.016
β_2^*	0.803	0.909	1.010	β_9^*	0.000	0.000	0.024
β_3^*	0.789	0.904	1.019	β_{10}^*	0.000	0.000	0.018
β_4^*	0.793	0.904	1.023	\vdots	\vdots	\vdots	\vdots
β_5^*	0.796	0.907	1.017	β_{23}^*	0.000	0.000	0.021
β_6^*	0.000	0.000	0.020	β_{24}^*	0.000	0.000	0.016
β_7^*	0.000	0.000	0.021	β_{25}^*	0.000	0.000	0.005

task to select among 25 candidate variables, one of them being endogenous. Indeed, classical selection procedures like the BIC would require to solve 2^{25} least squares problems. Our methods are much less numerically intensive. They are based on linear and conic programming, and their computational cost is polynomial in the dimension. We study both the setting with all the 50 instruments and the setting where we estimate the linear projection instrument.

Consider first the case where we use all the instruments. Set for brevity

$$\hat{w} \triangleq \left(1 - \frac{r}{\kappa_{1,\hat{J}}^*} - \frac{r^2}{\kappa_{1,\hat{J}}}\right)_+^{-1}, \quad w(5) \triangleq \left(1 - \frac{r}{\kappa_1^*(5)} - \frac{r^2}{\kappa_1(5)}\right)_+^{-1}.$$

These are the quantities appearing in (10.4) and (10.5). As above, we take \hat{J} equal to the set of the first five coordinates; $w(5)$ corresponds to the sparsity certificate approach with $s = 5$. Computing the exact coordinate-wise sensitivities we obtain the bound $\hat{w} \leq 1.6277$. The sparsity certificate approach with $s = 5$ yields $w(5) \leq 1.6306$. We obtain $\hat{\sigma} = 0.2970$ and the estimates in Table 5. The values $\hat{\beta}_{l,\hat{J}}$ and $\hat{\beta}_{u,\hat{J}}$ are the lower and upper confidence limits respectively obtained from (10.4); $\hat{\beta}_{l,SC}$ and $\hat{\beta}_{u,SC}$ are the lower and upper confidence limits obtained from (10.5) (sparsity certificate approach with $s = 5$). The thresholds $\omega_{k,\hat{J}}$ and $\omega_k(5)$ are computed from the formulas

$$\omega_{k,\hat{J}} = \frac{2 \cdot 1.6277 \hat{\sigma} r}{x_{k*} \kappa_{k,\hat{J}}^*}, \quad \omega_k(5) = \frac{2 \cdot 1.6306 \hat{\sigma} r}{x_{k*} \kappa_k^*(5)}.$$

Table 5 shows that in this example thresholding works well: The true support of β^* is recovered exactly by selecting the variables, for which the estimated coefficient is larger than the threshold. Note that the threshold for the endogenous variable is very close to the estimate of the first coefficient $\hat{\beta}_1$ since the confidence intervals are wider for the endogenous variable.

We now consider the case where we use only 25 instruments; the 24 exogenous variables serve as their own instruments and the Square-root Lasso estimator of the linear projection instrument is

TABLE 5. Confidence intervals and selection of variables, $n = 8000$

	$\hat{\beta}_{l,SC}$	$\hat{\beta}_{l,\hat{\mathcal{J}}}$	$\hat{\beta}$	$\hat{\beta}_{u,\hat{\mathcal{J}}}$	$\hat{\beta}_{u,SC}$	$\kappa_{k,\hat{\mathcal{J}}}^*$	$\kappa_k^*(5)$	$\omega_{k,\hat{\mathcal{J}}}$	$\omega_{k,SC}$
β_1^*	0.131	0.135	1.048	1.960	1.965	0.134	0.134	0.912	0.917
β_2^*	0.795	0.804	0.995	1.185	1.195	0.897	0.855	0.191	0.200
β_3^*	0.824	0.829	1.004	1.179	1.185	0.796	0.775	0.175	0.180
β_4^*	0.817	0.822	0.998	1.173	1.178	0.858	0.833	0.175	0.181
β_5^*	0.833	0.834	1.001	1.168	1.168	0.793	0.790	0.167	0.168
β_6^*	-0.163	-0.160	0.003	0.166	0.169	0.807	0.791	0.163	0.166
β_7^*	-0.173	-0.168	0.002	0.172	0.177	0.846	0.823	0.170	0.175
β_8^*	-0.173	-0.170	0.001	0.173	0.175	0.789	0.779	0.172	0.174
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
β_{23}^*	-0.190	-0.188	0.003	0.194	0.197	0.802	0.793	0.191	0.193
β_{24}^*	-0.171	-0.166	0.001	0.168	0.172	0.842	0.821	0.167	0.172
β_{25}^*	-0.172	-0.169	-0.005	0.158	0.162	0.828	0.811	0.163	0.167

TABLE 6. Estimates of the coefficients in the linear projection instrument

$\hat{\zeta}_1$	$\hat{\zeta}_2$	$\hat{\zeta}_3$	$\hat{\zeta}_4$	$\hat{\zeta}_5$	$\hat{\zeta}_6$	$\hat{\zeta}_7$	$\hat{\zeta}_8$	$\hat{\zeta}_9$	$\hat{\zeta}_{10}$	$\hat{\zeta}_{11}$	$\hat{\zeta}_{12}$	$\hat{\zeta}_{13}$	$\hat{\zeta}_{14}$
0.142	0.145	0.134	0.136	0.137	0.135	0.139	0.139	0.134	0.140	0.146	0.140	0.134	0.136
$\hat{\zeta}_{15}$	$\hat{\zeta}_{16}$	$\hat{\zeta}_{17}$	$\hat{\zeta}_{18}$	$\hat{\zeta}_{19}$	$\hat{\zeta}_{20}$	$\hat{\zeta}_{21}$	$\hat{\zeta}_{22}$	$\hat{\zeta}_{23}$	$\hat{\zeta}_{24}$	$\hat{\zeta}_{25}$	$\hat{\zeta}_{26}$		
0.137	0.138	0.141	0.128	0.142	0.137	0.133	0.135	0.135	0.142	0.137	0.138		

We only show the non-zero coefficients (keeping only three digits).

used for the endogenous variable. This time, we apply the Square-root Lasso with the recommended choice $c_{\sqrt{\text{Lasso}}} = 1.1$. We get $\sqrt{\hat{Q}(\hat{\beta})} = 0.3012$. The estimates of $\hat{\zeta}_l$ are given in Table 6. Next, we use (10.4) and (10.5) to obtain the confidence intervals. Computing the exact coordinate-wise sensitivities we get the bound $\hat{w} \leq 1.0941$. The sparsity certificate approach with $s = 5$ yields $w(5) \leq 1.0990$. We also get $\hat{\sigma} = 0.2970$. The thresholds $\omega_{k,\hat{\mathcal{J}}}$ and $\omega_k(5)$ are obtained from the formulas

$$\omega_{k,\hat{\mathcal{J}}} = \frac{2 \cdot 1.0941 \hat{\sigma} r}{x_{k*} \kappa_{k,\hat{\mathcal{J}}}^*}, \quad \omega_k(5) = \frac{2 \cdot 1.0990 \hat{\sigma} r}{x_{k*} \kappa_k^*(5)}.$$

The results are summarized in Table 7. Note that the confidence intervals and the thresholds are sharper than in the approach including all the instruments. The particularly good news is that the confidence interval for the coefficient of the endogenous variable becomes much tighter.

In conclusion, when the sample size is large, the coordinate-wise sensitivities based on the sparsity certificate work remarkably well for estimation, confidence intervals, and variable selection.

TABLE 7. Confidence intervals and selection of variables, $n = 8000$

	$\hat{\beta}_{l,SC}$	$\hat{\beta}_{l,\hat{J}}$	$\hat{\beta}$	$\hat{\beta}_{u,\hat{J}}$	$\hat{\beta}_{u,SC}$	$\kappa_{k,\hat{J}}^*$	$\kappa_k^*(5)$	$\omega_{k,\hat{J}}$	$\omega_{k,SC}$
β_1^*	0.901	0.909	1.048	1.187	1.194	0.556	0.531	0.139	0.146
β_2^*	0.872	0.883	0.995	1.106	1.118	0.968	0.882	0.111	0.123
β_3^*	0.896	0.905	1.004	1.103	1.112	0.888	0.821	0.099	0.108
β_4^*	0.885	0.893	0.998	1.102	1.110	0.907	0.848	0.105	0.113
β_5^*	0.899	0.902	1.001	1.100	1.103	0.843	0.823	0.099	0.102
β_6^*	-0.098	-0.092	0.003	0.099	0.104	0.868	0.822	0.095	0.101
β_7^*	-0.103	-0.098	0.002	0.102	0.107	0.907	0.869	0.100	0.105
β_8^*	-0.099	-0.095	0.001	0.098	0.102	0.886	0.853	0.096	0.101
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
β_{23}^*	-0.115	-0.109	0.003	0.115	0.121	0.862	0.825	0.112	0.118
β_{24}^*	-0.104	-0.099	0.001	0.101	0.106	0.888	0.848	0.100	0.105
β_{25}^*	-0.109	-0.104	-0.005	0.093	0.098	0.870	0.830	0.098	0.103

We also get a significant improvement from using the two-stage procedure with estimated linear projection instrument.

11. APPENDIX

11.1. **Lower Bounds on $\kappa_{p,J}^{(c,I)}$ for Square Matrices $\Psi_n^{(I)}$.** In this section we consider the case where all regressors are exogenous and we take $\mathbf{D}_Z^{(I)} = \mathbf{D}_X$. We therefore drop the exponent I . The following propositions establish lower bounds on $\kappa_{p,J}^{(c)}$ when Ψ_n is a square $K \times K$ matrix. For any $J \subseteq \{1, \dots, K\}$ we define the following restricted eigenvalue (RE) constants

$$\kappa_{\text{RE},J}^{(c)} \triangleq \inf_{\Delta \in \mathbb{R}^K \setminus \{0\}: \Delta \in C_J^{(c)}} \frac{|\Delta^T \Psi_n \Delta|}{|\Delta_J|_2^2}, \quad \kappa'_{\text{RE},J}^{(c)} \triangleq \inf_{\Delta \in \mathbb{R}^K \setminus \{0\}: \Delta \in C_J^{(c)}} \frac{|J| |\Delta^T \Psi_n \Delta|}{|\Delta_J|_1^2}.$$

Proposition 11.1. *For any $J \subseteq \{1, \dots, K\}$ and c in $(0, 1)$ we have*

$$\kappa_{1,J}^{(c)} \geq \frac{(1-c)^2}{4|J|} \kappa'_{\text{RE},J}^{(c)} \geq \frac{(1-c)^2}{4|J|} \kappa_{\text{RE},J}^{(c)}.$$

Proof. For such that $|\Delta_{J^c}|_1 \leq \frac{1+c}{1-c} |\Delta_J|_1$ we have $|\Delta|_1 \leq \frac{2}{1-c} |\Delta_J|_1$. Thus,

$$\frac{|\Delta^T \Psi_n \Delta|}{|\Delta_J|_1^2} \leq \frac{|\Delta|_1 |\Psi_n \Delta|_\infty}{|\Delta_J|_1^2} \leq \frac{4}{(1-c)^2} \frac{|\Psi_n \Delta|_\infty}{|\Delta|_1}.$$

This proves the first inequality of the proposition. The second inequality is obvious. \square

Proposition 11.2. *Let $J \subseteq \{1, \dots, K\}$ and c in $(0, 1)$ be such that*

$$(11.1) \quad \inf_{\Delta \in \mathbb{R}^K \setminus \{0\}: \Delta \in C_J^{(c)}} \frac{|\mathbf{X}\mathbf{D}_\mathbf{X}\Delta|_2}{\sqrt{n}|\Delta_J|_2} \geq \tilde{\kappa}^{(c)}$$

for some $\tilde{\kappa} > 0$, and let there exist $0 < \delta < 1$ such that

$$(11.2) \quad \left| \frac{1}{n} (\mathbf{X}\mathbf{D}_\mathbf{X} - \mathbf{Z}\mathbf{D}_\mathbf{Z})^T \mathbf{X}\mathbf{D}_\mathbf{X} \right|_\infty \leq \frac{\delta(1-c)^2(\tilde{\kappa}^{(c)})^2}{4|J|}.$$

Then

$$\kappa_{1,J}^{(c)} \geq \frac{(1-\delta)(1-c)^2(\tilde{\kappa}^{(c)})^2}{4|J|}.$$

Proof. We have

$$\begin{aligned} |\Psi_n \Delta|_\infty |\Delta|_1 &\geq |\Delta^T \Psi_n \Delta| \\ &\geq \left| \Delta^T \frac{1}{n} \mathbf{D}_\mathbf{X} \mathbf{X}^T \mathbf{X} \mathbf{D}_\mathbf{X} \Delta \right| - \left| \Delta^T \frac{1}{n} (\mathbf{X}\mathbf{D}_\mathbf{X} - \mathbf{Z}\mathbf{D}_\mathbf{Z})^T \mathbf{X}\mathbf{D}_\mathbf{X} \Delta \right| \end{aligned}$$

where

$$\begin{aligned} \left| \Delta^T \frac{1}{n} (\mathbf{X}\mathbf{D}_\mathbf{X} - \mathbf{Z}\mathbf{D}_\mathbf{Z})^T \mathbf{X}\mathbf{D}_\mathbf{X} \Delta \right| &\leq \left| \frac{1}{n} (\mathbf{X}\mathbf{D}_\mathbf{X} - \mathbf{Z}\mathbf{D}_\mathbf{Z})^T \mathbf{X}\mathbf{D}_\mathbf{X} \right|_\infty |\Delta|_1^2 \\ &\leq \frac{\alpha(1-c)^2 \tilde{\kappa}^2}{4|J|} |\Delta|_1^2. \end{aligned}$$

Combining these inequalities and using that $|\Delta|_1^2 \leq \frac{4}{(1-c)^2} |J| |\Delta_J|_2^2$ for all $\Delta \in C_J^{(c)}$ (cf. proof of Proposition 11.2) we get the result. \square

Relation (11.2) accounts for the closeness between the normalized instruments and the normalized regressors. In the case where there is only one endogenous regressor in the structural equation and all exogenous regressors are used as their own instrument, $\mathbf{X}\mathbf{D}_\mathbf{X}$ and $\mathbf{Z}\mathbf{D}_\mathbf{Z}$ only differ through one column. They are equal if there is no endogeneity. In that case we are left with the usual condition (11.1). It is the restricted eigenvalue condition of Bickel, Ritov and Tsybakov (2009) for the Gram matrix of X -variables, up to the normalization by $\mathbf{D}_\mathbf{X}$.

We now obtain bounds for sensitivities $\kappa_{p,J}^{(c)}$ with $1 < p \leq 2$ and c in $(0, 1)$. For any $s \leq K$, we consider a uniform version of the restricted eigenvalue constant: $\kappa_{\text{RE}}(s) \triangleq \min_{|J| \leq s} \kappa_{\text{RE},J}^{(c)}$.

Proposition 11.3. *For any $s \leq K/2$ and $1 < p \leq 2$, we have*

$$\kappa_{p,J}^{(c)} \geq C(p) s^{-1/p} \kappa_{\text{RE}}^{(c)}(2s), \quad \forall J: |J| \leq s,$$

where $C(p) = 2^{-1/p-1/2} (1-c) \left(1 + \frac{1+c}{1-c} (p-1)^{-1/p}\right)^{-1}$.

Proof. For $\Delta \in R^K$ and a set $J \subset \{1, \dots, K\}$, let $J_1 = J_1(\Delta, J)$ be the subset of indices in $\{1, \dots, K\}$ corresponding to the s largest in absolute value components of Δ outside of J . Define $J_+ = J \cup J_1$. If $|J| \leq s$ we have $|J_+| \leq 2s$. It is easy to see that the k th largest absolute value of elements of Δ_{J^c} satisfies $|\Delta_{J^c}|_{(k)} \leq |\Delta_{J^c}|_1/k$. Thus,

$$|\Delta_{J_+^c}|_p^p \leq |\Delta_{J^c}|_1^p \sum_{k \geq s+1} \frac{1}{k^p} \leq \frac{|\Delta_{J^c}|_1^p}{(p-1)s^{p-1}}.$$

For $\Delta \in C_J^{(c)}$, this implies

$$|\Delta_{J_+^c}|_p \leq \frac{|\Delta_{J^c}|_1}{(p-1)^{1/p}s^{1-1/p}} \leq \frac{c_0|\Delta_J|_1}{(p-1)^{1/p}s^{1-1/p}} \leq \frac{c_0|\Delta_J|_p}{(p-1)^{1/p}},$$

where $c_0 = \frac{1+c}{1-c}$. Therefore, for $\Delta \in C_J^{(c)}$,

$$(11.3) \quad |\Delta|_p \leq (1 + c_0(p-1)^{-1/p})|\Delta_{J_+}|_p \leq (1 + c_0(p-1)^{-1/p})(2s)^{1/p-1/2}|\Delta_{J_+}|_2.$$

Using (11.3) and the fact that $|\Delta|_1 \leq \frac{2}{1-c}|\Delta_J|_1 \leq \frac{2\sqrt{s}}{1-c}|\Delta_J|_2$ for $\Delta \in C_J^{(c)}$, we get

$$\begin{aligned} \frac{|\Delta^T \Psi_n \Delta|}{|\Delta_{J_+}|_2^2} &\leq \frac{|\Delta|_1 |\Psi_n \Delta|_\infty}{|\Delta_{J_+}|_2^2} \\ &\leq \frac{2\sqrt{s} |\Psi_n \Delta|_\infty}{(1-c) |\Delta_{J_+}|_2} \\ &\leq \frac{s^{1/p} |\Psi_n \Delta|_\infty}{C(p) |\Delta|_p}. \end{aligned}$$

Since $|J_+| \leq 2s$, this proves the proposition. \square

The lower bounds in Propositions 11.1 and 11.3 require to control from below $|\Delta^T \Psi_n \Delta|$ (where Ψ_n is a non-symmetric possibly non-positive definite matrix) by a quadratic form with many zero eigenvalues for vectors in a cone of dominant coordinates. This is potentially a strong restriction on the instruments that we can use. In other words, the sensitivity characteristics $\kappa_{p,J}^{(c)}$ can be much larger than the above bounds. The propositions of this section imply that, even in the case of symmetric matrices, these characteristics are more general and potentially lead to better results than the restricted eigenvalues $\kappa_{\text{RE}}^{(c)}(\cdot)$ appearing in the usual RE condition of Bickel, Ritov and Tsybakov (2009).

11.2. Moderate Deviations for Self-normalized Sums. Throughout this section X_1, \dots, X_n are independent random variables such that, for every i , $\mathbb{E}[X_i] = 0$. The following result is due to Efron (1969).

Theorem 11.4. *If X_i for $i = 1, \dots, n$ are symmetric, then for every r positive,*

$$\mathbb{P} \left(\frac{\left| \frac{1}{n} \sum_{i=1}^n X_i \right|}{\sqrt{\frac{1}{n} \sum_{i=1}^n X_i^2}} \geq r \right) \leq 2 \exp \left(-\frac{nr^2}{2} \right).$$

This upper bound is refined in Pinelis (1994) for i.i.d. random variables.

Theorem 11.5. *If X_i for $i = 1, \dots, n$ are symmetric and identically distributed, then for every r in $[0, 1)$,*

$$\mathbb{P} \left(\frac{\left| \frac{1}{n} \sum_{i=1}^n X_i \right|}{\sqrt{\frac{1}{n} \sum_{i=1}^n X_i^2}} \geq r \right) \leq \frac{4e^3}{9} \Phi(-\sqrt{nr}).$$

The following result is from Jing, Shao and Wang (2003).

Theorem 11.6. *Assume that $0 < \mathbb{E}[|X_i|^{2+\delta}] < \infty$ for some $0 < \delta \leq 1$ and set*

$$B_n^2 = \sum_{i=1}^n \mathbb{E}[X_i^2], \quad L_{n,\delta} = \sum_{i=1}^n \mathbb{E} \left[|X_i|^{2+\delta} \right], \quad d_{n,\delta} = B_n / L_{n,\delta}^{1/(2+\delta)}.$$

Then

$$\forall 0 \leq r \leq \frac{d_{n,\delta}}{\sqrt{n}}, \quad \mathbb{P} \left(\frac{\left| \frac{1}{n} \sum_{i=1}^n X_i \right|}{\sqrt{\frac{1}{n} \sum_{i=1}^n X_i^2}} \geq r \right) \leq 2\Phi(-\sqrt{nr}) \left(1 + A_0 \left(\frac{1 + \sqrt{nr}}{d_{n,\delta}} \right)^{2+\delta} \right)$$

where $A_0 > 0$ is an absolute constant.

Despite of its great interest to understand the large deviations behavior of self-normalized sums, the bound has limited practical use because A_0 is not an explicit constant.

The following result is a corollary of Theorem 1 in Bertail, Gauth rat and Harari-Kermadec (2009).

Theorem 11.7. *Assume that X_i for $i = 1, \dots, n$ are identically distributed and $0 < \mathbb{E}[|X_i|^4] < \infty$.*

Then

$$(11.4) \quad \forall r \geq 0, \quad \mathbb{P} \left(\frac{\left| \frac{1}{n} \sum_{i=1}^n X_i \right|}{\sqrt{\frac{1}{n} \sum_{i=1}^n X_i^2}} \geq r \right) \leq (2e + 1) \exp \left(-\frac{nr^2}{2 + \gamma_4 r^2} \right)$$

where $\gamma_4 = \frac{\mathbb{E}[X_i^4]}{\mathbb{E}[X_i^2]^2}$, while

$$\forall r \geq \sqrt{n}, \quad \mathbb{P} \left(\frac{\left| \frac{1}{n} \sum_{i=1}^n X_i \right|}{\sqrt{\frac{1}{n} \sum_{i=1}^n X_i^2}} \geq r \right) = 0.$$

Proof Bertail, Gauth erat and Harari-Kermadec (2009) obtain the upper bound for $r \geq \sqrt{n}$ and that for $0 \leq r < \sqrt{n}$

$$\mathbb{P} \left(\frac{|\frac{1}{n} \sum_{i=1}^n X_i|}{\sqrt{\frac{1}{n} \sum_{i=1}^n X_i^2}} \geq r \right) \leq \inf_{a>1} \left\{ 2e \exp \left(-\frac{nr^2}{2(1+a)} \right) + \exp \left(-\frac{n}{2\gamma_4} \left(1 - \frac{1}{a} \right)^2 \right) \right\}.$$

Because

$$\frac{1}{1+a} = \frac{1}{a} \frac{1}{1+\frac{1}{a}} \geq \frac{1}{a} \left(1 - \frac{1}{a} \right)$$

we obtain

$$-\frac{r^2}{1+a} \leq -\frac{r^2}{a} \left(1 - \frac{1}{a} \right).$$

This yields (11.4) by choosing a to equate the two exponential terms. \square

11.3. Proofs. Proof of Proposition 5.1. Parts (i), (ii) and (iv) of the proposition are straightforward. The upper bound in (5.8) follows immediately from (5.1). Next, obviously, $|\Delta|_p \leq |\Delta|_1^{1/p} |\Delta|_\infty^{1-1/p}$ and we get that, for $\Delta \neq 0$,

$$\frac{|\Psi_n^{(I)} \Delta|_\infty}{|\Delta|_p} \geq \frac{|\Psi_n^{(I)} \Delta|_\infty}{|\Delta|_\infty} \left(\frac{|\Delta|_\infty}{|\Delta|_1} \right)^{1/p}.$$

Furthermore, (5.1) implies $|\Delta|_1 \leq \frac{2}{1-c} |J| |\Delta|_\infty$ for $\Delta \in C_J$. Combining this with the above inequality we obtain the lower bound in (5.8). The sequence of inequalities in (5.9) follow from the fact that

$$|J_0| - 1 + 1/p |\Delta_{J_0}|_1 \leq |\Delta_{J_0}|_p \leq |J_0| 1/p |\Delta_{J_0}|_1.$$

\square

Proof of Proposition 5.2. For all $1 \leq k \leq K$ and $1 \leq l \leq L$,

$$\left| \left(\Psi_n^{(I)} \Delta \right)_l - (\Psi_n^{(I)})_{lk} \Delta_k \right| \leq |\Delta|_1 \max_{k' \neq k} |(\Psi_n^{(I)})_{lk'}|,$$

which yields

$$\left| (\Psi_n^{(I)})_{lk} \right| |\Delta_k| \leq |\Delta|_1 \max_{k' \neq k} |(\Psi_n^{(I)})_{lk'}| + \left| \left(\Psi_n^{(I)} \Delta \right)_l \right|.$$

The two inequalities of the assumption yield

$$\left| (\Psi_n^{(I)})_{l(k)k} \right| |\Delta_k| \leq |\Delta|_1 \frac{(1-\eta_2)(1-c)}{2|J|} |(\Psi_n^{(I)})_{l(k)k}| + \frac{1-c}{\eta_1} \left| \left(\Psi_n^{(I)} \Delta \right)_{l(k)} \right| \left| (\Psi_n^{(I)})_{l(k)k} \right|.$$

This inequality, together with the fact that $\left| \left(\Psi_n^{(I)} \Delta \right)_{l(k)} \right| \leq \left| \Psi_n^{(I)} \Delta \right|_\infty$, we obtain

$$(11.5) \quad |\Delta_j| \leq |\Delta|_1 \frac{(1-\eta_2)(1-c)}{2|J|} + \frac{1-c}{\eta_1} \left| \Psi_n^{(I)} \Delta \right|_\infty$$

Summing the inequalities over j in J , yields

$$|\Delta_J|_1 \leq \frac{(1-\eta_2)(1-c)}{2} |\Delta|_1 + \frac{|J|(1-c)}{\eta_1} \left| \Psi_n^{(I)} \Delta \right|_\infty.$$

This and the first inequality in (5.1) imply that

$$\frac{1-c}{2} |\Delta|_1 \leq \frac{(1-\eta_2)(1-c)}{2} |\Delta|_1 + \frac{|J|(1-c)}{\eta_1} \left| \Psi_n^{(I)} \Delta \right|_\infty$$

which yields

$$\frac{\eta_2(1-c)}{2} |\Delta|_1 \leq \frac{|J|(1-c)}{\eta_1} \left| \Psi_n^{(I)} \Delta \right|_\infty$$

and

$$(11.6) \quad \frac{\eta_1 \eta_2}{2|J|} |\Delta|_1 \leq \left| \Psi_n^{(I)} \Delta \right|_\infty.$$

We conclude, using the definition of the l_1 -sensitivity, that

$$(11.7) \quad \kappa_{1,J} = \frac{\eta_1 \eta_2}{2|J|}.$$

Next, plugging (11.6) into (11.5), we deduce

$$\begin{aligned} |\Delta_j| &\leq \left(\frac{1-\eta_2}{\eta_1 \eta_2} + \frac{1}{\eta_1} \right) (1-c) \left| \Psi_n^{(I)} \Delta \right|_\infty \\ &\leq \frac{1-c}{\eta_1 \eta_2} \left| \Psi_n^{(I)} \Delta \right|_\infty, \end{aligned}$$

which implies

$$\kappa_{\infty,J} \geq \frac{\eta_1 \eta_2}{1-c}.$$

This and the lower bound in (5.8) yield the result. \square

Proof of Theorem 6.5. Fix β in \mathcal{B}_s , denote by $u_i = y_i - x_i^T \beta$ and define the event

$$\mathcal{G} = \left\{ \forall l = 1, \dots, L, \left(\mathbf{D}_Z^{(I)} \right)_l \left| \frac{1}{n} \sum_{i=1}^n z_{li} u_i \right| \leq \max_{l \in I} \left(\mathbf{D}_Z^{(I)} \right)_l \sqrt{\widehat{Q}_l(\beta)} r \right\}.$$

Note that $\widehat{Q}_l(\beta) = \mathbb{E}_n[Z_l^2 U^2]$. We have

$$\mathcal{G}^c = \left(\bigcup_{l \in I^c} \left\{ \frac{1}{n} \left| \frac{\sum_{i=1}^n z_{li} u_i}{|z_{l*} \sqrt{\mathbb{E}_n[U^2]}|} \right| \geq r \right\} \right) \cup \left(\bigcup_{l \in I} \left\{ \frac{1}{n} \left| \frac{\sum_{i=1}^n z_{li} u_i}{\sqrt{\mathbb{E}_n[Z_l^2 U^2]}} \right| \geq r \right\} \right).$$

Under scenario 1, $\mathbb{P}(\mathcal{G}^c) = \alpha$. For the other scenarios, we use

$$(11.8) \quad \mathcal{G}^c \subset \bigcup_{l=1, \dots, L} \left\{ \left| \frac{\sum_{i=1}^n z_{li} u_i}{\sqrt{\sum_{i=1}^n (z_{li} u_i)^2}} \right| \geq \sqrt{nr} \right\}.$$

The union bound yields

$$\mathbb{P}(\mathcal{G}^c) \leq \sum_{l=1}^L \mathbb{P} \left(\left| \frac{\sum_{i=1}^n z_{li} u_i}{\sqrt{\sum_{i=1}^n (z_{li} u_i)^2}} \right| \geq \sqrt{nr} \right).$$

We conclude, using the moderate deviations result from Section 11.2 that the event \mathcal{G} holds with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5) under the respective choices of r . Because the event containing \mathcal{G}^c on the right hand side of (11.17) does not depend on I , we obtain statements which are uniform in I . Because \mathcal{G}^c does not depend on c we obtain statements which are uniform in c in $(0, 1)$ under all 5 scenarios. Set $\Delta \triangleq \mathbf{D}_{\mathbf{X}}^{-1}(\widehat{\beta}^{(c,I)} - \beta)$. On the event \mathcal{G} we have:

$$(11.9) \quad \left| \Psi_n^{(I)} \Delta \right|_{\infty} \leq \left| \frac{1}{n} \mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})} \mathbf{Z}^T (\mathbf{Y} - \mathbf{X} \widehat{\beta}^{(c,I)}) \right|_{\infty} + \left| \frac{1}{n} \mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})} \mathbf{Z}^T (\mathbf{Y} - \mathbf{X} \beta) \right|_{\infty}$$

$$(11.10) \quad \begin{aligned} &\leq r \widehat{\sigma}^{(c,I)} + \left| \frac{1}{n} \mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})} \mathbf{Z}^T \mathbf{U} \right|_{\infty} \\ &\leq r \left(\widehat{\sigma}^{(c,I)} + \max_{l \in I} (\mathbf{D}_{\mathbf{Z}})_{ll} \sqrt{\widehat{Q}_l(\beta)} \right). \end{aligned}$$

The inequality (11.10) holds because $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ belongs to the set $\widehat{\mathcal{I}}^{(I)}$ by definition. Notice that, on the event \mathcal{G} , the pair $(\beta, \max_{l=1, \dots, L} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\beta)})$ belongs to the set $\widehat{\mathcal{I}}^{(I)}$. On the other hand, $(\widehat{\beta}^{(c,I)}, \widehat{\sigma}^{(c,I)})$ minimizes the criterion $|\mathbf{D}_{\mathbf{X}}^{-1} \beta|_1 + c\sigma$ on the same set $\widehat{\mathcal{I}}^{(I)}$. Thus, on the event \mathcal{G} ,

$$(11.11) \quad \left| \mathbf{D}_{\mathbf{X}}^{-1} \widehat{\beta}^{(c,I)} \right|_1 + c \widehat{\sigma}^{(c,I)} \leq |\mathbf{D}_{\mathbf{X}}^{-1} \beta|_1 + c \max_{l \in I} (\mathbf{D}_{\mathbf{Z}})_{ll} \sqrt{\widehat{Q}_l(\beta)}.$$

This implies, again on the event \mathcal{G} ,

$$(11.12) \quad \begin{aligned} \left| \Delta_{J(\beta)^c} \right|_1 &= \sum_{k \in J(\beta)^c} \left| \mathbb{E}_n [X_k^2]^{1/2} \widehat{\beta}_k^{(c,I)} \right| \\ &\leq \sum_{k \in J(\beta)} \left(\left| \mathbb{E}_n [X_k^2]^{1/2} \beta_k \right| - \left| \mathbb{E}_n [X_k^2]^{1/2} \widehat{\beta}_k^{(c,I)} \right| \right) \\ &\quad + c \left(\max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\beta)} - \max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\widehat{\beta}^{(c,I)})} \right). \end{aligned}$$

For every l in I , $\gamma \mapsto (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\gamma)}$ is almost surely differentiable. Thus almost surely its subgradient is the gradient and thus the singleton

$$\partial \left((\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\cdot)} \right) (\beta) = \nabla \left((\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\cdot)} \right) (\beta) = \frac{\mathbb{E}_n [U X^T Z_l^2]}{\sqrt{\mathbb{E}_n [(U Z_l)^2]}}.$$

Because the function is also convex, almost surely,

$$(\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\beta)} - (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\widehat{\beta}^{(c,I)})} \leq \nabla \left((\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\cdot)} \right) (\beta)^T (\beta - \widehat{\beta}^{(c,I)})$$

$$\begin{aligned}
&\leq (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \left(\frac{\mathbb{E}_n[UX^T Z_l^2]}{\sqrt{\mathbb{E}_n[(UZ_l)^2]}} \right)^T \mathbf{D}_{\mathbf{X}} \mathbf{D}_{\mathbf{X}}^{-1} (\beta - \hat{\beta}^{(c,I)}) \\
&\leq -(\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \frac{\mathbb{E}_n[UX^T Z_l^2]}{\sqrt{\mathbb{E}_n[(UZ_l)^2]}} \mathbf{D}_{\mathbf{X}} \Delta.
\end{aligned}$$

By the Cauchy-Schwartz inequality,

$$\left| -(\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \frac{\mathbb{E}_n[UX^T Z_l^2]}{\sqrt{\mathbb{E}_n[(UZ_l)^2]}} \mathbf{D}_{\mathbf{X}} \right| \leq 1.$$

We will now use the same notation for \mathcal{G} and \mathcal{G} intersected with the probability 1 event where each mapping is differentiable and work on the later event. By the Dubovitsky-Milutin theorem (see, *e.g.*, Alekseev, Tikhomirov and Fomin (1987), Chapter 2)

$$\left(\max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\hat{Q}_l(\beta)} - \max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\hat{Q}_l(\hat{\beta}^{(c,I)})} \right) \leq |\Delta|_1.$$

Thus (11.12) implies

$$(11.13) \quad |\Delta_{J(\beta)^c}|_1 \leq \frac{1+c}{1-c} |\Delta_{J(\beta)}|_1.$$

Thus, $\Delta \in C_{J(\beta)}^{(c,I)}$ on the event \mathcal{G} . Using (11.9) and arguing as in (11.12) we find

$$\begin{aligned}
(11.14) \quad \left| \Psi_n^{(I)} \Delta \right|_{\infty} &\leq r \left(2\hat{\sigma}^{(c,I)} + \max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\hat{Q}_l(\beta)} - \hat{\sigma}^{(c,I)} \right) \\
&\leq r \left(2\hat{\sigma}^{(c,I)} + \max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\hat{Q}_l(\beta)} - \max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\hat{Q}_l(\hat{\beta}^{(c,I)})} \right) \\
&\leq r \left(2\hat{\sigma}^{(c,I)} + |\Delta|_1 \right).
\end{aligned}$$

This again uses the convexity for every l in I , of $\gamma \mapsto (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\hat{Q}_l(\gamma)}$. Using the definition of the sensitivities we get that, on the event \mathcal{G} ,

$$\left| \Psi_n^{(I)} \Delta \right|_{\infty} \leq r \left(2\hat{\sigma}^{(c,I)} + \frac{\left| \Psi_n^{(I)} \Delta \right|_{\infty}}{\kappa_{1,J(\beta)}^{(c,I)}} \right),$$

which implies

$$(11.15) \quad \left| \Psi_n^{(I)} \Delta \right|_{\infty} \leq 2\hat{\sigma}^{(c,I)} r \left(1 - \frac{r}{\kappa_{1,J(\beta)}^{(c,I)}} \right)_+^{-1}.$$

This inequality and the definition of the sensitivities yield (6.7) and (6.8).

To prove (6.9), it suffices to note that, by (11.11) and by the definition of $\kappa_{1,J(\beta),J(\beta)}^{(c,I)}$,

$$\begin{aligned} c\widehat{\sigma}^{(c,I)} &\leq |\Delta_{J(\beta)}|_1 + c \max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\beta)} \\ &\leq \frac{|\Psi_n^{(I)} \Delta|_\infty}{\kappa_{1,J(\beta),J(\beta)}^{(c,I)}} + c \max_{l \in I} (\mathbf{D}_{\mathbf{Z}}^{(\mathbf{I})})_{ll} \sqrt{\widehat{Q}_l(\beta)}, \end{aligned}$$

and to combine this inequality with (11.9). \square

Proof of Theorem 6.7. Take β in \mathcal{B}_s . Fix an arbitrary subset J of $\{1, \dots, K\}$. Acting as in (11.12) with J instead of $J(\beta)$, we get:

$$\begin{aligned} \sum_{k \in J^c} \left| \mathbb{E}_n[X_k^2]^{1/2} \widehat{\beta}_k^{(c,I)} \right| + \sum_{k \in J^c} \left| \mathbb{E}_n[X_k^2]^{1/2} \beta_k \right| &\leq \sum_{k \in J} \left(\left| \mathbb{E}_n[X_k^2]^{1/2} \beta_k \right| - \left| \mathbb{E}_n[X_k^2]^{1/2} \widehat{\beta}_k^{(c,I)} \right| \right) + 2 \sum_{k \in J^c} \left| \mathbb{E}_n[X_k^2]^{1/2} \beta_k \right| \\ &\quad + c \left(\sqrt{\widehat{Q}(\beta)} - \sqrt{\widehat{Q}(\widehat{\beta}^{(c,I)})} \right) \\ &\leq |\Delta_J|_1 + 2 |(\mathbf{D}_{\mathbf{X}}^{-1} \beta)_{J^c}|_1 + c |\Delta|_1. \end{aligned}$$

This yields

$$(11.16) \quad |\Delta_{J^c}|_1 \leq |\Delta_J|_1 + 2 |(\mathbf{D}_{\mathbf{X}}^{-1} \beta)_{J^c}|_1 + c |\Delta|_1.$$

Assume now that we are on the event \mathcal{G} . Consider the two possible cases. First, if $2 |(\mathbf{D}_{\mathbf{X}}^{-1} \beta)_{J^c}|_1 \leq |\Delta_J|_1$, then $\Delta \in \widetilde{C}_J^{(c,I)}$. From this, using the definition of the sensitivity $\widetilde{\kappa}_{p,J_0,J}^{(c,I)}$, we get that $|\Delta_{J_0}|_p$ is bounded from above by the first term of the maximum in (6.13). Second, if $2 |(\mathbf{D}_{\mathbf{X}}^{-1} \beta)_{J^c}|_1 > |\Delta_J|_1$, then for any $p \in [1, \infty]$ we have a simple bound

$$|\Delta_{J_0}|_p \leq |\Delta|_1 = |\Delta_{J^c}|_1 + |\Delta_J|_1 \leq \frac{6}{1-c} |(\mathbf{D}_{\mathbf{X}}^{-1} \beta)_{J^c}|_1.$$

In conclusion, $|\Delta|_p$ is smaller than the maximum of the two bounds. \square

Proof of Theorem 7.4. Part (i) of the theorem is a consequence of (6.9) and Assumptions 7.1 and 7.2. Parts (ii) and (iii) follow immediately from (6.7), (6.8), and Assumptions 7.1 and 7.2. Part (iv) is straightforward in view of (7.4). \square

Proof of Theorem 7.7. Fix β in \mathcal{B}_s . Let \mathcal{G}_j be the events of probabilities at least $1 - \gamma_j$ respectively appearing in Assumptions 7.1, 7.3, 7.6. Assume that all these events hold, as well as the event \mathcal{G} . Then, using Theorem 7.4 (i),

$$\omega_k^{(c,I)}(s) \leq \frac{2\sigma_* r}{c_k^{(c)*}(s) v_k} \left(1 + \frac{r}{c c_{1,J(\beta)}^{(c,I)}} \right) \left(1 - \frac{r}{c c_{1,J(\beta)}^{(c,I)}} \right)_+^{-1} \left(1 - \frac{r}{c_1^{(c,I)}(s)} \right)_+^{-1} \triangleq \omega_k^{(c)*}.$$

By assumption, $|\beta_k| > 2\omega_k^{(c)*}$ for $k \in J(\beta)$. Note that the following two cases can occur. First, if $k \in J(\beta)^c$ (so that $\beta_k = 0$) then, using (6.8) and Assumptions 7.1 and 7.6, we obtain $|\widehat{\beta}_k^{(c,I)}| \leq \omega_k^{(c,I)}$, which implies $\widetilde{\beta}_k^{(c,I)} = 0$. Second, if $k \in J(\beta)$, then using again (6.8) we get $\|\beta_k\| - |\widehat{\beta}_k^{(c,I)}| \leq |\beta_k - \widehat{\beta}_k^{(c,I)}| \leq \omega_k^{(c,I)} \leq \omega_k^{(c)*}$. Since $|\beta_k| > 2\omega_k^{(c)*}$ for $k \in J(\beta)$, we obtain that $|\widehat{\beta}_k^{(c,I)}| > \omega_k^{(c,I)}$, so that $\widetilde{\beta}_k^{(c,I)} = \widehat{\beta}_k^{(c,I)}$ and the signs of β_k and $\widehat{\beta}_k^{(c,I)}$ coincide. This yields the result. \square

Proof of Theorem 8.3. Fix β in \mathcal{B}_s and define the event

$$\mathcal{G} = \left\{ \left| \frac{1}{n} \mathbf{D}_Z \mathbf{Z}^T \mathbf{U} \right|_{\infty} \leq \sqrt{\widehat{Q}(\beta)r} \right\}.$$

Note that $\widehat{Q}(\beta) = \mathbb{E}_n[U^2]$. Under scenario 1, the event \mathcal{G} holds with probability at least $1 - \alpha$. For the other scenarios, the union bound yields

$$(11.17) \quad \begin{aligned} \mathbb{P}(\mathcal{G}^c) &\leq \sum_{l=1}^L \mathbb{P} \left(\frac{1}{n} \left| \frac{\sum_{i=1}^n z_{li} u_i}{z_{l*} \sqrt{\mathbb{E}_n[U^2]}} \right| \geq r \right) \\ &\leq \sum_{l=1}^L \mathbb{P} \left(\left| \frac{\sum_{i=1}^n z_{li} u_i}{\sqrt{\sum_{i=1}^n (z_{li} u_i)^2}} \right| \geq \sqrt{nr} \right). \end{aligned}$$

We conclude, using the moderate deviations result from Section 11.2 that the event \mathcal{G} holds with probability at least $1 - \alpha$ (approximately at least $1 - \alpha$ for scenario 4 and the two-stage procedure with scenario 5).

Take c in $(0, 1)$, set $\Delta \triangleq \mathbf{D}_X^{-1}(\widehat{\beta}^{(c)} - \beta)$. On the event \mathcal{G} we have:

$$(11.18) \quad \begin{aligned} |\Psi_n \Delta|_{\infty} &\leq \left| \frac{1}{n} \mathbf{D}_Z \mathbf{Z}^T (\mathbf{Y} - \mathbf{X} \widehat{\beta}^{(c)}) \right|_{\infty} + \left| \frac{1}{n} \mathbf{D}_Z \mathbf{Z}^T (\mathbf{Y} - \mathbf{X} \beta) \right|_{\infty} \\ &\leq r \widehat{\sigma}^{(c)} + \left| \frac{1}{n} \mathbf{D}_Z \mathbf{Z}^T \mathbf{U} \right|_{\infty} \\ &\leq r \left(\widehat{\sigma}^{(c)} + \sqrt{\widehat{Q}(\beta)} \right). \end{aligned}$$

Notice that, on the event \mathcal{G} , the pair $(\beta, \sqrt{\widehat{Q}(\beta)})$ belongs to the set $\widehat{\mathcal{L}}$. On the other hand, $(\widehat{\beta}^{(c)}, \widehat{\sigma}^{(c)})$ minimizes the criterion $|\mathbf{D}_X^{-1} \beta|_1 + c\sigma$ on the same set $\widehat{\mathcal{L}}$. Thus, on the event \mathcal{G} ,

$$(11.19) \quad \left| \mathbf{D}_X^{-1} \widehat{\beta}^{(c)} \right|_1 + c \widehat{\sigma}^{(c)} \leq |\mathbf{D}_X^{-1} \beta|_1 + c \sqrt{\widehat{Q}(\beta)}.$$

This implies, again on the event \mathcal{G} ,

$$(11.20) \quad \left| \Delta_{J(\beta)^c} \right|_1 = \sum_{k \in J(\beta)^c} \left| x_{k*} \widehat{\beta}_k^{(c)} \right|$$

$$\begin{aligned}
&\leq \sum_{k \in J(\beta)} \left(|x_{k*} \beta_k| - |x_{k*} \hat{\beta}_k^{(c)}| \right) + c \left(\sqrt{\widehat{Q}(\beta)} - \sqrt{\widehat{Q}(\hat{\beta}^{(c)})} \right) \\
&\leq |\Delta_{J(\beta)}|_1 + c \left(\sqrt{\widehat{Q}(\beta)} - \sqrt{\widehat{Q}(\hat{\beta}^{(c)})} \right) \\
&\leq |\Delta_{J(\beta)}|_1 + c \left| \frac{\mathbb{E}_n[UX^T] \mathbf{D}_X \Delta}{\sqrt{\mathbb{E}_n[U^2]}} \right| \quad (\text{by convexity of } \beta \mapsto \sqrt{\widehat{Q}(\beta)}) \\
&\leq |\Delta_{J(\beta)}|_1 + c \left| \frac{\mathbb{E}_n[UX^T] \mathbf{D}_X}{\sqrt{\mathbb{E}_n[U^2]}} \right|_\infty |\Delta|_1 \\
&\leq |\Delta_{J(\beta)}|_1 + c |\Delta|_1 \quad (\text{by the Cauchy-Schwarz inequality}).
\end{aligned}$$

Note that (11.20) can be re-written as a cone condition:

$$(11.21) \quad |\Delta_{J(\beta)^c}|_1 \leq \frac{1+c}{1-c} |\Delta_{J(\beta)}|_1.$$

Thus, $\Delta \in C_{J(\beta)}^{(c)}$ on the event \mathcal{G} . Using (11.9) and arguing as in (11.12) we find

$$\begin{aligned}
(11.22) \quad |\Psi_n \Delta|_\infty &\leq r \left(2\hat{\sigma}^{(c)} + \sqrt{\widehat{Q}(\beta)} - \hat{\sigma}^{(c)} \right) \\
&\leq r \left(2\hat{\sigma}^{(c)} + \sqrt{\widehat{Q}(\beta)} - \sqrt{\widehat{Q}(\hat{\beta}^{(c)})} \right) \quad (\text{since } \sqrt{\widehat{Q}(\hat{\beta}^{(c)})} \leq \hat{\sigma}^{(c)}) \\
&\leq r \left(2\hat{\sigma}^{(c)} + \left| \frac{\mathbb{E}_n[UX^T] \mathbf{D}_X \Delta}{\sqrt{\mathbb{E}_n[U^2]}} \right| \right) \\
&\leq r \left(2\hat{\sigma}^{(c)} + \max_{j \in J_{\text{exo}}^c} \left| \frac{\mathbb{E}_n[UX_j]}{\sqrt{\mathbb{E}_n[X_j^2 U^2]}} \right| |\Delta_{J_{\text{exo}}^c}|_1 + \max_{j \in J_{\text{exo}}} \left| \frac{\mathbb{E}_n[UX_j]}{\sqrt{\mathbb{E}_n[X_j^2 U^2]}} \right| |\Delta_{J_{\text{exo}}}|_1 \right) \\
&\leq r \left(2\hat{\sigma}^{(c)} + |\Delta_{J_{\text{exo}}^c}|_1 + \max_{j \in J_{\text{exo}}} \left| \frac{\mathbb{E}_n[UX_j]}{\sqrt{\mathbb{E}_n[X_j^2 U^2]}} \right| |\Delta_{J_{\text{exo}}}|_1 \right) \quad (\text{by the Cauchy-Schwarz inequality}).
\end{aligned}$$

Since the exogenous variables serve as their own instruments, we obtain that, on the event \mathcal{G} ,

$$\max_{j \in J_{\text{exo}}} \left| \frac{\mathbb{E}_n[UX_j]}{\sqrt{\mathbb{E}_n[X_j^2 U^2]}} \right| \leq r.$$

Combining this with (11.22) and using the definition of the block sensitivity $\kappa_{1, J_0, J(\beta)}^{(c)}$ with $J_0 = J_{\text{exo}}^c$, $J_0 = J_{\text{exo}}$, we get that, on the event \mathcal{G} ,

$$(11.23) \quad |\Psi_n \Delta|_\infty \leq r \left(2\hat{\sigma}^{(c)} + \sqrt{\widehat{Q}(\beta)} - \hat{\sigma}^{(c)} \right)$$

$$\leq r \left(2\widehat{\sigma}^{(c)} + \frac{|\Psi_n \Delta|_\infty}{\kappa_{1, J_{\text{exo}}, J(\beta)}^{(c)}} + r \frac{|\Psi_n \Delta|_\infty}{\kappa_{1, J_{\text{exo}}, J(\beta)}^{(c)}} \right),$$

which implies

$$(11.24) \quad |\Psi_n \Delta|_\infty \leq 2\widehat{\sigma}^{(c)} r \left(1 - \frac{r}{\kappa_{1, J_{\text{exo}}, J(\beta)}^{(c)}} - \frac{r^2}{\kappa_{1, J_{\text{exo}}, J(\beta)}^{(c)}} \right)_+^{-1}.$$

This inequality and the definition of the sensitivities yield (8.5) and (8.6).

To prove (8.7), it suffices to note that, by (11.19) and by the definition of $\kappa_{1, J(\beta), J(\beta)}^{(c)}$,

$$\begin{aligned} c\widehat{\sigma}^{(c)} &\leq |\Delta_{J(\beta)}|_1 + c\sqrt{\widehat{Q}(\beta)} \\ &\leq \frac{|\Psi_n \Delta|_\infty}{\kappa_{1, J(\beta), J(\beta)}^{(c)}} + c\sqrt{\widehat{Q}(\beta)}, \end{aligned}$$

and to combine this inequality with (11.18). \square

Proof of Theorem 8.6. Take β in \mathcal{B}_s . Note that

$$\begin{aligned} |z_i^T \widehat{\zeta}^{(c_{RF})}| &\geq |z_i^T \zeta| - \left| (\mathbf{D}_Z)^{-1} (\widehat{\zeta}^{(c_{RF})} - \zeta) \right|_1 \\ &\geq |z_i^T \zeta| - C_1(r, c_{RF}, s_{RF}) \end{aligned}$$

thus

$$\max_{i=1, \dots, n} |z_i^T \widehat{\zeta}^{(c_{RF})}| + C_1(r, c_{RF}, s_{RF})r \geq \max_{i=1, \dots, n} |z_i^T \zeta|.$$

As well, we have on E_α ,

$$\begin{aligned} \frac{1}{n} |\zeta^T \mathbf{Z}^T \mathbf{U}| &\leq \left| (\mathbf{D}_Z)^{-1} (\widehat{\zeta}^{(c_{RF})} - \zeta) \right|_1 \sqrt{\widehat{Q}(\beta)}r + \frac{1}{n} |\zeta^T \mathbf{Z}^T \mathbf{U}| \\ &\leq C_1(r, c_{RF}, s_{RF}) \sqrt{\widehat{Q}(\beta)}r + \frac{1}{n} |\zeta^T \mathbf{Z}^T \mathbf{U}|, \end{aligned}$$

thus

$$\begin{aligned} \frac{\frac{1}{n} |\widehat{\zeta}^{(c_{RF})T} \mathbf{Z}^T \mathbf{U}|}{\max_{i=1, \dots, n} |z_i^T \widehat{\zeta}^{(c_{RF})}| + C_1(r, c_{RF}, s_{RF})} &\leq \frac{C_1(r, c_{RF}, s_{RF}) \sqrt{\widehat{Q}(\beta)}r}{\max_{i=1, \dots, n} |z_i^T \widehat{\zeta}^{(c_{RF})}| + C_1(r, c_{RF}, s_{RF})} + \frac{\frac{1}{n} |\zeta^T \mathbf{Z}^T \mathbf{U}|}{\max_{i=1, \dots, n} |z_i^T \zeta|} \\ &\leq \frac{C_1(r, c_{RF}, s_{RF}) \sqrt{\widehat{Q}(\beta)}r}{\max_{i=1, \dots, n} |z_i^T \widehat{\zeta}| + C_1(r, c_{RF}, s_{RF})} + \sqrt{\widehat{Q}(\beta)}r \\ &\leq \sqrt{\widehat{Q}(\beta)}r \left(1 + \frac{C_1(r, c_{RF}, s_{RF})}{\max_{i=1, \dots, n} |z_i^T \widehat{\zeta}^{(c_{RF})}| + C_1(r, c_{RF}, s_{RF})} \right). \end{aligned}$$

This yields

$$\frac{\frac{1}{n}|\widehat{\zeta}^{(CRF)T}\mathbf{Z}^T\mathbf{U}|}{\max_{i=1,\dots,n}|z_i^T\widehat{\zeta}^{(CRF)}| + 2C_1(r, c_{RF}, s_{RF})} \leq \sqrt{\widehat{Q}(\beta)r}.$$

The rest of the proof is the same as for Theorem 8.3. \square

Proof of Theorem 9.1. Throughout the proof, we assume that we are on the event of probability at least $1 - \alpha_2$ where (9.4) holds. It follows easily from (9.4) that

$$(11.25) \quad \left| \frac{1}{n}\overline{\mathbf{Z}}^T\mathbf{X}(\widehat{\beta} - \beta^*) \right|_{\infty} \leq \widehat{b}\overline{z}_*.$$

Next, an argument similar to (11.17) and Theorem 11.6 yield that, with probability at least $1 - \alpha_1$,

$$(11.26) \quad \left| \frac{1}{n}\overline{\mathbf{Z}}^T\mathbf{U} - \theta^* \right|_{\infty} \leq r_1 \max_{l=1,\dots,L_1} \sqrt{\frac{1}{n} \sum_{i=1}^n (\overline{z}_{li}u_i - \theta_l^*)^2} = r_1 F(\theta^*, \beta^*).$$

In what follows, we assume that we are on the event of probability at least $1 - \alpha_1 - \alpha_2$ where both (11.25) and (11.26) are satisfied.

We will use the properties of $F(\theta, \beta)$ stated in the next lemma that we prove in Section 11.4.

Lemma 11.8. *We have*

$$(11.27) \quad F(\theta^*, \widehat{\beta}) - F(\widehat{\theta}, \widehat{\beta}) \leq |\widehat{\theta} - \theta^*|_1,$$

$$(11.28) \quad |F(\theta^*, \widehat{\beta}) - F(\theta^*, \beta^*)| \leq \overline{z}_* \left| \mathbf{D}_{\mathbf{X}}^{-1}(\widehat{\beta} - \beta^*) \right|_1 \leq \widehat{b}\overline{z}_*.$$

We proceed now to the proof of Theorem 9.1. First, we show that the pair $(\theta, \sigma_1) = (\theta^*, F(\theta^*, \beta^*))$ belongs to the set $\widehat{\mathcal{I}}_1$. Indeed, from (11.25) and (11.26) we get

$$\begin{aligned} \left| \frac{1}{n}\overline{\mathbf{Z}}^T(\mathbf{Y} - \mathbf{X}\widehat{\beta}) - \theta^* \right|_{\infty} &\leq \left| \frac{1}{n}\overline{\mathbf{Z}}^T\mathbf{U} - \theta^* \right|_{\infty} + \left| \frac{1}{n}\overline{\mathbf{Z}}^T\mathbf{X}(\widehat{\beta} - \beta^*) \right|_{\infty} \\ &\leq r_1 F(\theta^*, \beta^*) + \widehat{b}\overline{z}_*. \end{aligned}$$

Thus, the pair $(\theta, \sigma_1) = (\theta^*, F(\theta^*, \beta^*))$ satisfies the first constraint in the definition of $\widehat{\mathcal{I}}_1$. It satisfies the second constraint as well, since $F(\theta^*, \widehat{\beta}) \leq F(\theta^*, \beta^*) + \widehat{b}\overline{z}_*$ by (11.28).

Take any c in $(0, 1)$, as $(\theta^*, F(\theta^*, \beta^*)) \in \widehat{\mathcal{I}}_1$ and $(\widehat{\theta}^{(c)}, \widehat{\sigma}_1^{(c)})$ minimizes $|\theta|_1 + c\sigma_1$ over $\widehat{\mathcal{I}}_1$, we have

$$(11.29) \quad |\widehat{\theta}^{(c)}|_1 + c\widehat{\sigma}_1^{(c)} \leq |\theta^*|_1 + cF(\theta^*, \beta^*),$$

which implies

$$(11.30) \quad |\overline{\Delta}_{J(\theta^*)^c}|_1 \leq |\overline{\Delta}_{J(\theta^*)}|_1 + c(F(\theta^*, \beta^*) - \widehat{\sigma}_1^{(c)}),$$

where $\bar{\Delta} = \widehat{\theta}^{(c)} - \theta^*$. Using the fact that $F(\widehat{\theta}^{(c)}, \widehat{\beta}) \leq \widehat{\sigma}_1^{(c)} + \widehat{b}\bar{z}_*$, (11.27), and (11.28) we obtain

$$(11.31) \quad \begin{aligned} F(\theta^*, \beta^*) - \widehat{\sigma}_1^{(c)} &\leq F(\theta^*, \beta^*) - F(\widehat{\theta}^{(c)}, \widehat{\beta}) + \widehat{b}\bar{z}_* \\ &\leq |\widehat{\theta}^{(c)} - \theta^*|_1 + 2\widehat{b}\bar{z}_*. \end{aligned}$$

This inequality and (11.30) yield

$$|\bar{\Delta}_{J(\theta^*)^c}|_1 \leq |\bar{\Delta}_{J(\theta^*)}|_1 + c|\widehat{\theta}^{(c)} - \theta^*|_1 + 2c\widehat{b}\bar{z}_*,$$

or equivalently,

$$(11.32) \quad |\bar{\Delta}_{J(\theta^*)^c}|_1 \leq \frac{1+c}{1-c} |\bar{\Delta}_{J(\theta^*)}|_1 + \frac{2c}{1-c} \widehat{b}\bar{z}_*.$$

Next, using (11.25), (11.26) and the second constraint in the definition of $(\widehat{\theta}^{(c)}, \widehat{\sigma}_1^{(c)})$, we find

$$\begin{aligned} |\widehat{\theta}^{(c)} - \theta^*|_\infty &\leq \left| \frac{1}{n} \bar{\mathbf{Z}}^T (\mathbf{Y} - \mathbf{X}\widehat{\beta}) - \widehat{\theta}^{(c)} \right|_\infty \\ &\quad + \left| \frac{1}{n} \bar{\mathbf{Z}}^T \mathbf{U} - \theta^* \right|_\infty + \left| \frac{1}{n} \bar{\mathbf{Z}}^T \mathbf{X}(\widehat{\beta} - \beta^*) \right|_\infty \\ &\leq r_1(\widehat{\sigma}_1^{(c)} + F(\theta^*, \beta^*)) + 2\widehat{b}\bar{z}_*. \end{aligned}$$

This and (11.31) yield

$$(11.33) \quad |\widehat{\theta}^{(c)} - \theta^*|_\infty \leq r_1(2\widehat{\sigma}_1^{(c)} + |\widehat{\theta}^{(c)} - \theta^*|_1) + 2(1+r_1)\widehat{b}\bar{z}_*.$$

On the other hand, (11.32) implies

$$(11.34) \quad \begin{aligned} |\widehat{\theta}^{(c)} - \theta^*|_1 &\leq \frac{2}{1-c} |\bar{\Delta}_{J(\theta^*)}|_1 + \frac{2c}{1-c} \widehat{b}\bar{z}_* \\ &\leq \frac{2|J(\theta^*)|}{1-c} |\widehat{\theta}^{(c)} - \theta^*|_\infty + \frac{2c}{1-c} \widehat{b}\bar{z}_*. \end{aligned}$$

Inequalities (9.6) and (9.7) follow from solving (11.33) and (11.34) with respect to $|\widehat{\theta}^{(c)} - \theta^*|_\infty$ and $|\widehat{\theta}^{(c)} - \theta^*|_1$ respectively. \square

Proof of Theorem 9.3. We first prove part (i). We will assume that we are on the event of probability at least $1 - \alpha_1 - \alpha_2 - \varepsilon$ where (11.26), (9.8), and (9.10) are simultaneously satisfied. From (11.29) and the fact that (9.8) can be written as $F(\theta^*, \beta^*) \leq \sigma_{1*}$ we obtain

$$(11.35) \quad \widehat{\sigma}_1^{(c)} \leq |\widehat{\theta}^{(c)} - \theta^*|_1/c + \sigma_{1*}.$$

Note also that the argument in the proof of Theorem 9.1 and the results of that theorem remain obviously valid with \widehat{b} replaced by b_* . Thus, we can use (9.7) with \widehat{b} replaced by b_* , and combining it with (11.35) we obtain

$$(11.36) \quad \widehat{\sigma}_1^{(c)} \leq \overline{\sigma}_*.$$

This and (9.6) yield (9.11).

We now prove part (ii) of the theorem. In the rest of the proof, we assume that we are on the event \mathcal{G}' of probability at least $1 - \alpha_1 - \varepsilon - \gamma$ where (11.26), (9.8), and the events $\mathcal{G}, \mathcal{G}_j$ defined in the proofs of Theorems 6.5, 7.4 are simultaneously satisfied. Then item (ii) of Theorem 7.4 with $p = 1$ implies (9.10) with b_* defined in (9.12). This and (9.11) easily give part (ii) of the theorem.

To prove part (iii), note that, by Theorem 7.4 (i) and Assumption 7.6,

$$(11.37) \quad \widehat{b} = \frac{2\widehat{\sigma}rs}{\kappa_1(s)} \left(1 - \frac{r}{\kappa_1(s)}\right)_+^{-1} \leq b_*$$

for b_* defined in (9.12). This and (11.36) imply that the threshold ω satisfies $\omega \triangleq V(\widehat{\sigma}_1^{(c)}, c, \widehat{b}, J(\widehat{\theta})) \leq V(\overline{\sigma}_*, c, b_*, s_1) \triangleq \omega^{(c)*}$ on the event \mathcal{G}' . On the other hand, (9.6) guarantees that $|\widehat{\theta}_l^{(c)} - \theta_l^*| \leq \omega^{(c)}$ and, by assumption, $|\theta_l^*| > 2\omega^{(c)*}$ for all $l \in J(\theta^*)$. In addition, by (6.7) and (9.6) for all $l \in J(\theta^*)^c$ we have $|\theta_l^*| < \omega^{(c)}$, which implies $\widetilde{\theta}_l^{(c)} = 0$. We finish the proof in the same way as the proof of Theorem 7.4. \square

11.4. Proof of Lemma 11.8. Set $f_l(\theta_l) \triangleq \sqrt{\widehat{Q}_l(\theta_l, \widehat{\beta})}$, and $f(\theta) \triangleq \max_{l=1, \dots, L_1} f_l(\theta_l) \equiv F(\theta, \widehat{\beta})$. The mappings $\theta \mapsto f_l(\theta_l)$ are convex, so that by the Dubovitsky-Milutin, the subdifferential of their maximum f is contained in the convex hull of the union of the subdifferentials of the f_l :

$$(11.38) \quad \partial f \subseteq \text{Conv} \left(\bigcup_{l=1}^{L_1} \partial f_l \right).$$

Since, obviously, $\partial f_l(\theta_l) \subseteq [-1, 1]$, we find that $\partial f(\theta) \subseteq \{w \in \mathbb{R}^{L_1} : |w|_\infty \leq 1\}$ for all $\theta \in \mathbb{R}^{L_1}$. Using this property and the convexity of f , we get

$$f(\theta^*) - f(\widehat{\theta}) \leq \langle w, \theta^* - \widehat{\theta} \rangle \leq |\widehat{\theta} - \theta^*|_1, \quad \forall w \in \partial f(\theta^*),$$

where $\langle \cdot, \cdot \rangle$ denotes the standard inner product in \mathbb{R}^{L_1} . This yields (11.27). The proof of (11.28) is based on similar arguments. Instead of f_l , we now introduce the functions g_l defined by $g_l(\beta) \triangleq \sqrt{\widehat{Q}_l(\theta_l^*, \beta)}$, and set $g(\beta) \triangleq \max_{l=1, \dots, L_1} g_l(\beta) \equiv F(\theta^*, \beta)$. Next, notice that the subdifferential of g_l

satisfies $\partial g_l(\beta) \subseteq \{w \in \mathbb{R}^K : |w_k| \leq a_{lk}, k = 1, \dots, K\}$ for all $\beta \in \mathbb{R}^K, l = 1, \dots, L_1$, where

$$a_{lk} = \frac{|\frac{1}{n} \sum_{i=1}^n \bar{z}_{li} x_{ki} (\bar{z}_{li} (y_i - x_i^T \beta) - \theta_l^*)|}{\sqrt{\frac{1}{n} \sum_{i=1}^n (\bar{z}_{li} (y_i - x_i^T \beta) - \theta_l^*)^2}}.$$

Consequently, by the Cauchy-Schwarz inequality, $\mathbf{D}_{\mathbf{X}} \partial g_l(\beta) \subseteq \{w \in \mathbb{R}^K : |w|_\infty \leq \bar{z}_*\}$ for all $\beta \in \mathbb{R}^K, l = 1, \dots, L_1$. This and (11.38) with g, g_l instead of f, f_l imply $\mathbf{D}_{\mathbf{X}} \partial g(\beta) \subseteq \{w \in \mathbb{R}^K : |w|_\infty \leq \bar{z}_*\}$ for all $\beta \in \mathbb{R}^K$. Using this property and the convexity of g , we get

$$g(\beta) - g(\beta') \leq \langle w, (\beta - \beta') \rangle \leq |\mathbf{D}_{\mathbf{X}} w|_\infty |\mathbf{D}_{\mathbf{X}}^{-1}(\beta - \beta')|_1 \leq \bar{z}_* |\mathbf{D}_{\mathbf{X}}^{-1}(\beta - \beta')|_1, \quad \forall w \in \partial g(\beta),$$

for any $\beta, \beta' \in \mathbb{R}^K$. This proves (11.28). \square

Proof of Theorem 8.2. The only difference with the proof of Theorem 6.5 is that, because we do not have the l_1 -norm in the objective function (8.1), we drop the discussion leading to (11.13). \square

REFERENCES

- [1] Alekseev, V.M, V. M. Tikhomirov, and S. V. Fomin (1987): *Optimal Control*. Consultants Bureau, New York.
- [2] Amemiya, T. (1974): “The Non-Linear Two-Stage Least Squares Estimator”. *Journal of Econometrics*, 2, 105–110.
- [3] Andrews, D. W. K. (1999): “Consistent Moment Selection Procedures for Generalized Method of Moments Estimation”. *Econometrica*, 67, 543–564.
- [4] Andrews, D. W. K., and J. H. Stock (2007): “Inference with Weak Instruments”, in: *Advances in Economics and Econometrics Theory and Applications, Ninth World Congress*, Blundell, R., W. K. Newey, and T. Persson, Eds, 3, 122–174, Cambridge University Press.
- [5] Angrist, J. D., and A. B. Krueger (1991): “Does Compulsory School Attendance Affect Schooling and Earnings?”. *Quarterly Journal of Economics*, 106, 979–1014.
- [6] Bahadur, R. R., and L. J. Savage (1956): “The Nonexistence of Certain Statistical Procedures in Nonparametric Problems”. *Annals of Mathematical Statistics*, 27, 1115–1122.
- [7] Bai J., and S. Ng (2009): “Selecting Instrumental Variables in a Data Rich Environment”. *Journal of Time Series Econometrics*, 1, 105–110.
- [8] Basmann, R. (1960): “On Finite Sample Distributions of Generalized Classical Linear Identifiability Test Statistics.”. *Journal of the American Statistical Association*, 55, 650–659.
- [9] Belloni, A., D. Chen, V. Chernozhukov, and C. Hansen (2010): “Sparse Models and Methods for Optimal Instruments with an Application to Eminent Domain”. Forthcoming *Econometrica*.
- [10] Belloni, A., and V. Chernozhukov (2010): “Least Squares After Model Selection in High-dimensional Sparse Models”. Forthcoming *Bernoulli*.
- [11] Belloni, A., V. Chernozhukov, and L. Wang (2011): “Square-Root Lasso: Pivotal Recovery of Sparse Signals Via Conic Programming”. *Biometrika*, 98, 791–806.

- [12] Belloni, A., and V. Chernozhukov (2011a): “L1-Penalized Quantile Regression in High-Dimensional Sparse Models”. *The Annals of Statistics*, 39, 82–130.
- [13] Belloni, A., and V. Chernozhukov (2011b): “High Dimensional Sparse Econometric Models: an Introduction”, in: *Inverse Problems and High Dimensional Estimation, Stats in the Château 2009*, Alquier, P., E. Gautier, and G. Stoltz, Eds., *Lecture Notes in Statistics*, 203, 127–162, Springer, Berlin.
- [14] Bertail, P., E. Gauthérat, and H. Harari-Kermadec (2005): “Empirical-Discrepancies and Quasi-Empirical Likelihood : Exponential Bounds”. Preprint CREST 2005-34.
- [15] Bertail, P., E. Gauthérat, and H. Harari-Kermadec (2009): “Exponential Inequalities for Self Normalized Sums”. *Electronic Communications in Probability*, 13, 628–640.
- [16] Bickel, P., J. Y. Ritov, and A. B. Tsybakov (2009): “Simultaneous Analysis of Lasso and Dantzig Selector”. *The Annals of Statistics*, 37, 1705–1732.
- [17] Blundell, R., W. Chen, and D. Kristensen (2007): “Semi-nonparametric IV Estimation of Shape-invariant Engel Curves”. *Econometrica*, 75, 1613–1669.
- [18] Bühlmann, P., and S. A. van de Geer (2011): *Statistics for High-Dimensional Data*. Springer, New-York.
- [19] Caner, M. (2009): “LASSO Type GMM Estimator”. *Econometric Theory*, 25, 1–23.
- [20] Candès, E., and T. Tao (2007): “The Dantzig Selector: Statistical Estimation when p is Much Larger Than n ”. *The Annals of Statistics*, 35, 2313–2351.
- [21] Carrasco, M., and J. P. Florens (2000): “Generalization of GMM to a Continuum of Moment Conditions”. *Econometric Theory*, 16, 797–834.
- [22] Carrasco, M. (2012): “A Regularization Approach to the Many Instruments Problem”. Forthcoming in *Journal of Econometrics*.
- [23] Chamberlain, G. (1987): “Asymptotic Efficiency in Estimation with Conditional Moment Restrictions”. *Journal of Econometrics*, 34, 305–334.
- [24] Chamberlain, G. and G. Imbens (2004): “Random Effects Estimators with Many Instrumental Variables”. *Econometrica*, 72, 295–306.
- [25] Dalalyan, A., and A. B. Tsybakov (2008): “Aggregation by Exponential Weighting, Sharp PAC-Bayesian Bounds and Sparsity”. *Journal of Machine Learning Research*, 72, 39–61.
- [26] Donald, S. G., and W. K. Newey (2001): “Choosing the Number of Instruments”. *Econometrica*, 69, 1161–1191.
- [27] Donoho, D. L., M. Elad, and V. N. Temlyakov (2006): “Stable Recovery of Sparse Overcomplete Representations in the Presence of Noise”. *IEEE Transactions on Information Theory*, 52, 6–18.
- [28] Dufour, J.-M. (1997): “Impossibility Theorems in Econometrics with Applications to Structural and Dynamic Models”. *Econometrica*, 65, 1365–1387.
- [29] Efron, B. (1969): “Student’s t-test Under Symmetry Conditions”. *Journal of American Statistical Society*, 64, 1278–1302.
- [30] Gleser, L. J., and J. T. Hwang (1987): “The Nonexistence of $100(1 - \alpha)\%$ Confidence Sets of Finite Expected Diameter in Errors-in-Variables and Related Models”. *Annals of Statistics*, 15, 1351–1362.
- [31] Hall, A. R., and F. P. M. Peixe (2003): “A Consistent Method for the Selection of Relevant Instruments”. *Econometric Reviews*, 22, 269–287.

- [32] Hahn, J. (2002): “Optimal Inference with Many Instruments”. *Econometric Theory*, 18, 140–168.
- [33] Hansen, L. P. (1982): “Large Sample Properties of Generalized Method of Moments Estimators”. *Econometrica*, 50, 1029–1054.
- [34] Jing, B.-Y., Q. M. Shao, and Q. Wang (2003): “Self-Normalized Cramér-Type Large Deviations for Independent Random Variables”. *The Annals of Probability*, 31, 2167–2215.
- [35] Kolesár, M., R. Chetty, J. Friedman, E. Glaeser, and G. W. Imbens (2011): “Identification and Inference with Many Invalid Instruments”. Preprint.
- [36] Koltchinskii, V. (2009): “The Dantzig Selector and Sparsity Oracle Inequalities”. *Bernoulli*, 15, 799–828.
- [37] Koltchinskii, V. (2011): *Oracle Inequalities in Empirical Risk Minimization and Sparse Recovery Problems*. Forthcoming in *Lecture Notes in Mathematics*, Springer, Berlin.
- [38] Liao, Z. (2010): “Adaptive GMM Shrinkage Estimation with Consistent Moment Selection”. Working Paper.
- [39] Lounici, K. (2008): “Sup-Norm Convergence Rate and Sign Concentration Property of the Lasso and Dantzig Selector”. *Electronic Journal of Statistics*, 2, 90–102.
- [40] Nelson, C. R., and Startz, R. (1990a): “Some Further Results on the Exact Small Sample Properties of the Instrumental Variables Estimator”. *Econometrica*, 58, 967–976.
- [41] Nelson, C. R., and Startz, R. (1990b): “The Distribution of the Instrumental Variable Estimator and Its t Ratio When the Instrument Is a Poor One”. *Journal of Business*, 63, S125–S140.
- [42] Newey, W. K. (1990): “Efficient Instrumental Variables Estimation of Nonlinear Models”. *Econometrica*, 58, 809–837.
- [43] Okui, R. (2008): “Instrumental Variable Estimation in the Presence of Many Moment Conditions”. *Journal of Econometrics*, forthcoming.
- [44] Pinelis, I. (1994): “Probabilistic Problems and Hotelling’s t^2 Test Under a Symmetry Condition”. *Annals of Statistics*, 22, 357–368.
- [45] Raskutti, G., M. J. Wainwright, and B. Yu, (2011): “Minimax Rates of Estimation for High-dimensional Linear Regression Over l_q -balls”. *IEEE Transactions on Information Theory*, 57, 6976–6994.
- [46] Rigollet, P., and A. B. Tsybakov (2011): “Exponential Screening and Optimal Rates of Sparse Estimation”. *The Annals of Statistics*, 35, 731–771.
- [47] Romano, J. P., and M. Wolf (2000): “Finite Sample Nonparametric Inference and Large Sample Efficiency”. *The Annals of Statistics*, 28, 756–778.
- [48] Rosenbaum, M., and A. B. Tsybakov (2010): “Sparse Recovery Under Matrix Uncertainty”. *The Annals of Statistics*, 38, 2620–2651.
- [49] Sargan, J. D. (1958): “The Estimation of Economic Relationships Using Instrumental Variables”. *Econometrica*, 26, 393–415.
- [50] Sala-i-Martin, X. (1997): “I Just Ran Two Million Regressions”. *The American Economic Review*, 87, 178–183.
- [51] Städler, N., P. Bühlmann and S. van de Geer (2010): “ l^1 -penalization for Mixture Regression Models”. *Test*, 19, 209–256.
- [52] Stock, J. H., J.H. Wright, and M. Yogo (2002): “A Survey of Weak Instruments and Weak Identification in Generalized Method of Moments”. *Journal of Business & Economic Statistics*, 20, 518–529.

- [53] Surm, J. F. (1999): “Using SeDuMi 1.02, a Matlab Toolbox for Optimization Over Symmetric Cones”. *Optimization Methods and Software*, 11, 625–653.
- [54] Verzelen, N. (2010): “Minimax Risks for Sparse Regressions: Ultra-high-dimensional Phenomenons”. *Electronic Journal of Statistics*, 6,38–90.
- [55] Ye, F., and C.-H. Zhang (2010): “Rate Minimality of the Lasso and Dantzig Selector for the l_q Loss in l_r Balls”. *Journal of Machine Learning Research*, 11, 3519–3540.

CREST, ENSAE PARISTECH, 3 AVENUE PIERRE LAROUSSE, 92 245 MALAKOFF CEDEX, FRANCE.

E-mail address: `eric.gautier@ensae-paristech.fr`

CREST, ENSAE PARISTECH, 3 AVENUE PIERRE LAROUSSE, 92 245 MALAKOFF CEDEX, FRANCE.

E-mail address: `alexandre.tsybakov@ensae-paristech.fr`