

CLOSED SUBSETS OF ROOT SYSTEMS AND REGULAR SUBALGEBRAS

ANDREW DOUGLAS^{1,2} AND WILLEM A. DE GRAAF³

ABSTRACT. We describe an algorithm for classifying the closed subsets of a root system, up to conjugation by the associated Weyl group. Such a classification for an irreducible root system is closely related to the classification of the regular subalgebras, up to inner automorphism, of the corresponding simple Lie algebra. We implement our algorithm to classify the closed subsets of the irreducible root systems of ranks 3 through 7. We present a complete description of the classification for the closed subsets of the rank 3 irreducible root system. We employ this root system classification to classify all regular subalgebras of the rank 3 simple Lie algebras. We present only summary data for the classifications in higher ranks due to the large size of these classifications. Our algorithm is implemented in the language of the computer algebra system GAP.

1. INTRODUCTION

Closed subsets of root systems appear in a number of contexts. They are integral to the classification of regular semisimple subalgebras of the exceptional Lie algebras [Dyn52, dGr09, LG72]. Closed subsets of root systems also find application to classifications of the reflection subgroups of finite and affine Weyl groups [DL11]. They appear in the theory of Chevalley groups [HV96]. And, the problem of decomposing a root system into a union of two closed subsets has aided in the classification of complex homogeneous spaces [Mal80].

The problem of classifying all closed subsets of root systems is an unsolved problem, but incomplete results exist. We mention three examples. Borevich [Bor77] determined the number of regular subalgebras of rank n in a simple Lie algebra of rank n . In [Wes05], we determined the number of regular subalgebras of rank n in a simple Lie algebra of rank n . In [Wes05], we determined the number of regular subalgebras of rank n in a simple Lie algebra of rank n .

up to inner automorphism, of the corresponding simple Lie algebra. We implement our algorithm to classify the closed subsets of the irreducible root systems of ranks 3 through 7. We present a complete description of the classification for the closed subsets of the rank 3 irreducible root systems. We employ this to classify all regular subalgebras of the rank 3 simple Lie algebras. This is part of a bigger project to classify the non semisimple subalgebras of small-dimensional simple Lie algebras, see [DR15, DR16a, DR16b, DR17, May16]. We present only summary data for the classifications in higher ranks due to the large size of these classifications.

The article is organized as follows. In Section 2, we review root systems and closed subsets of root systems. In Section 3, we describe our algorithm for classifying the closed subsets of root systems, along with its implementation in GAP4 [GAP], using the SLA package [dGr09]. The latest version of this package (version 1.5) contains this implementation. We present the runtimes of the main steps of the algorithm on some sample inputs. All timings in this paper have been obtained on a 2.0Ghz processor with 4GB of memory for sf GAP.

Next, in Section 4, we present the full classification of the closed subsets of the irreducible root system of rank 3 derived from the implementation of our algorithm. In this section we also present summary data for the classifications of closed subsets of the irreducible root systems of ranks 4 through 7. In Section 5, we discuss the relationship between closed subsets of irreducible root systems and regular subalgebras of the corresponding simple Lie algebras. In this section, we also employ the classification of closed subsets of rank 3 irreducible root systems to yield a classification of regular subalgebras of the corresponding simple Lie algebras.

The ground field for Lie algebras in this article is the complex numbers. Unless explicitly stated otherwise, “ \oplus ” denotes the direct sum of vector spaces, not of Lie algebras.

Acknowledgement: We thank Heiko Dietrich for his help with the computation of the closed subsets of the root system of type E_7 , for which his computer worked several weeks. AD would also like to thank WAdG and the Department of Mathematics at the University of Trento for their warm hospitality during a recent visit.

2. ROOTS SYSTEMS AND THEIR CLOSED SUBSETS

Let E be a Euclidean space with positive definite symmetric bilinear form denoted (α, β) , for $\alpha, \beta \in E$. A subset Φ of E is called a *root system* in E if the following axioms are satisfied:

- (R1) Φ is finite, spans E , and does not contain 0.
- (R2) If $\alpha \in \Phi$, the only multiples of α in Φ are $\pm\alpha$.
- (R3) If $\alpha \in \Phi$, the reflection determined by α leaves Φ invariant.
- (R4) If $\alpha, \beta \in \Phi$, then $\langle \beta, \alpha^\vee \rangle = \frac{2(\beta, \alpha)}{(\alpha, \alpha)} \in \mathbb{Z}$.

The root system Φ is called *irreducible* if it cannot be partitioned into the union of two proper subsets such that each root in one set is orthogonal to each root in the other. Let s_α denote the reflection determined by $\alpha \in \Phi$. The *Weyl group* of a root system is the subgroup of $GL(E)$ generated by the reflections s_α , $\alpha \in \Phi$.

A subset T of a root system Φ is *closed* if for any $\alpha, \beta \in T$, $\alpha + \beta \in \Phi$ implies $\alpha + \beta \in T$. Any closed set T is a disjoint union of its *symmetric* part $T^s = \{\alpha \in T \mid -\alpha \in T\}$, and its *special* part $T^u = \{\alpha \in T \mid -\alpha \notin T\}$.

Closed subsets T and T' of a root system Φ are *conjugate* if there exists an element $w \in W(\Phi)$ such that $w(T) = T'$. The following lemma is well-known (and easy to prove).

Lemma 2.1. *T^r is a closed root subsystem of Φ . For any two roots $\alpha \in T^u$ and $\beta \in T$ such that $\alpha + \beta$ is a root, we have that $\alpha + \beta \in T^u$. In particular, T^u is closed.*

2.1. Topologies on finite sets and closed subsets of the root systems of type A_{n-1} . Here we briefly describe the correspondence that exists between the closed sets in the root systems of type A_{n-1} and the topologies that a set of size n can have. To the best of our knowledge this correspondence has first been described by Borevich ([Bor76], [Bor77]). Various computer programs have been developed to enumerate the topologies of a finite set (see for example [BM02]).

Let $n \geq 1$ and $X = \{1, 2, \dots, n\}$. A topology on X is a collection τ of subsets of X such that $\emptyset, X \in \tau$ and the union and intersection of elements of τ is again in τ . The elements of τ are said to be *closed sets*. For $i \in X$ we denote by \bar{i} the smallest closed set containing i . The topology τ is called T_0 if for $i, j \in X$, $i \neq j$ we have that $i \notin \bar{j}$ or $j \notin \bar{i}$. Two topologies τ, τ' are called *homeomorphic* if there is a $\pi \in S_n$ with $\tau' = \{\pi(T) : T \in \tau\}$.

To a topology τ on X we associate an $n \times n$ -matrix σ^τ , where $\sigma_{ij}^\tau = 1$ if $i \in \bar{j}$, and $\sigma_{ij}^\tau = 0$ otherwise. It is clear that from σ^τ we can reconstruct τ . Furthermore, a matrix σ with coefficients 0, 1 corresponds to a topology if

$$(1) \quad \sigma_{ii} = 1 \text{ for all } i \text{ and } \sigma_{ij} = 1, \sigma_{jk} = 1 \text{ imply } \sigma_{ik} = 1.$$

Finally, we have that τ, τ' are homeomorphic if and only if there is a $\pi \in S_n$ with $\sigma_{ij}^{\tau'} = \sigma_{\pi(i), \pi(j)}^\tau$.

The root system Φ of type A_{n-1} consists of all α_{ij} for $1 \leq i \neq j \leq n$. Here $\alpha_{1,2}, \alpha_{2,3}, \dots, \alpha_{n-1,n}$ are the simple roots. If $i < j$ then $\alpha_{ij} = \alpha_{i,i+1} + \alpha_{i+1,i+2} + \dots + \alpha_{j-1,j}$. If $i > j$ then $\alpha_{ij} = -\alpha_{ji}$. This implies that $\alpha_{ij} + \alpha_{kl} \in \Phi$ if $j = k$, when the sum is α_{il} , or if $l = i$, when the sum is α_{kj} . The Weyl group of Φ is isomorphic to S_n , and for $\pi \in S_n$ we have $\pi \cdot \alpha_{ij} = \alpha_{\pi(i), \pi(j)}$.

Let τ be a topology on X , and set $S = \{\alpha_{ij} \in \Phi \mid \sigma_{ij}^\tau = 1\}$. Then by (1) we see that S is closed. Moreover, τ is T_0 if and only if for each $i, j \in X$, $i \neq j$, at least one of $\sigma_{ij}^\tau, \sigma_{ji}^\tau$ is 0. But this is the same as saying that S is special.

Conversely, let $S \subset \Phi$ be closed. Define the matrix σ such that $\sigma_{ii} = 1$ for all i and $\sigma_{ij} = 1$ if $\alpha_{ij} \in S$, and $\sigma_{ij} = 0$ otherwise. Then σ satisfies (1), and therefore defines a topology on X . It is also clear that two topologies are homeomorphic if and only if the corresponding closed sets are conjugate under S_n .

The conclusion is that topologies on X correspond to closed sets in Φ , whereas topologies up to homeomorphism correspond to closed sets up to conjugacy by the Weyl group. In this correspondence the T_0 topologies correspond to special closed sets. Therefore, the tables in [BM02] imply that the root system of type A_{15} contains 4483130665195086 special closed sets.

3. THE ALGORITHM

In this section we outline an algorithm for listing the closed sets of a root system Φ . This algorithm is split into two parts. In the first part we find all special closed subsets of Φ up to conjugacy. The second part is a procedure to list, up to conjugacy, all non-special closed sets whose special part is a given special closed set obtained by the first half of the algorithm.

3.1. Listing the special closed sets. We first give a simple algorithm for listing the special closed sets of a root system up to conjugacy by the Weyl group. Subsequently we describe some ideas that can be used to implement it reasonably efficiently.

The algorithm is based on three lemmas, the first of which is proved in [Bou68], Chapter VI, Proposition 22, see [Sop04] for an easier proof.

Lemma 3.1. *A special closed set is conjugate to a subset of Φ^+ .*

For a closed set T we let $T + T$ be the set of all $\gamma \in T$ such that $\gamma = \alpha + \beta$ for certain $\alpha, \beta \in T$. Let Φ^+ be a fixed subset of positive roots with set of simple roots Δ . If $\alpha = \sum_{\alpha \in \Delta} k_\alpha \alpha$, then $\sum_{\alpha \in \Delta} k_\alpha$ is called the *height* of α . Write $W = W(\Phi)$, the Weyl group of Φ .

Lemma 3.2. *Let T be a special closed set. Then $T + T$ is strictly contained in T .*

Proof. Suppose that $T \subset \Phi^+$ and let $\alpha \in T$ be of minimal height. Then α cannot be a sum of two elements of T . The general case follows by Lemma 3.1. \square

For a closed set T we denote its conjugacy class by $[T]$, that is,

$$[T] = \{w(T) \mid w \in W\}.$$

Let T be a special closed set. By Lemma 3.2 there is an $\alpha \in T \setminus (T + T)$. Set $T' = T \setminus \{\alpha\}$. Then T' is closed and we say that $[T']$ is a *successor* of $[T]$. Note that the set of successors of $[T]$ depends on the class $[T]$ and not on the choice of T . Indeed, let $S \in [T]$, then there is a $w \in W$ with $w(T) = S$. So $\beta = w(\alpha) \in S \setminus (S + S)$ and with $S' = S \setminus \{\beta\}$ we have $S' = w(T')$ and therefore $[S'] = [T']$.

Lemma 3.3. *Let T' be a special closed set of cardinality $n < |\Phi^+|$. Then there is a special closed set T of cardinality $n + 1$ such that $[T']$ is a successor of $[T]$.*

Proof. By Lemma 3.1 we may assume that $T' \subset \Phi^+$. Set

$$\mathcal{N}(T') = \{\alpha \in \Phi \mid \alpha + \beta \in T' \text{ for all } \beta \in T' \text{ such that } \alpha + \beta \in \Phi\},$$

which we call the *normalizer* of T' . We claim that there is a $\gamma \in \Phi^+$ with $\gamma \notin T'$, $\gamma \in \mathcal{N}(T')$. Let \mathfrak{g} be the semisimple Lie algebra over \mathbb{C} with Cartan subalgebra \mathfrak{h} such that the corresponding root system is isomorphic to Φ . In the sequel we identify this root system and Φ . Let \mathfrak{n}^+ be the subalgebra of \mathfrak{g} spanned by the root vectors corresponding to the roots in Φ^+ . Let \mathfrak{a} be the subspace of \mathfrak{n}^+ spanned by the root vectors corresponding to the roots in T' . Since T' is closed, \mathfrak{a} is a subalgebra. Because \mathfrak{a} is spanned by root vectors we have $[\mathfrak{h}, \mathfrak{a}] \subset \mathfrak{a}$. Consider the normalizer of \mathfrak{a} in \mathfrak{n}^+ , $\mathfrak{n}(\mathfrak{a}) = \{x \in \mathfrak{n}^+ \mid [x, \mathfrak{a}] \subset \mathfrak{a}\}$. Let $x \in \mathfrak{n}(\mathfrak{a})$ and $y \in \mathfrak{a}$, $h \in \mathfrak{h}$, then $[[h, x], y] = -[[x, y], h] - [[y, h], x]$. Now $[x, y] \in \mathfrak{a}$ and hence $[[x, y], h] \in \mathfrak{a}$. Secondly, $[y, h] \in \mathfrak{a}$ so that $[[y, h], x] \in \mathfrak{a}$ as well. We conclude that $[h, x] \in \mathfrak{n}(\mathfrak{a})$. It follows that $\mathfrak{n}(\mathfrak{a})$ is a subalgebra of \mathfrak{n}^+ , normalized by \mathfrak{h} . Therefore, $\mathfrak{n}(\mathfrak{a})$ is also spanned by root vectors. If $\mathfrak{n}(\mathfrak{a}) = \mathfrak{a}$ then \mathfrak{a} would be a Cartan subalgebra of \mathfrak{n}^+ . But the only Cartan subalgebra of \mathfrak{n}^+ is \mathfrak{n}^+ itself. We conclude that $\mathfrak{n}(\mathfrak{a})$ contains a root vector corresponding to a $\gamma \in \Phi^+$, not lying in \mathfrak{a} . Then $\gamma \notin T'$, $\gamma \in \mathcal{N}(T')$.

Set $T = T' \cup \{\gamma\}$. Then T is closed and $\gamma \in T \setminus (T + T)$. The lemma follows. \square

Remark 3.4. Let T be a special closed set and $W_T = \{w \in W \mid w(T) = T\}$, the stabilizer of T in W . Let $\alpha, \beta \in T \setminus (T + T)$ be such that $\beta = w(\alpha)$, for a certain $w \in W_T$. Then $w(T \setminus \{\alpha\}) = T \setminus \{\beta\}$.

This, together with the previous lemma, implies the correctness of the following algorithm. It takes as input a set M of representatives of the conjugacy classes of special closed sets of cardinality $n + 1$. It outputs a set of representatives of the conjugacy classes of special closed sets of cardinality n . The basic idea is to compute the successors of each element of M and weed out the conjugate copies. Remark 3.4 helps with the latter task. The algorithm takes the following steps:

- (1) Set $L = \emptyset$.
- (2) For each T in M do
 - (a) Compute W_T .
 - (b) Compute the orbits of W_T on $T \setminus (T + T)$.
 - (c) For each such orbit take a representative α and set $S = T \setminus \{\alpha\}$. If S is not conjugate to any element of L , then add S to L .
- (3) Return L .

Write $m = |\Phi^+|$. Now starting with the set M containing the one special closed subset of Φ^+ of maximal cardinality, that is, Φ^+ itself, we iterate this algorithm and obtain the special closed sets (up to conjugacy) of sizes $m - 1, m - 2, \dots, 1$. So in the end we obtain, by Lemma 3.1, all special closed sets of Φ up to conjugacy.

Remark 3.5. We give some technical details of the implementation. The algorithm has been implemented in the language of the computer algebra system GAP4 [GAP], as part of the package SLA [dGr11].

In our implementation we use a list $\alpha_1, \dots, \alpha_N$ of all the roots of the root system. A root is represented by its position in this list. So instead of α_i we just use i . This gives a memory efficient way to represent a closed set as a list of indices. The addition relations of the roots are stored in a table. This is a 2-dimensional table \mathcal{T} such that if $\mathcal{T}(i, j) = k$, then $\alpha_i + \alpha_j = \alpha_k$ if $k \neq 0$, and $\alpha_i + \alpha_j$ is not a root if $k = 0$.

The Weyl group is given as a permutation group on $\{1, \dots, N\}$. This allows us to use the functionality for permutation groups present in GAP4. Thus we can readily compute generators of W_T (the command for this is `Stabilizer(W, T, OnSets)`) and the orbits of W_T on $T \setminus (T + T)$ (the command here is `Orbits(H, D)`, where $H = W_T$ and $D = T \setminus (T + T)$).

For deciding whether two sets are conjugate under the Weyl group backtrack methods (see [Ser03], Chapter 9) are used. They are present in the GAP core system, the command for checking whether S_1, S_2 are conjugate is `RepresentativeAction(W, S1, S2, OnSets)`.

In order to make the implementation more efficient we make use of certain invariants of closed sets. An invariant of closed sets is a function $\chi : \{\text{closed sets}\} \rightarrow A$, where A is some set, such that $S = w(T)$ for a $w \in W$ implies that $\chi(S) = \chi(T)$. First we describe three invariants. In order to describe the first two of these we need some well known facts on weights (see [Hum72], Chapter 13), which we recall first.

Let $\Delta = \{\alpha_1, \dots, \alpha_\ell\}$ be a fixed set of simple roots of Φ . Define the elements λ_i of the real space E spanned by Φ by $\langle \lambda_i, \alpha_j^\vee \rangle = \delta_{i,j}$. These λ_i are called the fundamental weights. A weight is a \mathbb{Z} -linear combination of the fundamental weights. A weight $\lambda = m_1\lambda_1 + \dots + m_\ell\lambda_\ell$ is *dominant* if $m_i \geq 0$ for all i . The Weyl group W acts on the set of weights, and each weight λ is W -conjugate to a unique dominant weight ([Hum72], Lemma 13.2.A). This dominant weight is computed as follows. Set $\mu_1 = \lambda$. Let $i \geq 1$

and suppose that μ_i is defined. Write $\mu_i = n_1\lambda_1 + \cdots + n_\ell\lambda_\ell$. If $n_j \geq 0$ for all j then μ_i is the dominant weight that we are after. Otherwise there is a j with $n_j < 0$ and we set $\mu_{i+1} = s_{\alpha_j}(\mu_i)$. It is not difficult to show that this procedure terminates and returns the unique dominant weight conjugate to λ (see, for example, [dGr00], p. 319). Note that by recording the s_{α_j} that are used we also find a $w \in W$ such that $w(\lambda)$ is dominant.

Now let $S \subset \Phi$ then its sum vector is $\xi(S) = \sum_{\alpha \in S} \alpha$. The sum vectors of conjugate sets are obviously conjugate, but in general not the same. In order to get an invariant we compute the unique dominant weight that is conjugate to $\xi(S)$. We denote the result by $\sigma(S)$. Then σ is an invariant of closed sets.

For a second invariant let r be a rational number and consider the set $S^{(r)} = \{\alpha \in \Phi \mid (\alpha, \beta) \geq r \text{ for all } \beta \in S\}$. If $T = w(S)$ for a $w \in W$ then also $T^{(r)} = w(S^{(r)})$ because the Weyl group leaves the inner product invariant. Define $\delta^r(S) = \sigma(S^{(r)})$; then δ^r is an invariant of closed sets. In our implementation we normalize the inner product so that $(\alpha, \alpha) = 2$ for short roots α and we use the invariant δ^{-1} . We also considered other values of r , but $r = -1$ turned out to work best.

Let $T = \{\alpha_{i_1}, \dots, \alpha_{i_m}\}$ be a closed set. The matrix of its inner products is $M_T = ((\alpha_{i_k}, \alpha_{i_l}))_{1 \leq k, l \leq m}$, and let \widehat{M}_T be the matrix consisting of the sorted rows of M_T , which are listed in lexicographical order. Since the Weyl group leaves the inner product invariant, if T, S are conjugate, then M_T can be obtained from M_S by applying a permutation to both the rows and the columns. So in that case $\widehat{M}_T = \widehat{M}_S$. Hence we have a third invariant.

We have three applications of these invariants: to make the computation of the stabilizer W_T more efficient, to reduce the number of sets S that we attempt to insert in the list L in Step (2c), and reduce the number of conjugacy tests that we perform in Step (2c) (that is, the number of calls to `RepresentativeAction`).

The computation of the stabilizer. We can use the invariant $\sigma(T)$ to compute a subgroup $W' \subset W$ such that the $W'_T = W_T$, making the computation of the stabilizer more efficient. For this we consider the sum vector $\xi(T)$. We compute a $w \in W$ such that $w(\xi(T))$ is dominant. Let $\lambda_1, \dots, \lambda_\ell$ denote the fundamental weights as above and write $w(\xi(T)) = m_1\lambda_1 + \cdots + m_\ell\lambda_\ell$. Then the stabilizer $W_{w(\xi(T))}$ is generated by the s_{α_i} where i is such that $m_i = 0$ ([Hum72], Lemma 10.3B). Obviously $W_{\xi(T)} = w^{-1}W_{w(\xi(T))}w$. We see that it is straightforward to compute the group $W_{\xi(T)}$ and we can let this be W' . Furthermore, in practice it is useful to check whether the generators of W' leave T invariant. If this is the case then we have the stabilizer without any further computation.

Reducing the number of sets S in Step (2c). Note that the algorithm starts with the closed set Φ^+ and the other closed sets are constructed as subsets of that. Hence each closed set produced by the algorithm is contained in Φ^+ .

In Step (2c) of the algorithm different non-conjugate sets T of size $n + 1$ can produce conjugate successors S of size n (and usually this happens rather often). In order to reduce the number of closed sets S that go into the second part of the step (that is, the conjugacy testing) we first compute all sets $T' \subset \Phi^+$ having S as a successor, and only continue with S if $\sigma(T)$ is minimal among all $\sigma(T')$. (For this we can use any order on the weights, for example the lexicographical order on coordinate vectors.) If it is not, then we are certain that a set conjugate to S will be constructed as successor of another set of size $n + 1$. More precisely we replace Step (2c) by the following

(2c) For each such orbit take a representative α and set $S = T \setminus \{\alpha\}$.

- (i) Compute $\mathcal{N}^+(S) = \{\beta \in \Phi^+ \mid \beta \notin S, S \cup \{\beta\} \text{ is closed}\}$.
- (ii) For each $\beta \in \mathcal{N}^+(S)$ set $T' = S \cup \{\beta\}$ and compute $\sigma(T')$.
- (iii) If $\sigma(T)$ is minimal among all $\sigma(T')$ then check whether S is not conjugate to any element of L . If it is not then add S to L .

Reduce the number of conjugacy tests. Next we come to the conjugacy testing. In order to limit the necessary tests we use the invariants \widehat{M}_S and $\delta^{-1}(S)$. Only if a set S' in the list L has $\widehat{M}_S = \widehat{M}_{S'}$ and $\delta^{-1}(S) = \delta^{-1}(S')$ we do a conjugacy test, otherwise we already know that S, S' are not conjugate. Often no S' in L has $\widehat{M}_S = \widehat{M}_{S'}$ and $\delta^{-1}(S) = \delta^{-1}(S')$ and in those cases S can directly be added to L . This explains the relatively low number of conjugacy tests in the sixth column of Table 1.

Remark 3.6. • One other powerful invariant is the permanent of the matrix M_T .

Experiments show that it is very effective. However, it is difficult to compute the permanent of a matrix, and the time needed to compute the permanents outweighs the gains made from using this invariant.

- Of course it is possible to use the invariants $\delta^{-1}, \widehat{M}_T$ also for reducing the number of sets S in Step (2c)(ii). In fact, experiments show that these manage to reduce the number of sets even further. However, the number of sets T' for which these invariants have to be computed can be very large (see the fourth column of Table 1). Therefore the invariant σ works better because it is much easier to compute. Similarly, σ could also be used to limit the number of conjugacy tests. But here the extra time needed to compute the invariants \widehat{M}_T and δ^{-1} pays off: these invariants limit the number of conjugacy tests more effectively.
- Many alternatives to our algorithm are possible. Here we briefly go into two of them. Instead of our approach to conjugacy testing it is possible to compute a minimal image of each closed set; that is, a closed set T is replaced by the minimal element of its orbit (in some ordering, for example the lexicographical). Then the conjugacy problem boils down to testing equality. In [JJPW19] algorithms for computing minimal images are given which have been implemented in the GAP4 package `images` ([JJPW18]). However, it appears that computing a minimal image of each closed set in the output of our algorithm takes longer than listing the closed sets in the first place (for example, for the root system of type E_6 we find 94635 special closed sets which are listed in 149 seconds, whereas the computation of a minimal image of each of the 94635 closed sets took 443 seconds). Another more general purpose algorithm that can be used for our problem is described in [Bet18]. The rough structure of this algorithm is similar to ours. It starts with two sets \mathcal{A} and \mathcal{B} (in our case these are the special closed sets of sizes $n+1$ and n respectively) on which a group G acts (in our case the Weyl group). Also given is a relation $\mathcal{R} \subset \mathcal{A} \times \mathcal{B}$ (in our case this is the successor relation). Knowing the G -orbits in \mathcal{A} the algorithm computes the G -orbits in \mathcal{B} . We have not implemented this algorithm, but it seems to be hard to combine it with the invariants that we use and which are very successful at making our algorithm more efficient. Probably it would be possible to modify the algorithm in [Bet18] to also use those invariants, but then the result would be a method very similar to ours.
- There always is the obvious question as to the reliability of our implementations. In other words: how can we be sure that output of our computations is correct. Of

course we can never be absolutely sure. However, we do have some circumstantial evidence that our implementations are correct. Firstly, in the course of our work we also have tried completely different algorithms, and the output has always been the same. Secondly, for the root systems of type A_n the number of special closed sets that we get is confirmed by the data in [BM02].

3.2. Practical performance. Here we present some data on the practical performance of our algorithm. In Table 1 we list the running times of various parts of the algorithm. As test root systems we have taken F_4 , E_6 , B_6 , B_7 . The second column of this table has the total time needed to compute the stabilizers W_T . The third column lists the time taken to compute the orbits of W_T on $T \setminus (T + T)$. The fourth column has the time taken to compute $\sigma(T)$ for the various T (to limit the number of sets S that we try to insert in the list L). The fifth column displays the total time taken to compute the invariants $\widehat{M}_T, \delta^{-1}$. The sixth column has the total time taken to do the conjugacy tests (that is, the time taken in calls to `RepresentativeAction`). The integer following the running times in these columns is the number of times the algorithm entered the computation (so, for example, for B_6 there have been 1273371 computations of $\sigma(T)$). The seventh and eighth columns show, respectively, the number of closed sets in the output and the total time taken. The last column has the number of miliseconds taken for each closed set in the output (this is the quotient of the entries in the eighth and seventh columns).

We list a few obvious observations:

- The parts contributing most to the running time are the computation of the stabilizers (which is done for almost every final closed set), and the computation of the invariant σ (which is done a huge number of times).
- The conjugacy test is not performed very often (compared to the number of closed sets in the output).
- The time taken per closed set increases markedly with increasing rank.

type	Stab	Orb	Invariant σ	Invariants $\widehat{M}_T, \delta^{-1}$	Conjugacy	Number of closed sets	Time	ms per closed set
F_4	0.9	0.3	0.4	0.4	0.12	3579	2.2	0.61
	3577	3577	19889	3954	410			
E_6	56.7	9.6	36.6	24.2	17.8	94635	149	1.57
	94634	94634	828924	115652	30925			
B_6	90.2	16.9	50.5	23.5	13.4	151386	201	1.33
	151384	151384	1273371	174243	27234			
B_7	5009	659	4656	1506	1414	6115473	13803	2.26
	6115471	6115471	69416764	7223257	1718047			

TABLE 1. Running times (in seconds) of various parts of the algorithm. See the text for a description of the contents of the columns.

3.3. Listing the non-special closed sets. Now we turn to the second part of the procedure, that is, to finding the non-special closed sets, given the special ones. We will

outline an algorithm for obtaining the closed sets Q such that $Q^u = T$, where T is a given special closed set. Since we have the list of all special closed sets, by Lemma 2.1 this suffices to get all closed sets. We start with the following simple lemma.

Lemma 3.7. *Let T be a special closed set. Let $\alpha \in \Phi \setminus (T \cup -T)$. There is a closed set S such that $S^u = T$ and $\alpha, -\alpha \in S^r$ if and only if $T \cup \{\alpha, -\alpha\}$ is closed.*

Proof. If there is such an S , then Lemma 2.1 shows that $T \cup \{\alpha, -\alpha\}$ is closed. For the converse we can take $S = T \cup \{\alpha, -\alpha\}$. \square

The next lemma is similar to [Sop04], Proposition 5.

Lemma 3.8. *Let T be a special closed set. Set $S = \{\alpha \in \Phi \setminus (T \cup -T) \mid T \cup \{\alpha, -\alpha\} \text{ is closed}\}$. Then $R = S \cup T$ is a closed set with $R^r = S$, $R^u = T$.*

Proof. We first show that S is closed. Let $\alpha, \beta \in S$ be such that $\alpha + \beta \in \Phi$. We show that $T \cup \{\alpha + \beta, -\alpha - \beta\}$ is closed. Let $\gamma \in T$ be such that $\alpha + \beta + \gamma \in \Phi$. Then at least two of $\alpha + \beta$, $\alpha + \gamma$, $\beta + \gamma$ are roots. (If Φ is simply laced, this can be shown by using the property that for $\delta, \epsilon \in \Phi$ we have that $\delta + \epsilon \in \Phi$ if and only if $(\delta, \epsilon) < 0$. In general the intersection of Φ and the set of linear combinations of α, β, γ is a root system of rank at most 3. For those root systems it is easy to show the statement by a brute force verification.) So at least one of $\alpha + \gamma$, $\beta + \gamma$ lies in Φ . Suppose $\alpha + \gamma \in \Phi$. Since $T \cup \{\alpha, -\alpha\}$ is closed, $\alpha + \gamma \in T$. But then also $\beta + \alpha + \gamma \in T$. In the same way we see that if $-\alpha - \beta + \gamma \in \Phi$ then $-\alpha - \beta + \gamma \in T$. Hence $T \cup \{\alpha + \beta, -\alpha - \beta\}$ is closed and the same follows for S .

Now it is obvious that R is closed. Furthermore, if $\alpha \in S$ then also $-\alpha \in S$ so that $R^r = S$, implying also $R^u = T$. \square

Now fix a special closed set T . The algorithm first constructs the set S as in the previous lemma. By Lemma 3.7, S contains all closed sets that are the symmetric part of a closed set whose special part is T . So an immediate way of proceeding would be to enumerate all symmetric closed subsets of S , for each such set P form the closed set $P \cup T$, and get rid of conjugate copies. However, there is a much more efficient way of doing it.

First note that S is a root subsystem of Φ . Let $\alpha \in S$, then the reflection s_α sends T to itself. (Indeed, if $\beta \in T$ then the string $\{\beta + k\alpha \mid k \in \mathbb{Z}\}$ is contained in S as S is closed, and it has to be contained in T by Lemma 2.1.) So T is stabilized by the Weyl group W^S of S . Hence, if two closed symmetric subsets $P, P' \subset S$ are conjugate under W^S , then $P \cup T, P' \cup T$ are as well, so that they are conjugate under W . Now we use Dynkin's algorithm ([Dyn52], see [dGr17], §5.9 for a more recent description) to find all symmetric closed subsets of S up to conjugacy by W^S . For each symmetric closed set P found we form the set $P \cup T$. Let W_T be the stabilizer of T in W . Two closed subsets $P \cup T, P' \cup T$ are conjugate under W if and only if they are under W_T . So we erase the W_T -conjugate copies from the list and we are done. Note that in order to execute this we do not have to compute the stabilizers W_T as they have already been constructed by the algorithm of Section 3.1.

By comparing the running times for the root systems F_4, B_5, B_6, E_6 in Tables 1, 10 we see that computing the non-special closed sets does not weigh heavily on the running time. The main part of the computation time is taken for listing the special closed sets.

4. CLOSED SUBSETS OF IRREDUCIBLE ROOT SYSTEMS

In this section, we present the full classification of the closed subsets of the irreducible root systems of rank 3 derived from the implementation of our algorithm. This classification is contained in Tables 2 through 9. In this section we also present summary data for the classifications of closed subsets of the irreducible root systems of ranks 4 through 7 in Table 10.

For each root system, the classification is divided into three parts: The symmetric closed subsets T^r , for which the special part is empty $T^u = \emptyset$; the special closed subsets T^u , for which the symmetric part is empty $T^r = \emptyset$; and the Levi decomposable closed subsets $T^r \cup T^u$, for which $T^r \neq \emptyset$ and $T^u \neq \emptyset$.

For symmetric closed subsets, we identify the isomorphism classes. The symmetric closed subsets have been computed with the implementation of Dynkin's algorithm ([Dyn52]) in the SLA package ([dGr11]) of GAP4.

	Special closed subsets \mathbf{T}^u ($\mathbf{T}^r = \mathbf{0}$)
1	$\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
2	$\alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
3	$\alpha_1, \alpha_2, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
4	$\alpha_1, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
5	$\alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
6	$\alpha_1, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
7	$\alpha_2, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
8	$\alpha_3, \alpha_1 + \alpha_2, \alpha_1 + \alpha_2 + \alpha_3$
9	$\alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
10	$\alpha_3, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
11	$\alpha_1, \alpha_1 + \alpha_2, \alpha_1 + \alpha_2 + \alpha_3$
12	$\alpha_1 + \alpha_2, \alpha_2 + \alpha_3$
13	$\alpha_1 + \alpha_2, \alpha_1 + \alpha_2 + \alpha_3$
14	$\alpha_3, \alpha_1 + \alpha_2 + \alpha_3$
15	α_3
	Levi decomposable closed subsets $\mathbf{T}^r \oplus \mathbf{T}^u$ ($\mathbf{T}^r, \mathbf{T}^u \neq \mathbf{0}$)
16	$\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_1$
17	$\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_3$
18	$\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_2$
19	$\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_1, -\alpha_3$
20	$\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_1, -\alpha_2, -\alpha_1 - \alpha_2$
21	$\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_2, -\alpha_3, -\alpha_2 - \alpha_3$
22	$\alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_3$
23	$\alpha_1, \alpha_2, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_1$
24	$\alpha_1, \alpha_3, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_1$
25	$\alpha_1, \alpha_2, \alpha_1 + \alpha_2, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_2$
26	$\alpha_1 + \alpha_2, \alpha_2 + \alpha_3, -\alpha_1 - \alpha_2$
27	$\alpha_3, \alpha_1 + \alpha_2, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_3$
28	$\alpha_3, \alpha_1 + \alpha_2, \alpha_1 + \alpha_2 + \alpha_3, -\alpha_1 - \alpha_2$

TABLE 2. The special and Levi decomposable closed subsets of the root system A_3 .

	Symmetric closed subsets \mathbf{T}^r ($\mathbf{T}^u = \mathbf{0}$)	Isomorphism class
29	$\alpha_1, -\alpha_1$	A_1
30	$\alpha_1, \alpha_2, \alpha_1 + \alpha_2, -\alpha_1, -\alpha_2, -\alpha_1 - \alpha_2$	A_2
31	$\alpha_1, \alpha_3, -\alpha_1, -\alpha_3$	$A_1 \times A_1$
32	Φ	A_3

TABLE 3. The symmetric closed subsets of the root system A_3 .

	Symmetric closed subsets \mathbf{T}^r ($\mathbf{T}^u = \mathbf{0}$)	Isomorphism class
80	$\alpha_1, -\alpha_1$	A_1
81	$\alpha_3, -\alpha_3$	A_1
82	$\alpha_1, \alpha_3, -\alpha_1, -\alpha_3$	$A_1 \times A_1$
83	$\alpha_1, \alpha_1 + 2\alpha_2 + 2\alpha_3, -\alpha_1, -\alpha_1 - 2\alpha_2 - 2\alpha_3$	$A_1 \times A_1$
84	$\alpha_1, \alpha_2, \alpha_1 + \alpha_2, -\alpha_1, -\alpha_2, -\alpha_1 - \alpha_2$	A_2
85	$\alpha_1, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, \alpha_1 + 2\alpha_2 + 2\alpha_3,$ $-\alpha_1, -\alpha_2 - \alpha_3, -\alpha_1 - \alpha_2 - \alpha_3, -\alpha_1 - 2\alpha_2 - 2\alpha_3$	B_2
86	$\alpha_1, \alpha_3, \alpha_1 + 2\alpha_2 + 2\alpha_3, -\alpha_1, -\alpha_3, -\alpha_1 - 2\alpha_2 - 2\alpha_3$	$A_1 \times A_1 \times A_1$
87	$\alpha_1, \alpha_2, \alpha_1 + \alpha_2, \alpha_2 + 2\alpha_3, \alpha_1 + \alpha_2 + 2\alpha_3, \alpha_1 + 2\alpha_2 + 2\alpha_3,$ $-\alpha_1, -\alpha_2, -\alpha_1 - \alpha_2, -\alpha_2 - 2\alpha_3, -\alpha_1 - \alpha_2 - 2\alpha_3, -\alpha_1 - 2\alpha_2 - 2\alpha_3$	A_3
88	Φ	B_3

TABLE 6. The symmetric closed subsets of the root system B_3 .

	Symmetric closed subsets \mathbf{T}^r ($\mathbf{T}^u = \mathbf{0}$)	Isomorphism class
79	$\alpha_1, -\alpha_1$	A_1
80	$\alpha_3, -\alpha_3$	A_1
81	$\alpha_1, \alpha_3, -\alpha_1, -\alpha_3$	$A_1 \times A_1$
82	$\alpha_3, 2\alpha_2 + \alpha_3, -\alpha_3, -2\alpha_2 - \alpha_3$	$A_1 \times A_1$
83	$\alpha_1, \alpha_2, \alpha_1 + \alpha_2, -\alpha_1, -\alpha_2, -\alpha_1 - \alpha_2$	A_2
84	$\alpha_1, 2\alpha_2 + \alpha_3, \alpha_1 + 2\alpha_2 + \alpha_3, 2\alpha_1 + 2\alpha_2 + \alpha_3,$ $-\alpha_1, -2\alpha_2 - \alpha_3, -\alpha_1 - 2\alpha_2 - \alpha_3, -2\alpha_1 - 2\alpha_2 - \alpha_3$	B_2
85	$\alpha_3, 2\alpha_2 + \alpha_3, 2\alpha_1 + 2\alpha_2 + \alpha_3, -\alpha_3, -2\alpha_2 - \alpha_3, -2\alpha_1 - 2\alpha_2 - \alpha_3$	$A_1 \times A_1 \times A_1$
86	$\alpha_1, \alpha_3, 2\alpha_2 + \alpha_3, \alpha_1 + 2\alpha_2 + \alpha_3, 2\alpha_1 + 2\alpha_2 + \alpha_3,$ $-\alpha_1, -\alpha_3, -2\alpha_2 - \alpha_3, -\alpha_1 - 2\alpha_2 - \alpha_3, -2\alpha_1 - 2\alpha_2 - \alpha_3$	$A_1 \times B_2$
87	Φ	C_3

TABLE 9. The symmetric closed subsets of the root system C_3 .

Root system	Number of special closed subsets	Number of Levi decomposable closed subsets	Number of symmetric closed subsets	Total Number of closed subsets	Time (s)
A_3	15	13	4	32	0.02
B_3	46	33	9	88	0.05
C_3	44	34	9	87	0.05
A_4	62	70	6	138	0.07
B_4	429	311	19	759	0.4
C_4	401	334	19	754	0.4
D_4	110	102	11	223	0.4
F_4	3579	1242	23	4844	3.0
A_5	317	390	10	717	0.4
B_5	6267	3592	35	9894	7.6
C_5	5744	4074	35	9853	7.7
D_5	1145	877	15	2037	1.4
A_6	2044	2476	14	4534	3.3
B_6	151386	61001	64	212451	257
C_6	137456	74081	64	211601	239
D_6	20549	11310	31	31890	36
E_6	94635	29487	20	124142	172
A_7	16998	18959	21	35978	33
B_7	6115473	1648192	109	7763774	15569
C_7	5523891	2163473	109	7687473	13781
D_7	623510	228140	44	851694	1498
E_7	144937928	10347518	46	155285492	weeks

TABLE 10. Closed subsets for the ranks 3 through 7 irreducible root systems, up to conjugation by the respective Weyl group. We exclude the empty set.

5. REGULAR SUBALGEBRAS

Let \mathfrak{g} be a semisimple Lie algebra and \mathfrak{h} a (fixed) Cartan subalgebra of \mathfrak{g} , with corresponding root system Φ . For $\alpha \in \Phi$ we denote \mathfrak{g}_α the corresponding root space. Let $T \subseteq \Phi$ be a closed subsystem. Let \mathfrak{t} be a subspace of \mathfrak{h} containing $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}]$ for $\alpha \in T$ such that $-\alpha \in T$. Then

$$(2) \quad \mathfrak{s}_{T, \mathfrak{t}} = \mathfrak{t} \oplus \bigoplus_{\alpha \in T} \mathfrak{g}_\alpha$$

is a regular subalgebra of \mathfrak{g} . Moreover, all regular subalgebras of \mathfrak{g} arise in this manner.

Let G be the adjoint group of \mathfrak{g} . There are many ways to characterize G . It is the identity component of the automorphism group of \mathfrak{g} . It is the connected algebraic subgroup of $\mathrm{GL}(\mathfrak{g})$ with Lie algebra $\mathrm{ad}\mathfrak{g}$. Thirdly, it is the group generated by all $\exp(\mathrm{ad}x)$ for $x \in \mathfrak{g}_\alpha$, $\alpha \in \Phi$.

The Weyl group of Φ naturally acts on the dual space \mathfrak{h}^* . Further, by identifying \mathfrak{h} and \mathfrak{h}^* via the Killing form, the Weyl group also acts on \mathfrak{h} . Letting s_α denote the reflection corresponding to $\alpha \in \Phi$, this means that $s_\alpha(h) = h - \alpha(h)h_\alpha$, where h_α is the unique element of $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}]$ with $\alpha(h_\alpha) = 2$.

Proposition 5.1. *The regular subalgebras $\mathfrak{s}_{T_1, \mathfrak{t}_1}$ and $\mathfrak{s}_{T_2, \mathfrak{t}_2}$ are conjugate under G if and only if there is a $w \in W(\Phi)$ with $w(T_1) = T_2$ and $w(\mathfrak{t}_1) = \mathfrak{t}_2$.*

Proof. Let $g' \in G$ be such that $g'(\mathfrak{s}_{T_1, \mathfrak{t}_1}) = \mathfrak{s}_{T_2, \mathfrak{t}_2}$. Let $\mathfrak{n}_{\mathfrak{g}} = \{x \in \mathfrak{g} : [x, \mathfrak{s}_{T_2, \mathfrak{t}_2}] \subset \mathfrak{s}_{T_2, \mathfrak{t}_2}\}$ be the normalizer of $\mathfrak{s}_{T_2, \mathfrak{t}_2}$. This is the Lie algebra of the algebraic group $N_G(\mathfrak{s}_{T_2, \mathfrak{t}_2}) = \{g \in G : g(\mathfrak{s}_{T_2, \mathfrak{t}_2}) = \mathfrak{s}_{T_2, \mathfrak{t}_2}\}$. Since \mathfrak{h} and $g'(\mathfrak{h})$ are Cartan subalgebras of $\mathfrak{n}_{\mathfrak{g}}(\mathfrak{s}_{T_2, \mathfrak{t}_2})$, there is a $g'' \in N_G(\mathfrak{s}_{T_2, \mathfrak{t}_2})$ with $g''g'(\mathfrak{h}) = \mathfrak{h}$. Set $g_0 = g''g'$. Note that $g_0 \in N_G(\mathfrak{h})$. There is a canonical homomorphism $N_G(\mathfrak{h}) \rightarrow W$ with kernel $Z_G(\mathfrak{h})$ and let $w \in W(\Phi)$ be the image of g_0 . A short calculation shows that $g_0(\mathfrak{g}_\alpha) = \mathfrak{g}_\beta$ where $\beta(h) = \alpha(g_0^{-1}(h))$ for $h \in \mathfrak{h}$. But this implies that $\beta = w(\alpha)$. (This is easy to see if w is a reflection, and it then follows in general.) So we see that $w(T_1) = T_2$. Also $w(\mathfrak{t}_1) = g_0(\mathfrak{s}_{T_1, \mathfrak{t}_1} \cap \mathfrak{h}) = \mathfrak{s}_{T_2, \mathfrak{t}_2} \cap \mathfrak{h} = \mathfrak{t}_2$.

For the reverse implication, let $g_0 \in N_G(\mathfrak{h})$ map to w . Then $g_0(\mathfrak{g}_\alpha) = \mathfrak{g}_{w(\alpha)}$ and $g_0(\mathfrak{t}_1) = \mathfrak{t}_2$. It follows that $g_0(\mathfrak{s}_{T_1, \mathfrak{t}_1}) = \mathfrak{s}_{T_2, \mathfrak{t}_2}$. \square

Corollary 5.2. *Let $T \subseteq \Phi$ be a closed set and $W(\Phi)_T = \{w \in W(\Phi) : w(T) = T\}$ its stabilizer in $W(\Phi)$. The regular subalgebras $\mathfrak{s}_{T, \mathfrak{t}_1}$ and $\mathfrak{s}_{T, \mathfrak{t}_2}$ are conjugate under G if and only if there is a $w \in W(\Phi)_T$ such that $w(\mathfrak{t}_1) = \mathfrak{t}_2$.*

5.1. Regular subalgebras of the simple Lie algebra of type A_3 . The (complex) Lie algebra of type A_3 is the special linear algebra $\mathfrak{sl}(4, \mathbb{C})$ of traceless 4×4 complex matrices. The Cartan subalgebra \mathfrak{h} that we use consists of the diagonal elements of $\mathfrak{sl}(4, \mathbb{C})$. The Weyl group W is isomorphic to the symmetric group S_4 , acting on \mathfrak{h} by permuting the diagonal entries. More precisely, for $\pi \in S_4$ we have

$$(3) \quad \pi \cdot \mathrm{diag}(a_1, a_2, a_3, a_4) = \mathrm{diag}(a_{\pi(1)}, a_{\pi(2)}, a_{\pi(3)}, a_{\pi(4)}).$$

We have that W is generated by the simple reflections s_1, s_2, s_3 , and the isomorphism $W \rightarrow S_4$ maps $s_1 \mapsto (1, 2)$, $s_2 \mapsto (2, 3)$, $s_3 \mapsto (3, 4)$. A W -invariant scalar product on \mathfrak{h} is defined by

$$(4) \quad h \cdot h' = \mathrm{trace}(hh'), \quad h, h' \in \mathfrak{h}.$$

The simple roots of the root system of $\mathfrak{sl}(4, \mathbb{C})$ are $\alpha_1, \alpha_2, \alpha_3$. The other positive roots are $\alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$.

We now wish to classify the regular subalgebras of $\mathfrak{sl}(4, \mathbb{C})$. To do so, for each closed subset T of the root system of type A_3 , we will determine the possible $\mathfrak{t} \subset \mathfrak{h}$ such that $\mathfrak{s}_{T, \mathfrak{t}}$ is a regular subalgebra. The cases in which \mathfrak{t} is 0- or 3-dimensional are trivial, so we only consider the 1- and 2-dimensional cases.

In the following example, we explicitly describe the process of determining the regular subalgebras obtained from a given closed subset of A_3 . Determining the regular subalgebras for all other closed subsets of A_3 is obtained in a similar manner, and the complete results are summarized in Table 11 through Table 13.

Example 5.3. Consider the special closed subset of A_3

$$(5) \quad T = \{\alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3\},$$

which is located in row 2 of Table 2. A calculation shows that the stabilizer of T , denoted W_T , is generated by the single element s_1 .

Since T consists of positive root vectors, we may construct a regular subalgebra

$$(6) \quad \mathfrak{s}_{T, \mathfrak{t}} = \mathfrak{t} \oplus \bigoplus_{\alpha \in T} \mathfrak{g}_\alpha$$

by letting \mathfrak{t} be any subspace of the Cartan subalgebra \mathfrak{h} . Then, the possible 1-dimensional \mathfrak{t} are spanned by the non-trivial traceless elements $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1 - a_2 - a_3)$ for $a_1, a_2, a_3 \in \mathbb{C}$.

Two such regular subalgebras determined by h_{a_1, a_2, a_3} and h_{b_1, b_2, b_3} , respectively, are W -conjugate if and only if $s_1 \cdot h_{a_1, a_2, a_3}$ is a scalar multiple of h_{b_1, b_2, b_3} by Corollary 5.2. Considering that s_1 acts as the permutation $(1, 2)$ on the entries of h_{a_1, a_2, a_3} , as described in Eq. (3), these regular subalgebras are W -conjugate if and only if (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) or (b_2, b_1, b_3) .

To determine the regular subalgebras constructed from 2-dimensional \mathfrak{t} , we use the fact that a 2-dimensional subspace is uniquely determined by a nonzero element of \mathfrak{h} perpendicular to it. Since the inner product is W -invariant, two subspaces are W -conjugate if and only if their perpendicular vectors are. Thus, the 2-dimensional \mathfrak{t} are the spaces orthogonal to h_{a_1, a_2, a_3} , for any $a_1, a_2, a_3 \in \mathbb{C}$, all of which are not zero.

Using the same reasoning as in the 1-dimensional case, regular subalgebras with \mathfrak{t} given by the space orthogonal to h_{a_1, a_2, a_3} , is conjugate to the space orthogonal to h_{b_1, b_2, b_3} if and only if (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) or (b_2, b_1, b_3) .

The results of this example are summarized in row 2 of Table 11.

T	W_T	\mathfrak{t}
1	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
2	s_1	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1 - a_2 - a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) or (b_2, b_1, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.
3	s_3	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(-a_1 - a_2 - a_3, a_1, a_2, a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.
4	s_2	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(-a_1 - a_2 - a_3, a_1, a_2, a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) or (b_2, b_1, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.
5-9	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
10	s_1, s_2	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1 - a_2 - a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff $\{a_1, a_2, a_3\}$ is a scalar multiple of $\{b_1, b_2, b_3\}$. Note: W_T consists of every permutation of the first three coordinates. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.
11	s_2, s_3	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(-a_1 - a_2 - a_3, a_1, a_2, a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff $\{a_1, a_2, a_3\}$ is a scalar multiple of $\{b_1, b_2, b_3\}$. Note: W_T consists of every permutation of the last three coordinates. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.
12	$s_1 s_3$	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1 - a_2 - a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff they're equal or (a_1, a_2, a_3) is a scalar multiple of $(b_2, b_1, -b_1 - b_2 - b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.
13	s_3	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(-a_1 - a_2 - a_3, a_1, a_2, a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff they're equal or (a_1, a_2, a_3) is a scalar multiple of (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.
14	$s_1 s_2 s_1$	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1 - a_2 - a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff they're equal or (a_1, a_2, a_3) is a scalar multiple of (b_3, b_2, b_1) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.
15	s_1	1-dimensional: Spanned by $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1 - a_2 - a_3)$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) or (b_2, b_1, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} with the same conjugacy condition.

TABLE 11. Special closed subsets T of A_3 with rows numbered according to the row numbering of Table 2; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T .

T	W_T	\mathfrak{t}
16	s_1	1-dimensional: $\langle \text{diag}(1, -1, 0, 0) \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(a, a, b, -2a - b)$, and all are non-conjugate.
17	s_3	1-dimensional: $\langle \text{diag}(0, 0, 1, -1) \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(-a - 2b, a, b, b)$, and all are non-conjugate.
18	s_2	1-dimensional: $\langle \text{diag}(0, 1, -1, 0) \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(-a - 2b, a, a, b)$, and all are non-conjugate.
19	s_1, s_3	$\langle \text{diag}(1, -1, 0, 0), \text{diag}(0, 0, 1, -1) \rangle$
20	s_1, s_2	$\langle \text{diag}(1, -1, 0, 0), \text{diag}(0, 1, -1, 0) \rangle$
21	s_2, s_3	$\langle \text{diag}(0, 1, -1, 0), \text{diag}(0, 0, 1, -1) \rangle$
22	s_1, s_3	1-dimensional: $\langle \text{diag}(0, 0, 1, -1) \rangle$ 2-dimensional: Spaces orthogonal to $h_{a,b,b} = \text{diag}(-a - 2b, a, b, b)$. The space orthogonal to $h_{a,b,b}$ is conjugate to that orthogonal to $h_{a',b',b'}$ iff $(-a' - 2b', a', b')$ is a scalar multiple of $(-a - 2b, a, b)$ or $(a, -a - 2b, b)$.
23	s_1, s_3	1-dimensional: $\langle \text{diag}(1, -1, 0, 0) \rangle$ 2-dimensional: Spaces orthogonal to $h_{a,a,b} = \text{diag}(a, a, b, -2a - b)$. The space orthogonal to $h_{a,a,b}$ is conjugate to that orthogonal to $h_{a',a',b'}$ iff $(a', b', -2a' - b')$ is a scalar multiple of $(a, b, -2a - b)$ or $(a, -2a - b, b)$.
24	s_1	1-dimensional: $\langle \text{diag}(1, -1, 0, 0) \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(a, a, b, -2a - b)$, and all are non-conjugate.
25	s_2	1-dimensional: $\langle \text{diag}(0, 1, -1, 0) \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(-2a - b, a, a, b)$, and all are non-conjugate.
26	$s_1 s_2 s_1$	1-dimensional: $\langle \text{diag}(1, 0, -1, 0) \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(a, b, a, -2a - b)$, and all are non-conjugate.
27	s_3	1-dimensional: $\langle \text{diag}(0, 0, 1, -1) \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(-a - 2b, a, b, b)$, and all are non-conjugate.
28	$s_1 s_2 s_1$	1-dimensional: $\langle \text{diag}(1, 0, -1, 0) \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(a, b, a, -2a - b)$, and all are non-conjugate.

TABLE 12. Levi decomposable closed subsets T of A_3 with rows numbered according to the row numbering of Table 2; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T .

T	W_T	\mathfrak{t}
29	s_1, s_3	1-dimensional: $\langle \text{diag}(1, -1, 0, 0) \rangle$ 2-dimensional: Spaces orthogonal to $h_{a,a,b} = \text{diag}(a, a, b, -2a - b)$. The space orthogonal to $h_{a,a,b}$ is conjugate to that orthogonal to $h_{a',a',b'}$ iff $(a', b', -2a' - b')$ is a scalar multiple of $(a, b, -2a - b)$ or $(a, -2a - b, b)$.
30	s_1, s_2	$\langle \text{diag}(1, -1, 0, 0), \text{diag}(0, 1, -1, 0) \rangle$
31	$s_1, s_3, s_2 s_1 s_3 s_2$	$\langle \text{diag}(1, -1, 0, 0), \text{diag}(0, 0, 1, -1) \rangle$

TABLE 13. Symmetric closed subsets T of A_3 with rows numbered according to the row numbering of Table 3; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . We omit the case $T = \Phi$, for which there are no nonempty \mathfrak{t} .

5.2. Regular subalgebras of the simple Lie algebra of type B_3 . The (complex) Lie algebra of type B_3 is the special orthogonal algebra $\mathfrak{so}(7, \mathbb{C})$. It is the Lie algebra of 7×7 complex matrices satisfying

$$(7) \quad X^t M = -M X,$$

where

$$(8) \quad M = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 0 & I_3 \\ 0 & I_3 & 0 \end{pmatrix}.$$

A Cartan subalgebra \mathfrak{h} is given by elements of the form

$$(9) \quad h = \text{diag}(0, \lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3),$$

for $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{C}$. The positive roots are

$$(10) \quad \Phi^+ = \left\{ \begin{array}{lll} \alpha_1, & \alpha_2, & \alpha_3, \\ \alpha_1 + \alpha_2, & \alpha_2 + \alpha_3, & \alpha_1 + \alpha_2 + \alpha_3, \\ \alpha_2 + 2\alpha_3, & \alpha_1 + \alpha_2 + 2\alpha_3, & \alpha_1 + 2\alpha_2 + 2\alpha_3 \end{array} \right\},$$

where

$$(11) \quad \begin{aligned} \alpha_1(h) &= \lambda_1 - \lambda_2, \\ \alpha_2(h) &= \lambda_2 - \lambda_3, \\ \alpha_3(h) &= \lambda_3. \end{aligned}$$

Define a change of basis

$$(12) \quad \beta_1 = \alpha_1 + \alpha_2 + \alpha_3, \quad \beta_2 = \alpha_2 + \alpha_3, \quad \beta_3 = \alpha_3.$$

The fundamental reflections are

$$(13) \quad \begin{aligned} s_i(\beta_i) &= \beta_{i+1}, & s_i(\beta_{i+1}) &= \beta_i, & i &= 1, 2, \\ s_i(\beta_j) &= \beta_j, & & & j &\neq i, i+1, i=1, 2, \\ s_3(\beta_3) &= -\beta_3, & s_3(\beta_i) &= \beta_i, & i &\neq 3. \end{aligned}$$

The Weyl group generated by the fundamental reflections is then seen to be isomorphic to $\mathbb{Z}_2^3 \rtimes S_3$: The isomorphism is given by $s_1 \mapsto (1, 2) \in S_3$, $s_2 \mapsto (2, 3) \in S_3$, and $s_3 \mapsto (0, 0, 1) \in \mathbb{Z}_2^3$. The actions of s_1 , s_2 , and s_3 on the simple roots α_1 , α_2 , and α_3 are

$$(14) \quad \begin{aligned} s_1(\alpha_1) &= -\alpha_1, & s_2(\alpha_1) &= \alpha_1 + \alpha_2, & s_3(\alpha_1) &= \alpha_1, \\ s_1(\alpha_2) &= \alpha_1 + \alpha_2, & s_2(\alpha_2) &= -\alpha_2, & s_3(\alpha_2) &= \alpha_2 + 2\alpha_3, \\ s_1(\alpha_3) &= \alpha_3, & s_2(\alpha_3) &= \alpha_2 + \alpha_3, & s_3(\alpha_3) &= -\alpha_3. \end{aligned}$$

The correspondence, due to the Killing form, between \mathfrak{h}^* and \mathfrak{h} determines an action of the Weyl group on \mathfrak{h} . Specifically, a computation shows that we have the correspondence

$$(15) \quad \text{diag}(0, \lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3) \longleftrightarrow 2(\lambda_1\beta_1 + \lambda_2\beta_2 + \lambda_3\beta_3).$$

The actions of s_1, s_2 , and s_3 on \mathfrak{h} are given by:

$$(16) \quad \begin{aligned} s_1 \cdot \text{diag}(0, \lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3) &= \text{diag}(0, \lambda_2, \lambda_1, \lambda_3, -\lambda_2, -\lambda_1, -\lambda_3), \\ s_2 \cdot \text{diag}(0, \lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3) &= \text{diag}(0, \lambda_1, \lambda_3, \lambda_2, -\lambda_1, -\lambda_3, -\lambda_2), \\ s_3 \cdot \text{diag}(0, \lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3) &= \text{diag}(0, \lambda_1, \lambda_2, -\lambda_3, -\lambda_1, -\lambda_2, \lambda_3). \end{aligned}$$

We now describe a set of generators for $\mathfrak{so}(7, \mathbb{C})$. Number matrix rows and columns $0, 1, 2, 3, -1, -2, -3$. Then, positive root vectors e_α and negative root vectors f_α corresponding to the simple roots are

$$(17) \quad \begin{aligned} e_{\alpha_1} &= E_{12} - E_{-2,-1}, & f_{\alpha_1} &= E_{21} - E_{-1,-2}, \\ e_{\alpha_2} &= E_{23} - E_{-3,-2}, & f_{\alpha_2} &= E_{32} - E_{-2,-3}, \\ e_{\alpha_3} &= 2E_{30} - E_{0,-3}, & f_{\alpha_3} &= E_{03} - 2E_{-3,0}. \end{aligned}$$

These root vectors determine the following basis of the Cartan subalgebra \mathfrak{h} :

$$(18) \quad \begin{aligned} h_{\alpha_1} &= \text{diag}(0, 1, -1, 0, -1, 1, 0), \\ h_{\alpha_2} &= \text{diag}(0, 0, 1, -1, 0, -1, 1), \\ h_{\alpha_3} &= \text{diag}(0, 0, 0, 2, 0, 0, -2). \end{aligned}$$

A W -invariant inner product on \mathfrak{h} is defined by

$$(19) \quad h \cdot h' = \text{trace}(hh').$$

We now wish to classify the regular subalgebras of $\mathfrak{so}(7, \mathbb{C})$. To do so, for each closed subset T of the root system of type B_3 , we will determine the possible $\mathfrak{t} \subset \mathfrak{h}$ such that $\mathfrak{so}_{T,\mathfrak{t}}$ is a regular subalgebra. The cases in which \mathfrak{t} is 0- or 3-dimensional are trivial, so we only consider the 1- and 2-dimensional cases.

In the following example, we explicitly describe the process of determining the regular subalgebras obtained from a given closed subset of B_3 . Determining the regular subalgebras for all other closed subsets of B_3 is obtained in a similar manner, and the complete results are summarized in Tables 14 through 20.

Example 5.4. Consider the Levi decomposable closed subset of B_3

$$(20) \quad T = \left\{ \begin{array}{l} \alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, \\ 2\alpha_2 + \alpha_3, \alpha_1 + 2\alpha_2 + \alpha_3, 2\alpha_1 + 2\alpha_2 + \alpha_3, -\alpha_3 \end{array} \right\},$$

which is located in row 47 of Table 5. A calculation shows that the stabilizer of T , denoted W_T , is generated by the single element s_3 .

Since T contains precisely one negative/positive root pair, $\{\alpha_3, -\alpha_3\}$, \mathfrak{t} must contain h_{α_3} . Hence, there is only one 1-dimensional \mathfrak{t} , which is given by $\langle h_{\alpha_3} \rangle$.

We now determine the 2-dimensional \mathfrak{t} , each of which must contain h_{α_3} . Each such subspace is determined by a vector orthogonal to it, and since \mathfrak{t} contains h_{α_3} , the orthogonal vector must be of the form $\text{diag}(0, a_1, a_2, 0, -a_1, -a_2, 0)$. Since s_3 fixes the vector $\text{diag}(0, a_1, a_2, 0, -a_1, -a_2, 0)$, Corollary 5.2 implies that each such subspace is non-conjugate.

The results of this example are summarized in row 47 of Table 18.

T	W_T	\mathfrak{t}
1	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
2	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
3	s_1	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) or (b_2, b_1, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
4	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ is a scalar multiple of (b_1, b_2, b_3) or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
5	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
6, 7	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
8	s_1, s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(a_1, a_2, \pm a_3)$ or $(a_2, a_1, \pm a_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
9	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
10	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
11	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_3, b_2) , or (b_1, b_2, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
12	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
13	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
14	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.

TABLE 14. Special closed subsets T –row numbers 1 to 14–of B_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(0, a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
15	s_1, s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of any permutation of the elements of (b_1, b_2, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
16	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_3, b_2) , or (b_1, b_2, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
17, 18	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
19	s_1, s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(a_1, a_2, \pm a_3)$ or $(a_2, a_1, \pm a_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
20	$s_1 s_3$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (a_1, a_2, a_3) or $(a_2, a_1, -a_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
21	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
22	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
23	$s_1 s_2 s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_3, b_2, b_1) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
24	$s_2 s_3 s_2$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
25, 26	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
27	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
28	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.

TABLE 15. Special closed subsets T —row numbers 15 to 28—of B_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(0, a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
29	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
30	s_2, s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, \pm b_3)$, or $(b_1, \pm b_3, \pm b_2)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
31	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
32	$s_1 s_3$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or $(b_2, b_1, -b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
33	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
34	$s_2 s_3 s_2, s_1 s_2 s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, b_3)$, or $(b_3, \pm b_2, b_1)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
35	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
36	$s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, \pm b_3)$ (all sign combinations included). 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
37	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
38	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
39	s_1, s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of any permutation of the elements of (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
40	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
41	$s_3, s_1 s_2 s_3 s_2 s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(\pm b_1, b_2, \pm b_3)$ (all sign combinations included). 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
42	$s_2 s_3 s_2$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.

TABLE 16. Special closed subsets T —row numbers 29 to 42—of B_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(0, a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
43	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
44	s_2, s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, \pm b_3)$, or $(b_1, \pm b_3, \pm b_2)$ (all sign combinations included). 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
45	$s_2, s_1 s_2 s_3 s_2 s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(\pm b_1, b_2, b_3)$, or $(\pm b_1, b_3, b_2)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
46	$s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$; and the spaces spanned by $\langle h_{a_1, a_2, a_2} \rangle \sim \langle h_{b_1, b_2, b_2} \rangle$ are conjugate if and only if (a_1, a_2, a_2) is a scalar multiple of $(b_1, \pm b_2, \pm b_2)$ (all sign combinations are included).

TABLE 17. Special closed subsets T –row numbers 43 to 46–of B_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(0, a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
47	s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, 0, -a_1, -a_2, 0)$, and all of them yield non-conjugate regular subalgebras.
48	s_1	1-dimensional: $\langle h_{\alpha_1} \rangle$. 2-dimensional \mathfrak{t} are orthogonal to $\text{diag}(0, a_1, a_1, a_3, -a_1, -a_1, -a_3)$, and all of them yield non-conjugate regular subalgebras.
49	s_2	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$, and all of them yield non-conjugate regular subalgebras.
50	s_1, s_3	1-dimensional: $\langle h_{\alpha_1} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_1, a_3, -a_1, -a_1, -a_3)$. The 2-dimensional space determined by (a_1, a_1, a_3) is conjugate to that determined by (b_1, b_1, b_3) iff (a_1, a_1, a_3) is a scalar multiple of $(b_1, b_1, \pm b_3)$.
51	s_1, s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, 0, -a_1, -a_2, 0)$. The 2-dimensional space determined by $(a_1, a_2, 0)$ is conjugate to that determined by $(b_1, b_2, 0)$ iff $(a_1, a_2, 0)$ is a scalar multiple of $(b_1, b_2, 0)$ or $(b_2, b_1, 0)$.
52	s_1, s_3	1-dimensional: None 2-dimensional: $\langle h_{\alpha_1}, h_{\alpha_3} \rangle$.
53	s_2	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$, and each is non-conjugate.
54	s_1, s_2	None
55	s_1	1-dimensional: $\langle h_{\alpha_1} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_1, a_3, -a_1, -a_1, -a_3)$, and each is non-conjugate.
56	s_2	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$, and each is non-conjugate.
57	s_1, s_3	1-dimensional: $\langle h_{\alpha_1} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_1, a_3, -a_1, -a_1, -a_3)$. The 2-dimensional space determined by (a_1, a_1, a_3) is conjugate to that determined by (b_1, b_1, b_3) if and only if (a_1, a_1, a_3) is a scalar multiple of $(b_1, b_1, \pm b_3)$.

TABLE 18. Levi decomposable closed subsets T —row numbers 47 to 57—of B_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(0, a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
58	$s_1 s_2 s_1$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, 2a_1, -a_1, -a_2, -2a_1)$. The 2-dimensional space determined by $(a_1, a_2, 2a_1)$ is conjugate to that determined by $(b_1, b_2, 2b_1)$ if and only if $(a_1, a_2, 2a_1)$ is a scalar multiple of $(b_1, b_2, 2b_1)$ or $(2b_1, b_2, b_1)$.
59	s_2, s_3	$\langle h_{\alpha_2}, h_{\alpha_3} \rangle$
60	$s_2, s_3 s_2 s_3$	1-dimensional: $\langle h_{\alpha_2} \rangle$ 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$. The 2-dimensional space determined by (a_1, a_2, a_2) is conjugate to that determined by (b_1, b_2, b_2) if and only if (a_1, a_2, a_2) is a scalar multiple of $(b_1, \pm b_2, \pm b_2)$ (all sign combinations are included).
61	$s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, 0, -a_1, -a_2, 0)$. The 2-dimensional space determined by $(a_1, a_2, 0)$ is conjugate to that determined by $(b_1, b_2, 0)$ if and only if $(a_1, a_2, 0)$ is a scalar multiple of $(b_1, \pm b_2, 0)$.
62	s_2, s_3	$\langle h_{\alpha_2}, h_{\alpha_3} \rangle$.
63	s_2	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$, and each is non-conjugate.
64	$s_2, s_3 s_2 s_3$	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$ and the spaces determined by h_{a_1, a_2, a_2} and h_{b_1, b_2, b_2} are conjugate if and only if (a_1, a_2, a_2) is a scalar multiple of $(b_1, \pm b_2, \pm b_2)$ (all sign combinations are included).
65	s_2, s_3	$\langle h_{\alpha_2}, h_{\alpha_3} \rangle$
66	s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, 0, -a_1, -a_2, 0)$, and all of them yield non-conjugate regular subalgebras.
67	$s_2 s_3 s_2, s_1 s_2 s_1$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, 2a_1, -a_1, -a_2, -2a_1)$. The 2-dimensional space determined by $(a_1, a_2, 2a_1)$ is conjugate to that determined by $(b_1, b_2, 2b_1)$ if and only if $(a_1, a_2, 2a_1)$ is a scalar multiple of $(b_1, \pm b_2, 2b_1)$ or $(2b_1, \pm b_2, b_1)$.
68	$s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, 0, -a_1, -a_2, 0)$. The 2-dimensional space determined by $(a_1, a_2, 0)$ is conjugate to that determined by $(b_1, b_2, 0)$ if and only if $(a_1, a_2, 0)$ is a scalar multiple of $(b_1, \pm b_2, 0)$.

TABLE 19. Levi decomposable closed subsets T —row numbers 58 to 68—of B_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(0, a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
69	s_2	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$, and each is non-conjugate.
70	s_1, s_2	None. (We must have $\mathfrak{t} = \mathfrak{h}$).
71	s_1	1-dimensional: $\langle h_{\alpha_1} \rangle$. 2-dimensional: Spaces are orthogonal to $\text{diag}(0, a_1, a_1, a_3, -a_1, -a_1, -a_3)$, and each is non-conjugate.
72	s_2	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$, and each is non-conjugate.
73	$s_3, s_1 s_2 s_3 s_2 s_1$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2 + \alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, 0, a_2, a_3, 0, -a_2, -a_3)$ and the space determined by $(0, a_2, a_3)$ is conjugate to that determined by $(0, b_2, b_3)$ if and only if $(0, a_2, a_3)$ is a scalar multiple of $(0, b_2, \pm b_3)$.
74	s_2	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$, and each is non-conjugate.
75	$s_3 s_2 s_3, s_2 s_3 s_2 s_3$	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$ and the space determined by (a_1, a_2, a_2) is conjugate to that determined by (b_1, b_2, b_2) if and only if (a_1, a_2, a_2) is a scalar multiple of $(b_1, \pm b_2, \pm b_2)$ (the signs match).
76	s_2, s_3	$\langle h_{\alpha_2}, h_{\alpha_3} \rangle$
77	$W_{s_2, s_1 s_2 s_3 s_2 s_1}$	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, a_1, a_2, a_2, -a_1, -a_2, -a_2)$ and the space determined by (a_1, a_2, a_2) is conjugate to that determined by (b_1, b_2, b_2) if and only if (a_1, a_2, a_2) is a scalar multiple of $(\pm b_1, b_2, b_2)$.
78	$s_3, s_1 s_2 s_3 s_2 s_1$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2 + \alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(0, 0, a_2, a_3, 0, -a_2, -a_3)$ and the space determined by $(0, a_2, a_3)$ is conjugate to that determined by $(0, b_2, b_3)$ if and only if $(0, a_2, a_3)$ is a scalar multiple of $(0, b_2, \pm b_3)$
79	$s_3, s_1 s_2 s_3 s_2 s_1$	$\langle h_{\alpha_2}, 2h_{\alpha_1} + h_{\alpha_3} \rangle$

TABLE 20. Levi decomposable closed subsets T –row numbers 69 to 79–of B_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(0, a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
80	$s_1, s_3, s_2 s_1 s_3 s_2 s_1 s_3 s_2$	1-dimensional: $\langle h_{\alpha_1} \rangle$ 2-dimensional: Spaces orthogonal to $h_{a,a,b}$. The space orthogonal to $h_{a,a,b}$ is conjugate to that orthogonal to $h_{a',a',b'}$ iff (a', a', b') is a scalar multiple of $(a, a, \pm b)$.
81	$s_1, s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{\alpha_3} \rangle$ 2-dimensional: Spaces orthogonal to $h_{a,b,0}$. The space orthogonal to $h_{a,b,0}$ is conjugate to that orthogonal to $h_{a',b',0}$ iff (a', b') is a scalar multiple of $(a, \pm b)$ or $(b, \pm a)$.

TABLE 21. Symmetric closed subsets T of B_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . We include only those closed systems of rank 1, as the others cases are straightforward. In this table $h_{a_1, a_2, a_3} = \text{diag}(0, a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

5.3. **Regular subalgebras of the simple Lie algebra of type C_3 .** The (complex) Lie algebra of type C_3 is the symplectic algebra $\mathfrak{sp}(6, \mathbb{C})$. It is the Lie algebra of 6×6 complex matrices satisfying

$$(21) \quad X^t M = -MX,$$

where

$$(22) \quad M = \begin{pmatrix} 0 & I_3 \\ -I_3 & 0 \end{pmatrix}.$$

A Cartan subalgebra \mathfrak{h} is given by elements of the form

$$(23) \quad h = \text{diag}(\lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3),$$

for $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{C}$. The positive roots are

$$(24) \quad \Phi^+ = \left\{ \begin{array}{lll} \alpha_1, & \alpha_2, & \alpha_3, \\ \alpha_1 + \alpha_2, & \alpha_2 + \alpha_3, & \alpha_1 + \alpha_2 + \alpha_3, \\ 2\alpha_2 + \alpha_3, & \alpha_1 + 2\alpha_2 + \alpha_3, & 2\alpha_1 + 2\alpha_2 + \alpha_3 \end{array} \right\},$$

where

$$(25) \quad \begin{aligned} \alpha_1(h) &= \lambda_1 - \lambda_2, \\ \alpha_2(h) &= \lambda_2 - \lambda_3, \\ \alpha_3(h) &= 2\lambda_3. \end{aligned}$$

Define a change of basis

$$(26) \quad \beta_1 = \alpha_1 + \alpha_2 + \frac{1}{2}\alpha_3, \quad \beta_2 = \alpha_2 + \frac{1}{2}\alpha_3, \quad \beta_3 = \frac{1}{2}\alpha_3.$$

The fundamental reflections are the same as those for B_3 :

$$(27) \quad \begin{aligned} s_i(\beta_i) &= \beta_{i+1}, & s_i(\beta_{i+1}) &= \beta_i, & i &= 1, 2, \\ s_i(\beta_j) &= \beta_j, & & & j &\neq i, i+1, i=1, 2, \\ s_3(\beta_3) &= -\beta_3, & s_3(\beta_i) &= \beta_i, & i &\neq 3. \end{aligned}$$

The Weyl group generated by the fundamental reflections is then seen to be isomorphic to $\mathbb{Z}_2^3 \rtimes S_3$: The isomorphism is given by $s_1 \mapsto (1, 2) \in S_3$, $s_2 \mapsto (2, 3) \in S_3$, and $s_3 \mapsto (0, 0, 1) \in \mathbb{Z}_2^3$. The actions of s_1 , s_2 , and s_3 on the simple roots α_1 , α_2 , and α_3 are

$$(28) \quad \begin{aligned} s_1(\alpha_1) &= -\alpha_1, & s_2(\alpha_1) &= \alpha_1 + \alpha_2, & s_3(\alpha_1) &= \alpha_1, \\ s_1(\alpha_2) &= \alpha_1 + \alpha_2, & s_2(\alpha_2) &= -\alpha_2, & s_3(\alpha_2) &= \alpha_2 + \alpha_3, \\ s_1(\alpha_3) &= \alpha_3, & s_2(\alpha_3) &= 2\alpha_2 + \alpha_3, & s_3(\alpha_3) &= -\alpha_3. \end{aligned}$$

The correspondence, due to the Killing form, between \mathfrak{h}^* and \mathfrak{h} determines an action of the Weyl group on \mathfrak{h} . Specifically, a computation shows that we have the correspondence

$$(29) \quad \text{diag}(\lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3) \longleftrightarrow 2(\lambda_1\beta_1 + \lambda_2\beta_2 + \lambda_3\beta_3).$$

The actions of s_1 , s_2 , and s_3 on \mathfrak{h} are given by:

$$(30) \quad \begin{aligned} s_1 \cdot \text{diag}(\lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3) &= \text{diag}(\lambda_2, \lambda_1, \lambda_3, -\lambda_2, -\lambda_1, -\lambda_3), \\ s_2 \cdot \text{diag}(\lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3) &= \text{diag}(\lambda_1, \lambda_3, \lambda_2, -\lambda_1, -\lambda_3, -\lambda_2), \\ s_3 \cdot \text{diag}(\lambda_1, \lambda_2, \lambda_3, -\lambda_1, -\lambda_2, -\lambda_3) &= \text{diag}(\lambda_1, \lambda_2, -\lambda_3, -\lambda_1, -\lambda_2, \lambda_3). \end{aligned}$$

We now describe a set of generators for $\mathfrak{sp}(6, \mathbb{C})$. Number matrix rows and columns 1, 2, 3, -1, -2, -3. Then, positive root vectors e_α and negative root vectors f_α corresponding to the simple roots are

$$(31) \quad \begin{aligned} e_{\alpha_1} &= E_{12} - E_{-2,-1}, & f_{\alpha_1} &= E_{21} - E_{-1,-2}, \\ e_{\alpha_2} &= E_{23} - E_{-3,-2}, & f_{\alpha_2} &= E_{32} - E_{-2,-3}, \\ e_{\alpha_3} &= E_{3,-3}, & f_{\alpha_3} &= E_{-3,3}. \end{aligned}$$

These root vectors determine the following basis of the Cartan subalgebra \mathfrak{h} :

$$(32) \quad \begin{aligned} h_{\alpha_1} &= \text{diag}(1, -1, 0, -1, 1, 0), \\ h_{\alpha_2} &= \text{diag}(0, 1, -1, 0, -1, 1), \\ h_{\alpha_3} &= \text{diag}(0, 0, 1, 0, 0, -1). \end{aligned}$$

A W -invariant inner product on \mathfrak{h} is defined by

$$(33) \quad h \cdot h' = \text{trace}(hh').$$

We now wish to classify the regular subalgebras of $\mathfrak{sp}(6, \mathbb{C})$. To do so, for each closed subset T of the root system of type C_3 , we will determine the possible $\mathfrak{t} \subset \mathfrak{h}$ such that $\mathfrak{s}_{T,\mathfrak{t}}$ is a regular subalgebra. The cases in which \mathfrak{t} is 0- or 3-dimensional are trivial, so we only consider the 1- and 2-dimensional cases.

In the following example, we explicitly describe the process of determining the regular subalgebras obtained from a given closed subset of C_3 . Determining the regular subalgebras for all other closed subsets of C_3 is obtained in a similar manner, and the complete results are summarized in Tables 22 through 29.

Example 5.5. Consider the symmetric closed subset $T = \{\alpha_1, -\alpha_1\}$ of C_3 , which is located in row 79 of Table 9. A calculation shows that the stabilizer of T , denoted W_T , is generated by the elements s_1 , s_3 , and $s_2s_1s_3s_2s_1s_3s_2$.

Clearly there is only one 1-dimensional \mathfrak{t} , which is given by $\langle h_{\alpha_1} \rangle$. We now determine the 2-dimensional \mathfrak{t} , each of which must contain h_{α_1} . Each such subspace is determined by a vector orthogonal to it, and since \mathfrak{t} contains h_{α_1} , the orthogonal vector must be of the form $\text{diag}(a, a, b, -a, -a, -b)$.

The actions of the generators of W_T on $\text{diag}(a, a, b, -a, -a, -b)$ are

$$(34) \quad \begin{aligned} s_1 \cdot \text{diag}(a, a, b, -a, -a, -b) &= \text{diag}(a, a, b, -a, -a, -b), \\ s_3 \cdot \text{diag}(a, a, b, -a, -a, -b) &= \text{diag}(a, a, -b, -a, -a, b), \\ s_2s_1s_3s_2s_1s_3s_2 \cdot \text{diag}(a, a, b, -a, -a, -b) &= \text{diag}(-a, -a, b, a, a, -b). \end{aligned}$$

Hence, by Corollary 5.2, the space orthogonal to $\text{diag}(0, a, a, b, -a, -a, -b)$ is conjugate to that orthogonal to $\text{diag}(a', a', b', -a', -a', -b')$ if and only if (a', a', b') is a scalar multiple of $(a, a, \pm b)$.

The results of this example are summarized in row 79 of Table 29.

T	W_T	\mathfrak{t}
1	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
2	s_1	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_2, b_1, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
3	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
4	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
5, 6	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
7	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
8	s_1, s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$, or $(b_2, b_1, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
9, 10	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
11	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.

TABLE 22. Special closed subsets T —row numbers 1 to 11—of C_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
12	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
13	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
14	s_1, s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of any permutation of the elements of (b_1, b_2, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
15, 16, 17	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
18	$s_1 s_3$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or $(b_2, b_1, -b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
19	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
20	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
21	$s_1 s_2 s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_3, b_2, b_1) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
22	s_2, s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, \pm b_3)$, or $(b_1, \pm b_3, \pm b_2)$ (all sign combinations included). 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.

TABLE 23. Special closed subsets T —row numbers 12 to 22—of C_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
23	$s_2 s_3 s_2$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
24	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
25	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
26	$s_1 s_2 s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_3, b_2, b_1) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
27	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
28	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
29	s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_1, b_3, b_2) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
30	s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
31	$s_1 s_3$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or $(b_2, b_1, -b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
32	s_1, s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of any permutation of the elements of (b_1, b_2, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
33	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.

TABLE 24. Special closed subsets T —row numbers 23 to 33—of C_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
34	1	\mathfrak{t} can be any subspace of \mathfrak{h} , and all of them yield non-conjugate regular subalgebras.
35	$s_3, s_2s_3s_2$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, \pm b_3)$ (all sign combinations included). 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
36	$s_2s_3s_2, s_2s_1s_2$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, b_3)$, or $(b_3, \pm b_2, b_1)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
37	s_1, s_2	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of any permutation of the elements of (b_1, b_2, b_3) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
38	$s_3s_2s_3$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or $(b_1, -b_3, -b_2)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
39	$s_1s_2s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_3, b_2, b_1) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
40	s_1, s_3	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, b_2, \pm b_3)$, or $(b_2, b_1, \pm b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
41	$s_1s_2s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or (b_3, b_2, b_1) . 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
42	$s_2s_3s_2$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, b_3)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
43	$s_2s_3s_2, s_1s_2s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of $(b_1, \pm b_2, b_3)$, or $(b_3, \pm b_2, b_1)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.
44	$s_1s_2s_1, s_1s_3s_2s_1$	1-dimensional: $\langle h_{a_1, a_2, a_3} \rangle$; $\langle h_{a_1, a_2, a_3} \rangle \sim \langle h_{b_1, b_2, b_3} \rangle$ iff (a_1, a_2, a_3) is a scalar multiple of (b_1, b_2, b_3) , or $(\pm b_3, b_2, b_1)$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_3} , with the same conjugacy condition.

TABLE 25. Special closed subsets T —row numbers 34 to 44—of C_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
45	$s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, 0}$. The 2-dimensional space determined by $(a_1, a_2, 0)$ is conjugate to that determined by $(b_1, b_2, 0)$ iff $(a_1, a_2, 0)$ is a scalar multiple of $(b_1, \pm b_2, 0)$.
46	s_1	1-dimensional: $\langle h_{\alpha_1} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(a_1, a_1, a_3, -a_1, -a_1, -a_3)$, and all of them yield non-conjugate regular subalgebras.
47	s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(a_1, a_2, 0, -a_1, -a_2, 0)$, and all of them yield non-conjugate regular subalgebras.
48	s_2	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(a_1, a_2, a_2, -a_1, -a_2, -a_2)$, and all of them yield non-conjugate regular subalgebras.
49	s_1, s_3	1-dimensional: $\langle h_{\alpha_1} \rangle$. 2-dimensional: Spaces orthogonal to h_{a_1, a_1, a_3} . The 2-dimensional space determined by (a_1, a_1, a_3) is conjugate to that determined by (b_1, b_1, b_3) iff (a_1, a_1, a_3) is a scalar multiple of $(b_1, b_1, \pm b_3)$.
50	s_1, s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, 0}$. The 2-dimensional space determined by $(a_1, a_2, 0)$ is conjugate to that determined by $(b_1, b_2, 0)$ iff $(a_1, a_2, 0)$ is a scalar multiple of $(b_1, b_2, 0)$, or $(b_2, b_1, 0)$.
51	s_1, s_3	$\langle h_{\alpha_1}, h_{\alpha_3} \rangle$
52	s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, 0}$, and all of them yield non-conjugate regular subalgebras.
53	s_1, s_2	$\langle h_{\alpha_1}, h_{\alpha_2} \rangle$
54	s_1	1-dimensional: $\langle h_{\alpha_1} \rangle$. 2-dimensional: Spaces orthogonal to h_{a_1, a_1, a_3} , and all of them yield non-conjugate regular subalgebras.
55	s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, 0}$, and all of them yield non-conjugate regular subalgebras.
56	$s_1 s_2 s_1$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_1} , and all of them yield non-conjugate regular subalgebras.

TABLE 26. Levi decomposable closed subsets T —row numbers 45 to 56—of C_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
57	s_2, s_3	$\langle h_{\alpha_2}, h_{\alpha_3} \rangle$
58	$s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces are orthogonal to $h_{a_1, a_2, 0}$. The 2-dimensional space determined by $(a_1, a_2, 0)$ is conjugate to that determined by $(b_1, b_2, 0)$ if and only if $(a_1, a_2, 0)$ is a scalar multiple of $(b_1, \pm b_2, 0)$.
59	$s_2, s_3 s_2 s_3$	1-dimensional: $\langle h_{\alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_2} . The 2-dimensional space determined by (a_1, a_2, a_2) is conjugate to that determined by (b_1, b_2, b_2) if and only if (a_1, a_2, a_2) is a scalar multiple of $(b_1, \pm b_2, \pm b_3)$, or $(b_1, \pm b_3, \pm b_2)$ (all sign combinations included).
60	s_2, s_3	$\langle h_{\alpha_2}, h_{\alpha_3} \rangle$
61	$s_2 s_3 s_2$	1-dimensional: $\langle h_{2\alpha_2 + \alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, 0, a_3}$, and all of them yield non-conjugate regular subalgebras.
62	s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, 0}$, and all of them yield non-conjugate regular subalgebras.
63	$s_1 s_2 s_1$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_1} , and all of them yield non-conjugate regular subalgebras.
64	s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, 0}$, and all of them yield non-conjugate regular subalgebras.
65	$s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{2\alpha_2 + \alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, 0, a_3}$. The 2-dimensional space determined by $(a_1, 0, a_3)$ is conjugate to that determined by $(b_1, 0, b_3)$ if and only if $(a_1, 0, a_3)$ is a scalar multiple of $(b_1, 0, \pm b_3)$.
66	$s_3, s_2 s_3 s_2$	$\langle h_{\alpha_2}, h_{\alpha_3} \rangle$
67	$s_2 s_3 s_2, s_2 s_1 s_2$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_1} . The 2-dimensional space determined by (a_1, a_2, a_1) is conjugate to that determined by (b_1, b_2, b_1) if and only if (a_1, a_2, a_1) is a scalar multiple of $(b_1, \pm b_2, b_1)$.

TABLE 27. Levi decomposable closed subsets T —row numbers 57 to 67—of C_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
68	$s_2 s_3 s_2, s_2 s_1 s_2$	1-dimensional: $\langle h_{2\alpha_2 + \alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, 0, a_3}$. The 2-dimensional space determined by $(a_1, 0, a_3)$ is conjugate to that determined by $(b_1, 0, b_3)$ if and only if $(a_1, 0, a_3)$ is a scalar multiple of $(b_1, 0, b_3)$, or $(b_3, 0, b_1)$.
69	$s_2 s_3 s_2, s_2 s_1 s_2$	$\langle h_{\alpha_1 + \alpha_2}, h_{2\alpha_2 + \alpha_3} \rangle$.
70	$s_3 s_2 s_3$	1-dimensional: $\langle h_{\alpha_2 + \alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, -a_2}$, and all of them yield non-conjugate regular subalgebras.
71	s_1, s_3	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, 0}$. The 2-dimensional space determined by $(a_1, a_2, 0)$ is conjugate to that determined by $(b_1, b_2, 0)$ if and only if $(a_1, a_2, 0)$ is a scalar multiple of $(b_1, b_2, 0)$, or $(b_2, b_1, 0)$.
72	$s_1 s_2 s_1$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_1} , and all of them yield non-conjugate regular subalgebras.
73	$s_2 s_3 s_2$	1-dimensional: $\langle h_{2\alpha_2 + \alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, 0, a_3}$, and all of them yield non-conjugate regular subalgebras.
74	$s_2 s_3 s_2, s_1 s_2 s_1$	1-dimensional: $\langle h_{2\alpha_2 + \alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $\text{diag}(a_1, 0, a_3, -a_1, 0, -a_3)$. The 2-dimensional space determined by $(a_1, 0, a_3)$ is conjugate to that determined by $(b_1, 0, b_3)$ if and only if $(a_1, 0, a_3)$ is a scalar multiple of $(b_1, 0, b_3)$, or $(b_3, 0, b_1)$.
75	$s_1 s_2 s_1, s_1 s_3 s_2 s_1$	$\langle h_{\alpha_3}, h_{\alpha_1 + \alpha_2} \rangle$
76	$s_3, s_1 s_2 s_3 s_2 s_1$	1-dimensional: $\langle h_{\alpha_3} \rangle$. 2-dimensional: Spaces orthogonal to $h_{a_1, a_2, 0}$. The 2-dimensional space determined by $(a_1, a_2, 0)$ is conjugate to that determined by $(b_1, b_2, 0)$ if and only if $(a_1, a_2, 0)$ is a scalar multiple of $(\pm b_1, b_2, 0)$.
77	$s_1 s_2 s_1, s_1 s_3 s_2 s_1 s_3$	1-dimensional: $\langle h_{\alpha_1 + \alpha_2} \rangle$. 2-dimensional: Spaces orthogonal to h_{a_1, a_2, a_1} . The 2-dimensional space determined by (a_1, a_2, a_1) is conjugate to that determined by (b_1, b_2, b_1) if and only if (a_1, a_2, a_1) is a scalar multiple of (b_1, b_2, b_1) , or $(-b_1, b_2, -b_1)$.
78	$s_1 s_2 s_1, s_1 s_3 s_2 s_1$	$\langle h_{\alpha_3}, h_{2\alpha_1 + 2\alpha_2 + \alpha_3} \rangle$

TABLE 28. Levi decomposable closed subsets T —row numbers 68 to 78—of C_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . In this table $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

T	W_T	\mathfrak{t}
79	$s_1, s_3, s_2 s_1 s_3 s_2 s_1 s_3 s_2$	1-dimensional: $\langle h_{\alpha_1} \rangle$ 2-dimensional: Spaces orthogonal to $h_{a,a,b}$. The space orthogonal to $h_{a,a,b}$ is conjugate to that orthogonal to $h_{a',a',b'}$ iff (a', a', b') is a scalar multiple of $(a, a, \pm b)$.
80	$s_1, s_3, s_2 s_3 s_2$	1-dimensional: $\langle h_{\alpha_3} \rangle$ 2-dimensional: Spaces orthogonal to $h_{a,b,0}$. The space orthogonal to $h_{a,b,0}$ is conjugate to that orthogonal to $h_{a',b',0}$ iff (a', b') is a scalar multiple of $(a, \pm b)$ or $(b, \pm a)$.

TABLE 29. Symmetric closed subsets T of C_3 ; the generators of the stabilizer of T , denoted W_T ; and the 1- and 2-dimensional subalgebras \mathfrak{t} of the Cartan subalgebra of the regular subalgebras determined by T . We include only those closed systems of rank 1, as the others cases are straightforward. In this table $h_{a_1, a_2, a_3} = \text{diag}(a_1, a_2, a_3, -a_1, -a_2, -a_3)$.

REFERENCES

- [Bet18] A. Betten, “How Fast Can We Compute Orbits of Groups?”, in: *Mathematical Software ICMS 2018*, J.H. Davenport, M. Kauers, G. Labahn, J. Urban editors, LNCS, volume 10931, 62-70 (2018).
- [Bor76] Z.I. Borevich, “Description of the subgroups of the general linear group that contain the group of diagonal matrices”. (Russian) *Rings and modules. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 64 (1976), 1229.
- [Bor77] Z.I. Borevich, “On the question of the enumeration of finite topologies,” (Russian) *Modules and representations. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 71 (1977), 47-65.
- [Bou68] N. Bourbaki. “Groupes et Algèbres de Lie, Chapitres 4, 5 et 6”, Hermann, Paris, 1968.
- [BM02] G. Brinkmann and B. D. McKay, “Posets on up to 16 points”. *Order* 19 (2002), no. 2, 147179.
- [GAP] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.9.2*; 2018, (<https://www.gap-system.org>).
- [dGr00] W. A. de Graaf: *Lie Algebras: Theory and Algorithms* (North Holland Mathematical Library, Elsevier, Amsterdam, 2000).
- [dGr17] W. A. de Graaf: *Computation with Linear Algebraic Groups* (CRC Press Boca Raton, 2017).
- [dGr09] W.A. de Graaf: *SLA-computing with Simple Lie algebras. a GAP package*, Version 1.5, (2018), <http://www.science.unitn.it/~degraaf/sla.html>.
- [dGr11] W.A. de Graaf, “Constructing semisimple subalgebras of semisimple Lie algebras,” *J. Algebra*, 325, 416-430 (2011).
- [DCH94] D. Ž. Đoković and P. Check and J.-Y. Hée, “On closed subsets of root systems”, *Canad. Math. Bull.*, 37, 338-345 (1994).
- [DR15] A. Douglas and J. Repka, “The Levi decomposable subalgebras of C_2 ,” *J. Math. Phys.*, **56** 051703, (2015).
- [DR16a] A. Douglas and J. Repka, “The subalgebras of $\mathfrak{so}(4, \mathbb{C})$,” *Comm. Algebra*, 44(12), (2016) 5269-5286.
- [DR16b] A. Douglas and J. Repka, “A classification of the subalgebras of A_2 ,” *J. Pure Appl. Algebra*, 220(6), (2016) 2389-2413.
- [DR17] A. Douglas and J. Repka, “The subalgebras of the rank two symplectic Lie algebra,” *Linear Algebra Appl.*, 527, (2017) 303-348.
- [DL11] M.J. Dyer and G.I. Lehrer, “Reflection subgroups of finite and affine Weyl groups,” *Trans. Amer. Math. Soc.*, 363, 5971-6005 (2011).
- [Dyn52] E.B. Dynkin, “Semisimple subalgebras of semisimple Lie algebras,” *Matematicheskii Sbornik N.S.* 30(72) (1952), no. 2, 349-462, (Russian).
- [HV96] A.L. Harebov and H.A. Vavilov, “On the lattice of subgroups of Chevalley groups containing a split maximal torus,” *Commun. Algebra*, 24(1), 109-133 (1996).
- [Hum72] J.E. Humphreys: *Introduction to Lie algebras and representation theory* (Springer-Verlag, New York, 1972).
- [JJPW18] C. Jefferson, E. Jonauskyste, M. Pfeiffer and R. Waldecker, *images - Minimal and Canonical images, a GAP package*, Version 1.1.0, (2018), <http://gap-packages.github.io/images/>.
- [JJPW19] C. Jefferson, E. Jonauskyste, M. Pfeiffer and R. Waldecker, “Minimal and canonical images”. *J. Algebra* 521, 481506 (2019).
- [LG72] M. Lorente and B. Gruber, “Classification of semisimple subalgebras of simple Lie algebras,” *J. Math. Phys.* 13 (1972) 1639-1663.
- [Mal80] F.M. Malyshev, “Decompositions of root systems,” *Math. Notes Acad. Sci. USSR* 27(1980), 418-421.
- [May16] E. Mayanskiy, “The subalgebras of G_2 ,” arXiv:1611.04070v1.
- [Ser03] Ákos Seress. “Permutation group algorithms”, volume 152 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2003.
- [Sop04] E.A. Sopkina, “On the sum of roots of a closed set,” *J. Math. Sci.* 124(1) (2004), 4832-4836.