

Nonlinear Reconnection in Magnetized Turbulence

NUNO F. LOUREIRO¹ AND STANISLAV BOLDYREV^{2,3}

¹*Plasma Science and Fusion Center, Massachusetts Institute of Technology, Cambridge MA 02139, USA*

²*Department of Physics, University of Wisconsin at Madison, Madison, WI 53706, USA*

³*Space Science Institute, Boulder, Colorado 80301, USA*

(Received July 24, 2019; Revised –; Accepted –)

Submitted to ApJ

ABSTRACT

Recent analytical works on strong magnetized plasma turbulence have hypothesized the existence of a range of scales where the tearing instability may govern the energy cascade. In this paper, we estimate the conditions under which such tearing may give rise to full nonlinear magnetic reconnection in the turbulent eddies, thereby enabling significant energy conversion and dissipation. When those conditions are met, a new turbulence regime is accessed where reconnection-driven energy dissipation becomes common, rather than the rare feature that it must be when they are not.

Keywords: magnetic fields — magnetic reconnection — plasmas — turbulence

1. INTRODUCTION

Magnetized plasmas are abundant in the Universe. Examples include the Earth’s magnetosphere, the solar wind and the solar corona, as well as many others, more distant and often more exotic, such as accretion disks around massive central objects, astrophysical jets, pulsar wind nebulae, etc. Many such environments, including all those just listed, are turbulent — a natural consequence of large-scale (roughly system-size) energy injection, and relatively infrequent collisions between the particles constituting those plasmas.

Beyond its interest as a fundamental physics prob-

of current sheets in this range (e.g. TenBarge & Howes 2013; Wan et al. 2015; Grošelj et al. 2018). Why this should be so has not been established on general theoretical grounds, but Boldyrev & Loureiro (2019) have recently advanced a possible explanation applicable to plasmas where $\beta_i \sim 1 \gg \beta_e$, such as found, for example, in the Earth’s magnetosheath (i.e., for so-called inertial kinetic-Alfvén wave turbulence (Chen & Boldyrev 2017; Passot et al. 2017, 2018; Roytershteyn et al. 2019)).

Current sheets being traditionally associated with magnetic reconnection (Biskamp 2000; Priest & Forbes 2000; Zweibel & Yamada 2009; Yamada et al. 2010), it is unsurprising that this process has acquired significant prominence as a potential key mechanism in magnetized turbulence (e.g. Matthaeus & Lamkin 1986; Retinò et al. 2007; Sundkvist et al. 2007; Servidio et al. 2009; Osman et al. 2014; Wan et al. 2015; Cerri et al. 2017; Shay et al. 2018). Fundamentally, reconnection leads to the conversion and dissipation of magnetic energy; thus, one expects that if it is indeed active in turbulence it may qualitatively impact the dynamics and observational signatures.

An important point that needs to be introduced in our discussion at this stage is that of the relationship — and distinction — between magnetic reconnection and the tearing mode (Furth et al. 1963). The latter is an instability which manifests itself through the reconnection of magnetic field lines [and the consequent opening of magnetic islands (or flux ropes)]. Strictly speaking, therefore, the tearing mode *can* be called magnetic reconnection; however, the term “reconnection” is most commonly used to refer to a strongly nonlinear plasma phenomenon associated with significant magnetic energy transfer and dissipation, as already mentioned. According to this classification, the deep nonlinear stage of evolution of the (strongly unstable, i.e., large instability parameter Δ') tearing mode (Coppi et al. 1976; Waelbroeck 1989; Jemella et al. 2003; Loureiro et al. 2005) is appropriately referred to as reconnection; but not its linear and early nonlinear stages. Let us now see why this distinction matters.

Recent works (Loureiro & Boldyrev 2017a; Mallet et al. 2017a; Boldyrev & Loureiro 2017; Loureiro & Boldyrev 2017b; Mallet et al. 2017b; Loureiro & Boldyrev 2018; Boldyrev & Loureiro 2018, 2019) have presented analytical arguments for the inevitability of the onset of the tearing mode (in either its resistive or collisionless forms, as appropriate) below a certain (so-called, critical) turbulence scale, $\lambda_{cr} \ll L$, where L is the energy injection scale, in a wide variety of plasma regimes. It is argued by these authors that the effect of the tearing mode is to redefine the energy cascade rate (to become

the tearing mode growth rate, see Section 2), resulting in a different energy spectrum and eddy anisotropy at scales $\lambda \ll \lambda_{cr}$. Magnetohydrodynamic simulations performed subsequently appear to lend support to these ideas (Walker et al. 2018; Dong et al. 2018), as does a detailed analysis of solar wind data (Vech et al. 2018). Additional consistent numerical evidence has been reported by Arzamasskiy et al. (2019); Landi et al. (2019) (specifically, the measurement of linear anisotropy of the turbulent fluctuations, $k_{\parallel} \sim k_{\perp}$, in the sub-ion range, as predicted for tearing-mediated inertial kinetic-Alfvén wave turbulence (Boldyrev & Loureiro 2019)).

Tearing onset in turbulence thus appears to be in reasonably strong footing — prompting the important question of whether it can (and, if so, under what conditions) lead to a fully nonlinear reconnecting stage. Essentially, the reason this question is non-trivial is that the reconnection rate differs from the tearing mode growth rate (and, therefore, from the eddy turnover rate in the tearing-mediated turbulence range). The goal of this paper is to address this issue¹.

2. PRELIMINARIES

The key idea underlying the suggestion that the tearing mode is activated at turbulence scales $\lambda \ll \lambda_{cr}$ derives from the observation that, at such scales, the tearing mode growth rate, $\gamma_t(\lambda)$, exceeds the eddy turnover rate, $\tau_{nl}^{-1}(\lambda)$, that would otherwise pertain to those scales, i.e.,

$$\gamma_t \tau_{nl} \gg 1, \quad (1)$$

with λ_{cr} resulting from solving this condition in the case of approximate equality (Loureiro & Boldyrev 2017a; Mallet et al. 2017a). It is demonstrated in these references that the specific mode (wavenumber) that solves Eq. (1), amongst all possible tearing-unstable modes, is the fastest-growing tearing mode (often dubbed the ‘Coppi’ mode (Coppi et al. 1976)).

The onset of the tearing mode, *per se*, is not sufficient to interfere with the turbulent cascade. Its ability to be dynamically significant naturally hinges on whether it can attain a nonlinear amplitude. In this regard, the tearing mode is a somewhat peculiar instability in that

¹ Hopefully, the reason for the somewhat tautological title of this paper is now clear. Reconnection is usually implicitly understood to be a nonlinear phenomenon. The specific phrasing of the title aims to stress the distinction between the linear and early nonlinear evolution of the tearing mode, on one hand, and its late, strongly nonlinear evolution on the other — the latter being what is meant here by the proper, or nonlinear, reconnection stage. This distinction is key, since the linear and early nonlinear stages of the tearing mode reconnect insignificant amounts of flux, and lead to negligible energy dissipation and conversion.

it becomes nonlinear at very small amplitudes: i.e., as soon as the width of the magnetic island that it creates exceeds the thickness of the inner boundary layer (which is, forcefully, asymptotically smaller than the characteristic length scale of variation of the background magnetic profile; i.e., in this case, than the size of the eddy, λ).² As the tearing mode begins its nonlinear evolution, it continues to grow exponentially at the same rate as in the linear stage³ (Wang & Bhattacharjee 1993; Porcelli et al. 2002; Loureiro et al. 2005). These notions imply that Eq. (1) correctly represents the condition for the nonlinear tearing mode to affect the turbulent cascade (Loureiro & Boldyrev 2017a; Boldyrev & Loureiro 2017; Mallet et al. 2017a). Furthermore, the tearing mode onset implies that γ_t becomes the eddy turnover rate at those scales, with a consequent change in the turbulence spectrum and other properties.

However, and as we now explain, it is less clear — but, we will argue, critical — whether the (early) nonlinear stage of the tearing mode evolution has the chance to evolve towards the deep nonlinear (i.e., properly reconnecting) stage, whereupon a significant amount of the magnetic flux in the eddy is reconnected, and considerable magnetic energy dissipation and conversion occurs.

In the absence of background turbulence, the (strongly unstable, large Δ') tearing mode is known to transition to a fully nonlinear reconnecting state once its amplitude becomes sufficiently large (Waelbroeck 1989; Jemella et al. 2003; Loureiro et al. 2005, 2013). At this moment, the tearing rate will, in most cases, change to a different value, usually referred to as the (normalized) reconnection rate, \mathcal{R} . The current understanding of reconnection suggests the following. In resistive MHD, there are two possibilities for \mathcal{R} , depending on the value of the Lundquist number, $S = L_{CS}v_A/\eta$, where L_{CS} is the current sheet length, v_A the Alfvén speed based on the reconnecting component of the magnetic field, and

² To be specific: purely from geometric considerations and the definition of separatrix, one has that the full width of a magnetic island is given by $W = 4\sqrt{-\tilde{\psi}/\psi''_{eq}}$, where $\tilde{\psi}$ is the perturbed (reconnected) flux, and $\psi''_{eq} \approx B_{eq}/a$ is the equilibrium current at the rational layer. The tearing mode becomes nonlinear when $W \approx \delta_{in}$, where δ_{in} is the width of the inner boundary layer that arises in the tearing mode calculation (Furth et al. 1963; Coppi et al. 1976); it scales with resistivity in the MHD regime, or with electron inertia in kinetic calculations and is, by definition, asymptotically smaller than a , the width of the background equilibrium. One thus finds that, in the early non-linear regime, $\tilde{\psi}/\psi_{eq} \approx 1/16 (\delta_{in}/a)^2 \ll 1$; note however that, numerical pre-factor aside, this condition implies that, for the Coppi mode, the perturbed and background currents are comparable at this stage.

³ This is not generally true for all wavenumbers unstable to tearing, but it is true for the most unstable (Coppi) mode.

η is the magnetic diffusivity. If $S \lesssim S_{cr} \approx 10^4$ we have $\mathcal{R} = S^{-1/2}$ (i.e., the Sweet-Parker rate (Parker 1957; Sweet 1958)). This is the only case where, in fact, the reconnection rate is the same as the tearing rate (of the most unstable tearing mode). However, this result is of limited applicability as, generally, $S \gg S_{cr}$; in such cases, one instead has $\mathcal{R} = S_{cr}^{-1/2} \approx 0.01$ (Loureiro et al. 2007; Samtaney et al. 2009; Bhattacharjee et al. 2009; Huang & Bhattacharjee 2010; Uzdensky et al. 2010; Loureiro et al. 2012; Loureiro & Uzdensky 2016). For collisionless reconnection, though absent a theoretical explanation (see, however, Liu et al. (2017)), it is generally accepted that $\mathcal{R} \approx 0.1$ (Birn et al. 2001; Comisso & Bhattacharjee 2016; Cassak et al. 2017).

For use in what follows, let us define the reconnection time as

$$\tau_{rec} = \mathcal{R}^{-1}\tau_{A,\lambda}, \quad (2)$$

where $\tau_{A,\lambda} = \lambda/v_{A,\lambda}$. The physical meaning of τ_{rec} is that it is the time that it takes to reconnect the magnetic flux contained in an eddy of size λ and reconnecting field $B_\xi(\lambda)$. The question which we wish to address is whether this reconnection rate is larger than the tearing rate, that is, whether an eddy distorted by the tearing instability may end up reconnecting significant magnetic flux, thus leading to significant energy conversion and dissipation. We propose that this will only happen if

$$\gamma_t\tau_{rec} \ll 1. \quad (3)$$

In other words, a typical eddy at scales $\lambda \gg \lambda_{cr}$ exists for a time of order γ_t^{-1} . It, therefore, has a finite probability of reaching the deep nonlinear stage, whereupon it may transition to the reconnection regime. If condition (3) is met, then the reconnection time is much shorter than the eddy turnover time, and it is thus expected that full reconnection will occur. Otherwise, reconnection is slower, and the eddy will cease to exist without significant reconnection having taken place. We now proceed to compute this condition, and discuss its implications, in three different cases: the pure MHD case, Section 3; and the cases when tearing, and reconnection, are enabled by kinetic physics (electron inertia) instead of resistivity, and the eddies in which they happen are above (Section 4.1) or below (Section 4.2) the ion kinetic scales.

3. THE MAGNETOHYDRODYNAMIC CASE

The onset of tearing in MHD turbulence has been addressed by Loureiro & Boldyrev (2017a); Mallet et al. (2017a); Boldyrev & Loureiro (2017). These authors find

$$\lambda_{cr}/L \sim S_L^{-4/7}, \quad (4)$$

where $S_L = LV_{A,0}/\eta$ is the outer scale Lundquist number, and $V_{A,0}$ is the Alfvén velocity based on the background (mean) field B_0 . Below this scale, the eddy turnover time becomes the growth rate of the fastest growing tearing mode:

$$\gamma_t \sim \tau_{A,\lambda}^{-1} (\lambda v_{A,\lambda}/\eta)^{-1/2}. \quad (5)$$

where (Boldyrev & Loureiro 2017)

$$v_{A,\lambda} \sim \varepsilon^{2/5} \eta^{-1/5} \lambda^{3/5}, \quad (6)$$

with $\varepsilon = V_{A,0}^3/L$ the injected power.

Therefore, evaluation of Eq. (3) yields the requirement

$$\lambda/L \gg \mathcal{R}^{-5/4} S_L^{-3/4}. \quad (7)$$

It is necessary for the validity of this result that $\lambda_{cr} \gg \lambda \gg \lambda_{diss}$, where $\lambda_{diss} \sim S_L^{-3/4} L$ (Boldyrev & Loureiro 2017) is the dissipation scale. Since $\mathcal{R} < 1$, the second inequality is automatically satisfied. As to the first, we find that it implies

$$S_L \gg \mathcal{R}^{-7}. \quad (8)$$

As mentioned before, as long as $S_\xi = \xi v_{A,\lambda}/\eta \gtrsim S_{cr} \sim 10^4$,⁴ the reconnection rate is $\mathcal{R} \sim S_{cr}^{-1/2} \sim 0.01$. We thus arrive at the conclusion that significant reconnection is only possible if $S_L \gg 10^{14}$, a considerable demand even by the standards of astrophysical and space plasmas — and certainly one that direct numerical simulations cannot be imagined to meet anytime in the foreseeable future⁵.

4. THE COLLISIONLESS CASE

Let us now examine the same question in a plasma where collisions are sufficiently rare that the breaking of frozen flux condition required to enable the tearing mode and the subsequent nonlinear reconnection stage is

⁴ To check whether this is true, we use Eq. (6) and the scaling $\xi \sim L(\lambda_{cr}/L)^{1/4}(\lambda/\lambda_{cr})^{9/5}$ (Boldyrev & Loureiro 2017), both of which expressions are valid in the tearing-mediated turbulence range. Then, we find that $S_\xi \gg S_{cr} \implies \lambda/L \gg S_{cr}^{5/12} S_L^{-73/84}$. This is smaller than λ_{cr}/L if $S_L \gg S_{cr}^{7/5} \sim 10^{28/5}$. This condition is superseded by Eq. (8).

⁵ It is straightforward to repeat this calculation for the case when the profile of the reconnecting magnetic field in the eddy is better represented by $\sin(x)$, instead of the default Harris ($\tanh(x)$) profile that we consider here. In that case, $\gamma_t \sim \tau_{A,\lambda}^{-1} (\lambda v_{A,\lambda}/\eta)^{-3/7}$, with $v_{A,\lambda} \sim \varepsilon^{7/18} \eta^{-1/6} \lambda^{5/9}$ (Boldyrev & Loureiro 2017). Eq. (7) is then replaced by $\lambda/L \gg \mathcal{R}^{-3/2} S_L^{-3/4}$. The estimation of the dissipation scale is unchanged from that pertaining to the Harris sheet; but, in this case, $\lambda_{cr}/L \sim S_L^{-6/11}$. Therefore, Eq. (8) is instead $S_L \gg \mathcal{R}^{-22/3}$.

due to electron inertia (active at the electron skin-depth scale, $d_e = c/\omega_{pe}$), rather than resistivity as considered in the previous section; i.e., in this case $\mathcal{R} \approx 0.1$.

As documented in Loureiro & Boldyrev (2017b), there are two cases that need considering: the first, somewhat simpler to address, is when the critical scale at which the tearing mode onsets is in the MHD range (i.e., λ_{cr} is larger than the ion kinetic scales) — even though, to repeat, the tearing and reconnection themselves require kinetic effects. This is treated in Section 4.1. The second case is when the onset of tearing only occurs for scales smaller than the ion kinetic scales. This is discussed in Section 4.2.

4.1. Reconnection at fluid scales

Several cases are possible, depending on plasma parameters (Loureiro & Boldyrev 2017b). There is no need here to be exhaustive: for any particular case, the calculation proceeds in a qualitatively similar way. Therefore, let us consider, as an example, a low beta plasma (see Sections 2 & 3 of Loureiro & Boldyrev (2017b)) — we choose to analyze this particular case because of its potential relevance to solar wind observations (Vech et al. 2018), and perhaps also to the solar corona. In this case, in the tearing-mediated range, the eddy turnover rate becomes the growth rate of the fastest growing tearing mode as given by $\gamma_t \sim v_{A,\lambda} d_e \rho_s / \lambda^3$, where ρ_s is the ion sound Larmor radius. Evaluation of Eq. (3) then yields:

$$\lambda \gg \mathcal{R}^{-1/2} (d_e \rho_s)^{1/2}. \quad (9)$$

This expression only applies in the range of scales $\lambda_{cr} \gg \lambda \gg \rho_s$ where, for this case, $\lambda_{cr}/L \sim (d_e/L)^{4/9} (\rho_s/L)^{4/9}$ (Loureiro & Boldyrev 2017b; Mallet et al. 2017b); this translates into

$$\mathcal{R} \ll \frac{d_e}{\rho_s} \ll \mathcal{R}^9 \left(\frac{L}{\rho_s} \right)^2. \quad (10)$$

The left inequality is probably not satisfied in the (pristine) solar wind at ~ 1 AU (it requires $\beta_e \ll 2(m_e/m_i)\mathcal{R}^{-2} \approx 0.1$, which may be too low). In that case, one concludes that reconnection in current sheets should not be a main energy dissipation mechanism in that turbulent environment. The opposite situation, however, should pertain to the solar corona: using standard parameters there is no difficulty in concluding that both inequalities in Eq. (10) should hold comfortably.⁶ From the point of view of numerical simulations, this

⁶ Note that this is indeed the regime that we would expect to describe turbulence in the solar corona at these scales, rather

result, like Eq. (8), unfortunately places close to impossible demands⁷.

4.2. Reconnection at kinetic scales

Finally, we analyze the case when the tearing onset only occurs at scales below the ion kinetic scales. Let us consider here the analysis recently proposed by Boldyrev & Loureiro (2019) of sub-ion range turbulence in plasmas such that $\beta_i \sim 1 \gg \beta_e$. The relevant eddy turnover rate is

$$\gamma_t \sim \frac{V_{Ae,\lambda}}{\lambda} \left(\frac{d_e}{\lambda} \right)^2. \quad (11)$$

Therefore we find:

$$\gamma_t \tau_{rec} \ll 1 \implies \frac{\lambda}{d_e} \gg \mathcal{R}^{-1/2} \sim 3. \quad (12)$$

Repeating this derivation for the $\sin(x)$ magnetic field profile instead yields $\lambda/d_e \gg \mathcal{R}^{-2/3} \sim 5$.

Unlike the two cases considered previously, an estimation of λ_{cr} is not available for this situation (a reflection of the fact that a detailed understanding of sub-ion range turbulence is still lacking). The only known constraint that applies to λ_{cr} is that it be smaller than $\min(d_i, \rho_i, \rho_s)$, which simply follows from the range of validity of the equations that are used by Boldyrev & Loureiro (2019) to compute Eq. (11). At small scales, it is required that $\lambda \gg d_e$ which is (marginally) satisfied by Eq. (12).

It is interesting to analyze this result in light of recent MMS observations of so-called electron-only reconnection in the Earth's magnetosheath (Phan et al. 2018; Stawarz et al. 2019). In Boldyrev & Loureiro (2019) we have estimated that the decoupling of the ions in these events requires $\lambda/d_e \ll \sqrt{d_i/d_e}$ or $\lambda/d_e \ll (d_i/d_e)^{2/3}$ depending on whether one assumes a $\tanh(x/\lambda)$ or $\sin(x/\lambda)$ magnetic field profile for the reconnecting field $B_\xi(x)$ in the eddy. These estimates range from ~ 6 to ~ 12 suggesting, therefore, a rather narrow range of scales where nonlinear 'electron-only' reconnection in the eddies may be possible. Remarkably, Phan et al.

(2018) report a current sheet thickness of $\sim 4 d_e$, strikingly consistent with these numbers and with Eq. (12). This is certainly very encouraging, but one must also bear in mind that all our analytical results are only order of magnitude estimates which ignore order unity numerical prefactors.

Another observationally-based result which we interpret to be consistent with our analysis is the recent claim by Chen et al. (2019) that energy dissipation at kinetic scales in the magnetosheath is dominated by linear electron Landau damping (the energy dissipation rate via that channel being comparable to the energy cascade rate). Indeed, Eq. (12) demonstrates that full reconnection in sub-ion scale eddies is permitted for typical magnetosheath parameters; and previous investigations of heating in (strong guide-field) collisionless reconnection (Loureiro et al. 2013; Numata & Loureiro 2015) show that when $\beta_e \ll 1$ linear electron Landau damping is by far the dominant energy dissipation channel.

5. CONCLUSION

This paper builds on previous recent work on the onset of the tearing instability in strong magnetic plasma turbulence, establishing the conditions under which this instability may develop into a deep nonlinear reconnecting state. The ability to do so is intimately tied to whether or not significant energy dissipation and conversion is to be expected at such turbulent scales. We think this has profound implications for turbulent systems. For example, in weakly collisional plasmas, reconnection is a well known efficient particle acceleration mechanism (e.g. Guo et al. 2014; Sironi & Spitkovsky 2014; Dahlin et al. 2015; Werner et al. 2017), and heats different species at different rates (e.g. Numata & Loureiro 2015; Shay et al. 2018). Therefore, if reconnecting eddies are a common occurrence — the conditions for which are worked out in this paper — then one might expect turbulence to be efficient at generating non-thermal populations and different electron-to-ion temperature ratios, which are indeed observed or expected in different space and astrophysical plasmas (see, e.g., Schekochihin et al. 2019, and references therein). Moreover, the very observability of reconnecting turbulence depends, obviously, on whether truly reconnecting eddies are the norm or an exception.

Yet another consequential implication of the analysis carried out in this paper stems from the fact that neither Eq. (8) nor Eq. (10) have ever been met in computer simulations conducted to date, nor is that likely to happen in the near future. It thus follows that all observations of reconnecting current sheets in (three-

than the MHD case of Section 3: for typical coronal conditions, i.e., $S_L \approx 10^{14}$, $L \approx 10^4$ km, Eq. (15) of Loureiro & Boldyrev (2017a), describing the MHD case, yields an estimate of the width of the inner boundary layer of the tearing mode occurring on an eddy of width λ_{cr} of approximately 1 cm, smaller than the electron skin depth and showing, therefore that the tearing mode at such scales is collisionless, as we consider in this Section.

⁷ For the $\sin(x)$ profile, Eq. (9) becomes instead $\lambda \gg \mathcal{R}^{-2/3} d_e^{4/9} \rho_s^{5/9}$; whereas Eq. (10) is replaced by $\mathcal{R}^{3/2} \ll d_e/\rho_s \ll \mathcal{R}^{63/6} (L/\rho_s)^{9/4}$. The conclusions drawn above pertain equally for these estimates.

dimensional) numerical simulations of strong turbulence in the plasma regimes to which those equations pertain are bound to be relatively rare or transient events, with no significant impact on the nature of energy dissipation. One immediate consequence is, therefore, that the energy dissipation (heating or particle energization) rates obtained in simulations of magnetic turbulence may be severely underestimated with respect to the environments that such simulations aim to study. One way to remedy this situation might be to hardwire, in numerical simulations, energy dissipation prescriptions based

on the energetics of reconnection (specific to the particular plasma parameters under study) at scales where the turbulent cascade is tearing dominated.

NFL was partially funded by NSF CAREER award no. 1654168 and by the NSF-DOE Partnership in Basic Plasma Science and Engineering, award no. desc0016215. SB was partly supported by the NSF under grant no. NSF PHY-1707272, by NASA under grant no. NASA 80NSSC18K0646, and by DOE grant no. DE-SC0018266.

REFERENCES

- Arzamasskiy, L., Kunz, M. W., Chandran, B. D. G., & Quataert, E. 2019, *The Astrophysical Journal*, 879, 53. <https://doi.org/10.3847/2F1538-4357%2F20190401>
- Bhattacharjee, A., Huang, Y.-M., Yang, H., & Rogers, B. 2009, *Physics of Plasmas*, 16, 112102. <https://aip.scitation.org/doi/abs/10.1063/1.3264103>
- Birn, J., Drake, J. F., Shay, M. A., et al. 2001, *Journal of Geophysical Research*, 106, 3715. <http://adsabs.harvard.edu/abs/2001JGR...106.3715B>
- Biskamp, D. 2000, *Magnetic Reconnection in Plasmas* (Cambridge University Press). <http://adsabs.harvard.edu/abs/2000mrp..book.....B>
- . 2003, *Magnetohydrodynamic Turbulence* (Cambridge University Press). <https://ui.adsabs.harvard.edu/abs/2003matu.book.....B/abstract>
- Boldyrev, S. 2006, *Physical Review Letters*, 96, doi:10.1103/PhysRevLett.96.115002. <https://link.aps.org/doi/10.1103/PhysRevLett.96.115002>
- Boldyrev, S., & Loureiro, N. F. 2017, *The Astrophysical Journal*, 844, 125. <http://stacks.iop.org/0004-637X/844/i=2/a=125?key=crossref.24ef5661071057097813176372358200f>
- . 2018, *Journal of Physics: Conference Series*, 1100, 012003. <http://stacks.iop.org/1742-6596/1100/i=1/a=012003?key=crossref.091c84a748d1e11993504c350c>
- . 2019, arXiv:1901.10096 [physics], arXiv: 1901.10096. <http://arxiv.org/abs/1901.10096>
- Cassak, P. A., Liu, Y.-H., & Shay, M. A. 2017, *Journal of Plasma Physics*, 83, doi:10.1017/S0022377817000666. <https://www.cambridge.org/core/journals/journal-of-plasma-physics/abstract/5901-reconnection-rate-problem/CB5057FE46231591>
- Cerri, S., Franci, L., Califano, F., Landi, S., & Hellinger, P. 2017, *Journal of Plasma Physics*, 83, doi:10.1017/S0022377817000265. <https://www.cambridge.org/core/product/identifier/S0022377817000265/type/journal-article>
- Chandran, B. D. G., Schekochihin, A. A., & Mallet, A. 2015, *The Astrophysical Journal*, 807, 39. <http://stacks.iop.org/0004-637X/807/i=1/a=39?key=crossref.12fb6d1527072e1533667744d6b>
- Chen, C. H. K. 2016, *Journal of Plasma Physics*, 82, 535820602. <http://adsabs.harvard.edu/abs/2016JPIPh..82f5302C>
- Chen, C. H. K., & Boldyrev, S. 2017, *The Astrophysical Journal*, 842, 122. <http://stacks.iop.org/0004-637X/842/i=2/a=122?key=crossref.775298>
- Chen, C. H. K., Klein, K. G., & Howes, G. G. 2019, *Nature Communications*, 10, 740. <https://doi.org/10.1038/s41467-019-08435-3>
- Comisso, L., & Bhattacharjee, A. 2016, *Journal of Plasma Physics*, 82, doi:10.1017/S002237781600101X. <http://www.cambridge.org/core/journals/journal-of-plasma-physics/a>
- Coppi, B., Galvão, R., Pellat, R., Rosenbluth, M., & Rutherford, P. 1976, *Soviet Journal of Plasma Physics*, 2, 961. <http://adsabs.harvard.edu/abs/1976SvJPP...2..533C>
- Dahlin, J. T., Drake, J. F., & Swisdak, M. 2015, *Physics of Plasmas*, 22, 100704. <https://doi.org/10.1063/1.4933212>
- Davidson, P. A. 2016, *Introduction to Magnetohydrodynamics* (Cambridge University Press), doi:10.1017/9781316372358. <https://www.cambridge.org/core/books/introduction-to-magnetohydrodynamics>
- Dong, C., Wang, L., Huang, Y.-M., Comisso, L., & Bhattacharjee, A. 2018, *Physical Review Letters*, 121, 165101. <https://link.aps.org/doi/10.1103/PhysRevLett.121.165101>
- Furth, H. P., Killeen, J., & Rosenbluth, M. N. 1963, *Physics of Fluids*, 6, 459. <http://adsabs.harvard.edu/abs/1963PhFl...6..459F>
- Goldreich, P., & Sridhar, S. 1995, *The Astrophysical Journal*, 438, 763. <http://adsabs.harvard.edu/abs/1995ApJ...438..763G>
- Grošelj, D., Mallet, A., Loureiro, N. F., & Jenko, F. 2018, *Physical Review Letters*, 120, 105101. <http://stacks.iop.org/0004-637X/120j5101G>

- Guo, F., Li, H., Daughton, W., & Liu, Y.-H. 2014, *Physical Review Letters*, 113, 155005. <https://link.aps.org/doi/10.1103/PhysRevLett.113.155005>
- Huang, Y.-M., & Bhattacharjee, A. 2010, *Physics of Plasmas*, 17, 062104. <https://aip.scitation.org/doi/abs/10.1063/1.3420208>
- Jemella, B. D., Shay, M. A., Drake, J. F., & Rogers, B. N. 2003, *Physical Review Letters*, 91, doi:10.1103/PhysRevLett.91.125002. <https://link.aps.org/doi/10.1103/PhysRevLett.91.125002>
- Landi, S., Franci, L., Papini, E., et al. 2019, arXiv e-prints, arXiv:1904.03903
- Liu, Y.-H., Hesse, M., Guo, F., et al. 2017, *Physical Review Letters*, 118, doi:10.1103/PhysRevLett.118.085101. <https://link.aps.org/doi/10.1103/PhysRevLett.118.085101>
- Loureiro, N. F., & Boldyrev, S. 2017a, *Physical Review Letters*, 118, doi:10.1103/PhysRevLett.118.245101. <http://link.aps.org/doi/10.1103/PhysRevLett.118.245101>
- . 2017b, *The Astrophysical Journal*, 850, 182. <http://stacks.iop.org/0004-637X/850/i=2/a=182?key=crossref.ea205300ef557655b4445c42164c39be>
- . 2018, *The Astrophysical Journal*, 866, L14. <http://stacks.iop.org/2041-8205/866/i=1/a=L14?key=crossref.db028cfe529c4ba6af61aacbebed4a1>
- Loureiro, N. F., Cowley, S. C., Dorland, W. D., Haines, M. G., & Schekochihin, A. A. 2005, *Physical Review Letters*, 95, doi:10.1103/PhysRevLett.95.235003. <https://link.aps.org/doi/10.1103/PhysRevLett.95.235003>
- Loureiro, N. F., Samtaney, R., Schekochihin, A. A., & Uzdensky, D. A. 2012, *Physics of Plasmas*, 19, 042303. <http://aip.scitation.org/doi/10.1063/1.3703318>
- Loureiro, N. F., Schekochihin, A. A., & Cowley, S. C. 2007, *Physics of Plasmas*, 14, 100703
- Loureiro, N. F., Schekochihin, A. A., & Zocco, A. 2013, *Physical Review Letters*, 111, doi:10.1103/PhysRevLett.111.025002. <https://link.aps.org/doi/10.1103/PhysRevLett.111.025002>
- Loureiro, N. F., & Uzdensky, D. A. 2016, *Plasma Physics and Controlled Fusion*, 58, 014021. <http://stacks.iop.org/0741-3335/58/i=1/a=014021>
- Mallet, A., Schekochihin, A. A., & Chandran, B. D. G. 2015, *Monthly Notices of the Royal Astronomical Society: Letters*, 449, L77. <http://academic.oup.com/mnrasl/article/449/1/L77/1029314/Refined%20critical%20balance%20in%20strong%20Alfvic>
- . 2017a, *Monthly Notices of the Royal Astronomical Society*, 468, 4862. <https://academic.oup.com/mnras/article-lookup/doi/10.1093/mnras/stw070>
- . 2017b, *Journal of Plasma Physics*, 83, doi:10.1017/S0022377817000812. <https://www.cambridge.org/core/product/identifier/S0022377817000812>
- Maron, J., & Goldreich, P. 2001, *The Astrophysical Journal*, 554, 1175. <https://doi.org/10.1086%2F321413>
- Matthaeus, W. H., & Lamkin, S. L. 1986, *Physics of Fluids*, 29, 2513. <https://aip.scitation.org/doi/10.1063/1.866004>
- Numata, R., & Loureiro, N. F. 2015, *Journal of Plasma Physics*, 81, doi:10.1017/S002237781400107X. http://www.journals.cambridge.org/abstract_S002237781400107X
- Osman, K. T., Matthaeus, W. H., Gosling, J. T., et al. 2014, *Physical Review Letters*, 112, 215002. <https://link.aps.org/doi/10.1103/PhysRevLett.112.215002>
- Parker, E. N. 1957, *Journal of Geophysical Research*, 62, 509. <http://doi.wiley.com/10.1029/JZ062i004p00509>
- Passot, T., Sulem, P. L., & Tassi, E. 2017, *Journal of Plasma Physics*, 83, doi:10.1017/S0022377817000514. <http://www.cambridge.org/core/journals/journal-of-plasma-physics/a>
- . 2018, *Physics of Plasmas*, 25, 042107. <http://aip.scitation.org/doi/10.1063/1.5022528>
- Phan, T. D., Eastwood, J. P., Shay, M. A., et al. 2018, *Nature*, 557, 202. <http://www.nature.com/articles/s41586-018-0091-5>
- Porcelli, F., Borgogno, D., Califano, F., et al. 2002, *Plasma Physics and Controlled Fusion*, 44, B389. <https://doi.org/10.1088%2F0741-3335%2F44%2F12b%2F327>
- Priest, E., & Forbes, T. 2000, *Magnetic reconnection: MHD theory and applications* (Cambridge University Press). <http://adsabs.harvard.edu/abs/2000mrmt.conf.....P>
- Retinò, A., Sundkvist, D., Vaivads, A., et al. 2007, *Nature Physics*, 3, 235
- Roytershteyn, V., Boldyrev, S., Delzanno, G. L., et al. 2019, *The Astrophysical Journal*, 870, 103. <http://stacks.iop.org/0004-637X/870/i=2/a=103?key=crossref.439e40>
- Samtaney, R., Loureiro, N. F., Uzdensky, D. A., Schekochihin, A. A., & Cowley, S. C. 2009, *Physical Review Letters*, 103, 105004
- Schekochihin, A. A. 2018, *MHD Turbulence: A Biased Review*, (*in preparation*), <http://www-thphys.physics.ox.ac.uk/research/plasma/JPP/papers/17/>
- Schekochihin, A. A., Kawazura, Y., & Barnes, M. A. 2019, *Journal of Plasma Physics*, 85, doi:10.1017/S0022377819000345. <https://www.cambridge.org/core/product/identifier/S0022377819000345>
- Servidio, S., Matthaeus, W. H., Shay, M. A., Cassak, P. A., & Dmitruk, P. 2009, *Physical Review Letters*, 102, 115003. <https://link.aps.org/doi/10.1103/PhysRevLett.102.115003>
- Shay, M. A., Haggerty, C. C., Matthaeus, W. H., et al. 2018, *Physics of Plasmas*, 25, 012304. <https://doi.org/10.1063/1.4993423>
- Simon, L., & Spitkovsky, A. 2014, *The Astrophysical Journal*, 783, L21. <https://doi.org/10.1088%2F2041-8205%2F783%2F1%2F121>

- Stawarz, J. E., Eastwood, J. P., Phan, T. D., et al. 2019, *The Astrophysical Journal Letters*, 877, L37.
<https://iopscience.iop.org/article/10.3847/2041-8213/ab21c8/meta>
- Sundkvist, D., Retinò, A., Vaivads, A., & Bale, S. D. 2007, *Physical Review Letters*, 99, doi:10.1103/PhysRevLett.99.025004.
<https://link.aps.org/doi/10.1103/PhysRevLett.99.025004>
- Sweet, P. A. 1958, in *IAU Symposium, Vol. 6, Electromagnetic Phenomena in Cosmical Physics*, ed. B. Lehnert, 123
- TenBarge, J. M., & Howes, G. G. 2013, *The Astrophysical Journal Letters*, 771, L27.
<http://adsabs.harvard.edu/abs/2013ApJ...771L..27T>
- Uzdensky, D. A., Loureiro, N. F., & Schekochihin, A. A. 2010, *Physical Review Letters*, 105, doi:10.1103/PhysRevLett.105.235002.
<https://link.aps.org/doi/10.1103/PhysRevLett.105.235002>
- Vech, D., Mallet, A., Klein, K. G., & Kasper, J. C. 2018, *The Astrophysical Journal*, 855, L27.
<http://stacks.iop.org/2041-8205/855/i=2/a=L27?key=crossref.3c4651e4bdcf44d52570508441ec42aa>
- Waelbroeck, F. L. 1989, *Physics of Fluids B: Plasma Physics*, 1, 2372.
<https://aip-scitation-org.libproxy.mit.edu/doi/10.1063/1.859172>
- Walker, J., Boldyrev, S., & Loureiro, N. F. 2018, *Physical Review E*, 98, doi:10.1103/PhysRevE.98.033209.
<https://link.aps.org/doi/10.1103/PhysRevE.98.033209>
- Wan, M., Matthaeus, W. H., Roytershteyn, V., et al. 2015, *Physical Review Letters*, 114, 175002.
<https://link.aps.org/doi/10.1103/PhysRevLett.114.175002>
- Wang, X., & Bhattacharjee, A. 1993, *Physical Review Letters*, 70, 1627.
<https://link.aps.org/doi/10.1103/PhysRevLett.70.1627>
- Werner, G. R., Uzdensky, D. A., Begelman, M. C., Cerutti, B., & Nalewajko, K. 2017, *Monthly Notices of the Royal Astronomical Society*, 473, 4840.
<https://doi.org/10.1093/mnras/stx2530>
- Yamada, M., Kulsrud, R., & Ji, H. 2010, *Reviews of Modern Physics*, 82, 603.
<http://adsabs.harvard.edu/abs/2010RvMP...82..603Y>
- Zhdankin, V., Uzdensky, D. A., Perez, J. C., & Boldyrev, S. 2013, *The Astrophysical Journal*, 771, 124.
<http://stacks.iop.org/0004-637X/771/i=2/a=124>
- Zweibel, E. G., & Yamada, M. 2009, *Annual Review of Astronomy and Astrophysics*, 47, 291.
<http://adsabs.harvard.edu/abs/2009ARA%26A...47..291Z>