Constraining the ¹²C+¹²C astrophysical S-factors with the ¹²C+¹³C measurements at very low energies

N.T. Zhang^{a,b}, X.Y. Wang^{a,b}, H. Chen^{a,b}, Z.J. Chen^{a,b}, W.P. Lin^a, W.Y. Xin^{a,b}, S.W. Xu^a, D. Tudor^{c,d}, A.I. Chilug^{c,d}, I.C. Stefanescu^{c,d}, M. Straticiuc^c, I. Burducea^c, D.G. Ghita^c, R. Margineanu^c, C. Gomoiu^c, A. Pantelica^c, D. Chesneanu^c, L. Trache^c, X.D. Tang^{a,b,e}, B. Bucher^f, L.R. Gasques^g, K. Hagino^{h,i}, S. Kubono^{a,j}, Y.J. Li^k, C.J. Lin^k, A.S. Umar^l, Y. Xu^m

^aInstitute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China

^bSchool of Nuclear Science and Technology, University of Chinese Academy of Sciences, Beijing 100049, China

^cHoria Hulubei National Institute of Physics and Nuclear Engineering, IFIN-HH, Măgurele 077125, Romania

^dDoctoral School of Physics, University of Bucharest, Bucuresti-Măgurele 077125, Romania

^e Joint department for nuclear physics, Lanzhou University and Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China

^fIdaho National Laboratory, Idaho Falls, Idaho 83415, USA

^gDepartamento de Física Nuclear, Instituto de Física da Universidade de São Paulo, 05508-090, São Paulo, SP, Brazil

ⁱResearch Center for Electron Photon Science, Tohoku University, 1-2-1 Mikamine, Sendai 982-0826, Japan

^jRIKEN Nishina Center, RIKEN, 2-1 Hirosawa, Saitama 351-0198, Japan

^kChina Institute of Atomic Energy, Beijing 102413, China

¹Department of Physics and Astronomy, Vanderbilt University, Nashville, Tennessee 37235, USA

^mExtreme Light Infrastructure-Nuclear Physics, Măgurele 077125, Romania

Abstract

We use an underground counting lab with an extremely low background to perform an activity measurement for the ${}^{12}C+{}^{13}C$ system with energies down to $E_{c.m.}=2.323$ MeV, at which the ${}^{12}C({}^{13}C,p){}^{24}$ Na cross section is found to be 0.22(7) nb. The ${}^{12}C+{}^{13}C$ fusion cross section is derived with a statistical model calibrated using experimental data. Our new result of the ${}^{12}C+{}^{13}C$ fusion cross section is the first decisive evidence in the carbon isotope systems which rules out the existence of the astrophysical S-factor maximum predicted by the phenomenological hindrance model, while confirming the rising trend of the S-factor towards lower energies predicted by other models, such as CC-M3Y+Rep, DC-TDHF, KNS, SPP and ESW. After normalizing the model predictions with our data, a more reliable upper limit is established for the ${}^{12}C+{}^{12}C$ fusion cross sections at stellar energies.

Keywords: Fusion cross section, astrophysical S-factor, extrapolation models, hindrance

1. Introduction

The ¹²C+¹²C fusion reaction is one of the most important reactions in nuclear astrophysics. The crucial energy range extends from a few 10's keV to $E_{c.m.}=3$ MeV [1]. All the measurements, going back 60 years, have been limited to energies above $E_{c.m.}=2.1$ MeV. The situation is further complicated by the existence of resonance states with ~50 keV widths and a few hundred keV spacings, continuing down to the lowest energies for which measurements have been made [2]. Extrapolation is inevitably necessary to obtain the reaction rate for astrophysical applications.

Modeling ¹²C+¹²C heavy ion fusion cross sections at deep sub-barrier energies has been a long standing challenge for nuclear reaction theory [2]. The Gamow energy window for most astrophysical applications typically spans 500 keV or more, and with no reliable method for predicting the resonances at lower

Email addresses: dana.tudor@nipne.ro (D. Tudor),

livius.trache@nipne.ro (L. Trache), xtang@impcas.ac.cn (X.D. Tang)

energies, the standard reaction rate (CF88) was established by using constant modified S-factors $(S^*(E) = S(E)exp(0.46E))$ based on the square well penetration factor [3, 4]. This rising trend of S-factor towards lower energies is confirmed by various phenomenological and microscopic models, such as density-constrained time dependent Hartree-Fock method (DC-TDHF) [5], wave-packet dynamics (TDWP) [6], barrier penetration model based on the global São Paulo potentials (SPP) [1] or the Krappe-Nix-Sierk potential (KNS) [7], and coupled channel calculations such as CC-M3Y+Rep [8, 9] (Fig. 1(a)). However, the hindrance model, a global phenomenological model based on the systematics observed in systems with $64 \gtrsim A \gtrsim 30$, predicts that the ¹²C+¹²C S-factor reaches its maximum at $E_{c.m.}$ =3.68±0.38 MeV [10–12]. At lower energies, this model predicts a rapid drop in the S-factor leading to a reduced reaction rate that is many orders of magnitude smaller than the standard rate used for astrophysical modeling. Although the experimental data are available down to 2.1 MeV, no decisive conclusion can be made due to the poor fitting of the resonancelike structure in the ${}^{12}C+{}^{12}C$ cross section with the smooth extrapolating models [12]. Recently, the cross sections were determined indirectly for $E_{c.m.} = 0.8$ MeV to 2.7 MeV using the

^hDepartment of Physics, Tohoku University, Sendai 980-8578, Japan

^{*}Corresponding author

Trojan Horse Method [13], recommending a new S-factor with a rising slope faster than any models presented in Fig. 1(a), resulting in a new rate which is 1 or 2 orders of magnitude higher than the standard one. The large uncertainty of the ${}^{12}C+{}^{12}C$ rate drastically impacts a number of models such as late-time massive star evolution, the ignition dynamics of type Ia supernovae and x-ray superbursts [1, 14].



Figure 1: (Color online) S-factors of ${}^{12}C+{}^{12}C$ (upper panel) and ${}^{12}C+{}^{13}C$ (lower panel) at sub-barrier energies. The ${}^{12}C+{}^{12}C$ data from Ref. [15], [7], [16], and [12] with errors better than 40% are shown as red, green, blue and black open circles, respectively. The ${}^{12}C+{}^{13}C$ data from Ref. [17] and [18] are shown respectively as green and blue filled circles. Model calculations, DC-TDHF (dark blue), CC-M3Y+Rep (red), TDWP (black) and CF88 (magenta), are shown as solid lines; SPP using global density distribution (red) and the proposed hindrance signatures (black) are shown as the dashed lines. The KNS prediction for the ${}^{12}C+{}^{12}C$ non-resonant component and the THM measurement are shown as the light blue dashed-dotted line and dark green dashed line, respectively.

While the complicated resonance-like structure in ${}^{12}C+{}^{12}C$ and the lack of reliable measurements at lower energies prevent us from drawing a clear conclusion [12], the isotope fusion reaction ¹²C+¹³C offers an ideal opportunity to constrain the ${}^{12}C+{}^{12}C$ S-factor. It has been observed at energies below and above the Coulomb barrier that ${}^{12}C+{}^{13}C$ and ${}^{13}C+{}^{13}C$ cross sections upper bound the ¹²C+¹²C cross sections, and match the maxima of the resonance-like structure seen in ${}^{12}C+{}^{12}C$ in a wide range from 10^{-8} b to 1 b [19] (see Fig. 2). This strong correlation among the three systems has been well explained by a coupled channel calculations and the significantly different level densities of the compound states [20]. At sub-barrier energies, the valence neutron(s) in ${}^{12}C+{}^{13}C$ and ${}^{13}C+{}^{13}C$ increase the level densities of their compound states by at least one order of magnitude in comparison to ${}^{12}C+{}^{12}C$ and result in smooth cross sections. According to eq.5 in Ref. [20], the upper limit of the ${}^{12}C+{}^{12}C$ cross section is reached using the high level density limit and can be modeled consistently with the other C+C isotope systems [20]. The average of the ${}^{12}C+{}^{12}C$ fusion cross section is also predicted by modulating the upper limit with the averaged ratio of the level width $\langle \Gamma \rangle$ and the level spacing $\langle D \rangle$ of ²⁴Mg [20]. Since the effect of $\langle \Gamma/D \rangle$ is not sensitive to the energy, the shape of the averaged cross section is essentially determined by the model used for the upper limit prediction. Therefore, a model well constrained by ${}^{12}C+{}^{13}C$ is crucial for us to set an reliable upper limit and constrain the shape of the averaged cross section for ${}^{12}C+{}^{12}C$.

However, the large deviations among various global models exist in ${}^{12}C+{}^{13}C$ and affect the S-factor extrapolation as shown in Fig. 1(b). Very similar to ${}^{12}C+{}^{12}C$, the global models, CC-M3Y+Rep, São Paulo and DC-TDHF, predict a rising trend for the ${}^{12}C+{}^{13}C$ S-factor, while the hindrance model suggests that the S-factor reaches its maximum at $E_{c.m.}=3.45\pm0.37$ MeV [10, 11]. However, the current experimental data are insufficient to test these models. Therefore, it is crucially important to extend the fusion cross section measurement of ${}^{12}C+{}^{13}C$ down to lower energies and provide a strict test for models which can constrain on the ${}^{12}C+{}^{12}C$ cross section.



Figure 2: (Color online) Cross sections of ${}^{12}C({}^{13}C,p){}^{24}Na$ are obtained in Ref. [19] (black) and this work (red). The latter represent the first measurements to reach the energy region $E_{c.m.} < 2.6$ MeV. A 10% experimental systematical uncertainty is included in both data. The ${}^{12}C{}^{+13}C$ total fusion cross sections from Ref. [17], [18] are also shown as green and blue squares, respectively. The ${}^{12}C{}^{+12}C$ total fusion cross sections are also shown as open circles with symbols explained in the Fig. 1 caption.

Obtaining a precise measurement of such small fusion cross sections is a great experimental challenge due to extremely low yields and ambient backgrounds. Up to now, only ¹²C+¹²C has been measured down to 2 nb with errors \leq 40% while other measurements for light systems have stopped above 10 nb, which is not low enough to differentiate the various predictions [21, 22]. Notani *et al.* has extended the measurement of ¹²C+¹³C to 20 nb [19]. The current work further extends that effort by exploiting an underground counting lab with an ultra-low background, pushing the measurement of the ¹²C(¹³C,*p*)²⁴Na cross section down to 0.22(7) nb with which the ¹²C+¹³C fusion cross section is determined to be 0.9 nb with an uncertainty \leq 30%.

2. Experiment and results

The fusion reaction measurement of ${}^{12}C+{}^{13}C$ was performed by detecting the residual nucleus ${}^{24}Na$ ($T_{1/2}=15.0$ h) from

¹²C(¹³C,p)²⁴Na. A ¹³C beam with a typical current up to 15 $p\mu A$ was delivered on a 1.5-mm-thick natural carbon target by the 3 MV Tandem of IFIN-HH [23]. The accelerator GVM was calibrated with the ${}^{27}Al(p,\gamma){}^{28}Si$ reaction with a uncertainty less than 0.1%. The ¹²C(¹³C,p)²⁴Na reaction was studied in the energy range from E_{lab} =4.640 to 10.995 MeV with a energy step of ΔE_{lab} =0.2 MeV. At lower energies the target samples were irradiated and then transported in 3 hours to the underground counting lab (μ Bq) in the SLANIC salt mine, Romania for offline γ -ray measurements [24]. In the measurement, a wellshielded HPGe detector was used to detect two cascading γ rays (1369, 2754 keV) emitted by the ²⁴Na β -decay, with only 1369 keV γ -ray being used in the final yield determination. Some of the higher energy measurements ($E_{c.m.}$ >2.8 MeV) were also performed in the Low Background Gamma-Ray Spectroscopy Laboratory (GamaSpec) in a basement of IFIN-HH. The detector efficiencies were carefully calibrated with standard sources. The correction of the summing effect was evaluated by comparing the measurements of a ²⁴Na sample at two different distances, 15 cm from the detector and a very close geometry (less than 1 mm from the detector). The results are also compared with the Monte Carlo simulations. The total times for irradiation and measurements were 20 and 33 days, respectively. Fig. 3 shows the γ -ray spectrum obtained at the energy of $E_{\rm c.m.}$ =2.323 MeV with a ²⁴Na production cross section of 0.2 nb. The 1369 keV peak, used in our yield determinations, is clearly seen in both raw and background subtracted spectra.



Figure 3: (Color online) (a) Measurements in the underground lab with 3 samples irradiated at $E_{c.m.}$ =2.323 MeV. The irradiations and measurements took 3.4 and 3.9 days, respectively. (b) Spectrum after subtracting background.

The fusion cross sections of ${}^{12}C({}^{13}C,p){}^{24}Na$ are obtained by differentiating the thick target yield using the procedure given in Ref. [19]. Our measurement includes a 10% systematic uncertainty which primarily results from the uncertainties in the beam current (5%), beam energy (2%), detector efficiency (5%), and stopping power (7%) from Ref. [25]. The measurement by Notani *et al.* is a relative measurement. Their ${}^{12}C({}^{13}C,p){}^{24}Na$ cross sections are normalized to our data. Both results are shown in Fig. 2.

Two different methods, in-beam characteristic γ ray [17] and summing- γ -ray yield [18], have been used to obtain the total fu-

sion cross sections with the aid of the statistical model, making ${}^{12}C+{}^{13}C$ a good case to assess the model's systematic error. The ratio between these two measurements is $1.11\pm0.07(\sigma)$. The deviation of 0.11 from 1 and the fluctuation of 0.07 result from a combination of the statistical error and the systematic uncertainties of both the experiment and the statistical model. As a conservative value, we combine the central deviation (0.11) with the standard deviation (0.07) and estimate the systematic error of the statistical model as 13%. This error has been included in both total cross sections in the latter part of this paper.

The total fusion cross sections over the full energy range measured in this work are converted from the measured cross sections of the ${}^{12}C({}^{13}C,p){}^{24}Na$ channel based on the statistical model calculations, and hence a reliable branching ratio of the proton channel with a quantified uncertainty is needed. The experimental branching ratio is obtained by comparing the ${}^{12}C({}^{13}C,p){}^{24}Na$ cross sections to the two sets of measured total fusion cross sections [17, 18] (Fig. 4). Two different statistical model codes, Talys [26] and Empire [27], are used to calculate the branching ratio (see Fig. 4). In the Talys calculation, the spin population of the ²⁵Mg compound is set using the prediction by [9]. The potential parameters of the p and n channels are tuned to reproduce the partial cross sections in Ref. [17]. In the Empire calculation, the ²⁴Na channel branching ratio is predicted using the default parameters and normalized to the experimental data by a factor of 0.84. These two calculations are similar with a difference less than 7%. The fluctuation of data around the best fit is $14\%(1\sigma)$, which is related to the systematic uncertainties of the measurements and statistical model. In this work, the Talys result is used for the branching ratio correction. By including the systematic difference of the two models $(\pm 3.5\%)$, we adopt 14% as the systematic error for the ${}^{12}C({}^{13}C,p){}^{24}Na$ branching ratio obtained from the model. This error was overestimated as 20% by Notani et al. [19].



Figure 4: (Color online) Branching ratio of the ${}^{12}C({}^{13}C, p){}^{24}Na$ reaction channel. The red and black symbols correspond to our measurement and the normalized Notani *et al* measurement [19], respectively, with the branching ratios deduced from the total fusion cross sections of Ref. [17] (circles) and Ref. [18] (stars). Solid lines show the normalized theoretical branching ratios calculated with Empire (black) and Talys (blue), with dashed lines representing the 1σ limits from the Talys calculation.

3. Discussion

The experin shown in Fig. maximum at model, is obs decreases, th other models global theore DC-TDHF a up to 35%, normalizing reduced- χ^2 at $E_{\rm c.m.}=1$ viations as of the thre (Fig. 5(a)) Hindra rapifi dro and ¹²C· strong h But our perimer at lowe below tween range sive e with ih th A serv <u>30</u>] S-f the is ТĮ 12 f

> c+¹³C redictive ded for iction

lim

onances around 1.5 MeV to enhance the rate by more than 25 times and ignite the carbon burning at 0.5 GK [35]. With our upper limit, the maximum enhancement is about a factor 2 instead. It suggests more physics needs to be included in the current superburst model to solve the ignition problem.

4. Conclusion

As a conclusion, the strong correlation between the ${}^{12}C+{}^{12}C$ and ${}^{12}C+{}^{13}C$ fusion cross sections offers a great opportunity to constrain the upper limit for the ${}^{12}C+{}^{12}C$ fusion cross sections at lower energies. However, such an upper limit towards the stellar energies suffers from the large deviations among various models. Our new measurement of the ${}^{12}C+{}^{13}C$ fusion cross section down to $E_{c.m.}=2.323$ MeV disagrees with the prediction of the hindrance model and is the first decisive evidence in the carbon isotope systems that rules out the existence of the Sfactor maximum predicted for ${}^{12}C+{}^{13}C$ by this phenomenological model. It also confirms the exponentially rising trend of the S-factor towards lower energies predicted by CC-M3Y+Rep, DC-TDHF, KNS, SPP and ESW. After calibrating these model predictions using our data, a more reliable upper limit is established for the ${}^{12}C+{}^{12}C$ fusion cross sections at stellar energies.

5. Acknowledgments

X.D.T. thanks H. Esbensen, C.Y. Wong, Α. Mukhamedzhanov, D.Y. Pang, X. Fang, W.P. Tan, S.F. Zhu, D.L. Fang and X.D. Xu for the helpful discussions, A. Tumino for providing the THM data and A. Tang for proofreading. This project is supported in part by the National Key Research and Development program (MOST 2016YFA0400501) from the Ministry of Science and Technology of China and the grants PN 16420102 and PN-III-P4-ID-PCE-2016-0743 NUCASTRO2 from the Romanian Ministry for Research and Innovation. X.D.T. and N.T.Z. acknowledge supports from the National Natural Science Foundation of China under Grants No. 11021504, No. 11321064, No. 11475228, No. 11490564, 11405226. X.D.T. also acknowledges supports and No. from the key research program (XDPB09-2) and 100 talents Program of the Chinese Academy of Sciences. B.B. and S.U. acknowledge support by the U.S. DOE under Contract No. DE-AC07-05ID14517 and Grant No. DE-SC0013847.

References

References

- L. R. Gasques, E. F. Brown, A. Chieffi, C. L. Jiang, M. Limongi, C. Rolfs, M. Wiescher, D. G. Yakovlev, Implications of low-energy fusion hindrance on stellar burning and nucleosynthesis, Phys. Rev. C 76 (2007) 035802. doi:10.1103/PhysRevC.76.035802.
- [2] C. A. Barnes, S. Trentalange, S. C. Wu, Treatise on Heavy-Ion Science, Vol. 6, Plenum Preess, New York, 1985, Ch. 1, pp. 3–60.
- [3] J. R. Patterson, H. Winkler, C. S. Zaidins, Experimental investigation of the stellar nuclear reaction 12c-12c at low energies, Astrophys. J. 157 (1969) 367. doi:10.1086/150073.
- [4] G. R. Caughlan, W. A. Fowler, Thermonuclear reaction rates v, Atomic Data and Nuclear Data Tables 40 (2) (1988) 283–334.

- [5] A. S. Umar, V. E. Oberacker, C. J. Horowitz, Microscopic sub-barrier fusion calculations for the neutron star crust, Phys. Rev. C 85 (2012) 055801. doi:10.1103/PhysRevC.85.055801.
- [6] A. Diaz-Torres, M. Wiescher, Characterizing the astrophysical s factor for ¹²C +¹²C fusion with wave-packet dynamics, Phys. Rev. C 97 (2018) 055802. doi:10.1103/PhysRevC.97.055802.
- [7] E. F. Aguilera, P. Rosales, E. Martinez-Quiroz, G. Murillo, M. Fernández, H. Berdejo, D. Lizcano, A. Gómez-Camacho, R. Policroniades, A. Varela, E. Moreno, E. Chávez, M. E. Ortíz, A. Huerta, T. Belyaeva, M. Wiescher, New γ-ray measurements for ¹²C +¹² C sub-coulomb fusion: Toward data unification, Phys. Rev. C 73 (2006) 064601. doi:10.1103/PhysRevC.73.064601.
- [8] M. Assuncao, P. Descouvemont, Role of the hoyle state in c-12+c-12 fusion, Phys. Lett. B 723 (4-5) (2013) 355-359. doi:10.1016/j.physletb.2013.05.030.
- [9] H. Esbensen, X. Tang, C. L. Jiang, Effects of mutual excitations in the fusion of carbon isotopes, Phys. Rev. C 84 (2011) 064613. doi:10.1103/PhysRevC.84.064613.
- [10] B. B. Back, H. Esbensen, C. L. Jiang, K. E. Rehm, Recent developments in heavy-ion fusion reactions, Rev. Mod. Phys. 86 (2014) 317–360. doi:10.1103/RevModPhys.86.317.
- [11] C. L. Jiang, K. E. Rehm, B. B. Back, R. V. F. Janssens, Expectations for 12c and 16o induced fusion cross sections at energies of astrophysical interest, Phys. Rev. C 75 (2007) 015803. doi:10.1103/PhysRevC.75.015803.
- [12] C. L. Jiang, D. Santiago-Gonzalez, S. Almaraz-Calderon, K. E. Rehm, B. B. Back, K. Auranen, M. L. Avila, A. D. Ayangeakaa, S. Bottoni, M. P. Carpenter, C. Dickerson, B. DiGiovine, J. P. Greene, C. R. Hoffman, R. V. F. Janssens, B. P. Kay, S. A. Kuvin, T. Lauritsen, R. C. Pardo, J. Sethi, D. Seweryniak, R. Talwar, C. Ugalde, S. Zhu, D. Bourgin, S. Courtin, F. Haas, M. Heine, G. Fruet, D. Montanari, D. G. Jenkins, L. Morris, A. Lefebvre-Schuhl, M. Alcorta, X. Fang, X. D. Tang, B. Bucher, C. M. Deibel, S. T. Marley, Reaction rate for carbon burning in massive stars, Phys. Rev. C 97 (2018) 012801. doi:10.1103/PhysRevC.97.012801.
- [13] A. Tumino, C. Spitaleri, M. La Cognata, S. Cherubini, G. L. Guardo, M. Gulino, S. Hayakawa, I. Indelicato, L. Lamia, H. Petrascu, R. G. Pizzone, S. M. R. Puglia, G. G. Rapisarda, S. Romano, M. L. Sergi, R. Spartá, L. Trache, An increase in the 12c + 12c fusion rate from resonances at astrophysical energies, Nature 557 (2018) 687. doi:10.1038/s41586-018-0149-4.
- [14] K. Mori, M. A. Famiano, T. Kajino, M. Kusakabe, X. Tang, Impacts of the new carbon fusion cross-sections on type ia supernovae, Monthly Notices of the Royal Astronomical Society: Letters 482 (1) (2019) L70–L74. doi:10.1093/mnrasl/sly188.
- [15] H. W. Becker, K. U. Kettner, C. Rolfs, H. P. Trautvetter, The 12c-12c reaction at subcoulomb energies, Zeitschrift f
 ür Physik A 303 (4) (1981) 305–312. doi:10.1007/BF01421528.
- [16] T. Spillane, F. Raiola, C. Rolfs, D. Schürmann, F. Strieder, S. Zeng, H.-W. Becker, C. Bordeanu, L. Gialanella, M. Romano, J. Schweitzer, ¹²C +¹² C fusion reactions near the gamow energy, Phys. Rev. Lett. 98 (2007) 122501. doi:10.1103/PhysRevLett.98.122501.
- [17] R. Dayras, R. Stokstad, Z. Switkowski, R. Wieland, Gammaray yields from 12c-13c reactions near and below the coulomb barrier, Nuclear Physics A 265 (1) (1976) 153–188. doi:10.1016/0375-9474(76)90121-4.
- [18] B. Dasmahapatra, B. Čujec, F. Lahlou, Fusion cross for 12c-12c, 12c-13c 13c-13c low ensections and at ergies, Nuclear Physics A 384 (1) (1982) 257-272. doi:http://dx.doi.org/10.1016/0375-9474(82)90316-5.
- [19] M. Notani, H. Esbensen, X. Fang, B. Bucher, P. Davies, C. L. Jiang, L. Lamm, C. J. Lin, C. Ma, E. Martin, K. E. Rehm, W. P. Tan, S. Thomas, X. D. Tang, E. Brown, Correlation between the 12c-12c, 12c-13c, and 13c-13c fusion cross sections, Phys. Rev. C 85 (2012) 014607. doi:10.1103/PhysRevC.85.014607.
- [20] C. L. Jiang, B. B. Back, H. Esbensen, R. V. F. Janssens, K. E. Rehm, R. J. Charity, Origin and consequences of ¹²C +¹² C fusion resonances at deep sub-barrier energies, Phys. Rev. Lett. 110 (2013) 072701. doi:10.1103/PhysRevLett.110.072701.
- [21] A. Shrivastava, K. Mahata, S. Pandit, V. Nanal, T. Ichikawa, K. Hagino, A. Navin, C. Palshetkar, V. Parkar, K. Ramachandran, P. Rout, A. Kumar,

A. Chatterjee, S. Kailas, Evolution of fusion hindrance for asymmetric systems at deep sub-barrier energies, Phys. Lett. B 755 (2016) 332 – 336. doi:10.1016/j.physletb.2016.02.029.

- [22] G. Montagnoli, A. M. Stefanini, C. L. Jiang, K. Hagino, F. Galtarossa, G. Colucci, S. Bottoni, C. Broggini, A. Caciolli, P. Čolović, L. Corradi, S. Courtin, R. Depalo, E. Fioretto, G. Fruet, A. Gal, A. Goasduff, M. Heine, S. P. Hu, M. Kaur, T. Mijatović, M. Mazzocco, D. Montanari, F. Scarlassara, E. Strano, S. Szilner, G. X. Zhang, Fusion hindrance for the positive *q*-value system ¹²C +³⁰ Si, Phys. Rev. C 97 (2018) 024610. doi:10.1103/PhysRevC.97.024610.
- [23] I. Burducea, M. Straticiuc, D. Ghiţă, D. Moşu, C. Călinescu, N. Podaru, D. Mous, I. Ursu, N. Zamfir, A new ion beam facility based on a 3mv tandetronTM at ifin-hh, romania, Nucl. Instr. and Meth. in Phys. Res. B 359 (2015) 12 – 19. doi:10.1016/j.nimb.2015.07.011.
- [24] R. Margineanu, C. Simion, S. Bercea, O. Duliu, D. Gheorghiu, A. Stochioiu, M. Matei, The slanic-prahova (romania) underground lowbackground radiation laboratory, Applied Radiation and Isotopes 66 (10) (2008) 1501 – 1506. doi:10.1016/j.apradiso.2008.04.002.
- [25] J. F. Ziegler, Website, www.srim.org (accessed February 2014).
- [26] A. J. Koning, D. Rochman, Modern nuclear data evaluation with the talys code system, Nucl. Data Sheets 113 (12, SI) (2012) 2841–2934. doi:10.1016/j.nds.2012.11.002.
- [27] M. Herman, R. Capote, B. V. Carlson, P. Oblozinsky, M. Sin, A. Trkov, H. Wienke, V. Zerkin, Empire: Nuclear reaction model code system for data evaluation, Nucl. Data Sheets 108 (12) (2007) 2655–2715. doi:10.1016/j.nds.2007.11.003.
- [28] C. Simenel, A. S. Umar, K. Godbey, M. Dasgupta, D. J. Hinde, How the pauli exclusion principle affects fusion of atomic nuclei, Phys. Rev. C 95 (2017) 031601. doi:10.1103/PhysRevC.95.031601.
- [29] S. Mişicu, H. Esbensen, Hindrance of heavy-ion fusion due to nuclear incompressibility, Phys. Rev. Lett. 96 (2006) 112701. doi:10.1103/PhysRevLett.96.112701.
- [30] S. Mişicu, H. Esbensen, Signature of shallow potentials in deep sub-barrier fusion reactions, Phys. Rev. C 75 (2007) 034606. doi:10.1103/PhysRevC.75.034606.
- [31] K. Godbey, C. Simenel, A. S. Umar, Absence of hindrance in a microscopic ¹²C +¹² C fusion study, Phys. Rev. C 100 (2019) 024619. doi:10.1103/PhysRevC.100.024619.
- [32] A. Mukhamedzhanov, X. Tang, D. Pang, Comments on the 12c+12c fusion s-factor (2018). arXiv:1806.05921.
- [33] A. M. Mukhamedzhanov, D. Y. Pang, A. S. Kadyrov, Phys. Rev. C 99 (2019) 064618. doi:10.1103/PhysRevC.99.064618.
- [34] A. Tumino, et al., Reply to the comments on the 12c+12c fusion s-factor (2018). arXiv:1807.06148.
- [35] R. L. Cooper, A. W. Steiner, E. F. Brown, Possible resonances in the c-12+c-12 fusion rate and superburst ignition, ASTROPHYSICAL JOUR-NAL 702 (1) (2009) 660–671. doi:10.1088/0004-637X/702/1/660.