

On R -triviality of F_4

Maneesh Thakur

Abstract

It is known that simple algebraic groups of type F_4 defined over a field k are precisely the full automorphism groups of Albert algebras over k . We explore R -triviality for the group $\mathbf{Aut} A$ when A is an Albert algebra. In this paper, we consider the case when A is an Albert division algebra, that arises from the first Tits construction. We prove that $\mathbf{Aut} A$ is R -trivial, in the sense of Manin.

Keywords: Exceptional groups, Algebraic groups, Albert algebras, Structure group, Kneser-Tits conjecture, R -triviality.

1 Introduction

F_4 is a simple algebraic group of type F_4 defined over a field k . It is known that simple algebraic groups of type F_4 defined over a field k are precisely the full automorphism groups of Albert algebras over k . We explore R -triviality for the group $\mathbf{Aut} A$ when A is an Albert algebra. In this paper, we consider the case when A is an Albert division algebra, that arises from the first Tits construction. We prove that $\mathbf{Aut} A$ is R -trivial, in the sense of Manin.

2 Preliminaries

Let J be a finite dimensional algebra over a field k .
 A *cubic norm structure* on J is a map $N: J \rightarrow k$ satisfying:
 1. $N(1) = 1$ (identity element)
 2. $N(x) = x^3$ for all $x \in J$.

$N: J \rightarrow k$ is a cubic form on J .

N is a trace form on J .

The trace form $T: J \times J \rightarrow k$ is defined by $T(x, y) = \frac{1}{3}(x+y)^3 - x^3 - y^3$.

The adjoint map $J \rightarrow J$ is defined by $T(x^\#, y) = \frac{1}{3}(x+y)^3 - x^3 - y^3$.

$x^\# = 3x - x^3$.

$c^\# = 3c - c^3$.

$c \times x = T(c, x) - T(x, c)$ is the trace of x in the direction of c .

The logarithmic derivative $U_x: J \rightarrow J$ is defined by $U_x(y) = \frac{y}{x} \log f$ where $f = N(x+y)$.
 The directional derivative $\frac{y}{x} f$ is defined by $\frac{y}{x} f = \frac{1}{3}(x+y)^3 - x^3 - y^3$.

A quadratic Jordan algebra J is a finite dimensional algebra over a field k with a quadratic trace T and a set of U -operators U_x for $x \in J$.

The division algebra U_x is invertible if $x \neq 0$.
 The division algebra U_x is defined by $U_x(y) = \frac{y}{x} \log f$.

Example.

Let D be a division algebra over k .
 The Tits process $T_D: D \rightarrow D$ is defined by $T_D(x) = \frac{1}{3}(x+x^3)$.
 The division algebra B, α is defined by $B, \alpha = \{b \in B \mid \sigma(b) = b\}$.

Tits process :

$N_D: T_D \rightarrow k$ is defined by $N_D(x) = \lambda N_D(x) - \lambda^{-1} N_D(x) - T_D(xy)$.

$$N_D(x, y) = \lambda N_D(x) - \lambda^{-1} N_D(x) - T_D(xy)$$

$$x, y, z \#_{L} x\# - yz, \lambda^{-1}z\# - xy, \lambda y\# - zx.$$

first Tits process

$J D, \lambda$ D $J D, \lambda$ D_+

$\lambda \notin N_D$ D λ

K n k B

K K/k σ $x \mapsto \bar{x}$ K

admissible pair u, μ $u \in B, \alpha +$ N_B u $\mu \in K^\times$

$B, \alpha + \oplus B$ I

N $b, \alpha \#_{L}$ N_B $T_K \mu N_B$ $- T_B b x u \sigma$

$b, \alpha \#_{L}$ $b\# - x u \sigma$ $, \bar{\mu} \sigma \#_{L} u^{-1} - b \sigma$ $, \nu_{L}$ $\nu_{B, \Gamma}$.

second Tits process

$J B, \sigma, u, \mu$ $J B, \sigma, u, \mu$

$B, \alpha +$ B B, α, u μ

K $k \times k$ B $D \times D^\circ$ σ D°

D $J B, \sigma, u, \mu$

$J D, \lambda$

Tits constructions of Albert algebras :

D $\lambda \in k^\times$ *first Tits construction* A $J D, \lambda$ k

K k *second Tits construction* A $J B, \sigma, u, \mu$ k

k A $J B, \sigma, u, \mu$ $A \otimes_k K$ $J B, \mu$ K

K B A A *pure first construction*

pure second construction A

B, α K n k A

Norm similarities of Albert algebras :

D *norm similarity* A $A \rightarrow A$

N f ν f N $x \in A$ ν $f \in k^\times$ k

A k N $a \in A$ U U_a


U_a $y \#_{L}$ $T a, y$ $a - a\# \times y$ $y \in A$ $a \in A$ U_a

A $x \in A$ $N U_a$ $N^2 N$


N_D $aa\#$ $a\#a$ $\forall a \in D$ $xy \#_{L}$ $y\#x\#$ $a \mapsto a\#$

N $x\#$ N a^2


Algebraic groups arising from Albert algebras :



R-equivalence and *R*-triviality:



3 Automorphisms of Albert algebras



Proposition 3.1 *Let A be an Albert algebra over k .*

1. Suppose $S \cong D_+ \subset A$ for D a degree 3 central simple k -algebra. Write $A \cong J(D, \mu)$ for some $\mu \in k^\times$. Then $\text{Aut } A/S \cong \text{SL}_3(D)$, the algebraic group of norm 1 elements in D .

2. Let $S \simeq B, \alpha_+ \subset A$ for B a degree n central simple algebra of degree n^2 over a quadratic extension K of k with an involution σ of the second kind over K/k . Write $A \simeq J(B, \sigma, u, \mu)$ for suitable parameters. Then $\text{Aut } A/S \simeq \text{SU}(B, \sigma, \mu)$, where $\sigma_u \simeq \text{Int } u \circ \sigma$.

In particular, the subgroups described in (1) and (2) are simply connected, simple of type A_2 , defined over k and since these are rank- n groups, they are rational and hence are R-trivial.

Theorem 3.1 *Let A be an Albert division algebra over k and $\phi \in \text{Aut } A$ an automorphism of A . Then ϕ fixes a cubic subfield of A pointwise.*

Extending automorphisms : A $S \subset A$

$\{g \in B^\times \mid g\sigma(g) \in k^\times\}$ $x \mapsto pxp^{-1}$ $p \in U(B, \sigma)$ $\sigma_u \simeq \text{Int } u \circ \sigma$

Proposition 3.2 *Let $A \simeq J(B, \sigma, u, \mu)$ be an Albert algebra over k with $S \simeq B, \alpha_+$ a division algebra and $K \simeq Z(B)$. Let $\phi \in \text{Aut } S$ be given by $\phi(a) = g\sigma(g)^{-1}a$ for $g \in \text{Sim } B, \alpha$ with $g\sigma(g) = \lambda \in k^*$ and $\nu \in N_B \setminus \{0\} \in K^*$. Let $q \in U(B, \sigma)$ be arbitrary with $N_B(q) = \nu^{-1}\nu$. Then the map $\phi \simeq A \rightarrow A$, given by*

$$a, b \mapsto gag^{-1}, \lambda^{-1}\sigma(g)\#bq,$$

is an automorphism of A extending ϕ .

Corollary 3.1 *Let $A \simeq J(D, \mu)$ be an Albert algebra arising from the first Tits construction, where D is a degree n central division algebra. Let $\phi \in \text{Aut } D$ be given by $\phi(a) = g\sigma(g)^{-1}a$ for $g \in D^\times$ and $x \in D_+$. Then the map*

$$x, y, z \mapsto gxg^{-1}, gyh^{-1}, hzg^{-1},$$

for any $h \in D^\times$ with $N_D(h) = N_D(g)$, is an automorphism of A extending ϕ .

Proposition 3.3 *Let $A \simeq J(B, \sigma, u, \mu)$ be an Albert algebra with $S \simeq B, \alpha_+$ a division algebra and $K \simeq Z(B)$, a quadratic étale extension of k . Then any automorphism of A stabilizing S is of the form $a, b \mapsto pap^{-1}, pbq$ for $p \in U(B, \alpha)$ and $q \in U(B, \sigma)$ with $N_B(q) = N_B(p)$. We have*


$$\text{Aut } A, B, \alpha_+ \simeq U(B, \alpha) \times U(B, \sigma) \backslash \text{det} / K^{(1)},$$

where

$$U(B, \alpha) \times U(B, \sigma) \backslash \text{det} = \{p, q \in U(B, \alpha) \times U(B, \sigma) \mid N_B(p) = N_B(q)\},$$

and $K^{(1)}$ denotes the group of norm-1 elements in K , embedded diagonally.

4 Norm similarities and R -triviality



Theorem 4.1 *Let A be an Albert division algebra over k and $S \subset A$ be a n -dimensional subalgebra. Then every element $\psi \in \text{Str } S$ admits an extension $\tilde{\psi} \in \text{Str } A$. Let $S \simeq B, \alpha_+$ and $A \simeq J B, \sigma, u, \mu$ for suitable parameters. Let $\psi \in \text{Str } S$ be given by $b \mapsto \gamma g b \sigma g$ for $b \in B, \alpha_+$, $g \in B^\times$ and $\gamma \in k^\times$. Then for any $q \in U B, \sigma$ with $N_B q = N_B \sigma g^{-1} g$, the map $\tilde{\psi}$ given by*

$$\tilde{\psi}(b, \alpha) = \gamma g b \sigma g, \sigma g \# x q,$$

is a norm similarity of A extending ψ .

Theorem 4.2 *Let $A \simeq J B, \sigma, u, \mu$ be an Albert division algebra, written as a second Tits construction and $S \simeq B, \alpha_+$. The group $\text{Str } A, S$ consists of the maps $b, \alpha \mapsto \gamma g b \sigma g, \sigma g \# x q$ where $\gamma \in k^\times, g \in B^\times$ are arbitrary and $q \in U B, \sigma$ satisfies $N_B q = N_B \sigma g^{-1} g$. We have,*

$$\text{Str } A, B, \alpha_+ \simeq \frac{k^\times \times H_0}{K^\times},$$


where $H_0 = \{g, q \in B^\times \times U B, \sigma \mid N_B q = N_B \sigma g^{-1} g\}$ and $K^\times \hookrightarrow k^\times \times H_0$ via $\alpha \mapsto (N_K \alpha^{-1}, \alpha, \bar{\alpha}^{-1} \alpha)$.

Corollary 4.1 *Let $A \simeq J D, \mu$ be an Albert algebra arising from the first Tits construction, where D is a degree d central division algebra over k . Then the group $\text{Str } A, D$ consists of the maps*

$$x, y, z \mapsto \gamma a x b, b \# y c, c^{-1} z a \#,$$

where $a, b, c \in D^\times, \gamma \in k^\times$ with $N_D a = N_D b = N_D c$.

5 R-triviality results




Theorem 5.1 *Let A be an Albert (division) algebra. Let $S \subset A$ be a n -dimensional subalgebra. Then, with the notations as above, $\text{Str } A, \mathbf{S}$ is R-trivial.*

Corollary 5.1 *Let A be an Albert (division) algebra and S a n -dimensional subalgebra of A . Then $\text{Aut } A, \mathbf{S} \subset R \text{Str } A, \mathbf{S}$.*

Theorem 5.2. *Let A be an Albert division algebra over a field k of arbitrary characteristic. Then $\text{Str } A, \mathbf{S} \subset R \text{Str } A, \mathbf{S}$.*

Theorem 5.3. *Let A be an Albert division algebra arising from the first Tits construction, over a field k of arbitrary characteristic. Then $G, \mathbf{S} \subset \text{Aut } A, \mathbf{S}$ is R-trivial.*

Theorem 5.4. *Let A be an Albert division algebra arising from the first Tits construction over a field k and $L \subset A$ a cyclic cubic subfield. Let $\phi \in \text{Aut } A/L$. Then ϕ is a product of two automorphisms, each stabilizing a n -dimensional subalgebra.*



Theorem 5.5. *Let A be an Albert division algebra arising from the first Tits construction, over a field k of arbitrary characteristic. Let $S \subset A$ be a n -dimensional subalgebra with trivial mod- n invariant. Then $\text{Aut } A, \mathbf{S} \subset R \text{Aut } A, \mathbf{S}$.*

Proof. $S \cong D_+ \cong D \times D^{\text{op}}$, $\nu \in k^\times$, $\gamma \in \text{Aut } J(D, \nu)$, $\phi \in \text{Aut } A, \mathbf{S} \subset D$, $a, b \in D^\times$, $N_D(a) = N_D(b)$, $\phi(x, y, z) = (axa^{-1}, ayb^{-1}, bza^{-1})$, $\forall x, y, z \in D$.
 $D \subset \text{Aut } A, \mathbf{S} \subset R \text{Aut } A, \mathbf{S}$, $\phi \in \mathcal{I}_{ab^{-1}} \mathcal{I}_a$.

$$\mathcal{J}_c \in \text{Aut } A, c \in \mathcal{O}^*, D, \mathcal{J}_c(x, y, z) = x, yc, c^{-1}z$$

$$\mathcal{I}_d \in \text{Aut } A, d \in D^\times, \mathcal{I}_d(x, y, z) = dxd^{-1}, dyd^{-1}, dzd^{-1}, \forall x, y, z \in D.$$

$$\mathcal{J}_p(x, y, z) = x, yp, p^{-1}z, p = ab^{-1} \in \mathcal{O}^*, D, \mathcal{I}_a(x, y, z) = axa^{-1}, aya^{-1}, aza^{-1}, \forall x, y, z \in D.$$

$$\text{SL}_3(D) \curvearrowright \text{Aut } A, D, R_\nu \curvearrowright \theta \in \text{Aut } A_k^1 \rightarrow \text{Aut } A, D,$$

$$\theta(x, y, z) = a_t x a_t^{-1}, a_t y a_t^{-1}, a_t z a_t^{-1}, x, y, z \in D,$$

$$a_t \in \mathcal{O}^*, t \in D, \theta \in \text{Aut } A_k^1 \rightarrow \text{Aut } A, D, \mathcal{I}_a, \theta \curvearrowright \mathcal{J}_{ab^{-1}}, \mathcal{I}_a \curvearrowright \text{RAut } A, D, k_F$$

$$\phi \in \text{RAut } A, D, S_\sigma(B, \alpha) \curvearrowright K_F^n \curvearrowright B, \alpha \curvearrowright N_B(u, \mu) \curvearrowright A, S, D, \sigma \curvearrowright J(B, \sigma, u, \mu) \curvearrowright B, \alpha \times K^\times \curvearrowright N_B(u, \mu)$$

$$f_3(A) \curvearrowright f_3(J(B, \sigma, u, \mu)) \curvearrowright f_3(B, \sigma) \curvearrowright f_3(B, \alpha).$$

$$\sigma_u \curvearrowright w \in B^\times, \lambda \in k^\times, u \curvearrowright \lambda w \sigma u^{-1} \curvearrowright J(B, \sigma, \lambda w \sigma u^{-1}, \mu) \curvearrowright J(B, \sigma, \lambda, N u \sigma^{-1} \mu).$$

$$D \curvearrowright N \lambda \curvearrowright \lambda^3 \curvearrowright N u \sigma^{-1} \mu \overline{N u \sigma^{-1} \mu} \curvearrowright \lambda \curvearrowright N u \sigma^{-1} \mu \lambda^{-1} \overline{N u \sigma^{-1} \mu \lambda^{-1}}.$$

$$\gamma \in \text{Aut } A \curvearrowright J(B, \sigma, u, \mu) \curvearrowright J(B, \sigma, \lambda, N u \sigma^{-1} \mu) \curvearrowright J(B, \sigma, \mu)$$

$$\nu \curvearrowright \mu^{-2} N u \sigma^2 \lambda^3 \curvearrowright \gamma \phi \gamma^{-1} \curvearrowright J(B, \sigma, \mu, u) \curvearrowright J(B, \sigma, \mu) \curvearrowright B, \alpha \curvearrowright \phi \in \text{RAut } A, S, \phi \in \text{RAut } A', S', A' \curvearrowright J(B, \sigma, \mu, u) \curvearrowright S' \curvearrowright B, \alpha \curvearrowright A', D \curvearrowright \phi' \curvearrowright \psi_{p, q}, p, q \in B, \alpha \curvearrowright N_B(p) \curvearrowright N_B(q) \curvearrowright \phi'(b) \curvearrowright p b p^{-1}, p x q^{-1}, \forall b \in B, \alpha, x \in B.$$

$$N_B(p) \curvearrowright N_B(q) \curvearrowright N_B(p q^{-1}) \curvearrowright p q^{-1} \in \mathcal{O}^*, B, \alpha \curvearrowright \psi_{p, q} \curvearrowright \psi_{p, p} \psi_{1, p q^{-1}}.$$

$$D \curvearrowright \gamma \phi \gamma^{-1} \in \text{RAut } A', S' \curvearrowright \text{Aut } B, \alpha \curvearrowright \text{Aut } B, \alpha \curvearrowright \text{Aut } A, S \curvearrowright \square$$

Proposition 5.1. *Let A be an Albert division algebra arising from the first Tits construction. Let $U \subset A$ be the Zariski open set $U = \{a \in A \mid k[a] \text{ is étale}\}$. Then for $a \in U$, there exists $v \in v \circ a \in L^\perp$, $v \circ / \Gamma$ such that $a \mapsto \lambda \circ L \circ N \circ v \circ a$ is a morphism $U \rightarrow \mathbb{A}_k^1$ defined over k .*

Proof. $\delta \circ a \in U$
 $a \in U$
 k
 $a \mapsto \delta \circ a$
 $U \rightarrow \mathbb{A}_k^1$
 $L \circ k$
 A
 $\delta \circ a$
 $k \circ D$
 L
 $\langle \delta \rangle^\perp$
 n_{C_0}
 C
 $pure \ norm \ n_{C_0}$
 C
 k
 n_C
 $v \in C_0$
 $v \circ / \Gamma$
 n_{C_0}
 $q_L \ v \otimes \dots$
 $L \circ k \circ a$
 $k \circ a$
 $a \in k \circ n$
 $x \in L^\perp$
 $x^\#$
 L
 $quadratic \ trace \ form$
 A
 n
 L
 k
 n_C
 $q_L \ L$
 A
 A
 $v \circ a$
 $v \circ a \circ L$
 $v \otimes k \circ a$
 $v \circ a \circ / \Gamma$
 $v \circ a$
 a
 k
 $a \mapsto v \circ a$
 $I \ U \rightarrow A$
 N
 $f \ A$
 k
 \square

Theorem 5.6. *Let A be an Albert division algebra over a field k of arbitrary characteristic, arising from the first Tits construction and $S \subset A$ a a -dimensional subalgebra. Then $Aut \ A, S \subset RAut \ A \ k$.*

Proof. A S
 $\theta \in \mathbb{A}_k^1 \rightarrow Str \ A, S$
 k
 $\phi \in A, S$
 $\theta \circ a \in S$
 $L_t \circ k \circ a$
 $v_t \in L_t^\perp$
 $\lambda_t \circ N \circ v_t$
 $J \ L_t, \lambda_t$
 $\chi_t \in A, J \ L_t, \lambda_t$
 $R_{N_{L_t}(a_t)} U_{(0,0,1)} U_{(0, N_{L_t}(a_t)^{-1}, 0)}$
 $R_{N_{L_t}(a_t)}$
 A
 $N_{L_t} \ a_t$
 $J \ L_t, \lambda_t$
 $L_t \oplus L_t \oplus L_t$
 A
 $J \ L_t, \lambda_t$
 $\chi_t \theta \circ a_t \in A$
 $\eta \mid t \mapsto \chi_t \theta \circ a_t$
 $\mathbb{A}_k^1 \rightarrow Aut \ A$
 k
 η
 $\chi_0 \theta$
 ϕ_t
 θ
 ϕ
 $\phi_t \circ a_t$
 θ
 a_1
 θ
 \square

Corollary 5.2. *Let A be an Albert division algebra arising from the first Tits construction and $L \subset A$ be a cyclic cubic subfield. Then $Aut \ A/L \subset RAut \ A \ k$.*

Proof. $\rho \in \text{Gal}(L/k) \subset D_+ \subset A$. Let $\tilde{\rho} \in \text{RAut}(A/k)$ be a lift of ρ . Then $\tilde{\rho}^{-1}\rho \in \text{Aut}(A/L) \cap D_+ = \{1\}$. Hence $\tilde{\rho} = \rho$. Since $\tilde{\rho} \in \text{RAut}(A/k)$, we have $\tilde{\rho}^{-1}\rho \in \text{RAut}(A/k)$. \square

Corollary 5.3. *Let A be an Albert division algebra arising from the first Tits construction and $L \subset A$ be a cyclic cubic subfield. Then $\text{Aut}(A, L) \subset \text{RAut}(A/k)$.*

Proof. Let $\phi \in \text{Aut}(A, L)$. Then $\phi|_{L/k} \in \text{Gal}(L/k) \subset D_+ \subset A$. Let $\tilde{\rho} \in \text{RAut}(A/k)$ be a lift of $\phi|_{L/k}$. Then $\tilde{\rho}^{-1}\phi \in \text{Aut}(A/L) \cap D_+ = \{1\}$. Hence $\tilde{\rho} = \phi$. Since $\tilde{\rho} \in \text{RAut}(A/k)$, we have $\phi \in \text{RAut}(A/k)$. \square

Proposition 5.2 [1, Prop. 1.1]. *Let A be an Albert division algebra over a field k of arbitrary characteristic. Let $M \subset A$ be a cubic subfield and S_M denote the set of all 3-dimensional subalgebras of A that contain M . Then the group $\text{Str}(A, M)$ acts on S_M .*

Proof. Let $\psi \in \text{Str}(A, M)$ and $S \in S_M$. Then $\psi S \in S_M$. For any $\phi \in \text{Aut}(A/S)$, $\phi \psi S \in S_M$. Hence $\psi \phi S \in S_M$. Since $\psi \phi S \in S_M$ and $\psi S \in S_M$, we have $\psi \phi S = \psi S$. Hence $\phi S = S$. \square

Proposition 5.3 [1, Prop. 1.2]. *Let the notation be as above, $\psi \in \text{Str}(A, M)$ and $S \in S_M$. Then S and ψS are k -isomorphic as subalgebras of A .*

Proof. Let $f \in \text{Aut}(A/S)$. Then $f \psi S \in S_M$. Let $x \in S$. Then $\psi f x \in \psi S$. Hence $\psi f \psi^{-1} x \in \psi S$. \square

\square $\psi f \psi^{-1} x \in \psi S$. Let $f \in \text{Aut}(A/S)$. Then $f \psi S \in S_M$. Let $\phi \in \text{Aut}(A/\psi S)$. Then $\phi \psi S \in S_M$. Hence $\psi \phi S \in S_M$. Since $\psi \phi S \in S_M$ and $\psi S \in S_M$, we have $\psi \phi S = \psi S$. Hence $\phi S = S$. \square

$$A_2 \begin{matrix} \text{Aut } A/S & \text{Aut } A/\psi S \\ S \simeq C, \tau_0 & w \in C, \tau_0 \end{matrix} \quad \tau$$

$$Z_{\text{Aut}(A,S)} \text{Aut } A/S \simeq \text{SU } C, \tau \quad Z_{\text{Aut}(A,\psi(S))} \text{Aut } A/\psi S \simeq \text{SU } C, \tau_0.$$

$$\begin{matrix} \text{Aut } A/S & \text{Aut } A/\psi S \\ S \simeq C, \tau_0 & \psi S \simeq k \end{matrix} \quad \square$$

Proposition 5.4 *Let $L \subset A$ be a cyclic cubic subfield. Let $\psi \in \text{Str } A, \mathbf{L}$ and $S \in \mathcal{S}_L$, where $S \simeq B, \alpha_+$ for a degree 3 central division algebra B over a quadratic étale extension K/k and σ is a distinguished unitary involution on B . Then there exists $\phi \in \text{Aut } A, \mathbf{L}$ such that $\phi^{-1}\psi \in \text{Str } A, \mathbf{S}$.*

Proof. Let $v \in L$, $w \in LK^\times$, $\lambda \in k^\times$, $v = \lambda w w^* \sigma$.
 $L \subset B, \alpha_+ \xrightarrow{\eta} J LK, *, u, \mu \rightarrow B, \alpha_+$
 $L \xrightarrow{\eta} J LK, *, u', \mu' \rightarrow \psi S \simeq B, \sigma_+$
 $L \xrightarrow{\eta} J LK, *, u', \mu' \rightarrow \psi S \simeq B, \sigma_+$
 $w^\#, N_L u, \mu \rightarrow J LK, *, u, \mu$
 $w^\#, N_L u, \mu \rightarrow J LK, *, u, \mu$

$$u' = w^\# u \lambda w w^* \# = u \lambda^2 w^\# w^* = u w_0 w_0^*, \mu' = N_L u, \mu,$$

$$w_0 = \lambda w^\# \rightarrow \psi S \simeq B, \sigma_+$$

$$\phi_1 J LK, *, w^\#, N_L u, \mu \rightarrow J LK, *, u w_0 w_0^*, \mu \rightarrow J LK, *, u, N_{LK} w_0^{-1} \mu.$$

$$N_{LK} w_0 \overline{N_{LK} w_0} = N_{LK} \lambda w^\# \overline{N_{LK} \lambda w^\#} = \lambda^3 N_{LK} u^2 \lambda^3 \overline{N_{LK} u^2}$$

$$= \lambda^3 N_{LK} u \overline{N_{LK} u}^2 = N_L u^2 \nu.$$

$$w_1 \in LK \rightarrow w_1 w_1^* \rightarrow N_{LK} w_1 \overline{N_{LK} w_1}$$

$$\phi_2 J LK, *, u, N_{LK} w_0^{-1} \mu \rightarrow J LK, *, u, \mu$$

$$\begin{matrix} \phi_2 l, x w_1^{-1} \\ \phi_1^{-1} \phi_2^{-1} J LK, *, u, \mu \rightarrow J LK, *, u', \mu' \\ \phi_1^{-1} \psi S \simeq S \rightarrow \phi \rightarrow \phi \rightarrow A \end{matrix} \quad \square$$

Theorem 5.7. *Let A be an Albert division algebra arising from the first Tits construction and $L \subset A$ a cyclic cubic subfield. Then $\text{Str } A, \mathbf{L} \subset \text{RStr } A$.*

Proof. $\psi \in \text{Aut}(A, L) \implies D_+ \subset A$
 $k \subset L \subset D_+ \xrightarrow{\psi} D_+ \xrightarrow{\psi} D \times D^{op}, \psi \in \text{Aut}(A, L)$
 $\phi \in \text{Aut}(A, L) \implies \phi^{-1}\psi \in \text{Aut}(A, L) \implies \psi \in \text{Aut}(A, L)$

$$\psi(x, y, z) = \gamma axb, b^\#yc, c^{-1}za^\#$$

$$a, b \in L^\times, \gamma \in k^\times \implies c \in D^\times \implies N_D a \in N_D b N_D \implies \gamma \in \text{Aut}(A, L)$$

$$\psi(x, y, z) = axb, b^\#yc, c^{-1}za^\#$$

$$N_D ba^{-1} \in \text{Aut}(A, L) \implies c \in \text{Aut}(A, L) \implies ab^{-1}d \in \text{Aut}(A, L) \implies d \in \text{Aut}(A, L)$$

$$a, b, ab^{-1}d \in L^\times \times L^\times \times ab^{-1}d \in \text{Aut}(A, L)$$

$$R_{L/k} \mathbb{G}_m \times \text{SL}(n, D) \xrightarrow{\theta} R_{L/k} \mathbb{G}_m \times \text{SL}(n, D) \times N_D \implies \theta \in \text{Aut}(A, D, L)$$

$$\theta(x, y, z) = a_t x b_t, b_t^\# y c_t, c_t^{-1} z a_t^\#$$

$$a_{t_1} = -t_1 a_t, b_{t_1} = -t_1 b_t, c_{t_1} = a_t b_t^{-1} d_t$$

$$d_{t_1} = \gamma t_1$$

$$\theta(x, y, z) = axb, b^\#yab^{-1}d, d^{-1}ba^{-1}za^\# \implies axb, b^\#yc, c^{-1}b^{-1}aba^{-1}za^\# \implies \psi(x, y, z)$$

$$\theta(x, y, z) = x, y, c_1^{-1}z \implies x, y, d_1, d_1^{-1}z \implies x, y, z$$

$$\theta \in \text{Aut}(A, L) \implies \theta \in \text{Aut}(A, L) \implies \square$$

Theorem 5.8. *Let A be an Albert division algebra arising from the first Tits construction. Then $\text{Aut } A$ is R -trivial.*

Proof. $\phi \in \text{Aut}(A, L) \implies L \subset A$
 $M_L \phi \in \text{Aut}(A, L) \implies S = \langle L, M \rangle \implies \text{Aut}(A, L)$
 $\eta \in \text{Aut}(A, S) \implies \eta \in \text{Aut}(A, L) \implies \phi^{-1}\eta \in \text{Aut}(A, L)$
 $\pi \in \text{Aut}(A, L) \implies \pi^{-1}\psi \in \text{Aut}(A, S) \implies \psi \in \text{Aut}(A, L)$

$$S = \pi^{-1}\psi S = \pi^{-1}\phi^{-1}\eta S = \pi^{-1}\phi^{-1}S$$

$$\pi^{-1}\phi^{-1} \in \text{Aut}(A, S) \subset \text{Aut}(A, L) \implies \phi \in \text{Aut}(A, L) \implies \square$$

References

- , *R-triviality of groups of type F_4 arising from the first Tits construction*, *J. Algebra* **161** (1994), no. 1, 1–14.
- , *On the R-triviality of the groups $E_{7,1}^{78}$ and $E_{8,2}^{78}$* , *J. Algebra* **161** (1994), no. 1, 15–24.
- , *Linear Algebraic Groups*, Graduate Texts in Mathematics, vol. 42, Springer-Verlag, New York, 1975.
- , *R-equivalence and Special Unitary Group*, *J. Algebra* **209** (1998), no. 1, 1–14.
- , *The rationality problem for semisimple group varieties*, *Math. Ann.* **272** (1991), no. 4, 631–644.
- , *Kneser-Tits for a rank ν form of E_6 (After Veldkamp)*, *J. Algebra* **143** (1992), no. 1, 1–14.
- , *Outer automorphisms of algebraic groups and a Skolem-Noether theorem for Albert algebras*, *J. Algebra* **114** (1988), no. 1, 1–14.
- , *Le Problème de Kneser-Tits*, *Ann. Inst. Fourier (Grenoble)* **32** (1982), no. 3, 326–341.
- , *Spécialisation de la R-équivalence pour les groupes réductifs*, *Ann. Inst. Fourier (Grenoble)* **35** (1985), no. 3, 356–371.
- , *Algebras of odd degree with involution, trace forms and dihedral extensions*, *J. Algebra* **156** (1993), no. 1, 96–114.
- , *Lie algebras*, *J. Algebra* **156** (1993), no. 1, 115–124.
- , *Structure and representations of Jordan algebras*, *J. Algebra* **156** (1993), no. 1, 125–134.
- , *Finite-dimensional division algebras over fields*, *J. Algebra* **156** (1993), no. 1, 135–144.
- , *Some groups of transformations defined by Jordan algebras-I*, *J. Algebra* **201** (1998), no. 1, 1–14.
- , *The Book of Involutions*, *J. Algebra* **161** (1994), no. 1, 15–24.
- , *Cubic forms*, *J. Algebra* **161** (1994), no. 1, 25–34.
- , *A classification theorem for Albert algebras*, *J. Algebra* **350** (2010), no. 1, 1–14.

\mathbb{Z} Tits-index $E_{8,2}^{66}$ over an arbitrary field. 17

Structure theorems for Jordan algebras of degree three over fields of arbitrary characteristic. 32

A survey on Albert algebras. 41

(91) \mathbb{Z} Cubic subfields of exceptional simple Jordan algebras. 41

Springer forms and the first Tits construction of exceptional Jordan division algebras. 45

Albert algebras. 45

Classification of algebras arising from the Tits process. 98

98 \mathbb{Z} Jordan algebras of degree 3 and the Tits process. 98

273 2004 The étale Tits process of Jordan algebras revisited. 273

Algebraic groups and number theory. 273

\mathbb{Z} Tits-index $E_{8,2}^{66}$ over an arbitrary field. 17

83 On the Kneser-Tits problem for triality forms. 83

Octonions, Jordan Algebras and Exceptional Groups. 83

Linear Algebraic Groups. 83

Isotopy and invariants of Albert algebras. 74

Automorphisms of Albert algebras and a conjecture of Tits and Weiss. 365

II Automorphisms of Albert algebras and a conjecture of Tits and Weiss. 372

The cyclicity problem for Albert algebras. 372

Albert algebras and the Tits-Weiss conjecture

Algebraic Groups and Their Birational Invariants

...