

Multiplicity Dependence of Shear Viscosity, Isothermal Compressibility and Speed of Sound in pp collisions at $\sqrt{s} = 7$ TeV

Dushmanta Sahu, Sushanta Tripathy*, and Raghunath Sahoo†

Discipline of Physics, School of Basic Sciences, Indian Institute of Technology Indore, Simrol, Indore 453552, India

Archita Rani Dash

*Department of Physics, School of Advanced Sciences,
Vellore Institute of Technology, Vellore 632014, India*

(Dated: July 28, 2020)

In order to understand the detailed dynamics of systems produced in pp collisions, it is essential to know about the Equation of State (EoS) and various thermodynamic properties. In this work, we study the shear viscosity to entropy density ratio, isothermal compressibility and speed of sound of the system by considering a differential freeze-out scenario. We have used a thermodynamically consistent Tsallis non-extensive statistics to have a better explanation for the dynamics of pp collision systems. While the shear viscosity to entropy density ratio provides information about the measure of fluidity of a system formed in high energy collisions, the isothermal compressibility gives a clear idea about the deviation of the system from a perfect fluid. The speed of sound in the system as a function of $\langle dN_{\text{ch}}/d\eta \rangle$ gives us a vivid picture of the dynamics of the system. The results show quite an intuitive perspective on high multiplicity pp collisions and give us a limit of $\langle dN_{\text{ch}}/d\eta \rangle \gtrsim (10 - 20)$, after which a change in the dynamics of the system may be observed.

PACS numbers:

I. INTRODUCTION

Ultra-relativistic collisions of protons and heavy-ions at the Large Hadron Collider (LHC) at CERN, Switzerland, have been instrumental in understanding the sublime nature of the microscopic world at very high energies. One of the most astounding facets of the microscopic realm is a relatively new state of matter called Quark Gluon Plasma (QGP), which is expected to be formed in such collisions. QGP has partons (quarks and gluons) as the degrees of freedom, and exists at very high temperature and/or baryon density. Earlier, it was believed that there wouldn't be any QGP formation in pp collisions. However, recent studies indicate possible formation of QGP droplets in high multiplicity pp collisions [1–3]. Such reasons compel us to study high energy pp collisions with ever-increasing interest. Understanding the behavior of matter in the hadronic phase of these type of collisions, and having the knowledge about various thermodynamical quantities involved is very useful. The coefficient of shear viscosity to entropy density ratio (η/s), isothermal compressibility (κ_T) and speed of sound (c_s) are such thermodynamic quantities that tell us about the fascinating behaviours of the system.

The space-time evolution of a system formed in ultra-relativistic high energy collisions is governed by its dissipative properties such as the shear viscosity (η). To understand the behavior of a system formed in such col-

lisions, η/s is one of the most important properties. The elliptic flow measurements from heavy-ion collisions at RHIC [4] have found that the medium formed in such collisions gives η/s value closer to the KSS bound (Kovtun-Son-Starinets) [5], which might suggest that QGP is almost a perfect fluid [6, 7]. Thus η/s can be used to study the measure of fluidity of a system.

κ_T gives us information about how the volume of a system changes with the change in pressure at constant temperature [8]. It tells us about the deviation of a real fluid from a perfect fluid. For a perfect fluid, $\kappa_T = 0$; which means the fluid is incompressible. However, no such fluid exists in nature. But, recent findings have shown that QGP behaves as a nearly perfect fluid, having the lowest κ_T estimated till now [9]. This also complements previous findings of the ratio of shear viscosity and entropy (η/s) from ADS/CFT calculations which gives a lower bound (KSS bound) to the ratio [5].

The study of speed of sound will help us to have a proper idea about the equation of state (EOS) of the system. It plays a crucial role in the hydrodynamical evolution of the matter created in the collisions. It also affects the momentum distributions of the particles created in the collision systems. Observations from heavy-ion collisions have proved that the speed of sound is different in three different phases in the evolution of the collision systems, namely the QGP phase, the mixed phase and the hadronic phase. For a massless non-interacting gas, the value of the squared speed of sound, c_s^2 is 1/3, whereas for a hadron gas the value is around 1/5 [10]. The expansion time scale of the system is a measure of the speed of sound which is given by $\tau_{\text{exp}}^{-1} \sim \frac{1}{\epsilon} \frac{\partial \epsilon}{\partial \tau_p} = \frac{1+c_s^2}{\tau_p}$ [11]. Here, ϵ is the energy density of the system and τ_p is the proper time. While the collision time scale is given by

*Presently at: Instituto de Ciencias Nucleares, UNAM, Deleg. Coyoacán, Ciudad de México 04510

†Corresponding Author Email: Raghunath.Sahoo@cern.ch

$\tau_{\text{coll}}^{-1} \sim n\sigma v$, where n is density of particles, σ is the collision cross section and v is the particle velocity. For a system to be in thermal equilibrium, the expansion time scale must be greater than the collision time scale. However, we are taking a differential freeze-out scenario, so it will be interesting to see how the speed of sound varies for different hadronic species as a function of charged particle multiplicity.

A large number of particles are produced in high energy collisions, which demands us to take a statistical approach to study the QCD matter, the particle production and the thermodynamics of the systems. The transverse momentum (p_T) of the final state particles produced in high energy collisions are expected to follow a thermalized Boltzmann-Gibbs (BG) distribution. However, it is experimentally observed that the p_T -spectra in pp collisions at the RHIC [12, 13] and LHC [14–17] energies show a deviation from thermalized Boltzmann distribution. Higher contribution of pQCD effects are responsible for this deviation and the spectra are better described by a combination of Boltzmann-type exponential and pQCD inspired power-law distribution. Although a first principle derivation of Tsallis non-extensive distribution [18] is still a question, empirically it has been very successful in describing the p_T -spectra in hadronic collisions. This is used for getting the multiplicity and thus particle ratios in experimental papers. There are vast theoretical developments in this front to bring up the physics messages, which include considering systems formed in hadronic collisions as away from equilibrium, system thermodynamics etc. Although there are various forms of the Tsallis distribution function used in the literature, a thermodynamically consistent distribution function is used in this paper to describe the p_T -spectra in LHC pp collisions [19]. The deviation from equilibrium is denoted by a parameter q , with $q = 1$ denoting the equilibrium condition (BG scenario). At high charged particle multiplicity in high energy collisions, q tends to 1, which is an indication that the system has attained global equilibrium. The extracted thermodynamic parameters [20] such as temperature, T and non-extensivity parameter, q are obtained for different multiplicity classes, which are then used to have the estimation of shear viscosity to entropy density ratio, isothermal compressibility and the speed of sound in the medium (related to the equation of state).

In this paper, we study the shear viscosity to entropy density ratio, the isothermal compressibility and speed of sound in high energy pp collisions. In section II, we give a brief formulation for shear viscosity, isothermal compressibility and c_s^2 using non-extensive statistics. In section III, the results and discussions are given. Finally we summarize our findings in section IV.

II. FORMULATION

As discussed in the previous section, a finite degree of deviation from the equilibrium statistical description of transverse momentum spectra has been observed by experiments at RHIC and LHC [12–17, 21]. In addition, the matter produced in these experiments evolves rapidly in a non-homogeneous way. Hence, the spatial configuration becomes non-uniform and the global equilibrium in the system is not established [22–25]. Due to this, some thermodynamic observables become non-extensive. This may also happen if there are local temperature fluctuation and long-range correlations in the produced system [26]. Thus, for the calculation of shear viscosity, we use the relativistic non-extensive Boltzmann transport equation where we assume that a non-equilibrium system, which dissipates energy and produces entropy relaxes to a local q -equilibrium after a certain relaxation time. For a detailed description and calculation, one can follow the Refs. [27–29]. Here, we briefly describe the formulation.

The Boltzmann transport equation (BTE) is given by,

$$\frac{\partial f_p}{\partial t} + v_p^i \frac{\partial f_p}{\partial x^i} + F_p^i \frac{\partial f_p}{\partial p^i} = I(f_p), \quad (1)$$

where v_p^i is the velocity of the i th particle and F_p^i is the external force acting on that particle, whereas $I(f_p)$ is the collision integral which gives the rate of change of the non-equilibrium distribution function f_p when the system approaches a q -equilibrium.

Under the approximation of no external force and proceeding with the relaxation time approximation (RTA), the collision integral can be approximated as,

$$I(f_p) \simeq -\frac{(f_p - f_p^0)}{\tau(E_p)}, \quad (2)$$

where $\tau(E_p)$ is called the relaxation time or collision time, which can be interpreted as the mean time between collisions. Also, we consider the thermodynamically consistent Tsallis distribution function, having q value close to 1, as an asymptotic equilibrium function, which is called as a q -equilibrium. Thus, Tsallis distribution function is taken as f_p^0 near the local rest frame of the fluid. One should note here that the above RTA is made in order to simplify the collision term. However, the exact collision term calculation in the non-extensive BTE is more complicated than in the classical BTE and it is beyond the scope of this manuscript. Nevertheless, these calculations can be used to make reasonable first estimations for different transport coefficients [27].

The thermodynamically consistent Tsallis distribution function is given as,

$$f_p^0 = \frac{1}{\left[1 + (q-1) \left(\frac{E_p - \mathbf{p} \cdot \mathbf{u} - \mu}{T}\right)\right]^{\frac{q}{q-1}}}, \quad (3)$$

where \mathbf{u} is the fluid velocity, T and μ are the temperature and the chemical potential respectively.

The stress-energy tensor is given by,

$$T^{\mu\nu} = T_0^{\mu\nu} + T_{dissipative}^{\mu\nu}, \quad (4)$$

where $T_0^{\mu\nu}$ is the ideal part and $T_{dissipative}^{\mu\nu}$ is the dissipative part of the stress-energy tensor. When we provide hydrodynamical description of QCD, the shear viscosity (η) and bulk viscosity (ζ) are included in the dissipative part of stress-energy tensor, which can be written in the local Lorentz frame as,

$$T^{ij} = -\eta \left(\frac{\partial u^i}{\partial x^j} + \frac{\partial u^j}{\partial x^i} \right) - \left(\zeta - \frac{2}{3}\eta \right) \frac{\partial u^i}{\partial x^j} \delta^{ij}. \quad (5)$$

In terms of distribution function, the above expression becomes,

$$T^{ij} = \int \frac{d^3p}{(2\pi)^3} \frac{p^i p^j}{E_p} \delta f_p. \quad (6)$$

Here δf_p is the deviation of the distribution function from the q -equilibrium and is given by (from Eq. 1 and 2),

$$\delta f_p = -\tau(E_p) \left(\frac{\partial f_p^0}{\partial t} + v_p^i \frac{\partial f_p^0}{\partial x^i} \right). \quad (7)$$

Under the assumption of a steady flow of the form $u^i = (u_x(y), 0, 0)$ and space-time independent temperature, Eq. 5 becomes $T^{xy} = -\eta \partial u_x / \partial y$. From Eq. 6 and 7, by using $\mu = 0$ (for LHC energies), we get,

$$T^{xy} = \left\{ -\frac{1}{T} \int \frac{d^3p}{(2\pi)^3} \tau(E_p) \left(\frac{p_x p_y}{E_p} \right)^2 q (f_p^0)^{\frac{2q-1}{q}} \right\} \frac{\partial u_x}{\partial y}. \quad (8)$$

For a single component of hadronic matter, the coefficient of shear viscosity η can be expressed with the non-extensive parameters as,

$$\eta = \frac{1}{15T} \int \frac{d^3p}{(2\pi)^3} \tau(E_p) \frac{p^4 q}{E_p^2} (f_p^0)^{\frac{(2q-1)}{q}} \quad (9)$$

The energy dependent relaxation time is given by,

$$\tau^{-1}(E_a) = \sum_{bcd} \int \frac{d^3p_b d^3p_c d^3p_d}{(2\pi)^3 (2\pi)^3 (2\pi)^3} W(a, b \rightarrow c, d) f_b^0, \quad (10)$$

where E_a is the energy of the a th particle and f_b^0 is the distribution function for b th particle. $W(a, b \rightarrow c, d)$ is the transition rate defined as,

$$W(a, b \rightarrow c, d) = \frac{2\pi^4 \delta(p_a + p_b - p_c - p_d)}{2E_a 2E_b 2E_c 2E_d} |\mathcal{M}|^2, \quad (11)$$

where $|\mathcal{M}|$ is the transition amplitude. In the centre-of-mass frame Eq. 10 can be written as,

$$\begin{aligned} \tau^{-1}(E_a) &= \sum_b \int \frac{d^3p_b}{(2\pi)^3} \sigma_{ab} \frac{\sqrt{s-4m^2}}{2E_a 2E_b} f_b^0 \\ &\equiv \sum_b \int \frac{d^3p_b}{(2\pi)^3} \sigma_{ab} v_{ab} f_b^0 \end{aligned} \quad (12)$$

where v_{ab} is the relative velocity and \sqrt{s} is the centre-of-mass energy and σ_{ab} is the total scattering cross-section in the process. $\tau(E_a)$ can be further written approximately as the averaged relaxation time ($\bar{\tau}$), which can be obtained from Eq. 12 by averaging over f_a^0 as,

$$\begin{aligned} \bar{\tau}_a^{-1} &= \frac{\int \frac{d^3p_a}{(2\pi)^3} \tau^{-1}(E_a) f_a^0}{\int \frac{d^3p_a}{(2\pi)^3} f_a^0} = \sum_b \frac{\int \frac{d^3p_a}{(2\pi)^3} \frac{d^3p_b}{(2\pi)^3} \sigma_{ab} v_{ab} f_a^0 f_b^0}{\int \frac{d^3p_a}{(2\pi)^3} f_a^0} \\ &= \sum_b n_b \langle \sigma_{ab} v_{ab} \rangle. \end{aligned} \quad (13)$$

n_b is the number density of b th particle. Here, all the calculations are done for one single hadronic species at a time. The thermal average of the scattering of the same species of particles with constant cross-section and zero baryon density can be estimated as,

$$\langle \sigma_{ab} v_{ab} \rangle = \frac{\sigma \int d^3p_a d^3p_b v_{ab} e_q^{-\frac{E_a}{T}} e_q^{-\frac{E_b}{T}}}{\int d^3p_a d^3p_b e_q^{-\frac{E_a}{T}} e_q^{-\frac{E_b}{T}}}, \quad (14)$$

where $e_q^{(x)}$ is the q -exponential which is defined as,

$$e_q^{(x)} = [1 + (q-1)x]^{q/(q-1)}. \quad (15)$$

The momentum space volume elements are be written as,

$$d^3p_a d^3p_b = 8\pi^2 p_a p_b dE_a dE_b d\cos\theta. \quad (16)$$

On solving, Eq. 14 can be further written as,

$$\langle \sigma_{ab} v_{ab} \rangle = \frac{\sigma \int 8\pi^2 p_a p_b dE_a dE_b d\cos\theta e_q^{-\frac{E_a}{T}} e_q^{-\frac{E_b}{T}} \sqrt{(E_a E_b - p_a p_b \cos\theta)^2 - (m_a m_b)^2}}{\int 8\pi^2 p_a p_b dE_a dE_b d\cos\theta e_q^{-\frac{E_a}{T}} e_q^{-\frac{E_b}{T}}}. \quad (17)$$

σ is the hadronic collision cross-section and a constant value of 11.3 mb [28, 29] is used in our calculations. The thermodynamical quantities in non-extensive statistics are calculated as [30],

$$n = g \int \frac{d^3p}{(2\pi)^3} \left[1 + (q-1) \frac{E-\mu}{T} \right]^{\frac{-q}{q-1}} \quad (18)$$

$$\epsilon = g \int \frac{d^3p}{(2\pi)^3} E \left[1 + (q-1) \frac{E-\mu}{T} \right]^{\frac{-q}{q-1}} \quad (19)$$

$$P = g \int \frac{d^3p}{(2\pi)^3} \frac{p^2}{3E} \left[1 + (q-1) \frac{E-\mu}{T} \right]^{\frac{-q}{q-1}}. \quad (20)$$

n , ϵ and P are the number density, energy density and pressure of hadrons, respectively. g is the particle degeneracy. Also, the non-extensive entropy density is given by,

$$s = \frac{\epsilon + P - \mu n}{T} \quad (21)$$

From thermodynamics, the isothermal compressibility (κ_T) is defined as [28, 31, 32],

$$\kappa_T = - \frac{1}{V} \frac{\partial V}{\partial P} \Big|_T, \quad (22)$$

where V , P and T are the volume, pressure and temperature of the system. In terms of multiplicity fluctuations and average number, isothermal compressibility can be defined as,

$$\left\langle (N - \langle N \rangle)^2 \right\rangle = \text{var}(N) = \frac{T \langle N \rangle^2}{V} \kappa_T, \quad (23)$$

where N is the particle multiplicity. From basic thermodynamic relation, we have

$$\left\langle (N - \langle N \rangle)^2 \right\rangle = VT \frac{\partial n}{\partial \mu}. \quad (24)$$

Thus, from Eq.23 and Eq.24 we derive the expression,

$$\kappa_T = \frac{\partial n / \partial \mu}{n^2} \quad (25)$$

where,

$$\frac{\partial n}{\partial \mu} = \frac{gq}{T} \int \frac{d^3p}{(2\pi)^3} \left[1 + (q-1) \frac{E-\mu}{T} \right]^{\frac{1-2q}{q-1}} \quad (26)$$

At the LHC energies, the baryochemical potential of the system is almost zero and for our studies, we use $\mu = 0$ in the calculations. To have a better understanding about a system, it is important to know the Equation of State (EoS) which is given by the speed of sound in that system. Speed of sound squared is given by [33],

$$c_s^2 = \left(\frac{\partial P}{\partial \epsilon} \right)_{s/n}. \quad (27)$$

It can be further written as,

$$c_s^2 = \frac{\left(\frac{\partial P}{\partial T} \right)}{\left(\frac{\partial \epsilon}{\partial T} \right)}. \quad (28)$$

III. RESULTS AND DISCUSSION

The experimental data from ALICE for pp collisions at $\sqrt{s} = 7$ TeV [34] are used for this analysis. We have taken T and q values from Tsallis distribution function by fitting the p_T -spectra of produced identified particles in a differential freeze-out scenario [35]. The Eq. 9 and Eq. 21 are used to calculate the shear viscosity to entropy density ratio. Fig. 1 shows the coefficient of shear viscosity to entropy density ratio as a function of charged particle multiplicity. The final state charged particle multiplicity density in pseudorapidity, $\langle dN_{ch}/d\eta \rangle$ is used as an event classifier in pp collisions at the LHC energies in order to classify low-multiplicity and high-multiplicity events – like centrality classification in heavy-ion collisions. This is required to study various observables as a function of final state system multiplicity. We observe that η/s decreases with increase in $\langle dN_{ch}/d\eta \rangle$ and at higher multiplicity it becomes the lowest approaching the KSS bound value. As pion being the lightest particle among the studied particles, the density of pions is higher in a system formed in high energy collisions. Thus, pions exhibit lowest η/s at low multiplicity and all the particles approach to a minimum η/s at high-multiplicity pp collisions. At high multiplicity beyond $\langle dN_{ch}/d\eta \rangle \gtrsim (10 - 20)$, η/s becomes almost constant and remain the same for all the particles. Thus the limit $\langle dN_{ch}/d\eta \rangle \gtrsim (10 - 20)$ can be interpreted as the threshold after which the system seems to be going through a change in its dynamics. This is one of the significant observations in the current work due to the fact that we use an experimentally motivated and thermodynamically consistent Tsallis distribution to explain the experimental data and the obtained results indicate a possible formation of QGP droplets in high multiplicity pp collisions.




FIG. 1: (Color online) η/s as a function of charged particle multiplicity for pp collisions at $\sqrt{s} = 7$ TeV for different charged particles.

In order to look deeper into the possibility of a near perfect fluid behaviour of the produced system, let us now study another important observable in this direction, *i.e.* the isothermal compressibility. For this, we take various identified particles produced in pp collisions and use Eq.25 to estimate isothermal compressibility in which T and q values are taken by fitting Tsallis distribution function to the p_T -spectra [34, 35]. Figure 2 shows the variation of κ_T of the identified particles as a function of charged particle multiplicity. We observe that κ_T decreases with the increase in $\langle dN_{ch}/d\eta \rangle$. This result is in agreement with our previous findings [9]. For lighter particles, the values of κ_T are lower and the values increase as the mass of the particles increase till a certain $\langle dN_{ch}/d\eta \rangle$. Pion being the lightest meson, has the lowest κ_T as compared to the others. This is because, κ_T is inversely proportional to the number density of the system. As pion number density is highest in a collision system, its isothermal compressibility is the lowest. However, as it is clearly seen, for higher charged particle multiplicity ($\langle dN_{ch}/d\eta \rangle \gtrsim 10 - 20$), the κ_T of all the hadrons converge together and show only a slight variation from each other. Beyond this threshold limit, a QGP-like medium formation is expected regardless of the collision systems [36]. It is worth mentioning here that a differential kinetic freeze-out scenario becomes a single freeze-out as has been discussed in heavy-ion collisions for a multiplicity threshold of $\langle dN_{ch}/d\eta \rangle \gtrsim 10 - 20$, which is one of the important findings of the present study.

The isothermal compressibility of water at room temperature is reported to be $6.62 \times 10^{42} \text{ fm}^3/\text{GeV}$ [37]. We have estimated the κ_T of the hadron gases to be around $10 \text{ fm}^3/\text{GeV}$ at high charged particle multiplicity. This value is still larger than the isothermal compressibility found for QGP-like medium in our previous work, which was found to be $\sim 0.3 \text{ fm}^3/\text{GeV}$ [9].




FIG. 2: (Color online) κ_T as a function of charged particle multiplicity for pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for different final state particles.




FIG. 3: (Color online) Speed of sound squared vs charged particle multiplicity for pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for different final state particles.

The speed of sound squared (c_s^2) being related to the equation of state of the system (EoS), is one of the important thermodynamic observables. In Fig. 3, we have plotted the c_s^2 as a function of charged particle multiplicity, which is calculated using Eq.28. We observe that c_s^2 increases with the increase in $\langle dN_{ch}/d\eta \rangle$. Here we see a mass ordering between the identified particles. As pion is the most abundant particle in the system, the c_s^2 for pion is the highest. These observations complement the previous estimations [38, 39]. The other particles follow the same trend accordingly. At low $\langle dN_{ch}/d\eta \rangle$, the speed of sound is lower. This is because the c_s^2 depends on the density of the system. For low charged particle multiplicity, the density is lower, and it increases slowly with increase in $\langle dN_{ch}/d\eta \rangle$. However, after a certain charged particle multiplicity, the density of the system doesn't change [40]. We observe that there is a transition in the behavior of the c_s^2 plot at around $\langle dN_{ch}/d\eta \rangle \gtrsim 10 - 20$, and almost becomes constant after this limit. This gives us a hint about a change in the dynamics of the systems after certain charged particle multiplicity. The squared value of the speed of sound in air at room temperature is 1.3×10^{-12} and in distilled water at room temperature it is around 2.5×10^{-11} [41]. With comparison to these, the c_s^2 of the hadron gases we found here is very high. This indicates the fact that, the hadron gas systems formed in high energy collisions are very dense mediums and at higher charged particle multiplicities, the value of c_s^2 tend towards $1/3$, which means the medium behaves like almost an ideal gas.

IV. SUMMARY

In summary,

1. We have estimated the shear viscosity to entropy

density ratio of the system produced in pp collision at $\sqrt{s} = 7$ TeV energy and observed that at higher charged particle multiplicity, η/s becomes the lowest, approaching the KSS bound. This helps us to conclude that at higher charged particle multiplicities, the system could be non-dissipative in nature.

2. In order to look deeper into the possibility of a near perfect fluid behaviour of the produced system, we have also estimated the isothermal compressibility of the hadron gases in pp collision system by considering differential freeze-out scenario. We observed that κ_T of the systems decreases with the increase in the charged particle multiplicity. This suggests that at higher charged particle multiplicity, the system is less compressible.
3. The shear viscosity to entropy density ratio and isothermal compressibility becoming independent of particle species around $\langle dN_{ch}/d\eta \rangle \gtrsim 10-20$ is an indication of a transition from differential to single kinetic freeze-out, as is observed in heavy-ion collisions. This observed threshold in the final state charged particle density is an important finding in view of the scaling observed in the LHC energies across different collision species.
4. The speed of sound squared for different particles are estimated. We see a range of values from 0.15 to 0.32 for different particles, with pion having the highest c_s^2 .
5. In all our findings, we have observed a limit of

$\langle dN_{ch}/d\eta \rangle \gtrsim 10 - 20$ after which the system appears to be going through some change in its dynamics. This limit of charged particle multiplicity may suggest a requirement for the possible formation of QGP droplets in high multiplicity pp collisions [42, 43].

6. Although in this paper, a theoretical estimation of various important thermodynamic quantities are made as a function of final state multiplicity in pp collisions in order to carry out a systematic study, the physical interpretation of the low multiplicity system has to be done with caution.

It would be interesting to go to higher multiplicity classes by extending the same analysis to look for possible criticality in the system, through the inclusion of higher collision energies and collision species (p -Pb, Pb-Pb). However, it should be noted that higher final state multiplicity will lead the system towards a Boltzmann-Gibbs statistical description. A smooth transition in terms of system dynamics and statistical mechanics description from hadronic to nuclear collision has been an issue always.

Acknowledgement

The authors acknowledge the financial supports from ALICE Project No. SR/MF/PS-01/2014-IITI(G) of Department of Science & Technology, Government of India. R. S. acknowledges the financial supports from DAE-BRNS Project No. 58/14/29/2019-BRNS.

-
- [1] R. Sahoo, AAPPs Bull. **29**, 16 (2019).
 - [2] J. Adam, *et al.* [ALICE Collaboration], Nature Phys. **13**, 535 (2017).
 - [3] D. Velicanu [CMS Collaboration], J. Phys. G **38**, 124051 (2011).
 - [4] J. Adams *et al.* [STAR Collaboration], Nucl. Phys. A **757**, 102 (2005).
 - [5] P. Kovtun, D. T. Son and A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005).
 - [6] P. Romatschke and U. Romatschke, Phys. Rev. Lett. **99**, 172301 (2007).
 - [7] T. Hirano and M. Gyulassy, Nucl. Phys. A **769**, 71 (2006).
 - [8] M. Mukherjee, S. Basu, A. Chatterjee, S. Chatterjee, S. P. Adhya, S. Thakur and T. K. Nayak, Phys. Lett. B **784**, 1 (2018).
 - [9] D. Sahu, S. Tripathy, R. Sahoo and S. K. Tiwari, arXiv:2001.01252 [hep-ph].
 - [10] B. Mohanty and J. e. Alam, Phys. Rev. C **68**, 064903 (2003).
 - [11] J. D. Bjorken, Phys. Rev. D **27**, 140 (1983).
 - [12] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. C **75**, 064901 (2007).
 - [13] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. C **83**, 064903 (2011).
 - [14] K. Aamodt *et al.* [ALICE Collaboration], Eur. Phys. J. C **71**, 1655 (2011).
 - [15] B. Abelev *et al.* [ALICE Collaboration], Phys. Lett. B **717**, 162 (2012).
 - [16] B. Abelev *et al.* [ALICE Collaboration], Phys. Lett. B **712**, 309 (2012).
 - [17] S. Chatrchyan *et al.* [CMS Collaboration], Eur. Phys. J. C **72**, 2164 (2012).
 - [18] Tsallis, C. J Stat. Phys. **52**, 479 (1988).
 - [19] J. Cleymans and D. Worku, J. Phys. G **39**, 025006 (2012).
 - [20] C. Tsallis and E. P. Borges, cond-mat/0301521 [cond-mat.stat-mech].
 - [21] T. Bhattacharyya, J. Cleymans, A. Khuntia, P. Preeek and R. Sahoo, Eur. Phys. J. A **52**, 30 (2016).
 - [22] J. Randrup, Phys. Rev. C **79**, 054911 (2009).
 - [23] L. Palhares, E. Fraga and T. Kodama, J. Phys. G **38**, 085101 (2011).
 - [24] V. Skokov and D. Voskresensky, JETP Lett. **90**, 223 (2009).
 - [25] V. Skokov and D. Voskresensky, Nucl. Phys. A **828**, 401 (2009).
 - [26] G. Wilk and Z. Wlodarczyk, Phys. Rev. Lett. **84**, 2770 (2000).
 - [27] T. Biro and E. Molnar, Phys. Rev. C **85**, 024905 (2012).

- [28] S. K. Tiwari, S. Tripathy, R. Sahoo and N. Kakati, Eur. Phys. J. C **78**, 938 (2018).
- [29] G. P. Kadam and H. Mishra, Phys. Rev. C **92**, 035203 (2015).
- [30] J. Cleymans and D. Worku, Eur. Phys. J. A **48**, 160 (2012).
- [31] K. Huang, Statistical Mechanics, 2nd Edition, John Wiley, New York (1987).
- [32] L.D. Landau and E.M. Lifshitz, Statistical Physics, Pergamon, Oxford (1980).
- [33] J. Cleymans and D. Worku, Mod. Phys. Lett. A **26**, 1197 (2011).
- [34] S. Acharya *et al.* [ALICE Collaboration], Phys. Rev. C **99**, 024906 (2019).
- [35] A. Khuntia, H. Sharma, S. Kumar Tiwari, R. Sahoo and J. Cleymans, Eur. Phys. J. A **55**, 3 (2019).
- [36] D. Sahu, S. Tripathy, G. S. Pradhan and R. Sahoo, Phys. Rev. C **101**, 014902 (2020).
- [37] Rana A. Fine, Frank J. Millero, J. Chem. Phys. **59**, 5529 (1973).
- [38] S. Deb, G. Sarwar, R. Sahoo and J. e. Alam, arXiv:1909.02837 [hep-ph].
- [39] A. Khuntia, P. Sahoo, P. Garg, R. Sahoo and J. Cleymans, Eur. Phys. J. A **52**, 292 (2016).
- [40] N. Sharma, J. Cleymans and B. Hippolyte, Adv. High Energy Phys. **2019**, 5367349 (2019).
- [41] Samuel J Ling, Sanny Jeff, and Moebs William, University Physics: Volume 1, (2016).
- [42] R. Campanini and G. Ferri, Phys. Lett. B **703**, 237 (2011).
- [43] R. Sahoo, D. Thakur, S. De and S. Dansana, Springer Proc. Phys. **234**, 173 (2019).