

Locating Dominating Sets in local tournaments

Thomas Bellitto*, Caroline Brosse†, Benjamin Lévêque‡, Aline Parreau§

September 8, 2021

Abstract

A dominating set in a directed graph is a set of vertices S such that all the vertices that do not belong to S have an in-neighbour in S . A locating set S is a set of vertices such that all the vertices that do not belong to S are characterized uniquely by the in-neighbours they have in S , *i.e.* for every two vertices u and v that are not in S , there exists a vertex $s \in S$ that dominates exactly one of them. The size of a smallest set of a directed graph D which is both locating and dominating is denoted by $\gamma^{LD}(D)$. Foucaud, Heydarshahi and Parreau proved that any twin-free digraph D satisfies $\gamma^{LD}(D) \leq \frac{4n}{5} + 1$ but conjectured that this bound can be lowered to $\frac{2n}{3}$. The conjecture is still open. They also proved that if D is a tournament, *i.e.* a directed graph where there is one arc between every pair of vertices, then $\gamma^{LD}(D) \leq \lceil \frac{n}{2} \rceil$.

The main result of this paper is the generalization of this bound to connected local tournaments, *i.e.* connected digraphs where the in- and out-neighbourhoods of every vertex induce a tournament. We also prove $\gamma^{LD}(D) \leq \frac{2n}{3}$ for all quasi-twin-free digraphs D that admit a supervising vertex (a vertex from which any vertex is reachable). This class of digraphs generalizes twin-free acyclic graphs, the most general class for which this bound was known.

1 Introduction

In this paper we consider loopless and finite digraphs. Our terminology is consistent with [2]. Especially, we refer the reader to this book for further information about the classes of digraphs we consider. The *order* of a digraph is its number of vertices. A digraph D is *simple* if there is at most one arc between two

*Faculty of Mathematics, Informatics and Mechanics, University of Warsaw, Poland. This author is supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme Grant Agreement 714704.

†Université Clermont Auvergne, Clermont Auvergne INP, CNRS, Mines Saint-Etienne, LIMOS, F-63000 Clermont-Ferrand, France. This author is supported by the French government IDEX-ISITE initiative 16-IDEX-0001 (CAP 20-25).

‡CNRS, Laboratoire G-SCOP, Grenoble, France. This author is partially supported by the French ANR Projects GATO (ANR-16-CE40-0009-01)

§LIRIS, Université Claude Bernard Lyon 1, CNRS, France

vertices x and y . A digraph D is called *connected*, if for every vertices x and y of D there exists a (non necessarily directed) path from x to y . A digraph D is called *strongly connected* (or *strong* for short), if for every vertices x and y of D , there exists a directed path from x to y and a directed path from y to x .

Let x, y be distinct vertices of a digraph D . If there is an arc from x to y , we say that x *dominates* y and denote it by $x \rightarrow y$. We say that y is an *out-neighbour* of x , and that x is an *in-neighbour* of y . The *open in-neighbourhood* and the *open out-neighbourhood* of a vertex v , denoted by $N^-(v)$ and $N^+(v)$ respectively, are the set of in-neighbours of v and the set of out-neighbours of v , respectively. A *source* is a vertex with no in-neighbours, and a *sink* is a vertex with no out-neighbours. We use the notations $d^-(v) = |N^-(v)|$ and $d^+(v) = |N^+(v)|$ to denote the in-degree and the out-degree of v . Further, the *closed in-neighbourhood* of v is $N^-[v] = N^-(v) \cup \{v\}$ and the *closed out-neighbourhood* of v is $N^+[v] = N^+(v) \cup \{v\}$. Two vertices are called *twins* if $N^-(x) = N^-(y)$ or if $N^-[x] = N^-[y]$, and *quasi twins* if $N^-(x) = N^-[y]$ or $N^-(y) = N^-[x]$.

A digraph is *twin-free* (resp. *quasi-twin-free*) if it contains no twins (resp. no twins nor quasi-twins). A *dominating set* S of a digraph D is a set of vertices such that any vertex not in S is dominated by a vertex in S . The smallest size of a dominating set of D is called its domination number, and is denoted by $\gamma(D)$. A *locating set* S of D is a subset of vertices such that for every pair of vertices x and y not in S , there exists a vertex s in S that dominates exactly one vertex among x and y . The smallest size of a locating set of D is denoted by $\gamma^L(D)$. A *locating-dominating set* of D is a set of vertices that is both locating and dominating. The minimum size of a locating-dominating set of D is called the *location-domination number*, denoted by $\gamma^{LD}(D)$. Note that at most one vertex of D is not dominated by a locating set of D , ensuring that:

$$\gamma^L(D) \leq \gamma^{LD}(D) \leq \gamma^L(D) + 1$$

Locating-dominating sets in digraphs have been introduced in [3, 7] and further studied in [5] where some upper bounds on the location-domination number of digraphs have been proved. If D contains at least one edge, then for any vertex x that is not isolated, the set $V(D) \setminus \{x\}$ is locating-dominating and thus, if D has order n , then $\gamma^{LD}(D) \leq n - 1$. This bound is tight and a complete characterization of the digraphs reaching it is given in [5]. However, all these digraphs contain many pairs of twins. The authors of [5] showed that any twin-free digraph D of order n satisfies $\gamma^{LD}(D) \leq \frac{4n}{5} + 1$. Note that similar questions were first considered in the non-oriented case for which it is conjectured that any twin-free graph has a locating-dominating set of size $\frac{n}{2}$ [4, 6].

The authors of [5] lowered the general upper bound $\frac{4n}{5} + 1$ for special cases. A digraph D is a *tournament* if there is exactly one arc between every pair of distinct vertices of D . In particular, it is proven in [5] that for a tournament D , the upper bound on the location-domination number can be lowered to $\gamma^{LD}(D) \leq \lceil \frac{n}{2} \rceil$. We extend this last result to a larger class of digraphs. A digraph D is a *local tournament* if it is simple and if the in-neighbourhood and

the out-neighbourhood of every vertex of D induce tournaments. The main result of the current paper is the following theorem.

Theorem 1. *A connected local tournament D of order n satisfies $\gamma^{LD}(D) \leq \lceil \frac{n}{2} \rceil$.*

Note that an edgeless graph of order n is a local tournament for which $\gamma^{LD}(D) = n - 1$. So if one removes the "connected" hypothesis in Theorem 1, then the conclusion does not hold.

In [5], the authors asked if the general upper bound $\frac{4n}{5} + 1$ for twin-free graphs can be lowered to $\frac{2n}{3}$, for which some tight constructions are known (see Figure 1 for the tight strongly connected example given in [5]). They proved that it is the case for quasi-twin-free acyclic digraphs. We extend these results by proving that if a quasi-twin-free digraph D contains a *supervising vertex*, that is a vertex from which there exists a directed path to all the other vertices of the graph, then $\gamma^{LD}(D) \leq \frac{2n}{3}$. In particular, this bound is true (and asymptotically tight) for quasi-twin-free strongly connected digraphs and quasi-twin-free local-in semi-complete digraphs (i.e. digraphs, not necessarily simple, where two in-neighbours of a common vertex are connected).




Figure 1: A strongly connected twin-free and quasi-twin-free digraph of order n with location-domination number $\frac{2(n-2)}{3}$, with a locating-dominating set in gray.

After giving some preliminary results on local tournaments in Section 2, we prove Theorem 1 in two steps in Sections 3 and 4. In Section 5, we prove the general upper bound for digraphs that contain a supervising vertex.

2 Preliminaries

In order to prove the upper bound on the location-domination number of local tournaments, we start by exposing some useful properties of local tournaments.

Lemma 2. *A connected local tournament is twin-free.*

Proof. Consider a connected local tournament D and suppose by contradiction that there exist two distinct vertices x, y of D such that x, y are twins.

Suppose first that x, y have some in-neighbours, then they have a common in-neighbour z . Since x, y are both in the out-neighbourhood of z , there is an arc between them, a contradiction.

Suppose now that x, y have no in-neighbours. Since D is connected, we can consider a shortest (non necessarily directed) path $P = v_1, \dots, v_k$ between x, y , such that $v_1 = x$ and $v_k = y$. Note that $k \geq 3$. By assumption, we have $v_1 \rightarrow v_2$ and $v_k \rightarrow v_{k-1}$. Thus there exists $1 < i < k$ such that $v_{i-1} \rightarrow v_i$ and $v_{i+1} \rightarrow v_i$. So v_{i-1} and v_{i+1} are both in the in-neighbourhood of v_i and thus there is an arc between them, contradicting the fact that P is a shortest path. \square

The structure of local tournaments has been studied in [1]. In particular, the authors introduce the *round decomposition*, which will be crucial in our proofs. Therefore, let us have a closer look at the construction of such a decomposition.

We call a digraph D on r vertices *round* if its vertices can be labelled v_1, \dots, v_r in such a way that $N^-(v_i) = \{v_{i-d^-(v_i)}, \dots, v_{i-1}\}$ and $N^+(v_i) = \{v_{i+1}, \dots, v_{i+d^+(v_i)}\}$ (indices are understood modulo r). One can easily check that a simple round digraph is a local tournament.

Let R be a digraph on r vertices and L_1, \dots, L_r be a collection of r digraphs. Then $R[L_1, \dots, L_r]$ is the digraph obtained from R by replacing each vertex v_i of R with L_i , and adding an arc from every vertex of L_i to every vertex of L_j if and only if $v_i \rightarrow v_j$ is an arc of R . A local tournament D is said to be *roundable* if it can be written as $R[T_1, \dots, T_r]$ where R is a round simple digraph on $r \geq 2$ vertices, and T_1, \dots, T_r are (non empty) strong tournaments. The decomposition $R[T_1, \dots, T_r]$ is called a *round decomposition* of D .

Given a round decomposition $R[T_1, \dots, T_r]$ of a roundable local tournament D , if there is an arc between a vertex in T_i and a vertex in T_j , then all the arcs between T_i and T_j are present in D , and we write $T_i \Rightarrow T_j$. If $T_i \Rightarrow T_j$ and $i < j$, then for all $k \in \{i+1, \dots, j-1\}$, we have $T_i \Rightarrow T_k$ and $T_k \Rightarrow T_j$. If $T_i \Rightarrow T_j$ and $j < i$, then for all $k \in \{1, \dots, j-1\} \cup \{i+1, \dots, r\}$, we have $T_i \Rightarrow T_k$ and $T_k \Rightarrow T_j$.

Suppose that D is a roundable connected local tournament that is not strongly connected. Then, as proved in [1], there is a unique round decomposition $R[T_1, \dots, T_r]$ of D such that $T_i \Rightarrow T_j$ only happens if $i < j$. We often consider this particular round decomposition in this case, that we call the *canonical* round decomposition of D . When D is a roundable strong local tournament, there is a unique round decomposition up to cyclic permutation.

3 Roundable local tournaments

This section is devoted to the study of roundable connected local tournaments. Consider a roundable connected local tournament D and a round decomposition $R[T_1, \dots, T_r]$ of D that is canonical if D is not strong.

Lemma 3. *For $1 \leq i < r$, we have $T_i \Rightarrow T_{i+1}$.*

Proof. Suppose by contradiction that there exists $1 \leq i < r$ such that we don't have $T_i \Rightarrow T_{i+1}$. Then there is no arc $v_i \rightarrow v_{i+1}$ in R and so by definition of round, we have $d^+(v_i) = 0$. For $1 \leq k \leq i < \ell \leq r$, there is no arc $v_k \rightarrow v_\ell$ as such an arc implies an arc $v_i \rightarrow v_\ell$. Since v_i is a sink, the digraph D is not strong and the decomposition is canonical. So there is no edge between $V(T_1) \cup \dots \cup V(T_i)$ and $V(T_{i+1}) \cup \dots \cup V(T_r)$, contradicting the fact that D is connected. \square

We define the sequence of integers $(i_k)_{0 \leq k \leq t}$ as follows (see Figure 2). Let $i_0 = 0$. If i_k is defined and $i_k < r$, then we define i_{k+1} as the greatest integer $i_k + 1 < i_{k+1} \leq r$ such that $T_{i_{k+1}} \Rightarrow T_{i_k}$ (note that this integer exists by Lemma 3). This procedure stops when some i_t is defined equal to r . For $1 \leq k \leq t$, let $D_k = D[T_{i_{k-1}+1} \cup \dots \cup T_{i_k}]$. Note that D_k is a tournament for all k .




Figure 2: Illustration of the definition of the subgraphs D_k . The dashed arc means that there is no arc between the subgraphs, whereas thick arcs means that there are all the arcs between the two subgraphs.

Step by step, from $k = t$ to $k = 1$, we define a set S of vertices of D that will be our candidate for a locating-dominating set of D . We consider the following three different cases:

- *Case 1: $k = t$, or $k < t$ and $S \cap V(D_{k+1})$ is assumed to be a dominating set of D_{k+1} .*

If $|V(D_k)|$ is even, we add to S a minimum locating-dominating set of D_k , otherwise we add to S a minimum locating set of D_k .

- *Case 2: $k < t$, the set $S \cap V(D_{k+1})$ is not assumed to be a dominating set of D_{k+1} , and $|V(T_{i_k})| = 1$.*

Let v be the unique vertex of T_{i_k} and $D'_k = D_k \setminus \{v\}$. We add v to S . Moreover we add to S a minimum locating-dominating set of D'_k when

$|V(D'_k)|$ is even, or a minimum locating set of D'_k when $|V(D'_k)|$ is odd. (Note that the set of D_k that is added to S , including v , is a dominating set of D_k when $|V(D'_k)|$ is odd.)

- *Case 3: $k < t$, the set $S \cap V(D_{k+1})$ is not assumed to be a dominating set of D_{k+1} , and $|V(T_{i_k})| > 1$.*

We add to S a minimum locating-dominating set of D_k .

Note that in all the three cases we add to S a set of vertices of D_k that is a locating set of D_k , even in Case 2 since a locating set of D'_k plus v forms a locating set of D_k .

In the next lemmas we prove some properties of the set S .

Lemma 4. *For $1 \leq i \leq r$, the set $S \cap V(T_i)$ is a locating set of T_i , in particular if T_i has at least 2 vertices, then $S \cap V(T_i) \neq \emptyset$.*

Proof. Let $1 \leq i \leq r$. If $V(T_i) \setminus S$ has size 1, then $S \cap V(T_i)$ is a locating set of T_i (even if empty) and the conclusion holds. So we can assume that $V(T_i) \setminus S$ has size at least 2. Let x, y be two distinct vertices of $V(T_i) \setminus S$. Since S is a locating set of D_k , there exists a vertex s in $S \cap V(D_k)$ that dominates exactly one vertex among x and y . Since $R[T_1, \dots, T_r]$ is a round decomposition, all the vertices of D_k not in T_i dominates either all or none of the vertices of T_i . So $s \in V(T_i)$. Therefore $S \cap V(T_i)$ is a locating set of T_i and $S \cap V(T_i) \neq \emptyset$. \square

Lemma 5. *For $1 \leq k \leq t$, if $S \cap V(D_k)$ is a dominating set of D_k , then $S \cap V(T_{i_{k-1}+1}) \neq \emptyset$. Otherwise, there is exactly one vertex of D_k that is not dominated by $S \cap V(D_k)$ and this vertex is in $V(T_{i_{k-1}+1})$.*

Proof. Let $1 \leq k \leq t$. By definition of i_k , all the vertices of $T_{i_{k-1}+1}$ dominates $D_k \setminus V(T_{i_{k-1}+1})$, so there is no arc from $V(D_k) \setminus V(T_{i_{k-1}+1})$ to $V(T_{i_{k-1}+1})$ (since R is simple). Thus, if $S \cap V(D_k)$ is a dominating set of D_k , then S must contains a vertex of $V(T_{i_{k-1}+1})$.

Suppose now that there exists a vertex of D_k that is not dominated by $S \cap V(D_k)$. Since $S \cap V(D_k)$ is a locating set of D_k , there is at most one vertex that is not dominated by $S \cap V(D_k)$, and thus exactly one such vertex v . If there exists a vertex u in $S \cap V(T_{i_{k-1}+1})$, then u dominates $D_k \setminus V(T_{i_{k-1}+1})$, so $v \in V(T_{i_{k-1}+1})$. If there is no vertices in $S \cap V(T_{i_{k-1}+1})$, then the vertices of $T_{i_{k-1}+1}$ are not dominated by $S \cap V(D_k)$, so $V(T_{i_{k-1}+1}) = \{v\}$. \square

Lemma 6. *There is at most one vertex of D that is not dominated by S , and if it exists, it is a vertex of T_1 .*

Proof. Assume by contradiction that there exists a vertex x of D that is not dominated by S and x is not in D_1 . Let $2 \leq k \leq t$ such that $x \in D_k$. By Lemma 5, vertex x is in $T_{i_{k-1}+1}$. If $T_{i_{k-1}}$ contains a unique vertex v , then, at step $k-1$ of the construction of S we are in Case 2 and v is added to S . If $T_{i_{k-1}}$ contains at least two vertices, then, by Lemma 4, it contains a vertex of S . In both cases, $S \cap V(T_{i_{k-1}}) \neq \emptyset$. By Lemma 3, we have $T_{i_{k-1}} \Rightarrow T_{i_{k-1}+1}$, so x is dominated by S , a contradiction.

Thus all the vertices that are not dominated by S are in D_1 . By Lemma 5, there is at most one such vertex and if it exists it is a vertex of T_1 . \square

Lemma 7. *Suppose that there exists a pair $\{x, y\}$ of vertices of $V(D) \setminus S$ not located by S . Then, D is strongly connected, and one of x, y is in T_1 and is the only vertex of D_1 not dominated by $S \cap V(D_1)$.*

Proof. Let $1 \leq k \leq t$, $1 \leq \ell \leq t$ such that $x \in V(D_k)$ and $y \in V(D_\ell)$. Let $1 \leq i \leq r$, $1 \leq j \leq r$ such that $x \in V(T_i)$ and $y \in V(T_j)$. Recall that for all $1 \leq m \leq t$, $S \cap V(D_m)$ is a locating set of D_m , so $k \neq \ell$ and $i \neq j$. Since a local tournament is simple, we can assume, w.l.o.g., that y does not dominate x .

Suppose by contradiction that there exists $s \in S \cap V(D_\ell)$ that dominates y . Let m be such that $s \in T_m$. Since s, y are vertices of D_ℓ , we have $1 \leq m \leq j \leq r$. Since $k \neq \ell$, we know that T_m, T_j and T_i appear in this order along the cyclic order T_1, \dots, T_r (with maybe $m = j$). Since y does not dominate x , the vertex s does not dominate x by definition of round. Thus s separates x and y , a contradiction. So $S \cap V(D_\ell)$ does not dominate y and by Lemma 5, y is the only vertex of D_ℓ that is not dominated by $S \cap V(D_\ell)$ and $y \in T_{i_{\ell-1}+1}$.

Suppose by contradiction that $\ell > 1$. We use an argument that is similar to the proof of Lemma 6. If $T_{i_{\ell-1}}$ contains a unique vertex v , then, at step $\ell - 1$ of the construction of S we are in Case 2 and v is added to S . If $T_{i_{\ell-1}}$ contains at least two vertices, then, by Lemma 4, it contains a vertex of S . In both cases, $S \cap V(T_{i_{\ell-1}}) \neq \emptyset$. Let $s' \in S \cap V(T_{i_{\ell-1}})$. By Lemma 3, we have $T_{i_{\ell-1}} \Rightarrow T_{i_{\ell-1}+1}$, so s' dominates y . Since x and y are not located by S , vertex s' dominates x . Hence, $T_{i_{\ell-1}}, T_{i_{\ell-1}+1}, T_i$ appear in this order along the cyclic order T_1, \dots, T_r (with maybe $i_{\ell-1} = i$). Since x is dominated by s' and not by y , we have $i_{\ell-1} = i$. So $k = \ell - 1$. Thus both x and s' are in $V(T_{i_k})$ and y is in $V(T_{i_{k+1}})$. At step k of the construction of S , we are in Case 3, and so $S \cap V(D_k)$ is a dominating set of D_k . By Lemma 5, the set S contains a vertex of $V(T_{i_{k-1}+1})$. By definition of i_k , this vertex dominates $x \in V(T_{i_k})$ but not $y \in V(T_{i_{k+1}})$, a contradiction. So $\ell = 1$, $y \in T_1$ and y is the only vertex of D_1 that is not dominated by $S \cap V(D_1)$.

Suppose by contradiction that S is not strongly connected, then the round decomposition is canonical and there is no vertex of $D \setminus V(D_1)$ that can dominate y . So y is not dominated by S and by Lemma 6, vertex x is dominated by S , a contradiction. So S is strongly connected. \square

We now need to slightly modify S in a particular case. If $S \cap V(D_1)$ is not a dominating set of D_1 , then by Lemma 5, let $z \in T_1$ be the unique vertex of D_1 not dominated by $S \cap V(D_1)$. In this case, let $S^+ = S \cup \{z\}$, otherwise, let $S^+ = S$. We are now able to prove the locating-dominating property of S^+ .

Lemma 8. *The set S^+ is a locating-dominating set of D .*

Proof. Suppose first that $S \cap V(D_1)$ is a dominating set of D_1 . Then S^+ is a dominating set of D by Lemma 6 and a locating set of D by Lemma 7. Suppose now that $S \cap D_1$ is not a dominating set. Recall that z is the only vertex of D_1 not dominated by $S \cap D_1$. By Lemma 6, there is a unique vertex that is

not dominated by S and this vertex is in T_1 , so this vertex is z . So S^+ is a dominating set of D . By Lemma 7, vertex z is in all the pairs of vertices of $V(D) \setminus S$ not located by S . Thus S^+ is locating. \square

By Lemma 8, we have defined a locating-dominating set of D . We now have to bound its size. For that purpose we use the following theorem from [5]:

Theorem 9 ([5]). *A tournament D of order n satisfies $\gamma^{LD}(D) \leq \lceil \frac{n}{2} \rceil$ and $\gamma^L(D) \leq \lfloor \frac{n}{2} \rfloor$.*

We use Theorem 9 to first bound the size of S and then of S^+ . At each step k of the construction of S , since D_k is a tournament, the theorem gives an upper bound on the size of the set that is added to S :

Lemma 10. *For $1 \leq k \leq t$, the set of vertices of D_k that is added to S at step k of the construction of S has size at most $\lceil |V(D_k)|/2 \rceil$ when it is assumed to be a locating-dominating set of D_k and size at most $\lfloor |V(D_k)|/2 \rfloor$ when it is assumed to be a locating set of D_k .*

Proof. Let $1 \leq k \leq t$. The lemma is clear by Theorem 9 if we are in Cases 1 or 3 at step k of the construction of S . Consider now that we are in Case 2. If $|V(D'_k)|$ is even (i.e. $|V(D_k)|$ is odd), then $S \cap V(D_k)$ is composed of a minimum locating-dominating set of D'_k plus the unique vertex of T_{i_k} . Thus, by Theorem 9, $S \cap V(D_k)$ is a locating-dominating set of D_k of size at most $\lceil (|V(D_k)| - 1)/2 \rceil + 1 = \lceil |V(D_k)|/2 \rceil$. If $|V(D'_k)|$ is odd (i.e. $|V(D_k)|$ is even), then $S \cap V(D_k)$ is composed of a minimum locating set of D'_k plus the unique vertex of T_{i_k} . Thus, similarly, $S \cap V(D_k)$ is a locating set of D_k of size at most $\lfloor (|V(D_k)| - 1)/2 \rfloor + 1 = \lfloor |V(D_k)|/2 \rfloor$. \square

For every $1 \leq k \leq t$, let $n_k = |V(D_k)| + \dots + |V(D_t)|$. In the following lemma, we bound the size of a minimum locating-dominating set of a subgraph of D .

Lemma 11. *For $1 \leq k \leq t$, the size of $S \cap (V(D_k) \cup \dots \cup V(D_t))$ is at most $\lfloor \frac{n_k}{2} \rfloor$.*

Proof. We prove the lemma by induction on k from t down to 1.

At step t of the construction of S , we are in Case 1. If $|V(D_t)|$ is even, then $S \cap V(D_t)$ is a minimum locating-dominating set of D_t of size at most $\lceil |V(D_t)|/2 \rceil = \lfloor n_t/2 \rfloor$ by Theorem 9. If $|V(D_t)|$ is odd, then $S \cap V(D_t)$ is a minimum locating set of D_t of size at most $\lfloor n_t/2 \rfloor$ by Theorem 9.

Let us now fix $1 \leq k < t$. We assume that the lemma holds for all j with $k+1 \leq j \leq t$ and we prove that it holds for k . If $V(D_k)$ has even size or $S \cap V(D_k)$ is not assumed to dominate D_k , then by Lemma 10, $|S \cap V(D_k)| \leq \lfloor |V(D_k)|/2 \rfloor$ and it follows that the lemma holds for k . So we can now suppose that $V(D_k)$ has odd size and $S \cap V(D_k)$ is assumed to dominate D_k . Thus, we are in Case 2 or 3 at step k and $S \cap V(D_{k+1})$ is not assumed to dominate D_{k+1} . Then, at step $k+1$, we are either in Case 1 with $V(D_{k+1})$ has odd size, or in Case 2 with $V(D_{k+1})$ has even size, $k+1 < t$ and $S \cap V(D_{k+2})$ is not assumed to

dominate D_{k+2} . In the second case, again it means that at step $k+2$ we are in the same situation, i.e. we are either in Case 1 with $V(D_{k+2})$ has odd size, or in Case 2 with $V(D_{k+2})$ has even size, $k+2 < t$ and $S \cap V(D_{k+3})$ is not assumed to dominate D_{k+3} . We can repeat this argument until we actually encounter a step ℓ of the construction of S such that we are in Case 1 at step ℓ with $V(D_\ell)$ has odd size (we assume ℓ is the first such step from $k+1$ towards t). Note that for $k+1 \leq j \leq \ell$, $S \cap V(D_j)$ is not assumed to dominate D_j . To summarize, we have:

- $1 \leq k < \ell \leq t$
- $V(D_k)$ and $V(D_\ell)$ have odd sizes
- $V(D_j)$ has even size for $k+1 \leq j \leq \ell-1$,
- $S \cap V(D_k)$ is assumed to dominate D_k
- $S \cap V(D_j)$ is not assumed to dominate D_j for $k+1 \leq j \leq \ell$,

Thus, by Theorem 9, we obtain the following:

$$\begin{aligned} & |S \cap (V(D_k) \cup \dots \cup V(D_\ell))| \\ & \leq (|V(D_k)| + 1)/2 + |V(D_{k+1}) \cup \dots \cup V(D_{\ell-1})|/2 + (|V(D_\ell)| - 1)/2 \\ & \leq \lfloor |V(D_k) \cup \dots \cup V(D_\ell)|/2 \rfloor \end{aligned}$$

If $\ell = t$, then the lemma holds for k and we are done. If $\ell < t$, then we use the induction hypothesis that the lemma holds for $\ell+1$ to conclude. \square

A corollary of Lemma 11 is that the size of S is at most $\lfloor n/2 \rfloor$. We are now able to bound $|S^+|$.

Lemma 12. *The size of S^+ is at most $\lceil n/2 \rceil$.*

Proof. By Lemma 11, we have $|S| \leq \lfloor n/2 \rfloor$. Thus if $S = S^+$, then the lemma holds. Thus we can assume that $S \neq S^+$, i.e. $S \cap V(D_1)$ is not a dominating set of D_1 and $S^+ = S \cup \{z\}$. If n is odd, then $|S^+| = |S| + 1 \leq \lfloor n/2 \rfloor + 1 = \lceil n/2 \rceil$ and we are done. So we can assume that n is even.

By Lemma 10, the size of $S \cap V(D_1)$ is at most $\lfloor |V(D_1)|/2 \rfloor$

By construction of S , at step 1, we are either in Case 1 with $|V(D_1)|$ is odd, or in Case 2 with $|V(D_1)|$ is even, $1 < t$ and $S \cap V(D_2)$ is not assumed to dominate D_2 . We consider this two different cases below.

Consider first that we are in Case 1. Since $|V(D_1)|$ is odd and n is even, we have $1 < t$ and $n_2 = n - |V(D_1)|$ is odd. By Lemma 11, the size of $S \cap (V(D_2) \cup \dots \cup V(D_t))$ is at most $\lfloor n_2/2 \rfloor = (n_2 - 1)/2$. Moreover $S \cap V(D_1)$ has size at most $\lfloor |V(D_1)|/2 \rfloor = (|V(D_1)| - 1)/2$. So in total S has size at most $n/2 - 1$. Therefore S^+ has size at most $n/2$ and we are done.

Consider now that we are in Case 2 of the construction at step 1. As in the proof of Claim 11, let D_ℓ be the first step from 2 towards t such that we are in Case 1 at step ℓ of the construction of S with $V(D_\ell)$ has odd size. Then, we have:

- $2 \leq \ell \leq t$
- $V(D_j)$ has even size for $1 \leq j \leq \ell - 1$,
- $V(D_\ell)$ has odd size
- $S \cap V(D_j)$ is not assumed to dominate D_j for $1 \leq j \leq \ell$,

Thus, by Theorem 9, we obtain the following:

$$|S \cap (V(D_1) \cup \dots \cup V(D_\ell))| \leq |V(D_1) \cup \dots \cup V(D_{\ell-1})|/2 + (|V(D_\ell)| - 1)/2$$

Since n is even and $|V(D_1) \cup \dots \cup V(D_\ell)|$ is odd, we have $\ell < t$ and $n_{\ell+1}$ is odd. By Lemma 11, the size of $S \cap (V(D_{\ell+1}) \cup \dots \cup V(D_t))$ is at most $\lfloor (n_{\ell+1})/2 \rfloor = (n_{\ell+1} - 1)/2$. So again S has size at most $n/2 - 1$ and S^+ has size at most $n/2$. \square

By combining the results of this section we obtain the following:

Lemma 13. *The set S^+ is a locating dominating set of D of size at most $\lceil n/2 \rceil$. Moreover, if D is not strongly connected, then S is a locating set of D of size at most $\lfloor n/2 \rfloor$.*

Proof. The first part is a consequence of Lemmas 8 and 12. The second part is a consequence of Lemmas 7 and 11. \square

As a corollary, this proves Theorem 1 for connected roundable local tournaments:

Lemma 14. *A connected roundable local tournament D of order n satisfies $\gamma^{LD}(D) \leq \lceil \frac{n}{2} \rceil$. Moreover if D is not strongly connected, then $\gamma^L(D) \leq \lfloor \frac{n}{2} \rfloor$.*

4 Non-roundable local tournaments

In the previous section, we proved the upper bound $\gamma^{LD}(D) \leq \lceil n/2 \rceil$ when D is a connected roundable local tournament. By Theorem 9, this result is also true for tournaments. In order to prove Theorem 1, we can now restrict ourselves to connected local tournaments which are not tournaments and not roundable.

Consider a connected local tournament D that is not a tournament and not roundable. By [1, Corollary 3.2], every connected local tournament that is not strong is roundable, so D is strongly connected. Then, by [1, Corollary 3.2, Lemmas 3.4 and 3.5], there exists a set of vertices X of $V(D)$ such that:

- $D \setminus X$ is not strongly connected and X is minimal for this property
- $D \setminus X$ is a connected local tournament that is not a tournament
- $D[X]$ is a tournament




Figure 3: Decomposition of non-roundable local tournaments

- Let $R[T_1, \dots, T_r]$ be the canonical round decomposition of $D \setminus X$, then $r \geq 3$ and there are all the arcs from $V(T_r)$ to X , and from X to $V(T_1)$.

Let $Y = V(D) \setminus X$ and $Z = Y \setminus V(T_r)$. See Figure 3 for an illustration of the decomposition of D . Let $D_1 = D[Y]$ and $D_2 = D[Z]$. For $i \in \{1, 2\}$, let n_i be the order of D_i . Note that since $r \geq 3$, the digraph D_2 is also a connected local tournament that is not strong with canonical round decomposition $R[T_1, \dots, T_{r-1}]$. So we can apply on D_1 and D_2 the method and results of Section 3. For $i \in \{1, 2\}$, let S_i be the set that is defined on D_i exactly as S is defined on D in Section 3. By Lemma 13, the set S_i is a locating set of D_i of size $\lfloor n_i/2 \rfloor$. We will moreover need some particular properties obtained in Section 3, namely Lemmas 4 and 6.

Using these definitions of S_1 and S_2 we are now able to define a set of vertices S of D that will be our candidate for a locating-dominating set of D . We consider the following four cases (see Figure 4 for an illustration):

- *Case 1:* $|V(T_r)| = 1$ and $|X| = 1$
Let S be the union of $V(T_r)$ and a minimum locating-dominating set of D_2
- *Case 2:* $|V(T_r)| = 1$ and $|X| > 1$
Let S be the union of $V(T_r)$, S_2 , and a minimum locating set of $D[X]$.
- *Case 3:* $|V(T_r)| > 1$ and $|X| = 1$
Let S be the union of S_1 and X .
- *Case 4:* $|V(T_r)| > 1$ and $|X| > 1$
Let S be the union of S_1 and a minimum locating-dominating set of $D[X]$.

Lemma 15. *The set S is a locating-dominating set of D .*

Proof. (Case 1) Let x be the only vertex of X and t be the only vertex of T_r . The set S contains a locating-dominating set of D_2 so all the vertices of $Y \setminus S$ are dominated by S . Moreover x is dominated by t . So the set S is a dominating set of D .




Figure 4: Construction of a locating-dominating sets in a non-roundable local tournament. Gray vertices are in the locating-dominating sets.

Every pair of vertices of $Y \setminus S$ is separated by S . Since the round decomposition is canonical, there is no arc from t to Z , so t separates x from any vertex of $Y \setminus S$. Hence S is a locating set of D .

(Case 2) As previously, let t be the only vertex of T_r . By Lemma 6 applied on S_2 , there is at most one vertex of Z that is not dominated by S_2 , and if it exists, it is a vertex of T_1 . However, by assumption, $|X| > 1$, so a locating set of $D[X]$ contains at least one vertex of X . Thus S contains at least one vertex s of X . Since there are all the arcs from X to $V(T_1)$, all the vertices of T_1 are dominated by s . All the vertices of X are dominated by t . So S is a dominating set of D .

Every pair of vertices of $X \setminus S$ and every pair of vertices of $Z \setminus S$ are separated by S . There are all the arcs from t to X and no arc from t to $Y \setminus \{t\}$. Hence, t separates the vertices of X from the vertices of Z , and S is a locating set of D .

(Case 3) Let x be the only vertex of X . By Lemma 6 applied on S_1 , there is at most one vertex of Y that is not dominated by S_1 , and if it exists, it is a vertex of T_1 . Since $x \in S$ and there are all the arcs between x and T_1 , then every vertex of T_1 is dominated by x . Hence S is a dominating set of D .

By definition of S , all the pairs of vertices of $Y \setminus S$ are separated by S and $x \in S$. So S is a locating set of D .

(Case 4) Note that, since S contains a locating set of X and $|X| > 1$, the set S contains at least one vertex of X . Then the proof that S is a dominating set of D is exactly the same as in Case 3.

By definition of S , every pair of vertices of $X \setminus S$ and every pair of vertices of $Y \setminus S$ are separated by S . Now consider $x \in X \setminus S$ and $y \in Y \setminus S$ and let us show that x and y are separated by S . We consider the following two cases:

- *Case A* : $y \in T_r$

Since S contains a dominating set of $D[X]$, there is a vertex x' of $X \cap S$ that dominates x in D . Since there are all the arcs from $V(T_s)$ to X , there is an arc from y to x' . Since we are considering a simple digraph, there is no arc from x' to y , so x' separates x and y .

- *Case B* : $y \notin T_r$

By Lemma 4 applied on S_1 , there exists a vertex y' in $S \cap T_r$. Since the round decomposition is canonical, there is no arc from y' to y . Plus, there are all the arcs from $V(T_r)$ to X , and there is an arc from y' to x . Hence y' separates x and y .

Therefore S is a locating-dominating set of D . □

Lemma 16. *The size of S is at most $\lceil \frac{n}{2} \rceil$.*

Proof. (*Case 1*) By Lemma 14, a minimum locating-dominating set of D_2 has size at most $\lceil \frac{n_2}{2} \rceil$. So S has size at most $\lceil \frac{n_2}{2} \rceil + 1 = \lceil \frac{n-2}{2} \rceil + 1 = \lceil \frac{n}{2} \rceil$.

(*Case 2*) By Lemma 13, the size of S_2 is at most $\lfloor \frac{n_2}{2} \rfloor = \lfloor \frac{n-1-|X|}{2} \rfloor$. By Theorem 9, a minimum locating set of $D[X]$ has size at most $\lfloor \frac{|X|}{2} \rfloor$. So S has size at most $\lfloor \frac{n-1-|X|}{2} \rfloor + \lfloor \frac{|X|}{2} \rfloor + 1 \leq \lceil \frac{n}{2} \rceil$ (check all the parity cases for the last inequality).

(*Case 3*) By Lemma 13, the size of S_1 is at most $\lfloor \frac{n_1}{2} \rfloor$. So S has size at most $\lfloor \frac{n-1}{2} \rfloor + 1 \leq \lceil \frac{n}{2} \rceil$.

(*Case 4*) By Lemma 13, the size of S_1 is at most $\lfloor \frac{n_1}{2} \rfloor$. By Theorem 9, a minimum locating-dominating set of $D[X]$ has size at most $\lceil \frac{|X|}{2} \rceil$. So S has size at most $\lfloor \frac{n-|X|}{2} \rfloor + \lceil \frac{|X|}{2} \rceil \leq \lceil \frac{n}{2} \rceil$ (check all the parity cases for the last inequality). □

By combining the results of this section we obtain the following:

Lemma 17. *A connected local tournament D of order n that is not roundable satisfies $\gamma^{LD}(D) \leq \lceil \frac{n}{2} \rceil$.*

Proof. By Lemmas 15 and 16, the set S is a locating dominating set of D of size at most $\lceil n/2 \rceil$. □

Theorem 1 is a direct consequence of Lemmas 14 and 17.

5 Supervising vertex

From now on, we consider digraphs that are not necessarily simple. In a digraph D , a *supervising* vertex is a vertex s of D such that, for any vertex v , there exists a directed path from s to v . In this section we prove the following theorem.

Theorem 18. *Let D be a twin-free digraph on n vertices containing a supervising vertex, then $\gamma^{LD}(D) \leq \frac{3n}{4}$. Moreover, if D is quasi-twin-free, then $\gamma^{LD}(D) \leq \frac{2n}{3}$.*

To prove Theorem 18, we will adapt the method used in [4] and [5] to prove general upper bounds on γ^{LD} . Let S be a set of vertices of a digraph D . The S -partition of D , denoted \mathcal{P}_S , is the partition of $V(D) \setminus S$ where two vertices are in the same part if and only if they have the same set of in-neighbours in S . We have the following lemma.

Lemma 19. *Let D be a twin-free digraph on n vertices and S a dominating set of D such that $|\mathcal{P}_S| \geq |S| - 1$. Then, $\gamma^{LD}(D) \leq \frac{3n}{4}$. Moreover, if D is quasi-twin-free, then $\gamma^{LD}(D) \leq \frac{2n}{3}$.*

This result is proved in [5] when $|\mathcal{P}_S| \geq |S|$ (Theorem 8 for $x = 1$). The proof can be adapted if we only have $|\mathcal{P}_S| \geq |S| - 1$.

Proof. Let $\mathcal{P}_S = P_1 \cup \dots \cup P_{n_1} \cup Q_1 \cup \dots \cup Q_{n_2}$, where P_1, \dots, P_{n_1} are the parts of size 1 and Q_1, \dots, Q_{n_2} are the parts of size at least 2.

If $n_2 = 0$, then S is a locating-dominating set of D . Since all the parts of \mathcal{P}_S have size 1, $|V(D)| = |S| + |\mathcal{P}_S| \geq 2|S| - 1$. Thus $|S| \leq \frac{n+1}{2}$ and we are done. Thus in the following we assume that $n_2 > 0$.

We assume that S is maximal with the property that \mathcal{P}_S has at least $|S| - 1$ parts (this is ensured by adding vertices to S while this property holds).

Now, let $X_1 = S \cup P_1 \cup \dots \cup P_{n_1}$. We have the following property:

Claim 20. *Two vertices in $V(D) \setminus X_1$ are located by X_1 , unless they form a pair of quasi-twins.*

Proof of claim. If two vertices are in different parts of \mathcal{P}_S , they are located by some vertices in S . Thus, by contradiction, let q_1 and q_2 be two vertices of $V(D) \setminus X_1$ belonging to some part Q_i of \mathcal{P}_S that are not quasi-twins but are not located by X_1 . Since D is twin-free, there is a vertex q_3 in $V(D) \setminus S$ that can locate q_1 and q_2 : without loss of generality q_3 is an in-neighbour of q_1 but not q_2 . By our assumption $q_3 \notin X_1$. Now, consider $S' = S \cup \{q_3\}$ and the corresponding S' -partition $\mathcal{P}_{S'}$ of $V(D) \setminus S'$. Since $q_3 \in \bigcup_i Q_i$, any part of \mathcal{P}_S still correspond to some part in $\mathcal{P}_{S'}$. But Q_i has been split into two parts so $\mathcal{P}_{S'}$ has at least one more part than \mathcal{P}_S , and thus, $|\mathcal{P}_{S'}| \geq |S'| - 1$. This contradicts the choice of S , which we assumed to be maximal with this property. \square

Since X_1 is a dominating set, Claim 20 shows that in the absence of quasi-twins, X_1 is locating-dominating. Next claim is proved in [5, Claim 8.B] to deal with quasi-twins.

Claim 21 ([5]). *Any two pairs of quasi-twins in $V(D) \setminus X_1$ are disjoint.*

For each pair of quasi-twins in $V(D) \setminus X_1$, we add one of the vertices of the pair in X_1 . By Claims 20 and 21, the resulting set X'_1 is a locating-dominating set and has size at most $|S| + n_1 + (n - |S| - n_1)/2 = (n + |S| + n_1)/2$.

Consider now the set X_2 of size $n - n_1 - n_2$ consisting of $V(D)$ without one vertex from each part of \mathcal{P}_S . Then all the vertices of $V(D) \setminus X_2$ are located and dominated by S and thus X_2 is a locating dominating set.

Assume now that D has no quasi-twins. Then X_1 and X_2 are two locating-dominating sets of D . If $|X_2| \leq 2n/3$ we are done. Thus we assume that $|X_2| > 2n/3$ which means that $|\mathcal{P}_S| < n/3$. Therefore,

$$\begin{aligned} |X_1| &= |S| + n_1 \\ &\leq |\mathcal{P}_S| + n_1 + 1 \\ &\leq |\mathcal{P}_S| + (n_1 + n_2) \quad \text{since } n_2 \geq 1 \\ &\leq 2|\mathcal{P}_S| \\ &\leq 2n/3 \end{aligned}$$

and we are done.

If D has some quasi-twins, we use the locating-dominating sets X'_1 and X_2 . Again, if $|X_2| \leq 3n/4$, we are done. So, assume that $|X_2| > 3n/4$. Then, $|\mathcal{P}_S| < n/4$.

Therefore,

$$\begin{aligned} |X'_1| &= \frac{|S| + n + n_1}{2} \\ &\leq \frac{|\mathcal{P}_S| + 1 + n + n_1}{2} \\ &\leq \frac{|\mathcal{P}_S| + n + n_1 + n_2}{2} \\ &\leq |\mathcal{P}_S| + \frac{n}{2} \\ &\leq \frac{n}{4} + \frac{n}{2} \\ &= \frac{3n}{4} \end{aligned}$$

and we are done. □

To apply Lemma 19, we prove that such a set S exists when there is a supervising vertex.

Lemma 22. *If a digraph D contains a supervising vertex s , then, there exists a dominating set S such that $|\mathcal{P}_S| \geq |S| - 1$.*

Proof. Consider a supervising vertex s of D . For $i \geq 0$, let V_i be the set of vertices of D such that the shortest directed path from s to v has length i .

Let k be the smallest integer such that V_{k+1} is empty. Since s is supervising, V_0, \dots, V_k form a partition of $V(D)$ where the vertices are sorted according to their distance from s . If $k = 0$, then D contains s as a unique vertex and the set $S = \emptyset$ satisfies the lemma. So from now we assume that $k > 0$.

We build the set S by the following method. Let S be a set of vertices of V_{k-1} that dominates V_k and that is minimal for this property. Then, step by step, for $i = 1$ to $k - 1$, we assume that S is a set of vertices of $V_{k-1} \cup \dots \cup V_{k-i}$ that dominates $V_k \cup \dots \cup V_{k-i+1}$ and we add to S a set of vertices of V_{k-i-1} that dominates $V_{k-i} \setminus S$ and that is minimal for this property. We continue the process until $i = k - 1$.

At each step i , by minimality of the chosen set, when a vertex v of V_{k-i-1} is added to S one can choose a vertex $f(v)$ in $V_{k-i} \setminus S$ whose in-neighbourhood in $V_{k-i-1} \cap S$ is exactly v . By doing so, at the end of the procedure, each vertex of $f(S)$ has different in-neighbours in S . So $|\mathcal{P}_S| \geq |S|$.

Finally if s is not already dominated by S , then we add it to S so that S is a dominating set of $V(D)$. This might increase the cardinality of S by 1 and in the end S is a dominating set such that $|\mathcal{P}_S| \geq |S| - 1$. \square

Then Theorem 18 is direct consequence of Lemmas 19 and 22.

Since all the vertices in a strongly connected digraph are supervising, we have the following corollary:

Corollary 23. *Let D be a twin-free strongly connected digraph on n vertices, then $\gamma^{LD}(D) \leq \frac{3n}{4}$. Moreover, if D is quasi-twin-free, then $\gamma^{LD}(D) \leq \frac{2n}{3}$.*

Note that the second bound is asymptotically tight (see Figure 1). A digraph is called *semicomplete* if there is at least one arc between every pair of vertices. A digraph D is *locally in-semicomplete* if the in-neighbourhood of every vertex x of D induces a semicomplete digraph. Note that semicomplete digraphs are a generalization of tournaments and thus, locally in-semicomplete digraphs naturally generalize local tournaments.

Lemma 24. *A locally in-semicomplete digraph has a supervising vertex.*

Proof. Suppose by contradiction, that there exists a locally in-semicomplete digraph D with no supervising vertex. Consider a vertex of D , such that the set S of vertices v of $V(D)$ for which there exists a directed path from s to v has maximum size, i.e. s is *supervising* a set of vertices of maximum size. Note that $s \in S$ since s forms a directed path of length zero from itself to itself. Since D contains no supervising vertices, we know that $V(D) \setminus S$ is nonempty. Plus, all the edges between $V(D) \setminus S$ and S are oriented from $V(D) \setminus S$ to S . Since D is connected, there exists at least such an edge from a vertex u of $V(D) \setminus S$ to a vertex of S . Recall that u has no in-neighbour in S . Let v be a vertex of S that is an out-neighbour of u and such that the length of a shortest directed path P from s to v has minimum length. By the choice of v , no vertex of P distinct from v is an out-neighbour of u . Note that v is distinct from s , since otherwise u is supervising more vertices than s , contradicting the choice of s . Let w be the in-neighbour of v along P . Both w and u are in the in-neighbourhood of v .

Hence, by in-semicompleteness of D , there must be an arc between them, which is a contradiction. \square

A consequence of Lemma 24 and Theorem 18 is the following corollary.

Corollary 25. *Let D be a twin-free locally in-semicomplete digraph on n vertices, then $\gamma^{LD}(D) \leq \frac{3n}{4}$. Moreover, if D is quasi-twin-free, then $\gamma^{LD}(D) \leq \frac{2n}{3}$.*

Consider the digraph D obtained from a vertex whose out-neighbourhood is made of k disjoint oriented triangles (see Figure 5). On this example, D is a quasi-twin-free locally in-semicomplete digraph on $3k + 1$ vertices. Note that $\gamma^{LD}(D) = 2k$ and $\frac{2n}{3} = 2k + 1$. So we are at distance 1 from the bound given by Corollary 25 and thus asymptotically tight.




Figure 5: Disjoint oriented triangles forming the out-neighbourhood of an extra vertex. A locating dominating set of minimum size is given by gray vertices.

Note that a similar method cannot be applied for locally out-semicomplete digraphs (digraphs where all the out-neighbourhoods are semicomplete). Indeed, there are twin-free locally out-semicomplete digraphs for which the minimum dominating set has size $2(n - 1)/3$ (see for example the reverse of the digraph of Figure 5). Thus, there is no dominating set S such that $|\mathcal{P}_S| \geq |S| - 1$ and Theorem 19 cannot be applied.

References

- [1] J. Bang-Jensen, Y. Guo, G. Gutin, L. Volkmann, *A classification of locally semicomplete digraphs*, Discrete Mathematics 167-168 (1997) 101-114.
- [2] J. Bang-Jensen and G. Gutin. *Digraphs: Theory, Algorithms and Applications*. Springer-Verlag, London, 2nd edition, 2009.
- [3] I. Charon, O. Hudry, and A. Lobstein. *Identifying and locating-dominating codes: Np-completeness results for directed graphs*. IEEE Transactions on Information Theory,48(8):2192–2200, 2002
- [4] F. Foucaud, M. A. Henning, C. Löwenstein and T. Sasse. Locating-dominating sets in twin-free graphs. *Discrete Applied Mathematics* 200:52–58:2016.

- [5] F. Foucaud, S. Heydarshahi, A. Parreau, *Domination and location in twin-free digraphs*, *Discrete Applied Mathematics* 284 (2020) 42-52.
- [6] D. Garijo, A. Gonzalez and A. Marquez. The difference between the metric dimension and the determining number of a graph. *Applied Mathematics and Computation* 249:487–501, 2014.
- [7] R. D. Skaggs. *Identifying vertices in graphs and digraphs*. PhD thesis, University of South Africa, 2007