

Asteroseismology of hot subdwarf B stars observed with TESS: discovery of two new gravity mode pulsating stars

Murat UZUNDAG^{1,2,*}, Roberto SILVOTTI³, Andrzej S. BARAN^{4,5}, Maja VUČKOVIĆ¹,
Péter NÉMETH⁶, Sumanta K. SAHOO^{4,7} and Mike REED⁸

¹ Instituto de Física y Astronomía, Universidad de Valparaíso, Gran Bretaña 1111, Playa Ancha, Valparaíso 2360102, Chile

² European Southern Observatory, Alonso de Cordova 3107, Santiago, Chile

³ INAF-Osservatorio Astrofisico di Torino, strada dell'Osservatorio 20, 10025 Pino Torinese, Italy

⁴ ARDASTELLA Research Group

⁵ Astronomical Observatory, University of Warsaw, Al. Ujazdowskie 4, 00-478 Warszawa, Poland

⁶ Astroserver.org, Fő tér 1, 8533 Malomsok, Hungary

⁷ Nicolaus Copernicus Astronomical Centre of the Polish Academy of Sciences, ul. Bartycka 18, 00-716 Warsaw, Poland

⁸ Department of Physics, Astronomy and Materials Science, Missouri State University, 901 S. National, Springfield, MO 65897, USA

* Corresponding author: muratuzundag.astro@gmail.com

Abstract

TIC 033834484 and TIC 309658435 are long-period pulsating subdwarf B star, which were observed extensively (675 and 621 days, respectively) by the Transiting Exoplanet Survey Satellite (TESS). The high-precision photometric light curve reveals the presence of more than 40 pulsation modes including both stars. All the oscillation frequencies that we found are associated with gravity (g)-mode pulsations, with frequencies spanning from $\sim 80 \mu\text{Hz}$ (2 500 s) to $\sim 400 \mu\text{Hz}$ (12 000 s). We utilize the asteroseismic tools including asymptotic period spacings and rotational frequency multiplets in order to identify the pulsational modes. We found dipole ($l = 1$) mode sequences for both targets and calculate the mean period spacing of dipole modes ($\Delta P_{l=1}$), which allows us to identify the modes. Frequency multiplets provide a rotation period of ~ 64 d for TIC 033834484. From follow-up ground-based spectroscopy, we find that TIC 033834484 has an effective temperature of 24 210 K (140), a surface gravity of $\log g/\text{cms}^{-2} = 5.28$ (03) and TIC 309658435 has an effective temperature of 25 910 K (150), a surface gravity of $\log g/\text{cms}^{-2} = 5.48$ (03).

Keywords: asteroseismology — stars: oscillations (including pulsations) — stars: interiors — stars: evolution — stars: horizontal-branch — stars: subdwarfs — Stars: individual: TIC 033834484, TIC 309658435

1. Introduction

One of the major progresses in our understanding of sdB stars was initiated by Kilkenny et al. (1997) discovering rapid pulsations in hot sdBs known as V361 Hya stars (often referred to as short-period sdBV stars). The V361 Hya stars show multiperiodic pulsations with periods spanning from 60 s to 800 s. Green et al. (2003) discovered the long-period sdB pulsators known as V1093 Her stars. These stars show brightness variations with periods up to a few hours and have amplitudes smaller than 0.1 per cent of their mean brightness. In the first group, the pulsational modes correspond to low-degree, low-order pressure (p)-modes while in the second group, the pulsational modes correspond to low-degree, medium- to high-order gravity (g)-modes. Some sdB pulsators showing both g- and p-modes have been discovered among the two described families of pulsating sdB stars. These stars are referred to as hybrid sdB pulsating stars. These modes are excited by a classical κ

far, TESS has monitored thousand of sdB stars with 2-min and 20-sec cadence during 56 sectors. The stars at high latitude region, close to the ecliptic caps, have been observed continuously (CVZ) and several sdBs were found in this zone. In this work, we present the analysis of two new discoveries of hot subdwarf B pulsating stars. TIC 033834484 and TIC 309658435, were observed by TESS during the first and third year in the southern ecliptic hemisphere. We obtained a low-resolution spectra of each target and applied the model atmosphere models in order to derive the fundamental parameters of the stars. We performed the frequency analysis and detailed seismic mode identification for both targets.

2. Spectroscopy

Intermediate-resolution spectra ($R = 3500$) of TIC 033834484 and TIC 309658435 were collected during three half-nights of observation between 2015 January 28 and February 22 at Las Campanas Observatory, with the IMACS spectrograph at the focus of the Baade 6.5 m telescope. The instrument was used at $f/4$ in longslit mode, and the $0''.75$ -wide slit was employed. The 1200–17.5 grating was tilted by an angle of $16^\circ.8$ to cover the spectral range 3660–5250 Å on the CCD. Figure 1 shows the IMACS spectra together with their best-fit TLUSTY/XTGRID models. The determined spectroscopic parameters demonstrate that TIC 033834484 has $T_{\text{eff}} = 24210(\pm 140)$ (K) and $\log g = 5.28(\pm 0.03)$ (cm s^{-2}) while TIC 309658435 has $T_{\text{eff}} = 25910(\pm 150)$ (K) and $\log g = 5.47(\pm 0.03)$ (cm s^{-2}). The spectra of both sdB stars are dominated by Balmer-lines and only weak He I lines are seen. Both stars belong to the He-weak sdB spectroscopic group, which overlaps with g-mode sdBV pulsators. The atmospheric parameters are in good agreement with the earlier results by Moni Bidin et al. (2017).

3. Photometric data

TIC 033834484 and TIC 309658435 located in the TESS’ CVZ, were observed extensively during 675 and 621 days, respectively. We analyzed TESS observations using the short-cadence (SC) mode with 2-minutes sampling, allowing us to analyze the frequency range up to the Nyquist frequency at about $4167 \mu\text{Hz}$. The Fourier transforms (FT) of the light curves were computed to examine the periodicities present in the data, aiming at identifying the frequency of all pulsation modes, along with their phase and amplitude.

4. Asteroseismology

4.1. Rotational multiplets

In rotating stars, the pulsation frequencies are split into $2l+1$ azimuthal components due to rotation, revealing equally spaced multiplets of $2l+1$ components. This $2l+1$ configuration can be resolved with high-precision photometry if the star has no strong magnetic field and the rotational




Figure 1: Optical spectra of the two new pulsating hot subdwarf B pulsating stars (black lines). Overplotted are the best-fit atmospheric models (red lines). Identifications of Balmer-lines and weak He I lines are marked.

period is not longer than the duration of the observation. Also, in order to detect rotationally split modes, the rotational axis has to be aligned with the pulsation axis, otherwise the pulsation frequencies are split into nine components for $l = 1$ mode (Silvotti et al., 2022, and references therein). Lastly, the inclination angle has to be different from 0 ($i \neq 0$).

The following equation gives the rotation period (P_{rot}) in terms of frequency splitting $\Delta\nu_{n,l,m}$.

$$\nu_{n,l,m} = \nu_{n,l,0} + \Delta\nu_{n,l,m} = \nu_{n,l,0} + m \frac{1 - C_{n,l}}{P_{\text{rot}}}, \quad (1)$$

where $C_{n,l}$ is the Ledoux constant (Ledoux, 1951), which, in the asymptotic limit, depends on the modal degree as $C_{n,l} = 1/l(l+1)$. For dipole and quadrupole modes therefore we get $C_{n,1} \sim 0.5$ and $C_{n,2} \sim 0.17$, respectively.

For TIC 033834484, we were able to determine a frequency splitting of $\sim 0.0905 \mu\text{Hz}$ (Fig. 2), and therefore this star appears to be another slow-rotating single sdB pulsating star with a rotation period of ~ 64 days.




Figure 2: FT of TIC 033834484 showing the region around the frequency triplets near $137 \mu\text{Hz}$ (left panel) and $148 \mu\text{Hz}$ (right panel). From top to bottom we show the window function, the original FT, and the FT of the residuals after subtracting each sequence of frequencies. A black dot represents each prewhitened peak.

4.2. Asymptotic period spacing

In the asymptotic limit of stellar pulsations, i.e., for large radial orders ($k \gg \ell$), g -modes of consecutive radial order in sdBs are approximately uniformly spaced in period (Tassoul, 1980). The asymptotic period spacing is given by $P_{\ell,n} = \Delta P_0 / \sqrt{\ell(\ell+1)n} + \varepsilon_\ell$, where ΔP_0 is the asymptotic period spacing for g -modes, which is defined as $\Delta P_0 \propto [\int_{r_1}^{r_2} \frac{|N|}{r} dr]^{-1}$, N being the Brunt-Väisälä frequency, the critical frequency of nonradial g -modes (Tassoul, 1980), and ε_ℓ is a constant (Unno et al., 1989). The pulsation spectrum is displayed in units of period in Fig. 3 with the expected locations of the $\ell = 1$ modes for even period spacing indicated. We derived the mean period spacing of dipole modes ($\Delta P_{\ell=1}$) and found $263.84_{-1.4}^{+0.8}$ s for TIC 033834484, $257.06_{-1.4}^{+1.5}$ s for TIC 309658435. We used a bootstrap resampling analysis as described by Uzunoglu et al. (2021) in order to propagate the errors in the mean period spacing obtained in our analysis.

5. Results and conclusions

We presented the discovery of two new long-period sdB pulsating stars TIC 033834484 and TIC 309658435. The long dataset allows us to achieve a high frequency resolution of $\sim 0.02 \mu\text{Hz}$ and study the temporal stability of the pulsations, as well as to reveal the rotation period of TIC 033834484. We derived the atmospheric parameters for both stars by fitting synthetic models to the spectra. We investigated the potential variability of the two new sDBVs by examining their




Figure 3: Pulsation spectrum of TIC 033834484 (top panel) and TIC 309658435 (bottom panel) in the period space with the red dots indicating the expected locations of $\ell = 1$ modes from the asymptotic pulsation theory. The horizontal blue lines show the confidence level of 0.1% FAP.

short and ultra-short-cadence observations obtained with TESS. Our investigation of the detected frequencies reveals:

- 12 dipole ($l = 1$) modes for TIC 033834484 with the relative radial order between 20 and 44 and mean period spacing of dipole modes ($\Delta P_{l=1}$) = $263.84^{+0.8}_{-1.4}$ s.
- 9 dipole modes for TIC 309658435 with the relative radial order between 11 and 21 and mean period spacing of dipole modes ($\Delta P_{l=1}$) = $257.06^{+1.5}_{-1.4}$ s.
- Rotational triplets for TIC 033834484 with $\Delta \nu_{l=1}$ of $\sim 0.0905 \mu\text{Hz}$, corresponding to a stellar rotation period of about 64 days, making this star another slowly rotating single pulsating sdB among ten others (Silvotti et al., 2022). TIC 309658435 does not show any multiplets suggesting that it has either a very slow rotation period (longer than ~ 600 days) or a pole-on orientation of the pulsation axis, or the side components are not driven.

Acknowledgments

M.U. acknowledges financial support from CONICYT Doctorado Nacional in the form of grant number No: 21190886 and ESO studentship program. P.N. acknowledges support from the Grant Agency of the Czech Republic (GAČR 22-34467S). The Astronomical Institute in Ondřejov is supported by the project RVO:67985815. This research has used the Sandbox services of www.Astroserver.org. The authors warmly thank Christian MoniBidin, who provided the spectroscopic observation of the two targets. Financial support from the National Science Center in Poland under project No. UMO-2017/26/E/ST9/00703 is acknowledged.

Further Information

ORCID identifiers of the authors

0000-0002-6603-994X (Murat UZUNDAG)

0000-0003-0963-0239 (Péter NÉMETH)

Author contributions

M.U. and R.S. and A.S.B. carried out the light curve analyses as well as frequency solutions. P.N. performed the atmospheric analysis. The text was written by M.U. All authors contributed to the discussion and interpretation of the results and commented on the written draft of the paper.

Conflicts of interest

The authors declare no conflict of interest.

References

- Charpinet, S., Fontaine, G., Brassard, P., Chayer, P., Rogers, F. J., Iglesias, C. A. and Dorman, B. (1997) A Driving Mechanism for the Newly Discovered Class of Pulsating Subdwarf B Stars. *ApJ*, 483, L123–L126. <https://doi.org/10.1086/310741>.
- Charpinet, S., Fontaine, G., Brassard, P. and Dorman, B. (1996) The Potential of Asteroseismology for Hot, Subdwarf B Stars: A New Class of Pulsating Stars? *ApJ*, 471, L103. <https://doi.org/10.1086/310335>.
- Charpinet, S., Giammichele, N., Zong, W., Grootel, V. V., Brassard, P. and Fontaine, G. (2018) Rotation in sdb stars as revealed by stellar oscillations. *Open Astronomy*, 27(1), 112 – 119. <https://www.degruyter.com/view/journals/astro/27/1/article-p112.xml>.
- Fontaine, G., Brassard, P., Charpinet, S., Green, E. M., Chayer, P., Billères, M. and Randall, S. K. (2003) A Driving Mechanism for the Newly Discovered Long-Period Pulsating Subdwarf B Stars. *ApJ*, 597, 518–534. <https://doi.org/10.1086/378270>.
- Green, E. M., Fontaine, G., Reed, M. D., Callerame, K., Seitzzahl, I. R., White, B. A., Hyde, E. A., Østensen, R., Cordes, O., Brassard, P., Falter, S., Jeffery, E. J., Dreizler, S., Schuh, S. L., Giovanni, M., Edelmann, H., Rigby, J. and Bronowska, A. (2003) Discovery of A New Class of Pulsating Stars: Gravity-Mode Pulsators among Subdwarf B Stars. *ApJ*, 583, L31–L34. <https://doi.org/10.1086/367929>.
- Kilkenny, D., Koen, C., O’Donoghue, D. and Stobie, R. S. (1997) A new class of rapidly pulsating star - I. EC 14026-2647, the class prototype. *MNRAS*, 285, 640–644. <https://doi.org/10.1093/mnras/285.3.640>.
- Ledoux, P. (1951) The Nonradial Oscillations of Gaseous Stars and the Problem of Beta Canis Majoris. *ApJ*, 114, 373. <https://doi.org/10.1086/145477>.
- Moni Bidin, C., Casetti-Dinescu, D. I., Girard, T. M., Zhang, L., Méndez, R. A., Vieira, K., Korchagin, V. I. and van Altena, W. F. (2017) Young stars in the periphery of the Large Magellanic Cloud. *MNRAS*, 466(3), 3077–3087. <https://doi.org/10.1093/mnras/stw3242>.
- Reed, M. D., Baran, A., Quint, A. C., Kawaler, S. D., O’Toole, S. J., Telting, J., Charpinet, S., Rodríguez-López, C., Østensen, R. H., Provencal, J. L., Johnson, E. S., Thompson, S. E., Allen, C., Middour, C. K., Kjeldsen, H. and Christensen-Dalsgaard, J. (2011) First Kepler results on compact pulsators - VIII. Mode identifications via period spacings in g-mode pulsating subdwarf B stars. *MNRAS*, 414, 2885–2892. <https://doi.org/10.1111/j.1365-2966.2011.18532.x>.

- Silvotti, R., Németh, P. and et al. (2022) Filling the gap between synchronized and non-synchronized sdBs in short-period sdBV+dM binaries with TESS. *MNRAS*, 511(2), 2201–2217. <https://doi.org/10.1093/mnras/stac160>.
- Tassoul, M. (1980) Asymptotic approximations for stellar nonradial pulsations. *ApJS*, 43, 469–490. <https://doi.org/10.1086/190678>.
- Unno, W., Osaki, Y., Ando, H., Saio, H. and Shibahashi, H. (1989) Nonradial oscillations of stars.
- Uzundag, M., Vučković, M., Németh, P., Miller Bertolami, M., Silvotti, R., Baran, A. S., Telting, J. H., Reed, M., Shoaf, K. A., Østensen, R. H. and Sahoo, S. K. (2021) Asteroseismic analysis of variable hot subdwarf stars observed with TESS. I. The mean g-mode period spacings in hot subdwarf B stars. *A&A*, 651, A121. <https://doi.org/10.1051/0004-6361/202140961>.