

GROUPS ACTING DISTALLY AND MINIMALLY ON \mathbb{S}^2 AND \mathbb{RP}^2

ENHUI SHI AND HUI XU

ABSTRACT. Let X be the 2-sphere \mathbb{S}^2 or the real projective plane \mathbb{RP}^2 . We show that if Γ is a finitely generated group acting minimally and distally on X , then Γ contains a nonabelian free subgroup.

1. INTRODUCTION

The aim of the note is continuing the study of the following question:

*Given a discrete group G and a compact metric space X ,
can G act on X distally and minimally?*

The answer to this question involves the discussions around the algebraic structure of G and the topology of X . Here, we are mainly concerned on the case that X is a closed surface (a compact connected 2-manifold with no boundary). There have been several related investigations around this topic. A remarkable result due to Furstenberg says that if a non-trivial space X admits a distal minimal action by a locally compact abelian group, then X cannot be simply connected (see [4, Theorem 11.1] or [1, Chapter 7-Theorem 16]). In [9], Shi proved further that if a continuum X admits a distal minimal amenable group action, then the first Čech cohomology group $\check{H}^1(X)$ of X with integer coefficients is nontrivial; in particular, if X is a CW-complex, then the fundamental group of X cannot be finite; so, the 2-sphere \mathbb{S}^2 and the real projective plane \mathbb{RP}^2 admit no distal minimal actions by amenable groups. Bronšteĭn proved that if X is a connected and locally connected finitely dimensional compact metric space which admits a distal minimal group action, then X must be a manifold and the fundamental group $\pi_1(X)$ is virtually nilpotent ([3]); this implies that if X is a closed surface except for the sphere \mathbb{S}^2 , the real projective plane \mathbb{RP}^2 , the torus \mathbb{T}^2 , and the Klein bottle \mathbb{K}^2 , then it admits no distal minimal actions by any group. Shi showed that no closed surface admits a distal minimal action by $SL(n, \mathbb{Z})$ with $n \geq 3$ ([10]).

The following is the main theorem of this paper. Recall that a group G is a *small group* if it contains no free nonabelian subgroups.

2010 *Mathematics Subject Classification.* 37B05.

Key words and phrases. distality, amenable group, free group, minimality, cohomology.

Theorem 1.1. *Let X be the 2-sphere \mathbb{S}^2 or the real projective plane \mathbb{RP}^2 . If Γ is a finitely generated group acting minimally and distally on X , then Γ contains a nonabelian free subgroup. Equivalently, X admits no distal minimal actions by a small group.*

Here we remark that the class of small groups is strictly larger than that of amenable groups, so this theorem is not implied by the main theorem in [9]; and it is easy to construct distal minimal actions on \mathbb{S}^2 and \mathbb{RP}^2 by $\mathbb{Z} * \mathbb{Z}$. In addition, the theorem does not hold when X is either the torus \mathbb{T}^2 or the Klein bottle \mathbb{K}^2 , since they admit distal minimal actions by abelian groups (see the appendix).

Now we summarize all the known results around the existence of distal minimal group actions on closed surfaces in the following tabular.

Closed surfaces	Existence	Non-existence
$\mathbb{S}^2, \mathbb{RP}^2$	$\mathbb{Z} * \mathbb{Z}$	small groups, $SL(n, \mathbb{Z}) (n \geq 3)$
$\mathbb{T}^2, \mathbb{K}^2$	\mathbb{Z}	$SL(n, \mathbb{Z}) (n \geq 3)$
Others		Any groups

2. PRELIMINARIES

In this section, we just list the theorems that will be used in the proof of the main theorem without mentioning the related notions and notations already appeared in [9, 10].

Theorem 2.1. [1, p.98] *Let (X, G, ϕ) and (Y, G, ψ) be distal minimal actions, and let $f : X \rightarrow Y$ be a homomorphism. Then f is open.*

Theorem 2.2. [1, p.104] *Suppose X is not a single point. If (X, G, ϕ) is distal minimal, then it has a nontrivial equicontinuous factor.*

Theorem 2.3. [3, Theorem 3.17.12] *Let $\pi : (X, G, \phi) \rightarrow (Y, G, \psi)$ be a homomorphism between minimal systems. Suppose that π is open and G is finitely generated. If (Y, G, ψ) is equicontinuous and there is some $y \in Y$ such that $f^{-1}(y)$ is of 0-dimension. Then (X, G, ϕ) is also equicontinuous.*

Theorem 2.4. [1, p.52] *Let (X, G, ϕ) be equicontinuous. Then the closure $\overline{\phi(G)}$ in $C(X, X)$ with respect to the uniform convergence topology is a compact topological group.*

Theorem 2.5. [8] *Let (X, G, ϕ) and (Y, G, ψ) be distal minimal actions, and let $f : X \rightarrow Y$ be a homomorphism. Then for every $y \in Y$, we have $\dim(Y) + \dim(f^{-1}(y)) = \dim(X)$.*

Theorem 2.6. [3, Theorem 3.17.10] *Let (X, G, ϕ) be a distal minimal system with X being a connected and locally connected compact metric space of finite dimension, then X is a manifold.*

Theorem 2.7. [5, Corollary 4.25] *Let G be a compact Lie group and let \mathfrak{g} be the Lie algebra of G . Then $\mathfrak{g} = Z(\mathfrak{g}) \oplus [\mathfrak{g}, \mathfrak{g}]$, where $Z(\mathfrak{g})$ is the center of \mathfrak{g} and $[\mathfrak{g}, \mathfrak{g}]$ is semisimple.*

Theorem 2.8. [5, Corollary 1.103] *Let G be a compact connected commutative Lie group of dimension n . Then G is isomorphic to the n -torus \mathbb{T}^n .*

Theorem 2.9. [11, Theorem 3.50] *Let G be a connected Lie group with Lie algebra \mathfrak{g} . Then the center of G is a closed Lie subgroup of G with Lie algebra the center of \mathfrak{g} .*

Theorem 2.10. [6, p.65] *Let X be a compact metric space and let (X, G) be an action of group G on X . Suppose G is compact. Then for every $x \in X$, G/G_x is homeomorphic to Gx , where $G_x = \{g \in G : gx = x\}$.*

Theorem 2.11. [6, p.99] *Let G be a compact group and let U be an open neighborhood of the identity e . Then U contains a closed normal subgroup H of G such that G/H is isomorphic to a Lie group.*

Theorem 2.12. [6, p.61] *Let X be a compact metric space and let (X, G) be an action of group G on X . Suppose G is compact and H is a closed normal subgroup of G . Then G/H can act on X/H by letting $gH \cdot H(x) = H(gx)$ for $gH \in G/H$ and $H(x) \in X/H$.*

We use $\check{H}^1(X)$ to denote the first Čech cohomology group of X with integer coefficients.

Theorem 2.13. [9, Corollary 2.15] *Let $f : X \rightarrow Y$ be an open map from a continuum X onto a continuum Y . Then $f^* : \check{H}^1(Y) \rightarrow \check{H}^1(X)$ is injective.*

Theorem 2.14. [2, Theorem 1.3] *Let G be a connected non-solvable real Lie group of dimension d . Then any finitely generated dense subgroup of G contains a dense free subgroup of rank $2d$.*

It is well known that a compact connected Hausdorff space is locally connected metrizable if and only if it is a continuous image of the closed interval $[0, 1]$ ([7, Theorem 8.18]). Thus the following result is direct.

Theorem 2.15. *Let X be a compact metric space which is connected and locally connected and Y be a Hausdorff space. If there is a continuous surjection $f : X \rightarrow Y$, then Y is locally connected and metrizable.*

3. PROOF OF THE MAIN THEOREM

Lemma 3.1. *Let X be the 2-sphere \mathbb{S}^2 or the real projective plane \mathbb{RP}^2 . Let Γ be a finitely generated group and $\phi : \Gamma \rightarrow \text{Homeo}(X)$ be a distal minimal action on X . Then (X, Γ, ϕ) is equicontinuous.*

Proof. Assume to the contrary that ϕ is not equicontinuous. From Theorem 2.2, we let (Y, Γ, ψ) be the maximal equicontinuous factor of (X, Γ, ϕ) with a factor map π . Then Y is a compact manifold of dimension ≤ 2 by Theorem 2.5, Theorem 2.6, and Theorem 2.15. If $\dim(Y) = 2$, then it follows from Theorem 2.5 that for every $x \in X$, $\dim(\pi^{-1}(x)) = 0$.

This together with Theorem 2.3 and Theorem 2.1 implies that (X, Γ, ϕ) is equicontinuous, which contradicts the assumption. So, we may assume that $\dim(Y) = 1$; this means that Y is the circle \mathbb{S}^1 . Since π is open and $\check{H}^1(\mathbb{S}^1)$ is the integer group, from Theorem 2.1 and Theorem 2.13, we have that $\check{H}^1(X)$ is infinite, which is a contradiction. \square

Lemma 3.2. *Let G be a connected compact Lie group acting faithfully and transitively on a closed surface X with finite fundamental group. Then G is semisimple.*

Proof. Assume to the contrary that G is not semisimple. Then by Theorem 2.7, Theorem 2.8, and Theorem 2.9, the connected component $Z(G)_0$ of the center of G is isomorphic to some torus \mathbb{T}^n with $n > 0$. Set $K = Z(G)_0$. For $x \in X$, let $\text{Stab}(x) := \{k \in K : kx = x\}$ be the stabilizer of x in K . Then from Theorem 2.10, Kx is homeomorphic to $K/\text{Stab}(x)$ which is also a torus. Thus Kx is either a point or a circle. If for every $x \in X$, Kx is a circle, then similar to the arguments in Lemma 3.1, X/K is a circle and the Čech cohomology group $\check{H}^1(X)$ with integer coefficients is infinite. This is a contradiction. So, there is some $x_0 \in X$ with $Kx_0 = x_0$. Since K is in the center of G , we have $Kgx_0 = gKx_0 = gx_0$ for every $g \in G$. Noting that $Gx_0 = X$, the action of K on X is trivial, which contradicts the faithfulness of the action. So, G is semisimple. \square

Proof of the main theorem. Let $\phi : \Gamma \rightarrow \text{Homeo}(X)$ be the distal minimal action. From Lemma 3.1, we see that the (X, Γ, ϕ) is equicontinuous. Let K be the closure of $\phi(\Gamma)$ with respect to the uniform topology on $\text{Homeo}(X)$. It follows from Theorem 2.4 that K is a compact metric group acting transitively on X . By Theorem 2.11, we can take a closed normal subgroup N of K such that K/N is a Lie group and X/N is not a single point. Then it canonically induces an action of K on X/N and the natural quotient map $X \rightarrow X/N$ is an equicontinuous extension. Thus it follows from Theorem 2.5, Theorem 2.6, and Theorem 2.15 that X/N is a manifold of dimension ≤ 2 . Similar to the arguments in Lemma 3.1, we have $\dim(X/N) = 2$.

Set $p : \mathbb{S}^2 \rightarrow X$ be a covering and $q : X \rightarrow X/N$ be the quotient map. Let $\pi : Y \rightarrow X/N$ be the universal covering and $\tilde{q}p : \mathbb{S}^2 \rightarrow Y$ be the lifting of qp . Since π is open and p and q are local homeomorphisms, we see that $\tilde{q}p$ is open. Thus $\tilde{q}p(\mathbb{S}^2)$ is open and closed in Y . Thus $Y = \tilde{q}p(\mathbb{S}^2)$ by the connectedness. So Y is compact (homeomorphic to \mathbb{S}^2). Thus π is a finite cover and then the fundamental group of X/N is finite.

Now consider the natural action of K/N on X/N (see Theorem 2.12). Since the connected component $(K/N)_0$ has finite index in K/N , the $(K/N)_0$ action on X/N is still transitive and $\phi(\Gamma)N \cap (K/N)_0$ is dense in $(K/N)_0$. Applying Lemma 3.2, we see that a quotient group of $(K/N)_0$ is semisimple. This together with Theorem 2.14 implies the existence of free nonabelian subgroups in Γ . \square

4. APPENDIX

In this appendix, we show that there is a minimal distal homeomorphism on the Klein bottle.

Let the torus \mathbb{T}^2 be $\mathbb{R}^2/\mathbb{Z}^2$. Then there is a \mathbb{Z}_2 action on \mathbb{T}^2 by

$$h(x, y) = \left(x + \frac{1}{2}, 1 - y\right) \pmod{\mathbb{Z}^2},$$

where h is the nonidentity element in \mathbb{Z}_2 . It is easy to see that this action is free and properly discontinuous and the quotient space $\mathbb{T}^2/\mathbb{Z}_2$ is the Klein bottle \mathbb{K}^2 .

Now for a homeomorphism T of \mathbb{T}^2 , if it commutes with h , i.e., $Th = hT$, then it induces a homeomorphism \tilde{T} of \mathbb{K}^2 .

Let α be an irrational number and $\phi : \mathbb{T} \rightarrow \mathbb{T}$ be a continuous mapping. Further, define a homeomorphism $T : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ by

$$T(x, y) = (x + \alpha, y + \phi(x)) \pmod{\mathbb{Z}^2}.$$

The system (\mathbb{T}^2, T) is a skew product system and it is well known that this system is distal, since it is a group extension of a minimal equicontinuous system. The following theorem characterizes the minimality of such skew product system.

Theorem 4.1. [1, Chapter 5, Theorem 10] *The above defined system (\mathbb{T}^2, T) is minimal if and only if for each $k \in \mathbb{Z} \setminus \{0\}$, there is no continuous function $f : \mathbb{T} \rightarrow \mathbb{T}$ such that $f(x + \alpha) = f(x) + k\phi(x)$ for each $x \in \mathbb{T}$.*

Now we take $\phi : \mathbb{T} \rightarrow \mathbb{T}$ to be $\phi(x) = 1 - |1 - 2x|$ for $x \in [0, 1)$. By comparing the Fourier coefficients, there is no continuous function $f : \mathbb{T} \rightarrow \mathbb{T}$ such that $f(x + \alpha) = f(x) + k\phi(x)$ for each $k \in \mathbb{Z} \setminus \{0\}$. Thus the system (\mathbb{T}^2, T) is minimal by Theorem 4.1. It is straightforward to calculate that for each $(x, y) \in \mathbb{T}^2$, $Th(x, y) = (x + \alpha + \frac{1}{2}, 1 - y + \phi(x + \frac{1}{2}))$ and $hT(x, y) = (x + \alpha + \frac{1}{2}, 1 - y - \phi(x))$. Note that

$$\phi\left(x + \frac{1}{2}\right) = \begin{cases} 1 - 2x, & x \in [0, 1/2) \\ 2x - 1, & x \in [1/2, 1) \end{cases} \text{ and } -\phi(x) = \begin{cases} -2x, & x \in [0, 1/2) \\ 2x - 2, & x \in [1/2, 1) \end{cases}.$$

Therefore, it follows that T commutes with h and thus the induced homeomorphism \tilde{T} on \mathbb{K}^2 is also minimal and distal.

REFERENCES

- [1] J. Auslander, *Minimal flows and their extensions*. North-Holland Mathematics Studies, 153. Notas de Matemática [Mathematical Notes], 122. North-Holland Publishing Co., Amsterdam, 1988.
- [2] E. Breuillard, T. Gelander, *On dense free subgroups of Lie groups*. J. Algebra 261 (2003), 448-467.

- [3] I. Bronštejn, *Extensions of minimal transformation groups*, Suthoff & Noordhoff, 1979.
- [4] H. Furstenberg, *The Structure of distal flows*. Amer. J. Math. 85 (1963), 477-515.
- [5] A. Knapp, *Lie groups beyond an introduction* (Second Edition). PM 140, Birkhauser, 2002.
- [6] D. Montgomery, L. Zippin, *Topological transformation groups*. Interscience Publishers, New York-London, 1955.
- [7] S. Nadler, *Continuum theory*, volume 158 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1992. An introduction.
- [8] M. Rees, *On the structure of minimal distal transformation groups with manifolds as phase spaces*, thesis, University of Warwick, Coventry, England.
- [9] E. Shi, *Continua having distal minimal actions by amenable groups*. arXiv:2001.03755.
- [10] E. Shi, *Distal higher rank lattice actions on surfaces*. arXiv:2001.01183.
- [11] F. Warner, *Foundations of differentiable manifolds and Lie groups*. Springer-Verlag, GTM 94, 1983.

(E.H. Shi) SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU 215006, P. R. CHINA

Email address: ehshi@suda.edu.cn

(H. Xu) CAS WU WEN-TSUN KEY LABORATORY OF MATHEMATICS, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI, ANHUI 230026, CHINA

Email address: huixu2734@ustc.edu.cn