

*A procedure to obtain symmetric cycles of any odd length using directed Hajós constructions*

# A procedure to obtain symmetric cycles of any odd length using directed Hajós constructions

Juan Carlos García-Altamirano, Mika Olsen, Jorge Cervantes-Ojeda

Departamento de Matemáticas Aplicadas y Sistemas  
Universidad Autónoma Metropolitana - Cuajimalpa  
México City, México

email: carlos\_treze@ciencias.unam.mx, olsen@cua.uam.mx, jcervantes@cua.uam.mx

(Received January 17th, 2023)

## Abstract

The dichromatic number of a digraph  $D$  is the minimum number of colors of a vertex coloring of  $D$  such that  $D$  has no monochromatic cycles. The Hajós join were recently extended to digraphs (using the dichromatic number) by J. Bang-Jensen et. al. and Hajós (directed) operations is a tool to obtain  $r$ -(di)chromatic (di)graphs. J. Bang-Jensen et. al. posed in 2020 the problem of how to obtain the symmetric cycle of length 5 from symmetric cycles of length 3. We recently solved this problem by applying a genetic algorithm. In this article, a procedure is presented to construct any odd symmetric cycle by applying directed Hajós operations to symmetric cycles of length 3, thus, generalizing the known construction of the symmetric cycle of length 5. In addition, this procedure is analyzed to determine its computational complexity.

## 1 Introduction

The dichromatic number of a digraph  $D$  was introduced by V. Neumann-Lara in 1982 [20] as an extension of the chromatic number of a graph. The dichromatic number of a digraph is the minimum number of colors of a vertex coloring of  $D$  such that  $D$  has no monochromatic cycles and several concepts and results for the chromatic number of a graph have been extended to digraphs using the dichromatic number. For instance, [1, 2, 9, 12, 14, 15]. In 2020 J. Bang-Jensen et. al. [3] extended the well-known Hajós join for graphs to digraphs.

The digraph  $H$  obtained by **identifying** a non-empty set  $I$  of independent vertices is defined as the digraph  $H = D - I$  adding a new vertex  $v$  and adding all arcs from  $v$  to  $N_D^+(I) = \bigcup_{u \in I} N_D^+(u)$  and all arcs from  $N_D^-(I) = \bigcup_{u \in I} N_D^-(u)$  to  $v$ . The new vertex  $v$  may preserve the label of one of the vertices of the independent set  $I$ .

The Hajós join was defined for digraphs in 2020 by Bang-Jensen et. al. [3] as an extension of the well-known Hajós join [13, 17, 24] for graphs. We use Figure 1 to illustrate the definition of directed Hajós Join. Let  $D_1$  and  $D_2$  be two disjoint digraphs. Let  $u_1v_1 \in A(D_1)$  and  $v_2u_2 \in A(D_2)$ . The **directed Hajós join**  $D = (D_1, u_1, v_1) \nabla (D_2, v_2, u_2)$  or, briefly  $D = D_1 \nabla D_2$  of  $D_1$  and  $D_2$  is defined as the disjoint union of  $D_1$  and  $D_2$  and deleting both arcs  $u_1v_1$  and  $v_2u_2$ , identifying the vertices  $v_1$  and  $v_2$  to a new vertex  $v$  and adding the arc  $u_1u_2$ . The vertex  $v$  may be denoted by  $v_1$ ,  $v_2$  or  $v$ .

---

**Key words and phrases:** Hajós operation, dichromatic number, computational complexity.  
**05C20, 05C15**




Figure 1: Directed Hajós join of two directed triangles and the resulting digraph.

We call the directed Hajós join and the vertex identifications the **directed Hajós operations**. The class of **Hajós- $k$ -constructible digraphs** defined as the smallest family of digraphs that contains all complete digraphs of order  $k$  and is closed under directed Hajós operations. J. Bang-Jensen et. al. [3] proved that any  $k$ -critical digraph is Hajós- $k$ -constructible. Since odd symmetric cycles are 3-critical, any symmetric odd cycle can be constructed from the symmetric complete digraph on three vertices  $D(K_3)$  using a sequence of directed Hajós operations (directed Hajós joins and identifications of non-adjacent vertices). In the same paper they posed the following question (Question 19):

*How can a bidirected (symmetric)  $C_5$  ( $D(C_5)$ ) be constructed from copies of  $D(K_3)$  by only using directed Hajós operations?*

In a recent paper J. C. García-Altamirano et. al. [11] obtained a sequence of Hajós operations applied to  $D(K_3)$  in order to obtain the symmetric cycle  $D(C_5)$  using a Rank Genetic Algorithm. In this paper we generalize this sequence to obtain any odd symmetric cycle applying Hajós operations from  $D(K_3)$ . This procedure provides an upper bound to the Hajós number of any odd symmetric cycle and an upper bound of its computational complexity.

We consider finite digraphs without loops and multiple arcs. For all definitions not given here we refer the reader to the book of J. Bang-Jensen and G. Gutin [4]. Let  $D$  be a digraph with vertex set  $V(D)$  and arc set  $A(D)$ . The **in-neighborhood** of a vertex  $u$  is  $N^-(u) = \{v \in V(D) \mid vu \in A(D)\}$  and the **out-neighborhood** of a vertex  $u$  is  $N^+(u) = \{v \in V(D) \mid uv \in A(D)\}$ . Two vertices in a digraph  $D$  are **independent** if there are no arcs between them in  $D$ , a set of vertices  $X$  is independent in a digraph  $D$  if any pair of vertices of  $X$  are independent in  $D$ . An arc  $uv \in A(D)$  is **symmetric (asymmetric)** if  $vu \in A(D)$  ( $vu \notin A(D)$ ), and a digraph is symmetric (bidirected graph) if every arc of  $D$  is a symmetric arc. The **symmetric digraph**  $D(G)$ , of the graph  $G$ , is the digraph obtained by replacing each edge by a symmetric arc.

## 2 Construction of odd symmetric cycles

Let  $D$  and  $D'$  be two disjoint digraphs of the same order  $n$ , with vertex-set  $\{v_0, v_1, \dots, v_{n-1}\}$  and  $\{v'_0, v'_1, \dots, v'_{n-1}\}$  resp., and let  $v_i v_j \in A(D)$  and  $v'_k v'_l \in A(D')$ , such that  $i, j, k, l \in \{0, 1, \dots, n-1\}$ ,  $i \neq j$ ,  $k \neq l$  and  $j - i \neq k - l$ . We define the **cyclic of Hajós identification**  $H = (D, v_i, v_j) \otimes (D', v'_k, v'_l)$  or, simply  $H = D \otimes D'$ , as the digraph obtained by the directed Hajós Join  $\hat{H} := (D, v_i, v_j) \nabla (D', v'_k, v'_l)$ , and the following  $n - 1$  identifications: for  $r = 1, 2, \dots, n - 1$ , the vertices  $v'_{k+r}$  and  $v_{j+r}$  are identified in the vertex  $v_{j+r}$ , the indices are taken modulo  $n$ .

**Remark 1.** By definition, we identify the vertices  $v'_a, v_{a-k+j} \in V(\hat{H})$  in the vertex  $v_{a-k+j}$ . The condition  $j - i \neq k - l$  assures that the vertices  $v'_i$  and  $v_i$  are not identified, because in this case a loop is obtained in the vertex  $v_i$  by the arc that is added in  $\hat{H}$ . Note that two vertices in  $\hat{H}$  which were original consecutive vertices in  $D'$  are identified into two vertices which were originally two consecutive vertices in  $D$ . Moreover, if  $v'_b v'_a \in A(\hat{H})$ , in  $H$  this arc becomes the arc  $v_{b-k+j} v_{a-k+j}$  in  $H$ .

**Lemma 1.** Let  $n \geq 1$  and let  $D$  be a symmetric cycle of order  $2^{n+1} + 1$  with asymmetric arcs  $v_0 v_{2^n}, v_x v_a$ , where  $x = 2^n + a + 1$ . Let  $D'$  be a copy of  $D$  such that  $v'_i \in V(D')$  is the copy of the vertex  $v_i \in V(D)$ . Then  $H = (D, v_x, v_a) \otimes (D', v'_0, v'_{2^n})$  is a symmetric cycle of order  $2^{n+1} + 1$  with two asymmetric arcs  $v_0 v_{2^n}, v_{2^n+2a+1} v_{2a}$ .

*Proof.* Let  $H = (D, v_x, v_a) \otimes (D', v'_0, v'_{2^n})$ , by Remark 1,

- (i) the vertex  $v'_{2^n} \in \hat{H}$  is identified with the vertex  $v_{(2^n)-(0)+(a)} = v_{2^n+a} = v_{x-1}$ , so the arc  $v_x v'_{2^n} \in A(\hat{H})$  is transformed into the arc  $v_x v_{x-1} \in A(H)$ , that is, an arc between two vertices consecutive, therefore this arc is a symmetric arc.
- (ii) The arc  $v'_x v'_a$  de  $\hat{H}$  is transformed into the arc  $v_{(2^n+a+1)-0+a} v_{a-0+a} = v_{2^n+2a+1} v_{2a} = v_{x+a} v_{2a}$  of  $H$ .

Note that the digraphs  $D$  and  $H$  only differ in two arcs, namely  $v_x v_a = A(D) \setminus A(H)$  and the arc  $v_{x+a} v_{2a} = A(H) \setminus A(D)$ , thus  $H = D + \{v_{x+a} v_{2a}\} - \{v_x v_a\}$ . In Figure 2 the dotted arrow represents the deleted arc  $v_x v_a$ .




Figure 2:  $H = D + \{v_{x+a} v_{2a}\} - \{v_x v_a\}$ .

For  $r = 2, \dots, n + 1$ , the digraph  $H_r = (H_{r-1}, v_{2^n+2^{r-2}+1}, v_{2^{r-2}}) \otimes (H'_{r-1}, v'_0, v'_{2^n})$  is a symmetric cycle of order  $2^{n+1}$  with the asymmetric arcs  $v_0 v_{2^n}, v_{2^n+2^{r-1}+1} v_{2^{r-1}}$ .  $\square$

**Theorem 2.** *Let  $n \geq 1$ . The symmetric cycle of order  $2^{n+1} + 1$  can be constructed from the symmetric cycle of order  $2^n + 1$  using  $(n + 2)(2^{n+1} + 1)$  directed Hajós operations.*

*Proof.* Let  $n \geq 1$ , consider the symmetric cycle of order  $2^n + 1$ ,  $D = D(C_{2^n+1}) = (v_0, v_1, \dots, v_{2^n}, v_0)$  and let  $D' = (v'_0, v'_1, \dots, v'_{2^n}, v'_0)$  be a disjoint copy of  $D$ , where  $v'_i$  is the copy of  $v_i$  for  $i \in \{0, 1, \dots, 2^n\}$ . We define the digraph  $H_0 = (D, v_{2^n}, v_0) \nabla (D', v'_0, v'_1)$ , see Figure 3.




Figure 3: The directed Hajós joint of two symmetric cycles and the resulting digraph  $H_0 = (D, v_{2^n}, v_0) \nabla (D', v'_0, v'_1)$ .

Relabel the vertices of  $H_0$  as follows:

$$w = \begin{cases} v_i & \text{if } w = v_i \text{ for } i = 0, 1, \dots, 2^n, \\ v_{2^n+i} & \text{if } w = v'_i \text{ for } i = 1, 2, \dots, 2^n. \end{cases}$$

Note that  $H_0$  is a digraph of order  $2^{n+1} + 1$ , with asymmetric arcs  $v_{2^n}v_{2^n+1}$ ,  $v_0v_{2^n}$ ,  $v_{2^n+1}v_0$ , and the symmetric path  $v_{2^n+1}, v_{2^n+2}, \dots, v_{2^n+1}, v_0, v_1, \dots, v_{2^n}$ . Let  $H'_0$  be a disjoint copy of  $H_0$ , with vertex-set  $\{v'_0, v'_1, \dots, v'_{2^n+1}\}$  where  $v'_i$  is the copy of the vertex  $v_i$  for  $i = 0, 1, \dots, 2^{n+1}$ .

Let  $H_1 = (H_0, v_0, v_{2^n}) \otimes (H'_0, v'_{2^n+1}, v'_0)$ . In Figure 4, we consider the directed Hajós union and  $2^{n+1}$  vertex identifications. The digraph on the left indicates the directed union of Hajós, where the dashed arrows must be removed, the thick arrow added, the black vertices are identified, and the shades of gray indicate the four pairwise vertex identifications. In each figure, the digraph on the right is the result of the  $2^{n+1} + 1$  Hajós operations.

By Remark 1:

- (i) The vertex  $v'_0 \in V(\hat{H}_1)$  is identified with the vertex  $v_{(0)-(2^n+1)+(2^n)} = v_{2^n+1}$ , hence, the arc  $v_0v'_0 \in A(\hat{H}_1)$ , becomes the arc  $v_0v_{2^n+1} \in A(H_1)$ .
- (ii) The arc  $v'_0v'_{2^n} \in A(\hat{H}_1)$  becomes the arc  $v_{(0)-(2^n+1)+(2^n)}v_{(2^n)-(2^n+1)+(2^n)} = v_{2^n+1}v_{2^n-1}$  in  $H_1$ .

Thus,  $H_1$  is a symmetric cycle of order  $2^{n+1} + 1$  with asymmetric arcs  $v_{2^n+1}v_{2^n-1}, v_{2^n+1}v_0$ .




Figure 4: The directed Hajós join of two copies of the digraph  $H_0$  and the resulting digraph  $H_1 = (H_0, v_0, v_{2^n}) \otimes (H'_0, v'_{2^n+1}, v'_0)$ .

In order to simplify the writing we relabel (in cyclic order) the vertices of  $H_1$ , for  $i = 0, 1, \dots, 2^{n+1}$ ,  $v_i := v_{i+1}$  the index are taken modulo  $2^{n+1} + 1$ .

For  $k = 1, \dots, n-1$ , let  $H_{k+1} = H_k \otimes H'_k$ . By Lemma 1,  $H_{k+1}$  is a symmetric cycle of order  $2^{n+1} + 1$  with asymmetric arcs  $v_0 v_{2^n}$  and  $v_{2^{n+2k+1}} v_{2^k}$ . Applying Lemma 1, the digraph  $H_{n+1}$  is a symmetric cycle of order  $2^{n+1} + 1$  with a unique asymmetric arc  $v_0 v_{2^n}$  because  $2^n + 2^n + 1 \equiv 0 \pmod{2^{n+1} + 1}$ .

Finally, let  $H_{n+2} = (H_{n+1}, v_0, v_{2^n}) \otimes (H'_{n+1}, v'_0, v'_{2^n})$ . By Remark 1, the vertex  $v'_{2^n} \in V(\hat{H}_{n+2})$  is identified with the vertex  $v_{2^n - (0) + 2^n} = v_{2^{n+1}}$ , thus the arc  $v_0 v'_{2^n} \in A(\hat{H}_{n+2})$  is transformed into the arc  $v_0 v_{2^{n+1}} \in A(H_{n+2})$ . Therefore,  $H_{n+2} = D(C_{2^{n+1}+1})$ .

Observe that we use only one directed Hajós operation (a directed Hajós join) to obtain the digraph  $H_0$ . For each of the digraphs  $H_1, H_2, \dots, H_{n+2}$  we use  $2^{n+1} + 1$  directed Hajós operations (a directed Hajós join and  $2^{n+1}$  identifications), thus we use  $(n+2)(2^{n+1} + 1) + 1$  directed Hajós operations.  $\square$

The Hajós number of an  $r$ -dichromatic digraph  $H$  was defined in [19] as the minimum number of Hajós operations needed to obtain  $H$  from  $D(K_r)$ .

**Theorem 3.** *Let  $n \geq 2$ . The Hajós number of a symmetric cycle of order  $2^n + 1$  is at most*

$$\frac{n(2^{n+2} + n + 5)}{2} - 7.$$

*Moreover, the Hajós number of a symmetric cycle of order  $2m + 1$ , where  $2^{n-1} + 1 < 2m + 1 < 2^n + 1$ , is at most*

$$\frac{n(2^{n+2} + n + 5)}{2} - 5.$$

*Proof.* By recursively applying Theorem 2, the number of steps to build the symmetric cycle of order  $2^n + 1$  from  $D(K_3)$  is at most:

$$\sum_{i=2}^n (i+1)(2^i + 1) + 1.$$

Note that

$$\begin{aligned}\sum_{i=1}^n i2^i &= n(2^{n+1} - 2) - \sum_{j=1}^{n-1} (2^{j+1} - 2) \\ &= (n-1)(2^{n+1}) + 2.\end{aligned}$$

Thus,

$$\begin{aligned}\sum_{i=1}^n (i+1)(2^i+1) + 1 &= \sum_{i=1}^n i2^i + \sum_{i=1}^n i + \sum_{i=1}^n 2^i + \sum_{i=1}^n 1 + \sum_{i=1}^n 1 \\ &= (n-1)2^{n+1} + 2 + \frac{n(n+1)}{2} + (2^{n+1} - 2) + 2n \\ &= \frac{n}{2}(2^{n+2} + n + 5).\end{aligned}$$

Since

$$\sum_{i=2}^n (i+1)(2^i+1) + 1 = \sum_{i=1}^n (i+1)(2^i+1) + 1 - 7,$$

the first result follows.

Consider a symmetric cycle of order  $2^n+1$ . Let  $1 \leq m < 2^{n-1}$ , let  $V_{2m} = \{v_{2m}, v_{2m+2}, \dots, v_{2^n}\}$  and let  $V_{2m+1} = \{v_{2m+1}, v_{2m+3}, \dots, v_{2^n+1}\}$ , note that  $V_{2m}$  and  $V_{2m+1}$  are both independent sets of vertices. Identifying the independent set of vertices  $V_{2m}$  into the vertex  $v_{2m}$  and identifying the independent set of vertices  $V_{2m+1}$  into the vertex  $v_{2m+1}$ , we obtain the symmetric cycle of order  $2m+1$ . Thus, we can construct the symmetric cycle of order  $2m+1$  using  $\frac{n(2^{n+2} + n + 5)}{2} - 5$  directed Hajós operations and the result follows.  $\square$

### 3 Computational Complexity

In this section we determine the complexity of our procedure and which is an upper bound for the complexity of constructing a symmetric odd cycle using directed Hajós operations.

**Theorem 4.** *Let  $n \geq 5$  be an odd integer. The complexity of the procedure to obtain  $D(C_n)$  from  $D(K_3)$  is  $\Theta(n \ln(n))$ .*

*Proof.* Let  $m \geq 2$  and let  $n$  be an odd integer such that  $n_1 < n \leq n_2$  with

$$n_1 = 2^{m-1} + 1 \text{ and } n_2 = 2^m + 1.$$

With our procedure, one needs

$$X = \frac{m}{2}(2^{m+2} + m + 5) - \alpha \tag{3.1}$$

directed Hajós operations to obtain a symmetric cycle of order  $n$  from  $D(K_3)$ , where

$$\alpha = \begin{cases} 5 & \text{if } n < n_2, \\ 7 & \text{if } n = n_2. \end{cases}$$

Since  $n_1 = 2^{m-1} + 1$ , then

$$\begin{aligned}n_1 - 1 &= 2^{m-1} \\ \log_2(n_1 - 1) &= m - 1,\end{aligned}$$

therefore

$$m = \log_2(n_1 - 1) + 1. \quad (3.2)$$

In equation 3.1 we have that

$$\begin{aligned} X &= \frac{\log_2(n_1 - 1) + 1}{2} (2^{(\log_2(n_1 - 1) + 1) + 2} + (\log_2(n_1 - 1) + 1) + 5) - \alpha \\ &= \frac{\log_2(n_1 - 1) + 1}{2} (8n_1 + \log_2(n_1 - 1) - 2) - \alpha \\ &= 4n_1 \log_2(n_1 - 1) + \frac{(\log_2(n_1 - 1))^2}{2} - \log_2(n_1 - 1) \\ &\quad + 4n_1 + \frac{\log_2(n_1 - 1)}{2} - 1 - \alpha \\ &< 9n_1 \log_2(n_1 - 1) < 9n \log_2(n) = \left( \frac{9}{\ln(2)} \right) n \ln(n) < 13n \ln(n). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} X &= 4n_1 \log_2(n_1 - 1) + \frac{(\log_2(n_1 - 1))^2}{2} - \log_2(n_1 - 1) \\ &\quad + 4n_1 + \frac{\log_2(n_1 - 1)}{2} - 1 - \alpha \\ &> 4n_1 \log_2(n_1 - 1) \\ &> 2n_1 \log_2((n_1 - 1)^2) = (2^m + 2) \log_2(2^{m+1}) \\ &> (2^m + 1) \log_2(2^m + 1) = n_2 \log_2(n_2) \\ &\geq n \log_2(n) > n \ln(n). \end{aligned}$$

Therefore  $n \ln(n) < X < 13n \ln(n)$  and the complexity of our procedure is  $\Theta(n \ln(n))$ .  $\square$

## 4 Conclusions

Although it was proved in [3] that any 3-critical digraph can be constructed by a sequence of Hajós operations, it is not a trivial task to obtain such a sequence even for simple digraphs such as symmetric cycles of odd length and in particular, the symmetric cycle of length 5. Using genetic algorithm, we obtained in [11], a sequence of Hajós operations for the symmetric cycle of length 5, and generalizing this particular result permitted us to construct any symmetric odd cycle using Hajós operations.

We believe that these ideas can be used in order to obtain the 3- and 4-critical tournaments characterized in [21] and hopefully a improvement of the upper bound for the minimum order of a 5-critical tournament, which is known to be 19 (see 4.6. An application [22]).

## References

- [1] D. Andres, W. Hochstattler. Perfect digraphs *J. Graph Theory* **79**, 21–29 (2015).
- [2] G. Araujo-Pardo, J. J. Montellano-Ballesteros, M. Olsen, C. Rubio-Montiel. The dichromatic number of digraphs. *Electr. J. Comb.* **25**(3), #P3.51 (2018).
- [3] J. Bang-Jensen, T. Bellitto, T. Schweser, M. Stiebitz. Hajós and Ore constructions for digraphs. *Electr. J. Comb.* **27**(1), #P1.63 (2020).
- [4] J. Bang-Jensen and G. Gutin. *Digraphs: Theory, Algorithms, and Applications*, 2nd edn, Springer-Verlag, London, 2009.
- [5] D. Bokal, G. Fijavz, M. Juvan, P.M. Kayll, B. Mohar. The circular chromatic number of a digraph. *J. Graph Theory* **46**, 227–240 (2004).
- [6] K. Bouazzi, M. Hammami, S. Bouamama. Application of an improved genetic algorithm to Hamiltonian circuit problem, *Procedia Comput. Sci.* **192**, 4337–4347 (2021).
- [7] J. Cervantes-Ojeda, M. Gómez-Fuentes, D. González-Moreno, and M. Olsen. Rainbow Connectivity Using a Rank Genetic Algorithm: Moore Cages with Girth Six. *Hindawi, J. Appl. Math.* **2019**, Article ID 4073905, 7 pages (2019).
- [8] J. Cervantes, C. R. Stephens, Limitations of existing mutation rate heuristics and how a rank GA overcomes them. *IEEE Transactions on Evolutionary Computation* **13**(2), 369–397 (2009).
- [9] N. Cordero-Michel, H. Galeana-Sánchez. New Bounds for the Dichromatic Number of a Digraph. *Discrete Math. Theor. Comput. Sci.* **21**(1), Paper No. 7, 14 pp (2019).
- [10] D. Dasgupta, Z. Michalewicz. *Evolutionary algorithms in engineering applications*. Springer Science and Business Media (2013).
- [11] J. C. García-Altamirano, M. Olsen, J. Cervantes-Ojeda. How to construct the symmetric cycle of length 5 using Hajós construction with an adapted Rank Genetic Algorithm (submitted) <https://doi.org/10.48550/arXiv.2210.05080>.
- [12] D. González-Moreno, R. Hernández-Ortiz, B. Llano, M. Olsen. The dichromatic polynomial of a digraph, *Graphs and Comb.* **38**:85 (2022).
- [13] G. Hajós. Über eine Konstruktion nicht  $n$ -färbbarer Graphen. *Wiss. Z. Martin Luther Univ. Halle-Wittenberg, Math. Natur. Reihe* **10**, 116–117 (1961).
- [14] A. Harutyunyan, B. Mohar. Strengthened brooks theorem for digraphs of girth at least three. *Electronic J. Comb.* **18**(1), #P195 (2011).
- [15] W. Hochstättler. A flow theory for the dichromatic number. *European J. of Comb.* **66**, 160–167 (2017).
- [16] S. Jakobs. On genetic algorithms for the packing of polygons. *Eur. J. Oper. Res.* **88**(1), 165–181 (1996).
- [17] T. R. Jensen, G. F. Royle. Hajós constructions of critical graphs, *J. Graph Theory* **30**(1), 37–50 (1999).



- [18] K. A. De Jong, W. M. Spears. Using genetic algorithms to solve NP-complete problems. In ICGA pp. 124–132 (1989, June).
- [19] A. J. Mansfield, D. J. A. Welsh. Some colouring problems and their complexity, Graph theory (Cambridge, 1981), North-Holland Math. Stud. **62**, Amsterdam: North-Holland, pp. 159–170 (1982).
- [20] V. Neumann-Lara. The dichromatic number of a digraph. J. Combin. Theory Ser. B **33**, 265–270 (1982).
- [21] V. Neumann-Lara. The 3- and 4-chromatic tournaments of minimum order. Discrete Math. **135**, 233–243 (1994).
- [22] V. Neumann-Lara. Dichromatic number, circulant tournaments and Zykov sums of digraphs. Discuss. Math. Graph Theory **20**(2) 197–207 (2000).
- [23] A. Pourrajabian, R. Ebrahimi, M. Mirzaei, M. Shams. Applying genetic algorithms for solving nonlinear algebraic equations. Appl. Math. Comput. **219**(24), 11483–11494 (2013).
- [24] A. Urquhart. The graph constructions of Hajós and Ore. J. Graph Theory **26** 211–215 (1997).