

**ON A FAMILY OF 2-AUTOMATIC SEQUENCES GENERATING  
ALGEBRAIC CONTINUED FRACTIONS IN  
CHARACTERISTIC 2**

YINING HU

ABSTRACT. We present family of automatic sequences that define algebraic continued fractions in characteristic 2. This family is constructed from ultimately period words and contains the period-doubling sequence.

1. INTRODUCTION

Ultimately periodic sequences are the simplest of all sequences. In this article we study a family  $\mathcal{S}$  of automatic sequences generated by ultimately periodic sequences and the continued fractions defined by them.

Each sequence  $\mathbf{s}$  in  $\mathcal{S}$  is built from an ultimately periodic sequence  $(\varepsilon_n)_n = p_0 \dots p_{l-1}(e_0 \dots e_{d-1})^\infty$  in the following way. Starting from the empty word  $W_0$ , we consider the sequence of words  $(W_n)_{n \geq 0}$  such that  $W_{n+1} = W_n, \varepsilon_n, W_n$  (note the comma is for concatenation and it will be omitted when it is suitable). Hence, we have  $W_1 = \varepsilon_0$ ,  $W_2 = \varepsilon_0, \varepsilon_1, \varepsilon_0$ , etc. Note that, for all  $n \geq 1$ ,  $W_n$  is a palindrome, centred in  $\varepsilon_n$ , of length  $2^n - 1$ .

By construction, the word  $W_{n+1}$  starts by  $W_n$  and and we define  $\mathbf{s}(\varepsilon)$  to be the limit of  $(W_n)$ . Hence, we have

$$\mathbf{s}(\varepsilon) = \varepsilon_0, \varepsilon_1, \varepsilon_0, \varepsilon_2, \varepsilon_0, \varepsilon_1, \varepsilon_0, \varepsilon_3, \varepsilon_0, \varepsilon_1, \varepsilon_0, \varepsilon_2, \dots$$

We illustrate the construction of  $\mathbf{s}(\varepsilon)$  with two basic cases.

1)  $\varepsilon = a, b, (c)^\infty$  then

$$\mathbf{s}(\varepsilon) = a, b, a, c, a, b, a, c, \dots = (a, b, a, c)^\infty.$$

Here  $\varepsilon$  is ultimately constant and  $\mathbf{s}(\varepsilon)$  is periodic. Consequently a basic property on continued fractions shows that  $\text{CF}(\mathbf{s}(\varepsilon))$  is quadratic over  $\mathbb{F}_2(t)$ .

2)  $\varepsilon = (a, b)^\infty$  then

$$\mathbf{s}(\varepsilon) = a, b, a, a, a, b, a, b, a, b, \dots$$

Here  $\mathbf{s}(\varepsilon)$  is the celebrated period-doubling sequence. It has been proved that  $\text{CF}(\mathbf{s}(\varepsilon))$  satisfies an algebraic equation of degree 4 (see [7, 8]).

Each sequence in  $\mathcal{S}$  is 2-automatic. The definition and relevant results about automatic sequences are briefly recalled in Section 1.3.

Continued fractions defined by two classical example of automatic sequences in characteristic 2 were studied in [7, 5, 8], where it was established that the Thue-Morse and period-doubling sequences both define algebraic continued fractions of

---

2010 *Mathematics Subject Classification.* 11B85, 11J70, 11B50, 11Y65, 05A15, 11T55.

*Key words and phrases.* algebraicity, automatic sequence, continued fraction, period-doubling sequence.

This work is supported by the National Science Foundation of China grant 12001216.

degree 4. On the other hand, both sequences define quadratic formal power series. The situation is different from the real case: Bugeaud [4] proved in 2013 that the continued fraction expansion of an algebraic real number of degree at least 3 is not automatic.

Our family  $\mathcal{S}$  is a natural generalization of the period-doubling sequence. We prove in this article that every sequence from this family defines an algebraic continued fraction in characteristic 2.

In Section 1.1 we give the notation necessary for stating our main result in 1.2. In Section 2 we prove the main theorem, and in Section 3 we give three examples. Finally, in Section 4 we make a comment on the link between continued fractions and Riccati differential equations.

**1.1. Continued Fractions in Power Series Rings.** Let  $A = \{a_0, \dots, a_k\}$  be a finite alphabet. We treat  $a_j$  as formal variables and define

$$\begin{aligned}\mathbb{F}_2[A] &:= \mathbb{F}_2[a_0, \dots, a_k] \\ \mathbb{F}_2(A) &:= \mathbb{F}_2(a_0, \dots, a_k) \\ \mathbb{F}_2((A)) &:= \mathbb{F}_2\left(\frac{1}{a_0}, \dots, \frac{1}{a_k}\right).\end{aligned}$$

Here  $\mathbb{F}_2\left(\frac{1}{a_0}, \dots, \frac{1}{a_k}\right)$  denotes the ring of power series of the form

$$(1.1) \quad \sum_{n_0, \dots, n_k \geq N} c_{n_0, n_1, \dots, n_k} a_0^{-n_0} \dots a_k^{-n_k},$$

where  $N$  is an integer and  $c_{n_0, n_1, \dots, n_k} \in \mathbb{F}_2$ .

Let  $(u_n)_{n \geq 0}$  be a sequence taking values in  $A$ . It defines a formal power series  $\sum_{n \geq 0} u_n z^n$  in  $\mathbb{F}_2[A][[z]]$ .

We define a norm on  $\mathbb{F}_2((A))$  by assigning a series of the form (1.1) the number  $2^{-m}$ , where  $m = \min\{n_0 + n_1 + \dots + n_k \mid c_{n_0, n_1, \dots, n_k} \neq 0\}$  with the convention that  $\min \emptyset = \infty$ . This norm makes  $\mathbb{F}_2((A))$  an ultrametric space.

The continued fraction  $\text{CF}(\mathbf{u}) = [u_0, u_1, \dots]$  is defined as the limit of the sequence  $([u_0, u_1, \dots, u_n])_n$ :

$$\begin{aligned}[u_0] &= u_0, \\ [u_0, u_1, \dots, u_n] &= u_0 + \frac{1}{[u_1, \dots, u_n]} \in \mathbb{F}_2((A)),\end{aligned}$$

for  $n \geq 1$ . For example,

$$\begin{aligned}[u_0, u_1, u_2] &= u_0 + \frac{1}{u_1 + \frac{1}{u_2}} \\ &= u_0 + \frac{u_1^{-1}}{1 + (u_1 u_2)^{-1}} \\ &= u_0 + u_1^{-1} + u_1^{-2} u_2^{-1} + u_1^{-3} u_2^{-2} + \dots\end{aligned}$$

The convergence of this sequence is proved in the same way as in the case of classical continued fraction for real numbers.

We may also choose to specialize the letters in  $A$  to non-constant polynomials in  $\mathbb{F}_2[z]$  (this gives a continuous map from  $\mathbb{F}_2((A))$  to  $\mathbb{F}_2((1/z))$ ), and then  $\text{CF}(\mathbf{u})$  can be seen as formal power series in  $\mathbb{F}_2((1/z))$ . For basic information on continued fractions, particularly in power series rings, the reader is referred to [9].

1.2. **Main result.** We prove the following theorem.

**Theorem 1.1.** *Let  $\varepsilon = p_0 \dots p_{l-1}(e_0 \dots e_{d-1})^\infty$  be an ultimately periodic sequence over the alphabet  $A$ .*

(i) *The continued fraction  $\text{CF}(\mathbf{s}(\varepsilon))$  is algebraic of degree at most  $2^d$  over  $\mathbb{F}_2(A)$ . It is of degree  $2^d$  if  $e_0$  is different from the other letters.*

(ii) *The formal power series*

$$F := \sum_{j \geq 0} s_j z^j \in \mathbb{F}_2(A)[[z]].$$

*is algebraic of degree at most  $2^{d-1}$  over  $\mathbb{F}_2(A)(z)$ . It is of degree  $2^{d-1}$  if  $e_0$  is different from the other letters.*

**Remark 1.** For the formal power series in in Theorem 1.1 and in the examples, we chose to keep the letters of an automatic sequence as formal variables instead of mapping them to elements in  $\mathbb{F}_p$ , because in the latter case, the formal power series and consequently its minimal polynomial depend on the mapping. For an automatic sequence  $\mathbf{u}$  taking values  $a_1, \dots, a_k$ , it is always possible to treat the letters as formal variables and get a minimal polynomial in  $\mathbb{F}_p[a_1, \dots, a_k, x][y]$ . This is because  $F$  can be decomposed into a finite sum  $\sum a_n F_n$ , where each  $F_n$  is algebraic over  $\mathbb{F}_p(x)$ . Therefore each  $a_n F_n$  is algebraic over  $\mathbb{F}_p(a_1, \dots, a_k)(x)$  and so is  $F$ .

1.3. **Automatic sequences and algebraicity.** Here we recall briefly the definition and properties of automatic sequences. We refer the readers to [2] for a comprehensive exposition on the subject. A sequence  $(u_n)_n$  is said to be  $k$ -automatic if it can be generated by a finite  $k$ -automaton. In this article we will use an equivalent definition. The  $k$ -kernel of a sequence  $(u_n)_n$  is defined as the smallest set containing  $(u_n)_n$  and stable under the operations  $(v_n)_n \mapsto (v_{kn+r})_n$  for  $r = 0, 1, \dots, k-1$ . The sequence  $\mathbf{u}$  is said to be  $k$ -automatic if its  $k$ -kernel is finite.

The well-known Theorem of Christol, Kamae, Mendès France and Rauzy [6] states that a formal power series in  $\mathbb{F}_p[[x]]$  is algebraic over  $\mathbb{F}_p(x)$  if and only if the sequence of its coefficients is  $p$ -automatic. This theorem is generalized by Salon [10] to multivariate cases: We define the  $k$ -kernel of a multi-dimensional sequence  $\mathbf{u} = (u_{n_1, \dots, n_d})_{n_1, \dots, n_d}$  to be the smallest set containing  $\mathbf{u}$  and stable under the operations

$$(v_{n_1, \dots, n_d})_{n_1, \dots, n_d} \mapsto (v_{k \cdot n_1 + r_1, \dots, k \cdot n_d + r_d})_{n_1, \dots, n_d}$$

for  $0 \leq r_1, \dots, r_d < k$ . The sequence  $\mathbf{u}$  is said to be  $k$ -automatic if its  $k$ -kernel is finite. When  $\mathbf{u}$  takes values in  $\mathbb{F}_p$ , then the formal power series  $\sum u_{n_1, \dots, n_d} x_1^{n_1} \cdot x_d^{n_d}$  is algebraic over  $\mathbb{F}_p(x_1, \dots, x_d)$  if and only if  $\mathbf{u}$  is  $k$ -automatic.

For  $r = 0, 1, \dots, p-1$ , the Cartier operators  $\Lambda_r$  are defined as

$$\begin{aligned} \Lambda_r : \mathbb{F}_p[[x]] &\rightarrow \mathbb{F}_p[[x]] \\ \sum_{n=0}^{\infty} u_n x^n &\mapsto \sum_{n=0}^{\infty} u_{pn+r} x^n. \end{aligned}$$

Thus for  $f \in \mathbb{F}_p[[x]]$ , we have

$$f = \sum_{r=0}^{p-1} x^r (\Lambda_r f)^p.$$

In the multivariate case, the Cartier operators are

$$\begin{aligned} \Lambda_{r_1, \dots, r_d} : \mathbb{F}_p[[x_1, \dots, x_d]] &\rightarrow \mathbb{F}_p[[x_1, \dots, x_d]] \\ \sum u_{n_1, \dots, n_d} x_1^{n_1} \cdots x_d^{n_d} &\mapsto \sum u_{pn_1+r_1, \dots, pn_d+r_d} x_1^{n_1} \cdots x_d^{n_d}. \end{aligned}$$

For  $f \in \mathbb{F}_p[[x_1, \dots, x_d]]$ , we have

$$f = \sum_{r_1, \dots, r_d=0}^{p-1} x_1^{r_1} \cdots x_d^{r_d} (\Lambda_{r_1, \dots, r_d} f)^p.$$

The following lemma will be used in the proof of theorem 1.1:

**Lemma 1.2.** *Let  $\mathbf{u} = (u_{n_1, \dots, n_d})_{n_1, \dots, n_d}$  be a  $p$ -automatic sequence taking value in  $\mathbb{F}_p$ . Let  $\mathbf{v}$  be an element from the  $p$ -kernel of  $\mathbf{u}$ , then the degree of algebraicity of  $\sum v_{n_1, \dots, n_d} x_1^{n_1} \cdots x_d^{n_d}$  is less than or equal to that of  $\sum u_{n_1, \dots, n_d} x_1^{n_1} \cdots x_d^{n_d}$ .*

*Proof.* Let  $f = \sum u_{n_1, \dots, n_d} x_1^{n_1} \cdots x_d^{n_d}$ . We only need to prove that for all  $(r_1, \dots, r_d) \in \{0, 1, \dots, p-1\}^d$ , for  $\Lambda := \Lambda_{r_1, \dots, r_d}$ , the degree of algebraicity of  $\Lambda f$  is less than or equal to that of  $f$ . From the proof of Christol's theorem we know that there exists  $n \geq 0$  and  $a_1, a_2, \dots, a_n \in \mathbb{F}_p(x_1, \dots, x_d)$  such that

$$f = a_1 f^{p^1} + a_2 f^{p^2} + \cdots + a_n f^{p^n}.$$

We make use of the following property of  $\Lambda$ : for two power series  $g$  and  $h$ , we have

$$\Lambda(g \cdot h^p) = \Lambda(g) \cdot h,$$

and

$$\Lambda \frac{g}{h} = \Lambda \frac{g \cdot h^{p-1}}{h^p} = \frac{\Lambda(g \cdot h^{p-1})}{h}.$$

Therefore

$$\Lambda f = \Lambda(a_1) f + \Lambda(a_2) f^p + \cdots + \Lambda(a_n) f^{p^{n-1}} \in \mathbb{F}_p(x_1, \dots, x_d)[f].$$

We conclude that the degree of  $\Lambda f$  is less than or equal to that of  $f$ .  $\square$

## 2. PROOF OF THEOREM 1.1

To prove of the first part of theorem 1.1, we consider the truncation  $[s_0, \dots, s_{2^n-2}]$  of the continued fraction  $\text{CF}(\mathbf{s}(\varepsilon))$  for  $n \geq 1$ . This corresponds to the palindrome  $W_n$ . For  $n \geq 1$ , we set  $u_n/v_n = [s_0, \dots, s_{2^n-2}]$ . We have  $(u_1, v_1) = (s_0, 1)$  and  $(u_2, v_2) = (s_0 s_1 s_2 + s_0 + s_2, s_1 s_2 + 1)$ , etc. The first step of the proof was introduced in the author's previous work with Lasjaunias concerning the particular case of the period-doubling sequence. In Lemma 3.2 from [8], using basic properties of continuants, we proved that the pair  $(u_n, v_n)$  satisfies a simple recurrence relation. Indeed, for  $n \geq 1$ , we have

$$(R) \quad u_{n+1} = \varepsilon_n u_n^2 \text{ and } v_{n+1} = \varepsilon_n u_n v_n + 1.$$

From (R) we get immediately

$$\frac{v_{n+1}}{u_{n+1}} = \frac{v_n}{u_n} + \frac{1}{u_n}$$

and therefore we obtain

$$\frac{v_n}{u_n} = \sum_{1 \leq k \leq n} \frac{1}{u_k}$$

for  $n \geq 1$ . Consequently

$$(\star) \quad \frac{1}{\text{CF}(\mathbf{s}(\varepsilon))} = \lim_{n \rightarrow \infty} \frac{v_n}{u_n} = \sum_{n \geq 1} \frac{1}{u_n}.$$

It is easier to work with the following series which differs by a rational fractional from  $1/\text{CF}(\mathbf{s}(\varepsilon))$ :

$$G := \sum_{n > l} \frac{1}{u_n}.$$

By regrouping every  $d$  terms and using (R), we decompose  $G$  as follows

$$(2.1) \quad G = \sum_{n=0}^{d-1} \frac{G_n^2}{e_n},$$

where

$$G_n = \sum_{k=0}^{\infty} \frac{1}{u_{l+n+kd}}$$

for  $n = 0, 1, \dots, d-1$ . It follows from their definition and (R) that  $\{G_n\}$  satisfy the following relations

$$(2.2) \quad G_0 = \frac{1}{u_l} + \frac{G_{d-1}^2}{e_{d-1}}$$

and

$$(2.3) \quad G_n = \frac{G_{n-1}^2}{e_{n-1}}$$

for  $n = 1, 2, \dots, d-1$ , with the convention  $u_0 = 1$ , from which we deduce

$$(2.4) \quad G_0 = \frac{1}{u_l} + \frac{G_0^{2^d}}{e_{d-1}e_{d-2}^2 \cdots e_0^{2^{d-1}}}.$$

Using (2.1), (2.3) and (2.4), we can express  $G$  in terms of  $G_0$

$$G = \frac{1}{u_l} + G_0 + \frac{G_0^2}{e_0} + \frac{G_0^4}{e_0^2 e_1} + \cdots + \frac{G_0^{2^{d-1}}}{e_0^{2^{d-2}} e_1^{2^{d-3}} \cdots e_{d-2}}.$$

This means that the degree algebraicity of  $G$  (and that of  $\text{CF}(\mathbf{s}(\varepsilon))$ ) is at most that of  $G_0$ . This proves that the degree of algebraicity of  $\text{CF}(\mathbf{s}(\varepsilon))$  is at most  $2^d$ .

When  $e_0$  is different from all other letters,  $G_0$  is in the 2-kernel of  $G$ , so that by lemma 1.2, the degree of  $G$  is at least that of  $G_0$ . To show that degree of algebraicity of  $G_0$  is at least  $2^d$ , consider the series

$$g = \sum_{k=0}^{\infty} \frac{1}{e_0^{2^{k \cdot d}}}$$

that satisfies

$$e_0 \cdot g^{2^d} + e_0 \cdot g + 1 = 0.$$

This is an irreducible polynomial by Eisenstein's criterion. If we project all the letters in  $G_0$  to  $e_0$ , then we get the series  $e_0 \cdot \sum_{k=l}^{\infty} \frac{1}{e_0^{2^{k \cdot d}}}$ , which has the same degree of algebraicity as  $g$ . Since projection cannot increase degree, we conclude that in this case, the degree of algebraicity of  $G_0$ , and therefore that of  $\text{CF}(\mathbf{s}(\varepsilon))$  is  $2^d$ .

To prove the second part of the theorem, we need the following lemma

**Lemma 2.1.** *Suppose that the letters  $e_j$  ( $0 \leq j < d$ ) are all distinct and that they are different from the letters  $p_j$  ( $0 \leq j < l$ ). For  $j = 0, 1, \dots, d-1$ , define*

$$P_j := \{k \mid s_k = e_j\}.$$

Then

$$(2.5) \quad P_{j+1} = \{2k+1 \mid k \in P_j\} \quad \text{for } j = 0, 1, \dots, d-2$$

and

$$(2.6) \quad P_0 = \{2k+1 \mid k \in P_{d-1}\} \cup \{(2k+1) \cdot 2^l - 1 \mid k = 1, 2, \dots\}.$$

*Proof.* To simplify notation, we write  $e_n$  instead of  $e_{n \pmod{d}}$  when  $n \geq d$ . By definition,  $\mathbf{s}(\varepsilon)$  can be written in two ways:

$$(2.7) \quad W_l e_0 W_l e_1 W_l e_0 W_l e_2 W_l e_0 W_l e_1 W_l e_0 W_l \dots$$

and

$$(2.8) \quad W_{l+1} e_1 W_{l+1} e_2 W_{l+1} e_1 W_{l+1} e_3 W_{l+1} e_1 W_{l+1} e_2 W_{l+1} e_1 W_{l+1} \dots$$

Letter  $e_0$  appears at the same places in (2.7) as  $e_1$  in (2.8), except that the length of the block  $W_l$  is  $2^l - 1$ , while that of  $W_{l+1}$  is  $2^{l+1} - 1$ . By the assumption on  $(e_j)$  and  $(p_j)$ , the letter  $e_j$  does not appear in  $W_{l+j}$  ( $0 \leq j < d$ ). The above observation allows us to conclude that

$$P_1 = \{2k+1 \mid k \in P_0\}.$$

Similar argument shows that

$$P_{j+1} = \{2k+1 \mid k \in P_j\} \quad \text{for } j = 1, \dots, d-2.$$

To prove (2.6), we also write  $\mathbf{s}(\varepsilon)$  in two ways:

$$(2.9) \quad W_{l+d-1} e_{d-1} W_{l+d-1} e_0 W_{l+d-1} e_{d-1} W_{l+d-1} e_1 W_{l+d-1} e_{d-1} W_{l+d-1} e_0 \dots$$

and

$$(2.10) \quad W_{l+d} e_0 W_{l+d} e_1 W_{l+d} e_0 W_{l+d} e_2 W_{l+d} e_0 W_{l+d} e_1 W_{l+d} e_0 W_{l+d} \dots$$

As with previous cases, the letter  $e_0$  appears at the same places in (2.10) as  $e_{d-1}$  in (2.9), except that the length of the block  $W_{l+d}$  is  $2^{l+d} - 1$ , while that of  $W_{l+d-1}$  is  $2^{l+d-1} - 1$ . Unlike previous cases, the letter  $e_0$  also occurs in  $W_{l+d}$  at indices  $(2k+1) \cdot 2^l - 1$  for  $k = 1, 2, \dots$ . That's how we get the additional term in (2.6).  $\square$

We consider another sequence  $\varepsilon' = p'_0 \dots p'_{l-1} (e'_0 \dots e'_{d-1})^\infty$ , where the letters  $p'_j$  ( $0 \leq j < l$ ) and  $e'_j$  ( $0 \leq j < d$ ) are all distinct, so that we can apply lemma 2.1 to obtain the sets  $P'_j$  for  $j = 0, 1, \dots, d-1$  corresponding to the sequence  $\mathbf{s}(\varepsilon')$ .

We decompose the series  $F$  as follows

$$(2.11) \quad F = R + e_0 F_0 + \dots + e_{d-1} F_{d-1},$$

where

$$R = \frac{\sum_{k=0}^{2^l-2} s_k z^k}{1 + z^{2^l}}$$

is a rational function which accounts for the preperiodic part of  $\varepsilon$  and

$$F_n = \sum_{j \in P_n} z^j$$

for  $n = 0, 1, \dots, d-1$ .

From lemma 2.1 we deduce the relations

$$(2.12) \quad F_n = zF_{n-1}^2$$

for  $n = 1, \dots, d-1$ . On the other hand, since the supports of  $R$  and  $F_0, \dots, F_{d-1}$  form a partition of  $\mathbb{N}$ , we have

$$(2.13) \quad F_0 = \frac{1}{1+z} - \frac{\sum_{k=0}^{2^l-2} z^k}{1+z^{2^l}} - \sum_{n=1}^{d-1} F_n = \frac{z^{2^l-1}}{1+z^{2^l}} - \sum_{n=1}^{d-1} F_n.$$

From (2.12) and (2.13) we deduce

$$(2.14) \quad F_0 = \frac{z^{2^l-1}}{1+z^{2^l}} - \sum_{n=1}^{d-1} z^{2^n-1} F_0^{2^n}.$$

Using (2.11) and (2.12) we can express  $F$  using  $F_1$

$$(2.15) \quad F = R + \sum_{n=0}^{d-1} e_n z^{2^n-1} F_0^{2^n}.$$

This shows that the degree of  $F$  is at most that of  $F_0$ .

The degree of  $F_0$  is  $2^{d-1}$ : if  $l$  were 0, in place of  $F_0$ , by 2.11, we would have a series  $f$  that satisfies

$$f = \frac{1}{1+z} - \sum_{n=1}^{d-1} z^{2^n-1} f^{2^n}.$$

Therefore the series  $h := zf$  satisfy

$$(1+z) \cdot \sum_{n=0}^{d-1} h^{2^n} + z = 0,$$

which is an irreducible polynomial in  $h$  by Eisenstein's criterion.

The support of  $f$  is the positions of  $e_0$  in

$$\mathbf{s}((e_0 e_1 \dots e_{d-1})^\infty) = W_0 e_0 W_0 e_1 W_0 e_0 W_0 e_2 W_0 e_0 W_0 e_1 W_0 e_0 W_0 \dots$$

Comparing the above expression with (2.7) we obtain

$$F_0 = z^{2^l-1} f^{2^l}.$$

Therefore  $f$  is in the 2-kernel of  $F_0$ . By lemma 1.2 the degree of  $F_0$  is at least that of  $f$ .

When  $e_0$  is different from the other letters, we can obtain  $F_0$  from  $F$  mapping  $e_0$  to 1 and the other letters to 0. This show that in this case the degree of  $F$  is at least that of  $F_0$ .

### 3. EXAMPLES

**Example 1.** Let  $\varepsilon = (ab)^\infty$ . Then  $\mathbf{s}(\varepsilon)$  is the period-doubling sequence. We have

$$\begin{aligned} \frac{1}{\text{CF}(\mathbf{s})} = G &= \frac{1}{a} + \frac{1}{a^2 b} + \frac{1}{a^4 b^2 a} + \frac{1}{a^8 b^4 a^2 b} + \frac{1}{a^{16} b^8 a^4 b^2 a} \dots \\ &= \frac{G_0^2}{a} + \frac{G_1^2}{b}, \end{aligned}$$

with

$$G_0 = 1 + \frac{1}{a^2 b} + \frac{1}{a^8 b^4 a^2 b} + \dots$$

$$G_1 = \frac{1}{a} + \frac{1}{a^4 b^2 a} + \frac{1}{a^{16} b^8 a^4 b^2 a} \cdots$$

The series  $G_0$  and  $\text{CF}(\mathbf{s})$  satisfy

$$G_0 = 1 + \frac{G_0^4}{a^2 b},$$

and

$$\frac{1}{\text{CF}(\mathbf{s})} = G = 1 + G_0 + \frac{G_0^2}{a}.$$

We get as expected (see [8])

$$(ab + b^2 + 1) + (a^2 b + ab^2) \cdot G + ab \cdot G^2 + G^4 = 0$$

The formal power series defined by  $\mathbf{s}(\varepsilon)$  is

$$F = a + bz + az^2 + az^3 + az^4 + bz^5 + az^6 + \cdots = aF_0 + bF_1.$$

The series  $F_0$  and  $F$  satisfy

$$\begin{aligned} F_0 &= \frac{1}{1+z} - zF_0^2, \\ F &= aF_0 + bzF_0^2. \end{aligned}$$

We deduce

$$(z^3 + z) \cdot F^2 + (az^2 + bz^2 + a + b) \cdot F + (a^2 z + abz + b^2 z + a^2 + ab) = 0.$$

**Remark 2.** Here we have a simple example of an algebraic formal power series whose continued fraction has unbounded partial quotients. If we let  $a = b = x$  in  $G_0$ , we get a series

$$g = 1 + \frac{1}{x} + \frac{1}{x^5} + \frac{1}{x^{21}} + \cdots,$$

which satisfies the algebraic equation

$$\frac{g^4}{x} + g + 1 = 0.$$

It can be shown that

$$g = [1, x, x^3, x, x^{11}, x, x^3, x, x^{43}, \dots] = [1, x^{c_0}, x^{c_1}, x^{c_2}, \dots],$$

where  $c_{2n} = 1$  and  $c_{2n+1} = 4c_n - 1$  for all  $n \geq 0$ . The exponents of partial quotients are not bounded, but they form nonetheless a 2-regular sequence (see [1]).

**Example 2.** Let  $\varepsilon = a(bc)^\infty$ . Then  $\mathbf{s}(\varepsilon) = abacabababacabacabac \cdots$ . We have

$$\frac{1}{\text{CF}(\mathbf{s})} = \frac{1}{a} + G,$$

where

$$\begin{aligned} G &= \frac{1}{a^2 b} + \frac{1}{a^4 b^2 c} + \frac{1}{a^8 b^4 c^2 b} + \frac{1}{a^{16} b^8 c^4 b^2 c} + \frac{1}{a^{32} b^{16} c^8 b^4 c^2 b} + \cdots \\ &= \frac{G_0^2}{b} + \frac{G_1^2}{c}, \end{aligned}$$

with

$$\begin{aligned} G_0 &= \frac{1}{a} + \frac{1}{a^4 b^2 c} + \frac{1}{a^{16} b^8 c^4 b^2 c} + \cdots \\ G_1 &= \frac{1}{a^2 b} + \frac{1}{a^8 b^4 c^2 b} + \frac{1}{a^{32} b^{16} c^8 b^4 c^2 b} + \cdots \end{aligned}$$



The series  $G_0$ , and  $\text{CF}(\mathbf{s}(\varepsilon))$  satisfy

$$(3.1) \quad G_0 = \frac{1}{a} + \frac{G_0^4}{b^2c},$$

$$(3.2) \quad \frac{1}{\text{CF}(\mathbf{s}(\varepsilon))} = G_0 + \frac{G_0^2}{b}.$$

We deduce from (3.1) and (3.2) that the minimal polynomial of  $\text{CF}(\mathbf{s})$  is

$$(3.3) \quad (ab^2c + abc^2 + c^2)x^4 + (a^2b^2c + a^2bc^2)x^3 + a^2bcx^2 + a^2.$$

The formal power series defined by  $\mathbf{s}(\varepsilon)$  is

$$\begin{aligned} F &= a + bz + az^2 + cz^3 + az^4 + bz^5 + az^6 + bz^7 + az^8 + bz^9 + az^{10} + \dots \\ &= \frac{a}{1+z^2} + bF_0 + cF_1. \end{aligned}$$

The series  $F_0$  and  $F$  satisfy

$$\begin{aligned} F_0 &= \frac{z}{1+z^2} - zF_0^2, \\ F &= \frac{a}{1+z^2} + bF_0 + czF_0^2. \end{aligned}$$

From these we deduce that the minimal polynomial for  $F$  is

$$Ax^2 + Bx + C = 0,$$

where

$$\begin{aligned} A &= z^5 + z, \\ B &= bz^4 + cz^4 + b + c, \\ C &= b^2z^3 + bcz^3 + c^2z^3 + abz^2 + acz^2 + a^2z + b^2z + bcz + ab + ac. \end{aligned}$$

**Example 3.** Let  $\varepsilon = (aabb)^\infty$ . Then  $\mathbf{s}(\varepsilon) = aaabaaabaabaaaaaab\dots$ . We have

$$\frac{1}{\text{CF}(\mathbf{s}(\varepsilon))} = G = \frac{1}{a} + \frac{1}{a^2a} + \frac{1}{a^4a^2b} + \frac{1}{a^8a^4b^2b} + \frac{1}{a^{16}a^8b^4b^2a} + \frac{1}{a^{32}a^{16}b^8b^4a^2a} + \dots$$

and

$$G = \frac{G_0^2}{a} + \frac{G_1^2}{a} + \frac{G_2^2}{b} + \frac{G_3^2}{b}.$$

with

$$\begin{aligned} G_0 &= 1 + \frac{1}{a^8a^4b^2b} + \dots \\ G_1 &= \frac{1}{a} + \frac{1}{a^{16}a^8b^4b^2a} + \dots \\ G_2 &= \frac{1}{a^2a} + \frac{1}{a^{32}a^{16}b^8b^4a^2a} \dots \\ G_3 &= \frac{1}{a^4a^2b} + \frac{1}{a^{64}a^{32}b^{16}b^8a^4a^2b} \dots \end{aligned}$$

The series  $G$  and  $G_0$  satisfy

$$\begin{aligned} G_0 &= 1 + \frac{G_0^{16}}{a^{12}b^3} \\ G &= 1 + G_0 + \frac{G_0^2}{a} + \frac{G_0^4}{a^3} + \frac{G_0^8}{a^6b}. \end{aligned}$$

We deduce

$$G^{16} + c_8 G^8 + c_2 G^2 + c_1 G + c_0 = 0.$$

where

$$\begin{aligned} c_0 &= a^{11}b^3 + a^{10}b^4 + a^3b^{11} + a^2b^{12} + a^6b^6 + a^4b^8 + a^2b^{10} + b^{12} + a^6b^2 + \\ &\quad a^4b^4 + a^2b^6 + b^8 + 1, \\ c_1 &= a^{12}b^3 + a^{11}b^4 + a^4b^{11} + a^3b^{12}, \\ c_2 &= a^{11}b^3 + a^{10}b^4 + a^8b^6 + a^6b^8 + a^4b^{10} + a^3b^{11}, \\ c_8 &= a^6b^2 + a^4b^4 + a^2b^6, \end{aligned}$$

This polynomial is irreducible.

The formal power series defined by  $\mathbf{s}(\varepsilon)$  is

$$F = a(F_0 + F_1) + a(F_2 + F_3),$$

where

$$\begin{aligned} F_0 &= 1 + z^2 + z^4 + z^6 + z^8 + z^{10} + z^{12} + z^{14} + z^{15} + z^{16} + z^{18} + \dots \\ F_1 &= z + z^5 + z^9 + z^{13} + z^{17} + z^{21} + z^{25} + z^{29} + z^{31} + z^{33} + z^{37} + \dots \\ F_2 &= z^3 + z^{11} + z^{19} + z^{27} + z^{35} + z^{43} + z^{51} + z^{59} + z^{63} + z^{67} + \dots \\ F_3 &= z^7 + z^{23} + z^{39} + z^{55} + z^{71} + z^{87} + z^{103} + z^{119} + z^{127} + z^{135} + \dots \end{aligned}$$

The series  $F_0$  and  $F$  satisfy

$$(3.4) \quad F_0 = \frac{1}{1+z} + zF_0^2 + z^3F_0^4 + z^7F_0^8,$$

$$(3.5) \quad F = aF_0 + azF_0^2 + bz^3F_0^4 + bz^7F_0^8.$$

Here  $F$  is not of degree 8, but 4, because we can write (3.4) and (3.5) as

$$(3.6) \quad \frac{1}{1+z} + (F_0 + zF_0^2) + z^3(F_0 + zF_0^2)^4 = 0$$

$$(3.7) \quad F = a(F_0 + zF_0^2) + bz^3(F_0 + zF_0^2)^4.$$

From (3.6) and (3.7) we find minimal polynomial of  $F$  to be

$$c_4x^4 + c_1x + c_0 = 0,$$

where

$$\begin{aligned} c_0 &= a^4z^3 + a^3bz^3 + a^2b^2z^3 + ab^3z^3 + b^4z^3 + a^4z^2 + a^3bz^2 + a^2b^2z^2 + \\ &\quad ab^3z^2 + a^4z + a^3bz + a^2b^2z + ab^3z + a^4 + a^3b + a^2b^2 + ab^3 \\ c_1 &= a^3z^4 + a^2bz^4 + ab^2z^4 + b^3z^4 + a^3 + a^2b + ab^2 + b^3 \\ c_4 &= z^7 + z^3 \end{aligned}$$

This example shows that the degree of the continued fraction is not always twice that of the formal power series.

## 4. CONTINUED FRACTIONS AND RICCATI DIFFERENTIAL EQUATIONS

In 1977, Baum and Sweet [3] considered the subset  $D$  of irrational continued fractions  $\alpha$  in  $\mathbb{F}_2((1/t))$  such that

$$\alpha = [0, a_1, a_2, \dots, a_n, \dots]$$

where  $\deg(a_i) = 1$  (i.e.,  $a_i = t$  or  $t+1$ ) for  $i \geq 1$ . Consider the subset  $P$  of  $\mathbb{F}_2((1/t))$

$$P := \{\alpha \in \mathbb{F}_2((1/t)) \mid \alpha = \sum_{i < 0} a_i t^i\}.$$

Hence  $P$  contains  $D$ . They proved the following theorem:

**Theorem 4.1.** [*Baum-Sweet, 1977*] *An element  $\alpha \in P$  is in  $D$  if and only if  $\alpha$  satisfies*

$$\alpha^2 + t\alpha + 1 = (1+t)\beta^2$$

for some  $\beta \in P$ .

This theorem has the following equivalent formulation (see [8] for a proof):

**Theorem 4.1'.** *An element  $\alpha \in P$  is in  $D$  if and only if  $\alpha$  satisfies*

$$(\alpha \cdot t(t+1))' = \alpha^2 + 1.$$

In [5] and [8], it was proved that the Thue-Morse and period-doubling continued fraction both satisfy the Riccati differential equation

$$(ab(a+b)f)' = (ab)'(1+f^2).$$

We prove that these are in fact particular cases of a more general result:

**Proposition 4.2.** *Let  $\mathbf{u}$  be a sequence taking value in  $\{a, b, a+b\}$ , where  $a, b$  and  $a+b$  (if it is in the image of  $\mathbf{u}$ ) are non-constant polynomials in  $\mathbb{F}_2[t]$ , then the continued fraction*

$$f = [u_0, u_1, \dots] \in \mathbb{F}_2((1/t))$$

*satisfies the Riccati differential equation*

$$(ab(a+b)f)' = (ab)'(1+f^2).$$

*Proof of Proposition 4.2.* Let  $P_n/Q_n$  ( $n \geq -1$ ) be the convergents of  $f$ . Recall that

$$(P_0, Q_0) = (u_0, 1), \quad (P_{-1}, Q_{-1}) = (1, 0)$$

and

$$(4.1) \quad P_n = u_n P_{n-1} + P_{n-2}, \quad Q_n = u_n Q_{n-1} + Q_{n-2} \quad \text{for } n \geq 1.$$

Then we only have to prove that

$$\lim_n (ab(a+b)f_n)' - (ab)'(1+f_n^2) = 0,$$

where  $f_n = \frac{P_n}{Q_n}$ .

Define

$$F_n := ab(a+b)P_n Q_n + ab(P_n^2 + Q_n^2),$$

then

$$(ab(a+b)f_n)' - (ab)'(1+f_n^2) = \frac{F_n'}{Q_n^2}.$$

We prove by induction that for all  $n \geq 0$ ,

$$(4.2) \quad F_n = ab + g_n^2$$

for some  $g_n \in \mathbb{F}_2[a, b]$ .

It is easy to check that the claim is true for  $n = -1, 0$ . Suppose now  $n > 1$  and that the claim holds for  $n - 1$  and  $n - 2$ . From the recurrence relations (4.1) we deduce

$$F_n = u_n^2 F_{n-1} + F_{n-2} + u_n ab(a+b)(P_{n-1}Q_{n-2} + P_{n-2}Q_{n-1}).$$

Notice that  $P_{n-1}Q_{n-2} + P_{n-2}Q_{n-1} = 1$  by a basic property of convergents, and if  $u_n = a$  or  $u_n = b$ ,  $u_n ab(a+b) = a^2 b^2 + u_n^2 ab$ , if  $u_n = a+b$ ,  $u_n ab(a+b) = u_n^2 ab$ . We have

$$F_n = ab + u_n^2 g_{n-1}^2 + g_{n-2}^2 + a^2 b^2$$

if  $u_n = a$  or  $u_n = b$ , and

$$F_n = ab + u_n^2 g_{n-1}^2 + g_{n-2}^2$$

if  $u_n = a+b$ . Thus we have proven (4.2).

Therefore

$$\frac{F'_n}{Q_n^2} = \frac{a'b + b'a}{Q_n^2}.$$

Since the valuation of  $1/Q_n$  in  $1/t$  is at least  $n$ , the above term converges to 0 in  $\mathbb{F}_2((1/t))$  as  $n$  goes to infinity.  $\square$

## 5. ACKNOWLEDGEMENT

I would like to thank Alain Lasjaunias (who has written a partial presentation on this subject intended for a private public) for valuable discussions. I would also like to thank Jean-Paul Allouche for his help.

## REFERENCES

- [1] Jean-Paul Allouche and Jeffrey Shallit. The ring of  $k$ -regular sequences. Theoretical Computer Science, 98(2):163–197, 1992.
- [2] Jean-Paul Allouche and Jeffrey Shallit. Automatic sequences. Theory, Applications, Generalizations. Cambridge University Press, Cambridge, 2003.
- [3] Leonard E. Baum and Melvin M. Sweet. Badly approximable power series in characteristic 2. Ann. of Math. (2), 105(3):573–580, 1977.
- [4] Yann Bugeaud. Automatic continued fractions are transcendental or quadratic. Ann. Sci. Éc. Norm. Supér. (4), 46(6):1005–1022, 2013.
- [5] Yann Bugeaud and Guo-Niu Han. The Thue-Morse continued fractions in characteristic 2 are algebraic. Acta Arith., 2022+.
- [6] Gilles Christol, Teturo Kamae, Michel Mendès France, and Gérard Rauzy. Suites algébriques, automates et substitutions. Bull. Soc. Math. France, 108(4):401–419, 1980.
- [7] Yining Hu and Guo-Niu Han. On the algebraicity of Thue-Morse and period-doubling continued fractions. Acta Arith., 203(4):353–381, 2022.
- [8] Yining Hu and Alain Lasjaunias. Period-doubling continued fractions are algebraic in characteristic 2. Ann. Inst. Fourier (Grenoble), forthcoming.
- [9] Alain Lasjaunias. Continued fractions. arXiv preprint arXiv:1711.11276, 2017.
- [10] Olivier Salon. Suites automatiques à multi-indices. Séminaire de théorie des nombres de Bordeaux, pages 1–36A, 1986.

(Yining Hu) SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, PR CHINA

*Email address:* huyining@protonmail.com