

# SELF-SIMILAR QUANTUM GROUPS

NATHAN BROWNLOWE AND DAVID ROBERTSON

ABSTRACT. We introduce the notion of self-similarity for compact quantum groups. For a finite set  $X$ , we introduce a  $C^*$ -algebra  $\mathbb{A}_X$ , which is the quantum automorphism group of the infinite homogeneous rooted tree  $X^*$ . Self-similar quantum groups are then certain quantum subgroups of  $\mathbb{A}_X$ . Our main class of examples are called finitely-constrained self-similar quantum groups, and we find a class of these examples that can be described as quantum wreath products by subgroups of the quantum permutation group.

## 1. INTRODUCTION

Self-similar groups are a class of groups acting faithfully on an infinite rooted homogeneous tree  $X^*$ . In particular, given an automorphism  $g \in \text{Aut}(X^*)$  and a vertex  $w \in X^*$ , by identifying  $wX^*$  with  $g(w)X^*$ , we get an automorphism  $g|_w \in \text{Aut}(X^*)$  which is uniquely determined by the identity

$$g \cdot (wv) = (g \cdot w)g|_w \cdot v \quad \text{for all } v \in X^*.$$

The automorphism  $g|_w$  is called the *restriction* of  $g$  by  $w$ , and a subgroup  $G \leq \text{Aut}(X^*)$  is *self-similar* if it is closed under restrictions. Self-similar groups are a significant class of groups that play an important role in geometric group theory, and have been a rich source of groups displaying interesting phenomena. Most notably, the Grigorchuk group [5] is a self-similar group which is an infinite, finitely generated periodic group and provided the first example of a group with intermediate growth, as well as the first known amenable group to not be elementary amenable.

When the group of automorphisms  $\text{Aut}(X^*)$  is equipped with the permutation topology, the closed self-similar groups are examples of compact, totally disconnected groups, and hence are profinite groups. A particular class of examples of interest are the self-similar groups of *finite type*, which are subgroups of automorphisms of  $X^*$  that act like elements of a given finite group locally around every vertex. Grigorchuk introduced this concept in [6], where he also showed that the closure of the Grigorchuk group is a self-similar group of finite type. Note that these groups are called finitely constrained self-similar groups in [9], and we will use that terminology.

The theory of compact quantum groups is by now a very substantial part of the wider field of quantum groups, and one which sits in the framework of operator algebras. The theory started with Woronowicz's introduction of the quantum  $\text{SU}(2)$  group in [12]. Woronowicz then defined compact matrix quantum groups in [13], before developing a general theory of compact quantum groups in [14]. An important class of compact matrix quantum groups was identified and studied by Wang through his quantum permutation

---

*Date:* February 6, 2023.

Brownlowe was supported by the Australian Research Council grant DP200100155, and both authors were supported by the Sydney Mathematical Research Institute.

groups in [11]. Wang was motivated by one of Connes' questions from his noncommutative geometry program: what is the *quantum* automorphism group of a space? Wang's work in [11] provided an answer for finite spaces; in particular, Wang formally defined the notion of a quantum automorphism group, and then showed that his quantum permutation group  $A_s(n)$  is the quantum automorphism group of the space with  $n$  points. For three or fewer points this algebra is commutative, and hence indicating no quantum permutations; but for four or more points, remarkably the algebra is noncommutative and infinite-dimensional.

Since the appearance of [11], follow-up work progressed in multiple directions, including the results of Bichon in [2] in which he introduced quantum automorphisms of finite graphs. These algebras are quantum subgroups of the quantum permutation groups. Bichon used this construction to define the quantum dihedral group  $D_4$ . Later still in [1], Banica and Bichon classified all the compact quantum groups acting on four points; that is, all the compact quantum subgroups of  $A_s(4)$ . Quantum automorphisms of infinite graphs have recently been considered by Rollier and Vaes in [8], and by Voigt in [15].

Our current work is the result of us asking the question: is there a reasonable notion of self-similarity for *quantum* groups? We answer this question in the affirmative for compact quantum groups. We do this by first constructing the quantum automorphism group  $\mathbb{A}_X$  of the homogeneous rooted tree  $X^*$ , and then identifying the quantum analogue of the restriction maps  $g \mapsto g|_w$  for  $g \in \text{Aut}(X^*)$ ,  $w \in X^*$ . We then define a self-similar quantum group to be any quantum subgroup  $A$  of  $\mathbb{A}_X$  for which the restriction maps factor through the quotient map  $\mathbb{A}_X \rightarrow A$ . We characterise self-similar quantum groups in terms of a certain homomorphism  $A \otimes C(X) \rightarrow C(X) \otimes A$ , which can be thought of as quantum state-transition function. The main class of examples we examine are quantum analogues of finitely constrained self-similar groups. In our main theorem about these examples we describe a class of finitely constrained self-similar groups as free wreath products by quantum subgroups of quantum permutation groups.

We start with a small preliminaries section in which we collect all the required definitions from the literature on compact quantum groups. In Section 3 we then identify a compact quantum group  $\mathbb{A}_X$  which we prove is the quantum automorphism group of the homogeneous rooted tree  $X^*$ . The  $C^*$ -algebra  $\mathbb{A}_X$  is a noncommutative, infinite-dimensional  $C^*$ -algebra whose abelianisation is the algebra of continuous functions on the automorphism group of the tree  $X^*$ . In Section 4 we introduce the notion of self-similarity for compact quantum groups, and we characterise self-similar quantum groups  $A$  in terms of morphisms  $A \otimes C(X) \rightarrow C(X) \otimes A$ , mimicking the fact that classical self-similar action are governed by the maps  $G \times X \rightarrow X \times G: (g, x) \mapsto (g \cdot x, g|_x)$ .

In Section 5 we define finitely constrained self-similar quantum groups, which are the quantum analogues of the classical finitely constrained self-similar groups studied in [4, 9]. In particular, we consider subalgebras  $\mathbb{A}_d$  of  $\mathbb{A}_X$ , which are the quantum automorphism groups of the finite subtrees  $X^{[d]}$  of  $X^*$  of depth  $d$ . To each quantum subgroup  $\mathbb{P}$  of  $\mathbb{A}_d$ , we construct a quantum subgroup  $A_{\mathbb{P}}$ , which we prove is a self-similar quantum group. We then build on the work of Bichon in [3] by constructing free wreath products of compact quantum groups by quantum subgroups of the quantum permutation group (which corresponds to the subalgebra  $\mathbb{A}_1$  of  $\mathbb{A}_X$ ), and we prove that every  $A_{\mathbb{P}}$  coming from a quantum subgroup  $\mathbb{P}$  of  $\mathbb{A}_1$  is canonically isomorphic to the free wreath product  $A_{\mathbb{P}} *_w \mathbb{P}$ .

## 2. PRELIMINARIES

In this section we collect some basics on compact quantum groups. We start with Woronowicz's definition of a compact quantum group [14].

**Definition 2.1.** A *compact quantum group* is a pair  $(A, \Phi)$  where  $A$  is a unital  $C^*$ -algebra and  $\Phi : A \rightarrow A \otimes A$  is a unital  $*$ -homomorphism such that

- (1)  $(\Phi \otimes \text{id})\Phi = (\text{id} \otimes \Phi)\Phi$
- (2)  $\overline{(A \otimes 1)\Phi(A)} = A \otimes A = \overline{(1 \otimes A)\Phi(A)}$ .

We call  $\Phi$  the *comultiplication* and (1) is called *coassociativity*.

*Remark 2.2.* It is proved in [14] that  $(A, \Phi)$  is a compact quantum group if and only if there is a family of matrices  $\{a_{\lambda} = (a_{ij}^{\lambda}) \in M_{d_{\lambda}}(A) : \lambda \in \Lambda\}$  for some indexing set  $\Lambda$  such that

- (1)  $\Phi(a_{ij}^{\lambda}) = \sum_{k=1}^{d_{\lambda}} a_{ik}^{\lambda} \otimes a_{kj}^{\lambda}$  for all  $\lambda \in \Lambda$  and  $1 \leq i, j \leq d_{\lambda}$ ,
- (2)  $a_{ij}^{\lambda}$  and its transpose  $(a_{ij}^{\lambda})^*$  are invertible elements of  $M_{d_{\lambda}}(A)$  for every  $\lambda \in \Lambda$ ,
- (3) the  $*$ -subalgebra  $\mathcal{A}$  of  $A$  generated by the entries  $\{a_{ij}^{\lambda} : 1 \leq i, j \leq d_{\lambda}, \lambda \in \Lambda\}$  is dense in  $A$ .

*Example 2.3.* A key example for us is Wang's *quantum permutation groups*  $(A_s(n), \Phi)$  from [11]. Here,  $n$  is a positive integer, and  $A_s(n)$  is the universal  $C^*$ -algebra generated by elements  $a_{ij}$ ,  $1 \leq i, j \leq n$ , satisfying

$$\begin{aligned} a_{ij}^2 &= a_{ij} = a_{ij}^* \text{ for all } 1 \leq i, j \leq n, \\ \sum_{i=1}^n a_{ij} &= 1 \text{ for all } 1 \leq j \leq n, \\ \sum_{j=1}^n a_{ij} &= 1 \text{ for all } 1 \leq i \leq n. \end{aligned}$$

The comultiplication  $\Phi$  satisfies  $\Phi(a_{ij}) = \sum_{k=1}^n a_{ik} \otimes a_{kj}$  for all  $1 \leq i, j \leq n$ .

**Definition 2.4.** If  $(A_1, \Phi_1)$  and  $(A_2, \Phi_2)$  are compact quantum groups, then a *morphism*  $\pi$  from  $(A_1, \Phi_1)$  to  $(A_2, \Phi_2)$  is a homomorphism of  $C^*$ -algebras  $\pi : A_1 \rightarrow A_2$  satisfying  $(\pi \otimes \pi) \circ \Phi_1 = \Phi_2 \circ \pi$ .

**Definition 2.5.** Let  $(A, \Phi)$  be a compact quantum group. A *Woronowicz ideal* is an ideal  $I$  of  $A$  such that  $\Phi(I) \subseteq \ker(q \otimes q)$ , where  $q$  is the quotient map  $A \rightarrow A/I$ . Then  $(A/I, \Phi')$ , where  $\Phi' : A/I \rightarrow A/I \otimes A/I$  satisfies  $\Phi' \circ q = (q \otimes q) \circ \Phi$  is a compact quantum group called a *quantum subgroup* of  $(A, \Phi)$ .

**Definition 2.6.** A *(left) coaction* of a compact quantum group  $(A, \Phi)$  on a unital  $C^*$ -algebra  $B$  is a unital  $*$ -homomorphism  $\alpha : B \rightarrow A \otimes B$  satisfying

- (1)  $(\text{id} \otimes \alpha)\alpha = (\Phi \otimes \text{id})\alpha$
- (2)  $\alpha(B)(A \otimes 1) = A \otimes B$ .

We refer to (1) as the *coaction identity* and (2) is known as the *Podleś condition*.

### 3. QUANTUM AUTOMORPHISMS OF A HOMOGENEOUS ROOTED TREE

In this section we introduce a compact quantum group  $\mathbb{A}_X$  which we prove is the quantum automorphism group of the infinite homogeneous rooted tree  $X^*$ . We start with the notion of an action of a compact quantum group on  $X^*$ . Note that for  $n \geq 0$  we write  $X^n$  for all the words in  $X$  of length  $n$ , and we then the tree  $X^*$  can be identified with  $\bigcup_{n \geq 0} X^n$ , where  $X^0 = \{\emptyset\}$  and  $\emptyset$  is the root of the tree.

**Definition 3.1.** Let  $X$  be a finite set and let  $(A, \Phi)$  be a compact quantum group. An action of  $A$  on the homogeneous rooted tree  $X^*$  is a system

$$\alpha = (\alpha_n : C(X^n) \rightarrow A \otimes C(X^n))$$

of left coactions, such that for any  $m < n$  the diagram

$$\begin{array}{ccc} C(X^m) & \xrightarrow{\alpha_{m,n}} & C(X^n) \\ \downarrow \alpha_m & & \downarrow \alpha_n \\ A \otimes C(X^m) & \xrightarrow{\text{id} \otimes \alpha_{m,n}} & A \otimes C(X^n) \end{array}$$

commutes, where  $i_{m,n} : C(X^m) \rightarrow C(X^n)$  is the injective homomorphism satisfying

$$i_{m,n}(p_w) = \sum_{w' \in X^{n-m}} p_{ww'}.$$

We now define the main object of interest in this section, the  $C^*$ -algebra  $\mathbb{A}_X$ , before proving that it is indeed a compact quantum group in Theorem 3.4. At some point in the later stages of this project we became aware of [8], and their notion of the quantum automorphism group  $\text{QAut } \Pi$  of a locally finite connected graph  $\Pi$ . A straightforward argument shows that  $\mathbb{A}_X$  is  $\text{QAut } \Pi$  for  $\Pi$  the homogeneous rooted tree, but we include the proof of Theorem 3.4 for completeness.

**Definition 3.2.** Let  $X$  be a finite set. Define  $\mathbb{A}_X$  to be the universal  $C^*$ -algebra generated by elements  $\{a_{u,v} : u, v \in X^n, n \geq 0\}$  subject to the following relations:

- (1)  $a_{\emptyset, \emptyset} = 1$ ,
- (2) for any  $n \geq 0, u, v \in X^n$ ,  $a_{u,v}^* = a_{v,u} = a_{u,v}$ ,
- (3) for any  $n \geq 0, u, v \in X^n$  and  $x \in X$

$$a_{u,v} = \sum_{y \in X} a_{ux} a_{vy} = \sum_{z \in X} a_{uz} a_{vx}.$$

*Remarks 3.3.* (i) For each  $d \in \mathbb{N}$  we denote by  $\mathbb{A}_d$  the subalgebra of  $\mathbb{A}_X$  generated by  $\{a_{u,v} : u, v \in X^d\}$ . Note that  $\mathbb{A}_1$  is the Wang's quantum permutation group  $A_s(|X|)$  from Example 2.3.

- (ii) We can interpret (3) as follows: each projection  $a_{u,v}$  decomposes as an  $|X| \times |X|$  square of projections  $\{a_{ux} a_{vy} : x, y \in X\}$  with a magic square type property where every row and column sums to  $a_{u,v}$ . For example, if  $X = \{0, 1, 2\}$  we have the following structure.

$$a_{u,v} \mapsto \begin{array}{|c|c|c|} \hline a_{u0} a_{v0} & a_{u0} a_{v1} & a_{u0} a_{v2} \\ \hline a_{u1} a_{v0} & a_{u1} a_{v1} & a_{u1} a_{v2} \\ \hline a_{u2} a_{v0} & a_{u2} a_{v1} & a_{u2} a_{v2} \\ \hline \end{array}$$

(iii) Repeated applications of (3) from Definition 3.2 show that for all  $u, u', v, v', w \in X^n, n \in \mathbb{N}$ , we have

$$u \neq u', v \neq v' \implies a_{u \cdot v} a_{u' \cdot v} = 0 = a_{u \cdot v} a_{u' \cdot v'},$$

and that for all  $u = u_1 \cdots u_n, v = v_1 \cdots v_n \in X^n, n \in \mathbb{N}$ , and  $x, y \in X$  we have

$$a_{x \cdot y} a_{u \cdot v} = a_{u \cdot v} a_{x \cdot y} = \begin{cases} a_{u \cdot v} & \text{if } u_1 = x, v_1 = y \\ 0 & \text{otherwise.} \end{cases}$$

We will freely use these two identities without comment throughout the rest of the paper.

**Theorem 3.4.** *The  $C^*$ -algebra  $\mathbb{A}_X$  is a compact quantum group with comultiplication  $\Delta : \mathbb{A}_X \rightarrow \mathbb{A}_X \otimes \mathbb{A}_X$  satisfying*

$$\Delta(a_{u \cdot v}) = \sum_{w \in X^n} a_{u \cdot w} \otimes a_{w \cdot v},$$

for all  $u, v \in X^n$  and  $n \geq 1$

*Proof.* To see that  $\Delta$  exists, it's enough to show that the elements

$$b_{u \cdot v} := \sum_{w \in X^n} a_{u \cdot w} \otimes a_{w \cdot v}$$

for  $u, v \in X^n$  and  $n \geq 1$  satisfy Definition 3.2.

Firstly,  $b_{\emptyset \cdot \emptyset} = \Delta(a_{\emptyset \cdot \emptyset}) = a_{\emptyset \cdot \emptyset} \otimes a_{\emptyset \cdot \emptyset} = 1 \otimes 1$ . For (2), we have

$$b_{u \cdot v}^* = \sum_{w \in X^n} a_{u \cdot w}^* \otimes a_{w \cdot v}^* = \sum_{w \in X^n} a_{u \cdot w} \otimes a_{w \cdot v} = b_{u \cdot v}$$

and

$$\begin{aligned} b_{u \cdot v}^2 &= \left( \sum_{w \in X^n} a_{u \cdot w} \otimes a_{w \cdot v} \right)^2 \\ &= \sum_{w, z \in X^n} a_{u \cdot w} a_{u \cdot z} \otimes a_{w \cdot v} a_{z \cdot v} \\ &= \sum_{w \in X^n} a_{u \cdot w}^2 \otimes a_{w \cdot v}^2 \\ &= \sum_{w \in X^n} a_{u \cdot w} \otimes a_{w \cdot v} \\ &= b_{u \cdot v}. \end{aligned}$$

For (3) fix  $u, v \in X^n$  and  $x \in X$ . Then

$$\begin{aligned} b_{u \cdot v} &= \sum_{w \in X^n} a_{u \cdot w} \otimes a_{w \cdot v} \\ &= \sum_{w \in X^n} \sum_{z \in X} a_{u \cdot x \cdot w z} \otimes a_{w \cdot v} \\ &= \sum_{w \in X^n} \sum_{z \in X} a_{u \cdot x \cdot w z} \otimes \sum_{y \in X} a_{w z \cdot y} \end{aligned}$$

$$\begin{aligned}
&= \sum_{y \in X} \sum_{w \in X^{n+1}} a_{u \cdot x \cdot w} \otimes a_{w \cdot x \cdot y} \\
&= \sum_{y \in X} b_{u \cdot x \cdot y}.
\end{aligned}$$

So by the universal property of  $\mathbb{A}_X$  there is a homomorphism  $\Delta : \mathbb{A}_X \rightarrow \mathbb{A}_X \otimes \mathbb{A}_X$  such that

$$\Delta(a_{u \cdot x}) = \sum_{w \in X^n} a_{u \cdot x \cdot w} \otimes a_{w \cdot x}.$$

For coassociativity, we have

$$\begin{aligned}
(\text{id} \otimes \Delta) \circ \Delta(a_{u \cdot x}) &= \sum_{w \in X^n} a_{u \cdot x \cdot w} \otimes \Delta(a_{w \cdot x}) \\
&= \sum_{w \in X^n} a_{u \cdot x \cdot w} \otimes \left( \sum_{z \in X^n} a_{w \cdot x \cdot z} \otimes a_{z \cdot x} \right) \\
&= \sum_{z \in X^n} \left( \sum_{w \in X^n} a_{u \cdot x \cdot w} \otimes a_{w \cdot x \cdot z} \right) \otimes a_{z \cdot x} \\
&= \sum_{z \in X^n} \Delta(a_{u \cdot x \cdot z}) \otimes a_{z \cdot x} \\
&= (\Delta \otimes \text{id}) \circ \Delta(a_{u \cdot x}).
\end{aligned}$$

Finally, we show that that the set of matrices

$$\{a_n = (a_{u \cdot x})_{u \cdot x \in X^n} \in M_{X^n}(\mathbb{A}_X) : n \geq 1\}$$

satisfy the conditions of Definition 2.2. Conditions (1) and (3) are clear. For (2) we show that given any  $n \geq 1$  the matrix  $a_n$  is invertible with inverse given by  $(a_n)^{\mathbf{f}}$ . Given  $u, v \in X^n$  we have

$$(a_n(a_n)^{\mathbf{f}})_{u \cdot x} = \sum_{w \in X^n} a_{u \cdot x \cdot w} a_{v \cdot x \cdot w} = \delta_{u \cdot x} \sum_{w \in X^n} a_{u \cdot x \cdot w} = \delta_{u \cdot x} 1_A.$$

Likewise, we can show  $((a_n)^{\mathbf{f}} a_n)_{u \cdot x} = \delta_{u \cdot x} 1_A$  and hence  $(a_n)^{\mathbf{f}} = a_n^{-1}$  as required.  $\square$

*Remark 3.5.* The canonical dense  $*$ -subalgebra of  $\mathbb{A}_X$  is the  $*$ -subalgebra generated by the projections  $\{a_{u \cdot x} : u, v \in X^n, n \geq 0\}$ . This is a Hopf  $*$ -algebra with counit  $\varepsilon : \mathbb{A}_X \rightarrow \mathbb{C}$  and coinverse  $\kappa : \mathbb{A}_X \rightarrow \mathbb{A}_X$  satisfying  $\varepsilon(a_{u \cdot x}) = \delta_{u \cdot x}$  and  $\kappa(a_{u \cdot x}) = a_{v \cdot x}$ , for  $u, v \in X^n$ ,  $n \in \mathbb{N}$ .

We now show that  $(\mathbb{A}_X, \Delta)$  is the quantum automorphism group (in the sense of [11, Definition 2.3]) of the homogeneous rooted tree.

**Proposition 3.6.** *There is an action  $\gamma = (\gamma_n)_{n=1}^\infty$  of  $\mathbb{A}_X$  on  $X^*$ . Moreover, if  $\alpha = (\alpha_n)_{n=1}^\infty$  is an action of a compact quantum group  $(A, \Phi)$  on  $X^*$  then there is a quantum group homomorphism  $\pi : \mathbb{A}_X \rightarrow A$  such that  $(\pi \otimes \text{id}) \circ \gamma_n = \alpha_n$  for any  $n \geq 1$ .*

*Proof.* For any  $n \geq 1$ , the elements

$$q_w := \sum_{w' \in X^n} a_{w \cdot x \cdot w'} \otimes p_{w'} \in \mathbb{A}_X \otimes C(X^n)$$

for each  $w \in X^n$  are mutually orthogonal projections and satisfy

$$\sum_{w \in X^n} q_w = \sum_{w, w' \in X^n} a_{w, w'} \otimes p_{w'} = 1 \otimes 1$$

Therefore there is a unital  $*$ -homomorphism  $\gamma_n : C(X^n) \rightarrow \mathbb{A}_X \otimes C(X^n)$  satisfying  $\gamma_n(p_w) = q_w$ . We have

$$\begin{aligned} (\Delta \otimes \text{id})\gamma_n(p_w) &= \sum_{w' \in X^n} \Delta(a_{w, w'}) \otimes p_{w'} \\ &= \sum_{w', z \in X^n} a_{w, z} \otimes a_{z, w'} \otimes p'_{w'} \\ &= \sum_{z \in X^n} a_{w, z} \otimes \alpha_n(p_z) \\ &= (\text{id} \otimes \gamma_n)\gamma_n(p_w), \end{aligned}$$

and so each  $\gamma_n$  satisfies the coaction identity.

For a fixed  $v \in X^n$  we have

$$\sum_{u \in X^n} \gamma_n(p_u)(a_{u, v} \otimes 1) = \sum_{u, w \in X^n} a_{u, w} a_{w, v} \otimes p_w = \sum_{u \in X^n} a_{u, v} \otimes p_v = 1 \otimes p_v.$$

Multiplying by any element  $a \otimes 1 \in \mathbb{A}_X \otimes 1$  shows that  $\gamma_n(C(X^n))(\mathbb{A}_X \otimes 1)$  contains the elements  $a \otimes p_v$  of  $\mathbb{A}_X \otimes C(X^n)$  and hence the required density is satisfied.

Finally, fix  $m < n$  and  $w \in X_l$ . Then

$$\begin{aligned} (\text{id} \otimes i_l)\gamma_l(p_w) &= \sum_{z \in X^m} a_{w, z} \otimes i_l(p_z) \\ &= \sum_{z \in X^m} \sum_{z' \in X^{n-m}} a_{w, z} \otimes p_{zz'} \\ &= \sum_{z \in X^m} \sum_{z' \in X^{n-m}} \sum_{w' \in X^{n-m}} a_{ww', z'} \otimes p_{zz'} \\ &= \sum_{w' \in X^{n-m}} \alpha_n(p_{ww'}) \\ &= \gamma_n(i_l(p_w)), \end{aligned}$$

and so the collection  $\gamma = (\gamma_n)_{n=1}^\infty$  defines an action of  $(\mathbb{A}_X, \Delta)$  on the homogeneous rooted tree  $X^*$ .

Now suppose  $(\alpha_n)_{n=1}^\infty$  is an action of a compact quantum group  $(A, \Phi)$  on  $X^*$ . Let  $b_{\emptyset, \emptyset} := 1 \in A$  and for  $n \geq 1$  and  $u, v \in X^n$  define  $b_{u, v} \in A$  to be the unique elements satisfying

$$\alpha_n(p_u) = \sum_{v \in X^n} b_{u, v} \otimes p_v.$$

The coaction identity for  $\alpha_n$  says that

$$(3.1) \quad \Phi(b_{u, v}) = \sum_{w \in X^n} b_{u, w} \otimes b_{w, v}$$

for any  $u, v \in X^n$ .

We claim that the collection  $\{b_{u\cdot X} : u, v \in X^n, n \geq 0\} \subseteq A$  satisfies Definition 3.2. Condition (1) is by definition. For (2) and (3), we appeal to the universal property of the quantum permutation groups  $A_s(|X|^n)$  for  $n \geq 1$ . Since for any  $n \geq 1$ ,  $\alpha_n$  defines a coaction of  $(A, \Phi)$  on  $C(X^n)$ , [11, Theorem 3.1] says that the elements  $\{b_{u\cdot X} : u, v \in X^n\}$  satisfy conditions (3.1)–(3.3) of [11, Section 3]. Condition (3.1) is precisely (2). Conditions (3.1) and (3.2) say that for any  $v \in X^n$  we have

$$\sum_{u \in X^n} b_{u\cdot X} = 1_A = \sum_{w \in X^n} b_{w\cdot X}.$$

For any  $u \in X^n$  and  $x \in X$  we have

$$p_{ux} \leq \sum_{y \in X} p_{uy} = i_{n\cdot X+1}(p_u),$$

and hence

$$\begin{aligned} \sum_{v \in X^n} \sum_{y \in X} b_{v\cdot X} \otimes p_{vy} &= \alpha_{n+1}(p_{ux}) \\ &\leq \alpha_{n+1}(i_{n\cdot X+1}(p_u)) \\ &= (\text{id}_A \otimes i_{n\cdot X+1})\alpha_n(p_u) \\ &= \sum_{v \in X^n} \sum_{y \in X} b_{v\cdot X} \otimes p_{vy}. \end{aligned}$$

It follows that  $b_{ux\cdot X} \leq b_{u\cdot X}$  for any  $x, y \in X$ . Therefore, for any  $u, v \in X^n$  and  $x \in X$  we have

$$b_{u\cdot X} = b_{u\cdot X} \left( \sum_{w \in X^n} \sum_{y \in X} b_{w\cdot X} \right) = \sum_{y \in X} b_{u\cdot X} \otimes p_{xy}.$$

Likewise for any  $y \in X$  we have  $b_{u\cdot X} = \sum_{x \in X} b_{ux\cdot X}$  and (3) holds.

Therefore, the universal property of  $\mathbb{A}_X$  provides a homomorphism  $\pi : \mathbb{A}_X \rightarrow A$  satisfying  $\pi(a_{u\cdot X}) = b_{u\cdot X}$ . It follows from (3.1) that  $(\pi \otimes \pi) \circ \Delta = \Phi \otimes \pi$  and so  $\pi$  is a compact quantum group homomorphism. The identity  $(\pi \otimes \text{id}) \circ \gamma_n = \alpha_n$  is immediate.  $\square$

**Proposition 3.7.** *For  $|X| \geq 2$  the  $C^*$ -algebra  $\mathbb{A}_X$  is non-commutative and infinite dimensional.*

*Proof.* Without loss of generality, assume  $X = \{0, 1\}$ . Let  $B$  be the universal unital  $C^*$ -algebra generated by two (non-commuting) projections  $p$  and  $q$ . It is known from [7] that  $B \cong C^*(\mathbb{Z}_2 * \mathbb{Z}_2)$ , which is non-commutative and infinite dimensional. Define the matrix

$$(b_{u\cdot X})_{u\cdot X \in X^2} = \begin{pmatrix} p & 1_B - p & 0 & 0 \\ 1_B - p & p & 0 & 0 \\ 0 & 0 & q & 1_B - q \\ 0 & 0 & 1_B - q & q \end{pmatrix} \in M_4(B).$$

Define  $b_{\emptyset\cdot X} = b_{0\cdot X} = b_{1\cdot X} = 1_B$ ,  $b_{0\cdot X} = b_{1\cdot X} = 0$  and for  $u, v \in X^2$  and  $w, w' \in X^*$  define  $b_{uw\cdot X} := \delta_{w\cdot X} b_{u\cdot X}$ . Then these elements satisfy the relations in Definition 3.2 and hence there is a surjective homomorphism  $\mathbb{A}_X \rightarrow B$ . Since  $B$  is non-commutative and infinite-dimensional so is  $\mathbb{A}_X$ .  $\square$



*Remark 3.8.* The group  $\text{Aut}(X^*)$  of automorphisms of a homogeneous rooted tree  $X^*$  is compact totally disconnected Hausdorff group under the permutation topology. A neighbourhood basis of the identity is given by the family of subgroups

$$\{G_u := \{g \in \text{Aut}(X^*) : g \cdot u = u\} : u \in X^*\},$$

and since the orbit of any  $u \in X^*$  is finite, each of these open subgroups is closed and hence compact. Cosets of these subgroups are of the form  $G_{u \cdot X} := \{g \in G : g \cdot v = u\}$ . Then  $\{G_{u \cdot X} : u, v \in X^*\}$  is a basis of compact open sets for the topology on  $\text{Aut}(X^*)$ . It follows that the indicator functions  $f_{u \cdot X} := 1_{G_{u \cdot X}}$  span a dense subset of  $C(\text{Aut}(X^*))$ . It is easily checked that the elements  $f_{u \cdot X}$  satisfy (1)–(3) of Definition 3.2 and the universal property of  $C(\text{Aut}(X^*))$  then implies that it is the abelianisation of  $\mathbb{A}_X$ .

#### 4. SELF-SIMILARITY

If  $g \in \text{Aut}(X^*)$  and  $x \in X$ , the *restriction*  $g|_x$  is the unique element of  $\text{Aut}(X^*)$  satisfying

$$g \cdot (xw) = (g \cdot x)g|_x \cdot w \quad \text{for all } w \in X^*.$$

A subgroup  $G \leq \text{Aut}(X^*)$  is called *self-similar* if  $G$  is closed under taking restrictions. That is, whenever  $g \in G$  and  $x \in X$ , the restriction  $g|_x$  is an element of  $G$ . With the topology inherited from  $\text{Aut}(X^*)$ , the restriction map  $G \rightarrow G : g \mapsto g|_x$  is continuous. If  $G$  is any group acting on  $X^*$  by automorphisms, we call the action *self-similar* if the image of  $G$  in  $\text{Aut}(X^*)$  is self-similar.

To have a reasonable notion of self-similarity for quantum subgroups of  $\mathbb{A}_X$ , we need to understand how restriction manifests itself in the function algebra  $C(\text{Aut}(X^*))$ . Given  $x \in X$  and  $u, v \in X^n$  we have

$$\begin{aligned} \{g : g|_x \cdot u = v\} &= \left( \bigcup_{y \in X} \{g : g \cdot x = y\} \right) \cap \{g : g|_x \cdot u = v\} \\ &= \bigcup_{y \in X} (\{g : g \cdot x = y\} \cap \{g : g|_x \cdot u = v\}) \\ &= \bigcup_{y \in X} \{g : g \cdot (xu) = yv\}, \end{aligned}$$

and hence the corresponding indicator functions satisfy

$$1_{\{g|_x \cdot u = v\}} = \sum_{y \in X} 1_{\{g \cdot (xu) = yv\}}.$$

This formula motivates the following result.

**Proposition 4.1.** *For each  $x \in X$  there is a homomorphism  $\rho_x : \mathbb{A}_X \rightarrow \mathbb{A}_X$  satisfying*

$$(4.1) \quad \rho_x(a_{u \cdot X}) = \sum_{y \in X} a_{y \cdot X} v,$$

for all  $u, v \in X^n$ .

We illustrate the formula for a restriction map in Figure 1 by considering  $X = \{0, 1, 2\}$  and looking at what the restriction map  $\rho_1$  does to the projection  $a_{1 \cdot X}$ .




FIGURE 1.  $\rho_1(a_{1\cdot X}) = a_{01\cdot X} + a_{11\cdot X} + a_{21\cdot X}$

*Proof of Proposition 4.1.* Fix  $x \in X$ . We show that the elements

$$\{b_{u\cdot X} := \rho_x(a_{u\cdot X}) : u, v \in X^n, n \geq 1\}$$

satisfy the conditions of Definition 3.2. For (1) we have

$$b_{\emptyset\cdot X} = \rho_x(a_{\emptyset\cdot X}) = \sum_{y \in X} a_{y\cdot X} = 1.$$

For (2), we have

$$b_{u\cdot X}^* = \left( \sum_{y \in X} a_{yu\cdot X} \right)^* = \sum_{y \in X} a_{yu\cdot X}^* = \sum_{y \in X} a_{yu\cdot X} = b_{u\cdot X}$$

and

$$b_{u\cdot X}^2 = \left( \sum_{y \in X} a_{yu\cdot X} \right)^2 = \sum_{y, z \in X} a_{yu\cdot X} a_{zu\cdot X} = \left( \sum_{y \in X} a_{yu\cdot X} \right) = b_{u\cdot X}.$$

For (3), fix  $y \in X$ . Then

$$\sum_{z \in X} b_{uy\cdot X} z = \sum_{z \in X} \sum_{w \in X} a_{wuy\cdot X} z = \sum_{w \in X} a_{wuy\cdot X} = b_{u\cdot X}.$$

A similar calculation shows  $\sum_{z \in X} b_{uz\cdot X} y = b_{u\cdot X}$ . Hence there is a homomorphism  $\rho_x$  with the desired formula.  $\square$

*Remark 4.2.* We define  $\rho_\emptyset$  to be the identity homomorphism  $\mathbb{A}_X \rightarrow \mathbb{A}_X$ , and for  $w = w_1 \cdots w_n \in X^n$  we define  $\rho_w$  to be the composition  $\rho_{w_1} \circ \cdots \circ \rho_{w_n}$ . A routine calculation shows that for all  $u, v \in X^n$  we have

$$\rho_w(a_{u\cdot X}) = \sum_{z \in X^n} a_{zu\cdot X} w.$$

*Remark 4.3.* A similar argument to the one in the proof of Proposition 4.1 shows that for each  $x \in X$  there is a homomorphism  $\sigma_x: \mathbb{A}_X \rightarrow \mathbb{A}_X$  satisfying

$$\sigma_x(a_{u\cdot X}) = \sum_{y \in X} a_{xu\cdot X} y$$

for all  $u, v \in X^n$ ,  $n \in \mathbb{N}$ . It is straightforward to see that  $\sigma_x = \kappa \circ \rho_x \circ \kappa$ , where  $\kappa$  is the coinverse.

We can now state the main definition of the paper.

**Definition 4.4.** We call  $\rho_w$  the *restriction by  $w$* . A quantum subgroup  $A$  of  $\mathbb{A}_X$  is *self-similar* if for each  $x \in X$  the restriction  $\rho_x$  factors through the quotient map  $q : \mathbb{A}_X \rightarrow A$ ; that is, if there exists a homomorphism  $\tilde{\rho}_x : A \rightarrow A$  such that the diagram

$$\begin{array}{ccc} \mathbb{A}_X & \xrightarrow{x} & \mathbb{A}_X \\ \downarrow q & & \downarrow q \\ A & \xrightarrow{\tilde{\rho}_x} & A \end{array}$$

commutes.

To motivate the main result of this section, let  $G$  be a group. To construct a self-similar action of  $G$  on  $X^*$ , it suffices to have a function  $f : G \times X \rightarrow X \times G$  such that  $f(e, x) = (x, e)$  for all  $x \in X$ , and such that the following diagram commutes

$$\begin{array}{ccc} G \times G \times X & \xrightarrow{G \times \text{id}_X} & G \times X \\ \downarrow \text{id}_G \times f & & \downarrow f \\ G \times X \times G & & \\ \downarrow f \times \text{id}_G & & \\ X \times G \times G & \xrightarrow{\text{id}_X \times G} & X \times G \end{array}$$

This data allows us to define an action of  $G$  on  $X^*$ , which is self-similar with  $g \cdot x$  and  $g|_x$  the unique elements of  $X$  and  $G$  satisfying  $(g \cdot x, g|_x) := f(g, x)$ .

Our next result is a compact quantum group analogue of the above result. We will be working with multiple different identity homomorphisms and units. For clarity we adopt the following notational conventions: we write  $\text{id}_A$  for the identity homomorphism on a  $C^*$ -algebra  $A$ , and for  $n \geq 1$  write  $\text{id}_n$  for the identity homomorphism on the commutative  $C^*$ -algebra  $C(X^n)$ . Likewise,  $1_A$  will denote the unit of  $A$ ,  $1$  and  $1_n$  will denote the units of  $C(X)$  and  $C(X^n)$  respectively.

**Theorem 4.5.** *Suppose  $(A, \Phi)$  is a compact quantum group equipped with a unital  $*$ -homomorphism  $\psi : C(X) \otimes A \rightarrow A \otimes C(X)$  satisfying*

$$(4.2) \quad (\Phi \otimes \text{id}_1)\psi = (\text{id}_A \otimes \psi)(\psi \otimes \text{id}_A)(\text{id}_1 \otimes \Phi)$$

and

$$(4.3) \quad \overline{\psi(C(X) \otimes 1_A)(A \otimes 1)} = A \otimes C(X).$$

Then  $(A, \Phi)$  acts on the homogeneous rooted tree  $X^*$  and moreover the image of  $\mathbb{A}_X$ , under the homomorphism  $\pi : \mathbb{A}_X \rightarrow A$  from Proposition 3.6, is a self-similar compact quantum group.

*Proof.* We begin by defining an action of  $(A, \Phi)$  on  $X^*$ . Identify  $C(X)$  with  $C(X) \otimes 1_A \subseteq C(X) \otimes A$  and let  $\alpha_1 := \psi|_{C(X) \otimes 1_A}$ . Then  $\alpha_1$  is clearly unital and the coaction identity and Podleś condition for  $\alpha_1$  follow from (4.2) and (4.3). Now inductively define  $\alpha_{n+1} := (\psi \otimes \text{id}_n)(\text{id}_1 \otimes \alpha_n) : C(X^{n+1}) \rightarrow A \otimes C(X^{n+1})$  for  $n \geq 1$ , where we are suppressing the canonical isomorphism  $C(X^{n+1}) \cong C(X) \otimes C(X^n)$ . Again,  $\alpha_{n+1}$  is clearly unital whenever  $\alpha_n$  is. If we assume  $\alpha_n$  satisfies the coaction identity, then

$$\begin{aligned} (\Phi \otimes \text{id}_{n+1})\alpha_{n+1} &= (\Phi \otimes \text{id}_{n+1})(\psi \otimes \text{id}_n)(\text{id}_1 \otimes \alpha_n) \\ &= (\text{id}_A \otimes \psi \otimes \text{id}_n)(\psi \otimes \text{id}_A \otimes \text{id}_n)(\text{id}_1 \otimes \Phi \otimes \text{id}_n)(\text{id}_1 \otimes \alpha_n) \end{aligned}$$

$$\begin{aligned}
&= (\text{id}_A \otimes \psi \otimes \text{id}_n)(\psi \otimes \text{id}_A \otimes \text{id}_n)(\text{id}_1 \otimes \text{id}_A \otimes \alpha_n)(\text{id}_1 \otimes \alpha_n) \\
&= (\text{id}_A \otimes \psi \otimes \text{id}_n)(\text{id}_A \otimes \text{id}_1 \otimes \alpha_n)(\psi \otimes \text{id}_n)(\text{id}_1 \otimes \alpha_n) \\
&= (\text{id}_A \otimes \alpha_{n+1})\alpha_{n+1},
\end{aligned}$$

and so  $\alpha_{n+1}$  also satisfies the coaction identity. Since  $\alpha_1$  is a coaction, we see that  $\alpha_n$  satisfies the coaction identity for any  $n \geq 1$ .

To see that each  $\alpha_n$  satisfies the Podleś condition, we argue by induction. We know it is satisfied for  $n = 1$ . Suppose for some  $n \geq 1$  that

$$\overline{\alpha_n(C(X^n))(A \otimes 1_n)} = A \otimes C(X^n).$$

Fix a spanning element  $a \otimes p_u \otimes p_x \in A \otimes C(X^{n+1})$  where  $u \in X^n$  and  $x \in X$ . By the inductive hypothesis we can approximate

$$a \otimes p_u \sim \sum_i \alpha_n(f_i)(a_i \otimes 1_n),$$

where  $f_i \in C(X^n)$  and  $a_i \in A$ . Then

$$(4.4) \quad a \otimes p_u \otimes p_x \sim \sum_i (\alpha_n(f_i) \otimes 1)(1_A \otimes 1_n \otimes p_x)(a_i \otimes 1_{n+1}).$$

By definition of  $\alpha_n$ , for any  $f \in C(X^n)$  we have

$$\begin{aligned}
\alpha_n(f) \otimes 1 &= ((\psi \otimes \text{id}_{n-1}) \dots (\text{id}_{n-1} \otimes \psi)(f \otimes 1_A)) \otimes 1 \\
&= (\psi \otimes \text{id}_n) \dots (\text{id}_{n-1} \otimes \psi \otimes \text{id}_1)(f \otimes 1_A \otimes 1) \\
&= (\psi \otimes \text{id}_n) \dots (\text{id}_{n-1} \otimes \psi \otimes \text{id}_1)(\text{id}_n \otimes \psi)(f \otimes 1 \otimes 1_A) \\
&= \alpha_{n+1}(f \otimes 1).
\end{aligned}$$

So we can write (4.4) as

$$\sum_i \alpha_{n+1}(f_i \otimes 1)(1_A \otimes 1_n \otimes p_x)(a_i \otimes 1_{n+1}).$$

Since  $\psi$  is unital, we have

$$1_A \otimes 1_n \otimes p_x = (\psi \otimes \text{id}_n)(1 \otimes 1_A \otimes 1_{n-1} \otimes p_x),$$

which can be approximated using the induction hypothesis by

$$\begin{aligned}
(\psi \otimes \text{id}_n)(1 \otimes 1_A \otimes 1_{n-1} \otimes p_x) &\sim (\psi \otimes \text{id}_n) \left( 1 \otimes \sum_i \alpha_n(g_i)(b_i \otimes 1_n) \right) \\
&= (\psi \otimes \text{id}_n) \left( \sum_i (\text{id}_1 \otimes \alpha_n)(1 \otimes g_i)(1 \otimes b_i \otimes 1_n) \right) \\
&= \sum_i \alpha_{n+1}(1 \otimes g_i)(\psi \otimes \text{id}_n)(1 \otimes b_i \otimes 1_n).
\end{aligned}$$

Finally, applying the Podleś condition for  $\alpha_1$  we can approximate

$$\psi(1 \otimes b_i) \sim \sum_j \alpha_1(h_j)(c_j \otimes 1) = \sum_j \psi(h_j \otimes 1_A)(c_j \otimes 1),$$

so

$$\begin{aligned} (\psi \otimes \text{id}_n)(1 \otimes b \otimes 1_n) &\sim \sum (\psi(h \otimes 1_A) \otimes 1_n)(c \otimes 1_{n+1}) \\ &= \sum \alpha_{n+1}(h \otimes 1_n)(c \otimes 1_{n+1}). \end{aligned}$$

Combining these approximations we can write

$$a \otimes p_u \otimes p_x \sim \sum_{\substack{X \\ X}} \alpha_{n+1}((f \otimes 1)(h \otimes g))(c \otimes a \otimes 1_{n+1}),$$

where  $f, g \in C(X^n)$ ,  $h \in C(X)$  and  $a, c \in A$ . Thus  $\alpha_{n+1}$  satisfies the Podleś condition and so by induction  $\alpha_n$  satisfies the Podleś condition for every  $n \geq 1$ .

It remains to show that  $\alpha_n \circ i_{\bullet, \mathbb{K}} = (\text{id}_A \otimes i_{\bullet, \mathbb{K}}) \circ \alpha$  for any  $m < n$ . As in the proof of Proposition 3.6, for any  $n \geq 1$  and  $u, v \in X^n$  we will let  $b_{u, \mathbb{K}} \in A$  be the unique elements satisfying

$$\alpha_n(p_u) = \sum_{v \in X^n} b_{u, \mathbb{K}} \otimes p_v.$$

We know from the same proof that for any  $n \geq 1$  and  $v \in X^n$  we have

$$\sum_{u \in X^n} b_{u, \mathbb{K}} = 1_A.$$

If  $m < n$ , for any  $u \in X^m$  we have

$$\begin{aligned} \alpha_n \circ i_{\bullet, \mathbb{K}}(p_u) &= (\psi \otimes \text{id}_{n-1}) \dots (\text{id}_{\bullet-1} \otimes \psi \otimes \text{id}_{n-\bullet}) (\text{id}_{\bullet} \otimes \alpha_{n-\bullet})(i_{\bullet, \mathbb{K}}(p_u)) \\ &= \sum_{w \in X^{n-m}} (\psi \otimes \text{id}_{n-1}) \dots (\text{id}_{\bullet-1} \otimes \psi \otimes \text{id}_{n-\bullet}) (\text{id}_{\bullet} \otimes \alpha_{n-\bullet})(p_u \otimes p_w) \\ &= \sum_{w \in X^{n-m}} (\psi \otimes \text{id}_{n-1}) \dots (\text{id}_{\bullet-1} \otimes \psi \otimes \text{id}_{n-\bullet}) (p_u \otimes b_{w, \mathbb{K}} \otimes p_z) \\ &= \sum_{z \in X^{n-m}} (\psi \otimes \text{id}_{\bullet-1}) \dots (\text{id}_{\bullet-1} \otimes \psi) \left( p_u \otimes \sum_{w \in X^{n-m}} b_{w, \mathbb{K}} \right) \otimes p_z \\ &= \sum_{z \in X^{n-m}} (\psi \otimes \text{id}_{\bullet-1}) \dots (\text{id}_{\bullet-1} \otimes \psi) (p_u \otimes 1_A) \otimes p_z \\ &= \sum_{z \in X^{n-m}} \alpha_{\bullet}(p_u) \otimes p_z \\ &= \sum_{v \in X^m} b_{u, \mathbb{K}} \otimes \sum_{z \in X^{n-m}} p_v \otimes p_z \\ &= \sum_{v \in X^m} b_{u, \mathbb{K}} \otimes i_{\bullet, \mathbb{K}}(p_v) \\ &= (\text{id}_A \otimes i_{\bullet, \mathbb{K}}) \circ \alpha_{\bullet}(p_u). \end{aligned}$$

So we have that  $(\alpha_n)_{n=1}^{\infty}$  defines an action of  $(A, \Phi)$  on  $X^*$ .

Finally, let  $\pi : \mathbb{A}_X \rightarrow A$  be the homomorphism from Proposition 3.6. We have  $\pi(a_{u \cdot \mathbb{X}}) = b_{u \cdot \mathbb{X}}$  for any  $u, v \in X^n$  and  $n \geq 1$ . For each  $x \in X$  define a homomorphism  $\tilde{\rho}_x : A \rightarrow A$  by

$$\psi(1 \otimes a) = \sum_{x \in X} \tilde{\rho}_x(a) \otimes p_x,$$

where  $a \in A$ . For any  $u \in X^n$  we have

$$\alpha_{n+1}(1 \otimes p_u) = \sum_{y \in X} \alpha_{n+1}(p_{yu}) = \sum_{v \in X^n} \sum_{x \cdot \mathbb{X} \in X} b_{y u \cdot \mathbb{X} v} \otimes p_x \otimes p_v.$$

On the other hand, we know  $\alpha_{n+1} = (\psi \otimes \text{id}_n)(\text{id}_1 \otimes \alpha_n)$  and

$$(\psi \otimes \text{id}_n)(\text{id}_1 \otimes \alpha_n)(1 \otimes p_u) = \sum_{v \in X^n} \psi(1 \otimes b_{u \cdot \mathbb{X}}) \otimes p_v = \sum_{v \in X^n} \sum_{x \in X} \tilde{\rho}_x(b_{u \cdot \mathbb{X}}) \otimes p_x \otimes p_v,$$

and by comparing tensor factors we see that  $\tilde{\rho}_x(b_{u \cdot \mathbb{X}}) = \sum_{y \in X} b_{y u \cdot \mathbb{X} v}$ . Hence, the diagram

$$\begin{array}{ccc} \mathbb{A}_X & \xrightarrow{x} & \mathbb{A}_X \\ \downarrow \iota & & \downarrow \iota \\ \pi(\mathbb{A}_X) & \xrightarrow{\tilde{x}} & \pi(\mathbb{A}_X) \end{array}$$

commutes, and so  $\pi(\mathbb{A}_X) \subseteq A$  is a self-similar quantum group.  $\square$

**Proposition 4.6.** *The following are equivalent*

- (1)  $(A, \Phi)$  is a quantum self-similar group, and
- (2)  $(A, \Phi)$  is a quantum subgroup of  $(\mathbb{A}_X, \Delta)$  and there is a homomorphism  $\psi : C(X) \otimes A \rightarrow A \otimes C(X)$  satisfying the hypotheses of Theorem 4.5.

*Proof.* Theorem 4.5 is the implication (2)  $\implies$  (1). To see (1)  $\implies$  (2) suppose  $(A, \Phi)$  is a quantum self-similar group. By definition there is a surjective quantum group morphism  $q : \mathbb{A}_X \rightarrow A$ . It is routine to check that there is a homomorphism  $\psi : C(X) \otimes A \rightarrow A \otimes C(X)$  satisfying

$$\psi(p_x \otimes q(a_{u \cdot \mathbb{X}})) = \sum_{y \in X} q(a_{x u \cdot \mathbb{X} y}) \otimes p_y.$$

Given  $u, v \in X^n$  we have

$$\begin{aligned} (\Phi \otimes \text{id}_1)\psi(p_x \otimes q(a_{u \cdot \mathbb{X}})) &= \sum_{y \in X} \Phi(q(a_{x u \cdot \mathbb{X} y})) \otimes p_y \\ &= \sum_{w \in X^n} \sum_{y \cdot \mathbb{X} \in X} q(a_{x u \cdot \mathbb{X} w}) \otimes q(a_{z w \cdot \mathbb{X} y}) \otimes p_y \\ &= (\text{id}_A \otimes \psi) \left( \sum_{w \in X^n} \sum_{z \in X} q(a_{x u \cdot \mathbb{X} w}) \otimes p_z \otimes q(a_{w \cdot \mathbb{X}}) \right) \\ &= (\text{id}_A \otimes \psi)(\psi \otimes \text{id}_A) \left( \sum_{w \in X^n} p_x \otimes q(a_{u \cdot \mathbb{X} w}) \otimes q(a_{w \cdot \mathbb{X}}) \right) \\ &= (\text{id}_A \otimes \psi)(\psi \otimes \text{id}_A)(\text{id}_1 \otimes \Phi)(p_x \otimes q(a_{u \cdot \mathbb{X}})), \end{aligned}$$

and so  $\psi$  satisfies (4.2). For (4.3) notice that for any  $q(a) \in A$  and  $z \in X$  we have

$$q(a) \otimes p_z = (1 \otimes p_z)(q(a) \otimes 1)$$

$$\begin{aligned}
 &= \left( \sum_{x \in X} q(a_{x \cdot X}) \otimes p_z \right) (q(a) \otimes 1) \\
 &= \left( \sum_{\substack{x \cdot X \\ w \in X}} q(a_{x \cdot X}) \otimes p_w \right) (q(a_{x \cdot X}) \otimes 1) (q(a) \otimes 1) \\
 &= \sum_{x \in X} \psi(p_x \otimes 1) (q(a_{x \cdot X} a) \otimes 1). \quad \square
 \end{aligned}$$

*Example 4.7.* If  $G$  is a closed subgroup of  $\text{Aut}(X^*)$  which is self-similar, then  $C(G)$  is a commutative self-similar quantum group. The quotient map  $\mathbb{A}_X \rightarrow C(\text{Aut}(X^*))$  takes a generator  $a_{u \cdot X}$  to the indicator function  $f_{u \cdot X}$  defined in Remark 3.8. For a function  $f \in C(\text{Aut}(X^*))$  and  $x \in X$  the restriction homomorphism  $\tilde{\rho}_x$  satisfies  $\tilde{\rho}_x(f)(g) = f(g|_x)$ , for any  $g \in G$ .

## 5. FINITELY CONSTRAINED SELF-SIMILAR QUANTUM GROUPS

**5.1. Classical finitely constrained self-similar groups.** Fix  $d \geq 1$ , and let  $X^{[d]} = \bigcup_{\leq d} X$  be the finite subtree of  $X^*$  of depth  $d$ . The group of automorphism  $\text{Aut}(X^{[d]})$  is a quotient of  $\text{Aut}(X^*)$ , and the quotient map is given by restriction to the finite subtree. We write  $r_d : \text{Aut}(X^*) \rightarrow \text{Aut}(X^{[d]})$  for this restriction map.

Fix a subgroup  $P \leq \text{Aut}(X^{[d]})$ . Define

$$G_P := \{g \in \text{Aut}(X^*) : r_d(g|_w) \in P \text{ for all } w \in X^*\}.$$

By the properties of restriction, if  $g, h \in G_P$ , then for any  $w \in X^*$

$$r_d((gh)|_w) = r_d(g|_{\cdot w} h|_w) = r_d(g|_{\cdot w}) r_d(h|_w) \in P.$$

Likewise,  $r_d(g^{-1}|_w) = r_d(g|_{\leftarrow 1 \cdot w})^{-1} \in P$ . Hence  $G_P$  is a self-similar group, called a *finitely constrained self-similar group*. More details for these groups can be found in [9].

**5.2. Finitely constrained self-similar quantum groups.** Consider the subalgebra  $\mathbb{A}_d \subseteq \mathbb{A}_X$  generated by the elements  $\{a_{u \cdot X} : |u| = |v| \leq d\}$ . Since  $\Delta : \mathbb{A}_d \rightarrow \mathbb{A}_d \otimes \mathbb{A}_d$  the subalgebra  $\mathbb{A}_d$  is a quotient quantum group. The abelianisation of  $\mathbb{A}_d$  is the algebra  $C(\text{Aut}(X^{[d]}))$  of continuous functions on the finite group  $\text{Aut}(X^{[d]})$ .

**Definition 5.1.** Suppose  $\mathbb{P}$  is a quantum subgroup of  $\mathbb{A}_d$ , where  $\mathbb{P} = \mathbb{A}_d/I$ . Denote by  $q_I : \mathbb{A}_d \rightarrow \mathbb{P}$  the quotient map; so  $I = \ker(q_I)$ . We denote by  $J$  the smallest closed 2-sided ideal of  $\mathbb{A}_X$  generated by  $\{\rho_w(I) : w \in X^*\}$ , and by  $A_{\mathbb{P}}$  the quotient  $A_{\mathbb{P}} := \mathbb{A}_X/J$ . In the next result we prove that  $A_{\mathbb{P}}$  is a self-similar quantum group, and we call it a *finitely constrained self-similar quantum group*.

**Proposition 5.2.** *Each  $A_{\mathbb{P}}$  is a self-similar quantum group.*

To prove Proposition 5.2 we need two lemmas. Recall that for  $g, h \in \text{Aut}(X^*), w \in X^*$  we have

$$(gh)|_w = g|_{\cdot w} h|_w.$$

In the first lemma, we establish an analogous relationship between the comultiplication  $\Delta$  on  $\mathbb{A}_X$  and the restriction maps  $\rho_w$ .

**Lemma 5.3.** *For any  $n \geq 1, w \in X^n$  and  $a \in \mathbb{A}_X$  we have*

$$(\Delta \circ \rho_w)(a) = \sum_{y \in X^n} (1 \otimes a_{y \cdot w})(\rho_y \otimes \rho_w)(\Delta(a)).$$

*Proof.* Let  $a_{u \cdot v}$  be a generator of  $\mathbb{A}_X$ , with  $|u| = |v| = k \geq 0$ . Then

$$\begin{aligned} (\Delta \circ \rho_w)(a_{u \cdot v}) &= \Delta \left( \sum_{\substack{X \\ \in X^n}} a_{u \cdot v} \right) \\ &= \sum_{y \in X^n} \sum_{\substack{X \\ \in X^k}} \sum_{\substack{X \\ \in X^n}} a_{u \cdot v} \otimes a_{y \cdot w} \\ &= \sum_{y \in X^n} \sum_{\substack{X \\ \in X^k}} \rho_y(a_{u \cdot X}) \otimes a_{y \cdot w} \\ &= \sum_{y \in X^n} \sum_{\substack{X \\ \in X^k}} \rho_y(a_{u \cdot X}) \otimes a_{y \cdot w} \rho_w(a_{\cdot v}) \\ &= \sum_{y \in X^n} (1 \otimes a_{y \cdot w})(\rho_y \otimes \rho_w) \left( \sum_{\substack{X \\ \in X^k}} a_{u \cdot X} \otimes a_{\cdot v} \right) \\ &= \sum_{y \in X^n} (1 \otimes a_{y \cdot w})(\rho_y \otimes \rho_w)(\Delta(a_{u \cdot v})). \end{aligned}$$

To see that this formula extends to  $\mathbb{A}_X$  it's enough to show that for any  $w \in X^*$  the map

$$a \mapsto \sum_{y \in X^n} (1 \otimes a_{y \cdot w})(\rho_y \otimes \rho_w)(\Delta(a))$$

is linear and multiplicative. Linearity is clear, and multiplicativity follows from the orthogonality of the projections  $1 \otimes a_{y \cdot w}$  and  $1 \otimes a_{z \cdot w}$  for  $y \neq z$ .  $\square$

**Lemma 5.4.** *Consider the quotient maps  $q_I : \mathbb{A}_d \rightarrow \mathbb{A}_d/I$  and  $q_J : \mathbb{A}_X \rightarrow \mathbb{A}_X/J$ . Then for any  $n \geq 1$  and  $y, w \in X^n$*

$$\ker(q_I \otimes q_I) \subseteq \ker((q_J \circ \rho_y) \otimes (q_J \circ \rho_w)).$$

*Proof.* By definition of  $J$  we have  $I \subseteq J \circ \rho_w$  for any  $w \in X^*$ . Therefore there is a commuting diagram

$$\begin{array}{ccc} \mathbb{A}_d & \hookrightarrow & \mathbb{A}_X \\ \downarrow q_I & & \downarrow q_J \circ \rho_w \\ \mathbb{A}_d/I & \xrightarrow{\tilde{w}} & \mathbb{A}_X / \ker(q_J \circ \rho_w) \end{array}$$

Then if  $c \in \ker(q_I \otimes q_I)$  we have

$$(q_J \circ \rho_y) \otimes (q_J \circ \rho_w)(c) = (\pi_y \circ q_I) \otimes (\pi_w \circ q_I)(c) = \pi_y \otimes \pi_w \circ (q_I \otimes q_I)(c) = 0,$$

as required.  $\square$

*Proof of Proposition 5.2.* To see that  $A_{\mathbb{P}}$  is a compact quantum group, it suffices to show that  $J$  is a Woronowicz ideal. In other words, we need to show that  $\Delta(J) \subset \ker(q_J \otimes q_J)$



where  $q_J : \mathbb{A}_X \rightarrow \mathbb{A}_X/J =: A_{\mathbb{P}}$  is the quotient map. Since  $J$  is generated as an ideal by  $\bigcup_{w \in X^*} \rho_w(I)$  it's enough to show that

$$(q_J \otimes q_J)(\Delta \circ \rho_w(i)) = 0$$

for any  $i \in I$  and  $w \in X^*$ . Because  $I$  is a Woronowicz ideal we know that  $\Delta(i) \in \ker(q_I \otimes q_I)$ . Then by Lemmas 5.3 and 5.4 we have

$$\begin{aligned} (q_J \otimes q_J)(\Delta \circ \rho_w(i)) &= (q_J \otimes q_J) \left( \sum_{y \in X^n} (1 \otimes a_{y \cdot \mathbb{X}})(\rho_y \otimes \rho_w)(\Delta(i)) \right) \\ &= \sum_{y \in X^n} (1 \otimes q_J(a_{y \cdot \mathbb{X}}))(q_J \circ \rho_y \otimes q_J \circ \rho_w)(\Delta(i)) \\ &= 0. \end{aligned}$$

Finally,  $A_{\mathbb{P}}$  is self-similar since by definition of  $J$  we have  $\rho_w(J) \subset J$  for any  $w \in X^*$ .  $\square$

**5.3. Free wreath products.** It is well known that for any  $d \geq 1$  the group  $\text{Aut}(X^{[d+1]})$  is isomorphic to the wreath product  $\text{Aut}(X^{[d]}) \wr \text{Sym}(X)$ . Since  $\text{Aut}(X^*)$  is the inverse limit over  $d$  of the groups  $\text{Aut}(X^{[d]})$ , it can be thought as the infinitely iterated wreath product  $\dots \wr \text{Sym}(X) \wr \text{Sym}(X)$ . It follows that  $\text{Aut}(X^*) \cong \text{Aut}(X^*) \wr \text{Sym}(X)$ . More generally, it is shown in [4] that if  $P \leq \text{Sym}(X) = \text{Aut}(X^{[1]})$ , then the finitely constrained self-similar group  $G_P$  is the infinitely iterated wreath product  $\dots \wr P \wr P$ . In this section we prove in Theorem 5.7 an analogue of this result for finitely constrained self-similar quantum groups.

In [3], Bichon constructs a free wreath product of a compact quantum group by the quantum permutation group  $\mathbb{A}_s(n)$ . Bichon also comments in Remark 2.4 of [3] that there is a natural analogue of this construction for free wreath products by quantum subgroups of  $\mathbb{A}_s(n)$ . In this section we formally extend this definition to take free wreath products by any quantum subgroup of  $\mathbb{A}_s(n)$ , and we prove that the finitely constrained self-similar quantum group  $A_{\mathbb{P}}$  induced from a quantum subgroup  $\mathbb{P}$  of  $A_s(n)$  is a free wreath product by  $\mathbb{P}$ . We begin by recalling the definition of the free wreath product from [3]; note that we use our notation  $\mathbb{A}_1$  instead of  $A_s(|X|)$ .

**Definition 5.5.** Let  $X$  be a set of at least two elements. Let  $(A, \Phi)$  be a compact quantum group, and  $\mathbb{P}$  a quantum subgroup of  $\mathbb{A}_1$ . For each  $x \in X$ , we denote by  $\nu_x$  the inclusion of  $A$  in the free product  $C^*$ -algebra  $(\ast_{x \in X} A) \ast \mathbb{P}$ . The *free wreath product* of  $A$  by  $\mathbb{P}$  is the quotient of  $(\ast_{x \in X} A) \ast \mathbb{P}$  by the two-sided ideal generated by the elements

$$\nu_x(a)q_I(a_{x \cdot \mathbb{X}}) - q_I(a_{x \cdot \mathbb{X}})\nu_x(a), \quad x, y \in X, a \in A.$$

The resulting  $C^*$ -algebra is denoted by  $A \ast_{X \cdot \mathbb{X}} \mathbb{P}$ , and the quotient map is denoted by  $q_w$ . If  $X$  is understood, we typically just write  $\hat{A} \ast_w \mathbb{P}$ .

**Theorem 5.6.** *Let  $(A, \Phi)$  be a compact quantum group, and  $\mathbb{P}$  a quantum subgroup of  $\mathbb{A}_1$ . The free wreath product  $A \ast_w \mathbb{P}$  from Definition 5.5 is a compact quantum group with comultiplication  $\Phi_w$  satisfying*

$$(5.1) \quad \Phi_w(q_w(q_I(a_{x \cdot \mathbb{X}}))) = \sum_{z \in X} q_w(q_I(a_{x \cdot \mathbb{X}})) \otimes q_w(q_I(a_{z \cdot \mathbb{X}}))$$

$$(5.2) \quad \Phi_w(q_w(\nu_x(a))) = \sum_{z \in X} (q_w \otimes q_w)((\nu_x \otimes \nu_z)(\Phi(a))(q_I(a_{x \cdot \mathbb{X}}) \otimes 1)),$$

for each  $x, y \in X$  and  $a \in A$ .

*Proof.* Since  $I$  is a Woronowicz ideal, we have  $\Delta|_I \subseteq \ker(q_I \otimes q_I)$ , and so the map  $(q_I \otimes q_I) \circ \Delta_{\mathbb{A}_1}$  descends to a map

$$\phi: \mathbb{P} \rightarrow \mathbb{P} \otimes \mathbb{P} \subseteq ((*_x \in X A) * \mathbb{P})^{\otimes 2}.$$

Then  $(q_w \otimes q_w) \circ \phi: \mathbb{P} \rightarrow (A *_w \mathbb{P})^{\otimes 2}$  satisfies

$$(q_w \otimes q_w) \circ \phi(q_I(a_{x \cdot y})) = \sum_{z \in X} q_w(q_I(a_{x \cdot z})) \otimes q_w(q_I(a_{z \cdot y})) \quad \text{for all } x, y \in X.$$

For each  $x \in X$ , consider the continuous linear map  $\phi_x: A \rightarrow (A *_w \mathbb{P})^{\otimes 2}$  given by

$$\phi_x(a) = (q_w \otimes q_w) \left( \sum_{z \in X} (\nu_x \otimes \nu_z)(\Phi(a))(q_I(a_{x \cdot z}) \otimes 1) \right).$$

We claim that  $\phi_x$  is a homomorphism. To see this, let  $\{a_\lambda = (a_{\lambda \cdot X}) \in M_{d_\lambda}(A) : \lambda \in \Lambda\}$  be a family of matrices satisfying (1)–(3) of Definition 2.2, and  $\mathcal{A}$  be the  $*$ -subalgebra of  $A$  spanned by the entries  $a_{\lambda \cdot X}$ . Let  $a, b \in \mathcal{A}$  and use Sweedler's notation to write  $\Phi(a) = a_{(1)} \otimes a_{(2)}$  and  $\Phi(b) = b_{(1)} \otimes b_{(2)}$ . We have

$$\phi_x(a)\phi_x(b) = \sum_{z, z' \in X} q_w(\nu_x(a_{(1)})q_I(a_{x \cdot z})\nu_x(b_{(1)})q_I(a_{x \cdot z'})) \otimes q_w(\nu_z(a_{(2)})\nu_{z'}(b_{(2)})),$$

and then since

$$\begin{aligned} q_w(\nu_x(a_{(1)})q_I(a_{x \cdot z})\nu_x(b_{(1)})q_I(a_{x \cdot z'})) &= q_w(\nu_x(a_{(1)}b_{(1)})q_I(a_{x \cdot z}a_{x \cdot z'})) \\ &= \delta_{z, z'} q_w(\nu_x(a_{(1)}b_{(1)})q_I(a_{x \cdot z})), \end{aligned}$$

we have

$$\begin{aligned} \phi_x(a)\phi_x(b) &= \sum_{z \in X} q_w(\nu_x(a_{(1)}b_{(1)})q_I(a_{x \cdot z})) \otimes q_w(\nu_z(a_{(2)})\nu_z(b_{(2)})) \\ &= (q_w \otimes q_w) \left( \sum_{z \in X} (\nu_x \otimes \nu_z)(a_{(1)}b_{(1)} \otimes a_{(2)}b_{(2)})(q_I(a_{x \cdot z}) \otimes 1) \right) \\ &= (q_w \otimes q_w) \left( \sum_{z \in X} (\nu_x \otimes \nu_z)(\Phi(ab))(q_I(a_{x \cdot z}) \otimes 1) \right) \\ &= \phi_x(ab). \end{aligned}$$

Since  $\mathcal{A}$  is dense in  $A$ , it follows that  $\phi_x$  is a homomorphism on  $A$ .

The universal property of  $(*_x \in X A) * \mathbb{P}$  now gives a homomorphism  $\tilde{\Phi}: (*_x \in X A) * \mathbb{P} \rightarrow (A *_w \mathbb{P})^{\otimes 2}$  satisfying

$$\begin{aligned} \tilde{\Phi}(q_I(a_{x \cdot y})) &= \sum_{z \in X} q_w(q_I(a_{x \cdot z})) \otimes q_w(q_I(a_{z \cdot y})), \\ \tilde{\Phi}(\nu_x(a)) &= \sum_{z \in X} (q_w \otimes q_w)((\nu_x \otimes \nu_z)(\Phi(a))(q_I(a_{x \cdot z}) \otimes 1)). \end{aligned}$$

For each  $a \in \mathcal{A}, x, y \in X$  we have

$$\tilde{\Phi}(\nu_x(a)q_I(a_{x \cdot y})) = \sum_{z, z' \in X} q_w(\nu_x(a_{(1)})q_I(a_{x \cdot z})q_I(a_{z \cdot y})) \otimes q_w(\nu_z(a_{(2)})q_I(a_{z \cdot y}))$$

$$\begin{aligned}
 &= \sum_{z \cdot \mathbf{X}' \in X} q_w \left( q_I(a_{x \cdot \mathbf{X}}) \nu_x(a_{(1)}) q_I(a_{x \cdot \mathbf{X}'}) \right) \otimes q_w \left( q_I(a_{z' \cdot \mathbf{X}'}) \nu_z(a_{(2)}) \right) \\
 &= \tilde{\Phi}(q_I(a_{x \cdot \mathbf{X}}) \nu_x(a)).
 \end{aligned}$$

It follows that  $\tilde{\Phi}(\nu_x(a) q_I(a_{x \cdot \mathbf{X}})) = \tilde{\Phi}(q_I(a_{x \cdot \mathbf{X}}) \nu_x(a))$  for each  $a \in A, x, y \in X$ , and hence  $\tilde{\Phi}$  descends to the desired  $\Phi_w: A *_w \mathbb{P} \rightarrow (A *_w \mathbb{P})^{\otimes 2}$ .

We now claim that  $(\text{id} \otimes \Phi_w) \circ \Phi_w = (\Phi_w \otimes \text{id}) \circ \Phi_w$ . Since  $\mathcal{A}$  is dense in  $A$ , to see that  $(\text{id} \otimes \Phi_w) \circ \Phi_w$  and  $(\Phi_w \otimes \text{id}) \circ \Phi_w$  agree on each  $q_w(\nu_x(A))$ , it suffices to show that

$$(\text{id} \otimes \Phi_w) \circ \Phi_w(q_w(\nu_x(a_{\cdot \mathbf{X}}))) = (\Phi_w \otimes \text{id}) \circ \Phi_w(q_w(\nu_x(a_{\cdot \mathbf{X}}))), \text{ for all } \lambda \in \Lambda, 1 \leq i, j \leq d .$$

Routine calculations using (5.1) and (5.2) show that both sides of this equation are equal to

$$\sum_{z \cdot \mathbf{X}' \in X} \sum_{1 \leq \mathbf{X} \leq d_\lambda} q_w^{\otimes 3}(\nu_x(a_{\cdot \mathbf{X}}) q_I(a_{x \cdot \mathbf{X}}) \otimes \nu_z(a_{\cdot \mathbf{X}}) q_I(a_{z \cdot \mathbf{X}'}) \otimes \nu_{z'}(a_{\cdot \mathbf{X}})),$$

and hence are equal. So we have  $(\text{id} \otimes \Phi_w) \circ \Phi_w(q_w(\nu_x(A))) = (\Phi_w \otimes \text{id}) \circ \Phi_w(q_w(\nu_x(A)))$  for each  $x \in X$ . It is straightforward to check that evaluating both  $(\text{id} \otimes \Phi_w) \circ \Phi_w$  and  $(\Phi_w \otimes \text{id}) \circ \Phi_w$  at  $q_I(a_{x \cdot \mathbf{X}})$  gives

$$\sum_{z \cdot \mathbf{X}' \in X} q_w^{\otimes 3} \left( q_I^{\otimes 3} (a_{x \cdot \mathbf{X}} \otimes a_{z \cdot \mathbf{X}'} \otimes a_{z' \cdot \mathbf{X}}) \right).$$

Hence we have  $(\text{id} \otimes \Phi_w) \circ \Phi_w = (\Phi_w \otimes \text{id}) \circ \Phi_w$ .

We now define the matrix  $a^X$  by  $a_{x \cdot \mathbf{X}}^X := q_w(q_I(a_{x \cdot \mathbf{X}}))$ , for  $x, y \in X$ ; and for each  $\lambda \in \Lambda, 1 \leq i, j \leq d, x, y \in X$ , the elements

$$a_{\left( \begin{smallmatrix} \mathbf{X} \\ \mathbf{X} \cdot \mathbf{X}' \cdot \mathbf{X} \end{smallmatrix} \right)} := q_w(\nu_x(a_{\cdot \mathbf{X}}) q_I(a_{x \cdot \mathbf{X}})) \in A *_w \mathbb{P},$$

define matrices  $a^{(\cdot \mathbf{X})} = (a_{\left( \begin{smallmatrix} \mathbf{X} \\ \mathbf{X} \cdot \mathbf{X}' \cdot \mathbf{X} \end{smallmatrix} \right)})$ . To finish the proof we have to show that these matrices satisfy (1)–(3) of Definition 2.2.

We have

$$\Phi_w(a_{\left( \begin{smallmatrix} \mathbf{X} \\ \mathbf{X} \cdot \mathbf{X}' \cdot \mathbf{X} \end{smallmatrix} \right)}) = \left( \sum_{z \in X} (q_w \otimes q_w)((\nu_x \otimes \nu_z)(\Phi(a_{\cdot \mathbf{X}}))(q_I(a_{x \cdot \mathbf{X}}) \otimes 1)) \right) (q_w \circ q_I)^{\otimes 2}(\Delta(a_{x \cdot \mathbf{X}})).$$

We know that (1) is satisfied for the matrix  $a^X$ . For each  $z \in X$  we have

$$\begin{aligned}
 &(q_w \otimes q_w)((\nu_x \otimes \nu_z)(\Phi(a_{\cdot \mathbf{X}}))(q_I(a_{x \cdot \mathbf{X}}) \otimes 1))(q_w \circ q_I)^{\otimes 2}(\Delta(a_{x \cdot \mathbf{X}})) \\
 &= (q_w \otimes q_w) \left( (\nu_x \otimes \nu_z) \left( \sum_{1 \leq \mathbf{X} \leq d_\lambda} a_{\cdot \mathbf{X}} \otimes a_{\cdot \mathbf{X}} \right) (q_I(a_{x \cdot \mathbf{X}}) \otimes 1) \right) (q_w \circ q_I)^{\otimes 2}(\Delta(a_{x \cdot \mathbf{X}})) \\
 &= \sum_{1 \leq \mathbf{X} \leq d_\lambda} (q_w(\nu_x(a_{\cdot \mathbf{X}})) \otimes q_w(\nu_z(a_{\cdot \mathbf{X}}))) (q_w(q_I(a_{x \cdot \mathbf{X}})) \otimes 1) (q_w \circ q_I)^{\otimes 2}(\Delta(a_{x \cdot \mathbf{X}})) \\
 &= \sum_{1 \leq \mathbf{X} \leq d_\lambda} (q_w(\nu_x(a_{\cdot \mathbf{X}})) \otimes q_w(\nu_z(a_{\cdot \mathbf{X}}))) \sum_{z' \in X} q_w(q_I(a_{x \cdot \mathbf{X}} a_{x \cdot \mathbf{X}'}) \otimes q_w(q_I(a_{z' \cdot \mathbf{X}}))) \\
 &= \sum_{1 \leq \mathbf{X} \leq d_\lambda} (q_w(\nu_x(a_{\cdot \mathbf{X}})) \otimes q_w(\nu_z(a_{\cdot \mathbf{X}}))) (q_w(q_I(a_{x \cdot \mathbf{X}})) \otimes q_w(q_I(a_{z \cdot \mathbf{X}}))) \\
 &= \sum_{1 \leq \mathbf{X} \leq d_\lambda} q_w(\nu_x(a_{\cdot \mathbf{X}}) q_I(a_{x \cdot \mathbf{X}})) \otimes q_w(\nu_z(a_{\cdot \mathbf{X}}) q_I(a_{z \cdot \mathbf{X}})).
 \end{aligned}$$

It follows that

$$\begin{aligned}\Phi_w(a_{(\bullet, \lambda)}^{(\lambda)})) &= \sum_{z \in X} \sum_{1 \leq i \leq d_\lambda} q_w(\nu_x(a_{\bullet, X})q_I(a_{x, \lambda})) \otimes q_w(\nu_z(a_{\bullet, X})q_I(a_{z, \lambda})) \\ &= \sum_{z \in X} \sum_{1 \leq i \leq d_\lambda} a_{(\bullet, \lambda)}^{(\lambda)} \otimes a_{(\bullet, \lambda)}^{(\lambda)},\end{aligned}$$

and so (1) holds for all matrices  $a^{(\lambda)}$ . To see that  $a^{(\lambda)}$  is invertible, we define  $b^{(\lambda)}$  by

$$b_{(\bullet, \lambda)}^{(\lambda)} := q_w(q_I(a_{y, \lambda})\nu_y((a_{\bullet, \lambda})^{-1})).$$

Then we have

$$\begin{aligned}(a^{(\lambda)}b^{(\lambda)})_{(\bullet, \lambda)} &= \sum_{z \in X} \sum_{1 \leq i \leq d_\lambda} a_{(\bullet, \lambda)}^{(\lambda)} b_{(\bullet, \lambda)}^{(\lambda)} \\ &= q_w \left( \sum_{z \in X} \sum_{1 \leq i \leq d_\lambda} \nu_x(a_{\bullet, X})q_I(a_{x, \lambda})q_I(a_{y, \lambda})\nu_y((a_{\bullet, \lambda})^{-1}) \right) \\ &= \delta_{x, \lambda} q_w \left( \sum_{1 \leq i \leq d_\lambda} \nu_x(a_{\bullet, X}) \left( \sum_{z \in X} q_I(a_{x, \lambda}) \right) \nu_x((a_{\bullet, \lambda})^{-1}) \right) \\ &= \delta_{x, \lambda} q_w \left( \nu_x \left( \sum_{1 \leq i \leq d_\lambda} a_{\bullet, X} (a_{\bullet, \lambda})^{-1} \right) \right) \\ &= \delta_{x, \lambda} q_w(\nu_x((a_{\bullet, \lambda})^{-1})) \\ &= \delta_{x, \lambda} \delta_{\bullet, \lambda} 1.\end{aligned}$$

A similar calculation shows that  $(b^{(\lambda)}a^{(\lambda)})_{(\bullet, \lambda)} = \delta_{x, \lambda} \delta_{\bullet, \lambda} 1$ , and so  $a^{(\lambda)}$  is invertible. Similar calculations also show that  $c^{(\lambda)}$  with entries

$$c_{(\bullet, \lambda)}^{(\lambda)} := q_w(q_I(a_{x, \lambda})\nu_x(((a_{\bullet, \lambda})^{-1}))$$

is the inverse of  $(a^{(\lambda)})^\mathbf{f}$ .

We also have

$$(a^X(a^X)^\mathbf{f})_{x, \lambda} = \sum_{z \in X} a_{x, \lambda}^X a_{y, \lambda}^X = q_w \left( q_I \left( \sum_{z \in X} a_{x, \lambda}^X a_{y, \lambda}^X \right) \right) = \delta_{x, \lambda} 1.$$

Similarly,  $(a^X)^\mathbf{f} a^X$  is the identity. So  $a^X$  and  $(a^X)^\mathbf{f}$  are mutually inverse, and (2) is satisfied.

We now claim that the entries of the matrices  $\{a^{(\lambda)} : \lambda \in \Lambda\} \cup \{a^X\}$  span a dense subset of  $A *_{X, \mathbb{P}} \mathbb{P}$ . For each  $x, y \in X$  we obviously have  $q_w(q_I(a_{x, \lambda}))$  in this span since they are the entries of  $a^X$ . For each  $x \in X$ ,  $\lambda \in \Lambda$  and  $1 \leq i, j \leq d_\lambda$  we have

$$\sum_{y \in X} a_{(\bullet, \lambda)}^{(\lambda)} = q_w \left( \nu_x(a_{\bullet, X})q_I \left( \sum_{y \in X} a_{x, \lambda} \right) \right) = q_w(\nu_x(a_{\bullet, X})),$$

and so each  $q_w(\nu_x(a_{\bullet, X}))$  is in the span of the entries. The claim follows, and so (3) holds.  $\square$

**Theorem 5.7.** *Let  $A_{\mathbb{P}}$  be a finitely-constrained self-similar quantum group in the sense of Definition 5.1. There is a unital quantum group isomorphism  $\pi : A_{\mathbb{P}} \rightarrow A_{\mathbb{P}} *_{w} \mathbb{P}$  satisfying*

$$(5.3) \quad \pi(q_J(a_{xu, \lambda}v)) = q_w(q_I(a_{x, \lambda})\nu_x(q_J(a_{u, \lambda})))$$

for all  $x, y \in X$ ,  $u, v \in X_l$ ,  $m \geq 0$ .

*Proof.* We define  $b_{\emptyset, \cdot}^*$  to be the identity of  $A_{\mathbb{P}} *_{\mathbb{P}} \mathbb{P}$ , and for each  $x, y \in X$ ,  $u, v \in X_l$ ,  $m \geq 0$ ,

$$b_{xu, \cdot}^* := q_w(q_I(a_{x, \cdot})\nu_x(q_J(a_{u, \cdot}))).$$

We claim that this gives a family of projections satisfying (1)–(3) of Definition 3.2. Condition (1) holds by definition. We have

$$\begin{aligned} b_{xu, \cdot}^* &= q_w(\nu_x(q_J(a_{u, \cdot}^*))q_I(a_{x, \cdot}^*)) \\ &= q_w(\nu_x(q_J(a_{u, \cdot}))q_I(a_{x, \cdot})) \\ &= q_w(q_I(a_{x, \cdot})\nu_x(q_J(a_{u, \cdot}))) \\ &= b_{xu, \cdot} \end{aligned}$$

and

$$b_{xu, \cdot}^2 = q_w(q_I(a_{x, \cdot})\nu_x(q_J(a_{u, \cdot}))q_I(a_{x, \cdot})\nu_x(q_J(a_{u, \cdot}))) = q_w(q_I(a_{x, \cdot}^2)\nu_x(q_J(a_{u, \cdot}^2))) = b_{xu, \cdot}.$$

So (2) holds. For each  $w \in X$  we have

$$\begin{aligned} \sum_{z \in X} b_{xuw, \cdot} b_{vz} &= \sum_{z \in X} q_w(q_I(a_{x, \cdot})\nu_x(q_J(a_{uw, \cdot}z))) \\ &= q_w(q_I(a_{x, \cdot})\nu_x(q_J(\sum_{z \in X} a_{uw, \cdot}z))) \\ &= q_w(q_I(a_{x, \cdot})\nu_x(q_J(a_{u, \cdot}))) \\ &= b_{xu, \cdot}, \end{aligned}$$

and

$$\begin{aligned} \sum_{z \in X} b_{xuz, \cdot} b_{vw} &= \sum_{z \in X} q_w(q_I(a_{x, \cdot})\nu_x(q_J(a_{uz, \cdot}w))) \\ &= q_w(q_I(a_{x, \cdot})\nu_x(q_J(\sum_{z \in X} a_{uz, \cdot}w))) \\ &= q_w(q_I(a_{x, \cdot})\nu_x(q_J(a_{u, \cdot}))) \\ &= b_{xu, \cdot}, \end{aligned}$$

and hence (3) holds. This proves the claim, and hence the universal property of  $\mathbb{A}_X$  now gives a homomorphism  $\tilde{\pi}: \mathbb{A}_X \rightarrow A_{\mathbb{P}} *_{\mathbb{P}} \mathbb{P}$  satisfying

$$\tilde{\pi}(a_{xu, \cdot}) = q_w(q_I(a_{x, \cdot})\nu_x(q_J(a_{u, \cdot}))),$$

for all  $x, y \in X$ ,  $u, v \in X_l$ ,  $m \geq 0$ .

We now claim that  $J$  is contained in  $\ker \tilde{\pi}$ . To see this, fix  $w \in X^n$ , with  $w = w_1 w'$  for  $w_1 \in X, w' \in X^{n-1}$ . We first prove the claim that for each  $x := a_{u_1, \cdot} \cdots a_{u_k, \cdot}$ , where  $k \geq 1$  and each pair  $u_i, v_i \in X_l$  for some  $m_i \geq 0$ , we have

$$(5.4) \quad \tilde{\pi}(\rho_w(x)) = \sum_{y \in X} q_w(q_I(a_{y, \cdot})\nu_y(q_J(\rho_{w'}(x)))).$$

Let  $k = 1$ . Then

$$\tilde{\pi}(\rho_w(a_{u_1, \cdot})) = \sum_{y \in X} \sum_{\in X^{n-1}} \tilde{\pi}(a_{y u_1, \cdot} v_1)$$

$$\begin{aligned}
&= \sum_{y \in X} \sum_{\in X^{n-1}} q_w(q_I(a_{y \bullet \mathfrak{X}_1}) \nu_y(q_J(a_{u_1 \bullet \mathfrak{X}' v_1}))) \\
&= \sum_{y \in X} q_w(q_I(a_{y \bullet \mathfrak{X}_1}) \nu_y(q_J(\sum_{\in X^{n-1}} a_{u_1 \bullet \mathfrak{X}' v_1}))) \\
&= \sum_{y \in X} q_w(q_I(a_{y \bullet \mathfrak{X}_1}) \nu_y(q_J(\rho_{w'}(a_{u_1 \bullet \mathfrak{X}_1}))),
\end{aligned}$$

and so (5.4) holds for  $k = 1$ . We now assume true for  $x$ , and prove for  $x_{+1}$ . Note that for  $y, y' \in X$  we have  $q_I(a_{y \bullet \mathfrak{X}_1}) q_I(a_{y' \bullet \mathfrak{X}_1}) = \delta_{y \bullet y'} q_I(a_{y \bullet \mathfrak{X}_1})$ , and hence

$$\begin{aligned}
&q_w(q_I(a_{y \bullet \mathfrak{X}_1}) \nu_y(q_J(\rho_{w'}(x))) q_I(a_{y' \bullet \mathfrak{X}_1}) \nu_{y'}(q_J(\rho_{w'}(a_{u_{k+1} \bullet \mathfrak{X}_{k+1}})))) \\
&= q_w(\nu_y(q_J(\rho_{w'}(x))) q_I(a_{y \bullet \mathfrak{X}_1}) q_I(a_{y' \bullet \mathfrak{X}_1}) \nu_{y'}(q_J(\rho_{w'}(a_{u_{k+1} \bullet \mathfrak{X}_{k+1}})))) \\
&= \delta_{y \bullet y'} q_w(\nu_y(q_J(\rho_{w'}(x))) q_I(a_{y \bullet \mathfrak{X}_1}) \nu_{y'}(q_J(\rho_{w'}(a_{u_{k+1} \bullet \mathfrak{X}_{k+1}})))) \\
&= \delta_{y \bullet y'} q_w(q_I(a_{y' \bullet \mathfrak{X}_1}) \nu_y(q_J(\rho_{w'}(x))) \nu_{y'}(q_J(\rho_{w'}(a_{u_{k+1} \bullet \mathfrak{X}_{k+1}})))) \\
&= \delta_{y \bullet y'} q_w(q_I(a_{y' \bullet \mathfrak{X}_1}) \nu_y(q_J(\rho_{w'}(x a_{u_{k+1} \bullet \mathfrak{X}_{k+1}})))).
\end{aligned}$$

It follows that

$$\tilde{\pi}(\rho_w(x_{+1})) = \tilde{\pi}(\rho_w(x)) \tilde{\pi}(\rho_w(a_{u_{k+1} \bullet \mathfrak{X}_{k+1}})) = \sum_{y \in X} q_w(q_I(a_{y \bullet \mathfrak{X}_1}) \nu_y(q_J(\rho_{w'}(x a_{u_{k+1} \bullet \mathfrak{X}_{k+1}})))) ,$$

and it follows that (5.4) holds for all  $k$ . Since linear combinations of products of the form  $x$  is a dense subalgebra of  $\mathbb{A}_X$ , it follows that

$$\tilde{\pi}(\rho_w(a)) = \sum_{y \in X} q_w(q_I(a_{y \bullet \mathfrak{X}_1}) \nu_y(q_J(\rho_{w'}(a))))$$

for all  $a \in \mathbb{A}_X$ . Now, if  $a \in I$ , then  $\rho_{w'}(a) \in J = \ker q_J$ , and hence the above equations shows that  $\tilde{\pi}(\rho_w(a)) = 0$ . Hence  $\rho_w(a) \in \ker \tilde{\pi}$  for all  $w \in X^n$  and  $a \in I$ , and hence  $J \subseteq \ker \tilde{\pi}$ . This means  $\tilde{\pi}$  descends to a homomorphism  $\pi: A_{\mathbb{P}} \rightarrow A_{\mathbb{P}} *_w \mathbb{P}$  satisfying

$$\pi(q_J(a_{xu \bullet \mathfrak{X} v})) = q_w(q_I(a_{x \bullet \mathfrak{X}}) \nu_x(q_J(a_{u \bullet \mathfrak{X}})))$$

for all  $x, y \in X$ ,  $u, v \in X_i$ ,  $m \geq 0$ .

We now show that  $\pi$  is an isomorphism by finding an inverse. For each  $x \in X$  consider the homomorphism  $q_J \circ \sigma_x: \mathbb{A}_X \rightarrow A_{\mathbb{P}}$ , where  $\sigma_x$  is the homomorphism from Remark 4.3. Since  $\sigma_x = \kappa \circ \rho_x \circ \kappa$ , and we know from [10, Remark 2.10] that  $\kappa(J) \subseteq J$ , it follows that  $q_J \circ \sigma_x$  descends to a homomorphism  $\phi_x: A_{\mathbb{P}} \rightarrow A_{\mathbb{P}}$  satisfying

$$\phi_x(q_J(a_{u \bullet \mathfrak{X}})) = q_J(\sigma_x(a_{u \bullet \mathfrak{X}})) = \sum_{y \in X} q_J(a_{xu \bullet \mathfrak{X} v}),$$

for all  $u, v \in X_i$ ,  $m \geq 0$ .

Each  $\phi_x$ , and the map  $q_I(a) \mapsto q_J(a)$  from  $\mathbb{P}$  to  $A_{\mathbb{P}}$ , now allow us to apply the universal property of the free product  $(*_x \in X A_{\mathbb{P}}) * \mathbb{P}$  to get a homomorphism  $\tilde{\phi}: (*_{x \in X} A_{\mathbb{P}}) * \mathbb{P} \rightarrow A_{\mathbb{P}}$  satisfying  $\tilde{\phi} \circ \nu_x = \phi_x$  for each  $x \in X$ , and  $\tilde{\phi}(q_I(a)) = q_J(a)$  for all  $a \in \mathbb{A}_1 \subseteq \mathbb{A}_X$ . We claim that

$$\tilde{\phi}(\nu_x(q_J(a_{u \bullet \mathfrak{X}})) q_I(a_{x \bullet \mathfrak{X}}) - q_I(a_{x \bullet \mathfrak{X}}) \nu_x(q_J(a_{u \bullet \mathfrak{X}}))) = 0,$$

for each  $x \in X$ ,  $u, v \in X_i$ ,  $m \in \mathbb{N}$ . We have

$$\tilde{\phi}(\nu_x(q_J(a_{u \bullet \mathfrak{X}})) q_I(a_{x \bullet \mathfrak{X}}) - q_I(a_{x \bullet \mathfrak{X}}) \nu_x(q_J(a_{u \bullet \mathfrak{X}})))$$

$$\begin{aligned}
 &= \phi_x(q_J(a_{u\dot{x}}))q_J(a_{x\dot{y}}) - q_G(a_{x\dot{y}})\phi_x(q_J(a_{u\dot{x}})) \\
 &= \sum_{y \in X} q_J(a_{xu\dot{y}v})q_J(a_{x\dot{y}}) - \sum_{y' \in X} q_J(a_{x\dot{y}})q_J(a_{xu\dot{y}'v}) \\
 &= q_J(a_{xu\dot{y}v}) - q_J(a_{xu\dot{y}v}) \\
 &= 0.
 \end{aligned}$$

It follows that  $\tilde{\phi}$  descends to a homomorphism  $\phi: A_{\mathbb{P}} *_w \mathbb{P} \rightarrow A_{\mathbb{P}}$  satisfying

$$\phi(q_w(\nu_x(q_J(a_{u\dot{x}})))) = q_J(\sigma_x(a_{u\dot{x}})) = \sum_{y \in X} q_J(a_{xu\dot{y}v})$$

for all  $x \in X$ ,  $u, v \in X_I$ ,  $m \geq 0$ , and

$$\phi(q_w(q_I(a_{x\dot{y}}))) = q_J(a_{x\dot{y}})$$

for all  $x, y \in X$ .

We claim that  $\pi$  and  $\phi$  are mutually inverse. For  $x, y \in X$ ,  $u, v \in X_I$ ,  $m \geq 0$ , we have

$$\phi(\pi(q_J(a_{xu\dot{y}v}))) = \phi(q_w(q_I(a_{x\dot{y}})\nu_x(q_J(a_{u\dot{x}})))) = q_J(a_{x\dot{y}}) \sum_{y \in X} q_J(a_{xu\dot{y}v}) = q_J(a_{xu\dot{y}v}),$$

and it follows that  $\phi \circ \pi$  is the identity on  $A_{\mathbb{P}}$ . For  $x \in X$ ,  $u, v \in X_I$ ,  $m \geq 0$ , we have

$$\begin{aligned}
 \pi(\phi(q_w(\nu_x(q_J(a_{u\dot{x}})))))) &= \pi\left(\sum_{y \in X} q_J(a_{xu\dot{y}v})\right) \\
 &= \sum_{y \in X} q_w(q_I(a_{x\dot{y}})\nu_x(q_J(a_{u\dot{x}}))) \\
 &= q_w\left(q_I\left(\sum_{y \in X} a_{x\dot{y}}\right)\nu_x(q_J(a_{u\dot{x}}))\right) \\
 &= q_w(\nu_x(q_J(a_{u\dot{x}}))),
 \end{aligned}$$

and for all  $x, w \in X$  we have

$$\pi(\phi(q_w(q_I(a_{x\dot{y}})))) = \pi(q_J(a_{x\dot{y}})) = q_w(q_I(a_{x\dot{y}})).$$

Hence  $\pi \circ \phi$  is the identity on  $A_{\mathbb{P}} *_w \mathbb{P}$ , and so  $\pi$  is an isomorphism.

We now need to show that  $\pi$  is a homomorphism of compact quantum groups, which means that  $\Delta_w \circ \pi = (\pi \otimes \pi) \circ \Delta_J$ , where  $\Delta_J$  is the comultiplication on  $A_{\mathbb{P}}$ . For  $x, y \in X$ ,  $u, v \in X_I$ ,  $m \geq 0$ , we have

$$\begin{aligned}
 &(\pi \otimes \pi) \circ \Delta_J(q_J(a_{xu\dot{y}v})) \\
 &= \sum_{m+1} \pi(q_J(a_{xu\dot{x}})) \otimes \pi(q_J(a_{\dot{y}v})) \\
 &= \sum_{z \in X} \sum_{\in X^m} \pi(q_J(a_{xu\dot{z}})) \otimes \pi(q_J(a_{z\dot{y}v})) \\
 &= \sum_{z \in X} \sum_{\in X^m} q_w(q_I(a_{x\dot{z}})\nu_x(q_J(a_{u\dot{x}}))) \otimes q_w(q_I(a_{z\dot{y}})\nu_x(q_J(a_{\dot{y}v}))).
 \end{aligned}$$

We have

$$\Delta_w \circ \pi(q_J(a_{xu\dot{y}v})) = \Delta_w(q_w(q_I(a_{x\dot{y}}))) \Delta_w(q_w(\nu_x(q_J(a_{u\dot{x}}))),$$

where

$$(5.5) \quad \Delta_w(q_w(q_I(a_{x\dot{X}}))) = \sum_{z \in X} q_w(q_I(a_{x\dot{X}})) \otimes q_w(q_{\mathbb{P}}(a_{z\dot{Y}})),$$

and

$$(5.6) \quad \begin{aligned} \Delta_w(q_w(\nu_x(q_J(a_{u\dot{X}})))) &= \sum_{z' \in X} (q_w \otimes q_w)((\nu_x \otimes \nu_{z'}) (\Delta_J(q_J(a_{u\dot{X}})))(q_I(a_{x\dot{X}}) \otimes 1)) \\ &= \sum_{z' \in X} \sum_{\epsilon \in X^m} q_w(\nu_x(q_J(a_{u\dot{X}}))q_I(a_{x\dot{X}})) \otimes q_w(\nu_{z'}(q_J(a_{\epsilon\dot{X}}))). \end{aligned}$$

A typical summand in the product of the expressions in (5.5) and (5.6) is

$$\begin{aligned} & q_w(q_I(a_{x\dot{X}})\nu_x(q_J(a_{u\dot{X}}))q_I(a_{x\dot{X}})) \otimes q_w(q_{\mathbb{P}}(a_{z\dot{Y}})\nu_{z'}(q_J(a_{\epsilon\dot{X}}))) \\ &= q_w(\nu_x(q_J(a_{u\dot{X}}))q_I(a_{x\dot{X}})q_I(a_{x\dot{X}})) \otimes q_w(q_{\mathbb{P}}(a_{z\dot{Y}})\nu_{z'}(q_J(a_{\epsilon\dot{X}}))) \\ &= \delta_{z\dot{Y}} q_w(\nu_x(q_J(a_{u\dot{X}}))q_I(a_{x\dot{X}})) \otimes q_w(q_{\mathbb{P}}(a_{z\dot{Y}})\nu_z(q_J(a_{\epsilon\dot{X}}))) \\ &= \delta_{z\dot{Y}} q_w(q_I(a_{x\dot{X}})\nu_x(q_J(a_{u\dot{X}}))) \otimes q_w(q_{\mathbb{P}}(a_{z\dot{Y}})\nu_z(q_J(a_{\epsilon\dot{X}}))). \end{aligned}$$

Hence

$$\begin{aligned} \Delta_w \circ \pi(q_J(a_{xu\dot{X}v})) &= \sum_{z \in X} q_w(q_I(a_{x\dot{X}})) \otimes q_w(q_{\mathbb{P}}(a_{z\dot{Y}})) \\ &= \sum_{z \in X} \sum_{\epsilon \in X^m} q_w(q_I(a_{x\dot{X}})\nu_x(q_J(a_{u\dot{X}}))) \otimes q_w(q_I(a_{z\dot{Y}})\nu_x(q_J(a_{\epsilon\dot{X}}))) \\ &= (\pi \otimes \pi) \circ \Delta_J(q_J(a_{xu\dot{X}v})), \end{aligned}$$

and it follows that  $\Delta_w \circ \pi = (\pi \otimes \pi) \circ \Delta_J$ .  $\square$

*Example 5.8.* An immediate consequence of Theorem 5.7 is that  $A_{\mathbb{P}}$  is noncommutative whenever  $\mathbb{P}$  is a noncommutative quantum subgroup of  $\mathbb{A}_1$ . A class of such examples comes from Banica and Bichon's [1, Theorem 1.1], in which they classify all the quantum subgroups  $\mathbb{P}$  of  $\mathbb{A}_1$  for  $|X| = 4$ ; the corresponding list of quantum groups  $A_{\mathbb{P}}$  gives us a list of potentially interesting self-similar quantum groups for further study.

## REFERENCES

- [1] T. Banica, J. Bichon, *Quantum groups acting on 4 points*, J. reine angew. Math. **626** (2009), 75–114.
- [2] J. Bichon, *Quantum automorphism groups of finite graphs*, Proc. Amer. Math. Soc. **131** (2003), 665–673.
- [3] J. Bichon, *Free wreath product by the quantum permutation group*, Algebr. Represent. Theory **7** (2004), 343–362.
- [4] I. Bondarenko, I. Samilovych, *On finite generation of self-similar groups of finite-type*, Internat. J. Algebra Comput. **23** (2013), 69–79.
- [5] R.I. Grigorchuk, *On Burnside's problem on periodic groups*, Func. Anal. Appl. **14** (1980), 41–43.
- [6] R.I. Grigorchuk, *Solved and unsolved problems around one group*, in *Infinite Groups: Geometric, Combinatorial and Dynamical Aspects*, Vol. 248, eds. L. Bartholdi, T. Ceccherini-Silberstein, T. Smirnova-Nagnibeda and A. Zuk, Progress in Mathematics (Birkhäuser Boston, 2005), 117–218.
- [7] I. Raeburn, A. M. Sinclair, *The  $C^*$ -algebra generated by two projections*, Math. Scand. **65** (1989), 278–290.
- [8] L. Rollier, S. Vaes, *Quantum automorphism groups of connected locally finite graphs and quantizations of discrete groups*, [arXiv:2209.03770](https://arxiv.org/abs/2209.03770).
- [9] Z. Šunić, *Pattern closure of groups of tree automorphisms*, Bull. Math. Sci. **1** (2011), 115–127.
- [10] S. Wang, *Free products of compact quantum groups*, Commun. Math. Phys. **167** (1995) 671–692.



- [11] S. Wang, *Quantum symmetry groups of finite spaces*, Commun. Math. Phys. **195** (1998) 195–211.
- [12] S.L. Woronowicz, *Twisted SU(2) group. An example of noncommutative differential calculus*, Publ. RIMS, Kyoto Univ. **23** (1987), 117–181.
- [13] S.L. Woronowicz, *Compact matrix pseudogroups*, Comm. Math. Phys. **111** (1987), 613–665.
- [14] S.L. Woronowicz, *Compact quantum groups*, Symétries quantiques (Les Houches, 1995), North-Holland, Amsterdam (1998) 845–884.
- [15] C. Voigt, *Infinite quantum permutations*, [arXiv:2208.01310](https://arxiv.org/abs/2208.01310).

NATHAN BROWLOWE, SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA.

*Email address:* `nathan.brownlowe@sydney.edu.au`

DAVID ROBERTSON, SCHOOL OF SCIENCE AND TECHNOLOGY, UNIVERSITY OF NEW ENGLAND, NSW 2351, AUSTRALIA.

*Email address:* `david.robertson@une.edu.au`