

Self-consistent effective-one-body theory for spinning binaries based on post-Minkowskian approximation

Jiliang Jing^{*,1,2} Weike Deng,¹ Sheng Long,¹ and Jieci Wang^{†1}

¹*Department of Physics, Key Laboratory of Low Dimensional
Quantum Structures and Quantum Control of Ministry of Education,
and Synergetic Innovation Center for Quantum Effects and Applications,
Hunan Normal University, Changsha, Hunan 410081, P. R. China*

²*Center for Gravitation and Cosmology,
College of Physical Science and Technology,
Yangzhou University, Yangzhou 225009, P. R. China*

Abstract

This paper extends the research on the self-consistent effective-one-body theory of a real spinless two-body system based on the post-Minkowskian approximation (Science China, 65, 100411, (2022)) to the case of a binary system for the spinning black holes. An effective rotating metric and an improved Hamiltonian for the spinning black hole binaries were constructed. The decoupled equation for the null tetrad component of the gravitational perturbed Weyl tensor ψ_4^B in the effective rotating spacetime is found with the help of the gauge transform characteristics of the Weyl tensors. The decoupled equation is then separated between radial and angular variables in the slowly rotating background spacetime, and a formal solution of ψ_4^B is obtained. On this basis, the formal expressions of the radiation reaction force and the waveform for the “plus” and “cross” modes of the gravitational wave are presented. These results, obtained in the same effective spacetime, constitute a self-consistent effective-one-body theory for the spinning black hole binaries based on the post-Minkowskian approximation.

PACS numbers: 04.25.Nx, 04.30.Db, 04.20.Cv

Keywords: real spin two-body system, decoupled equation with separable variables for ψ_4^B , effective-one-body theory

* jljing@hunnu.edu.cn

† jcwang@hunnu.edu.cn

Contents

I. Introduction	3
II. Effective rotating metric and Hamiltonian of the EOB theory for a real spin two-body system	
A. Effective rotating metric of the EOB theory for the real spin two-body system	6
B. Hamiltonian of the EOB theory for the real spin two-body system	7
III. Radiation reaction force and waveform for “plus” and “cross” modes of gravitational waves	
A. Decoupled equation of ψ_4^B for the gravitational perturbation in the effective rotating spacetime	10
B. Radial equation and solution of ψ_4^B for the gravitational perturbation in the effective rotating spacetime	
1. Separation of variables of ψ_4^B in the slowly rotating case	13
2. Tetrad components of the energy-momentum tensor	15
3. Solution of ψ_4^B in the effective rotating spacetime	18
C. Radiation reaction force and waveform for “plus” and “cross” modes of the gravitational wave	22
IV. Conclusions and discussions	23
Acknowledgments	25
References	25

I. INTRODUCTION

The gravitational waves (GWs) [1–6] generated by coalescing compact object binary systems are the most promising sources for the laser interferometer gravitational wave detectors, such as LIGO, Virgo, LISA, KAGRA, Taiji, and TianQin [7–10]. The gravitational waveform template plays a central role in the detection of the GW events generated by coalescing compact object binary systems. Tens of thousands of waveform templates may be needed to extract the GW signal from the noise by means of the matched filtering techniques. The high computational cost and the large parameter space for spinning binary black holes show that it will be impractical for numerical relativity alone to provide a gravitational waveform template bank [11].

To construct the gravitational waveform template for the gravitational radiation generated by coalescing compact object binary systems, the key point is to study the late dynamical evolution of the system. Buonanno and Damour [12] introduced the effective-one-body (EOB) theory based on the post-Newtonian (PN) approximation. The EOB theory is a novel approach to studying the general gravitational radiation generated by coalescing compact object binary systems. It [12–16] presents a theoretical framework that yields a resummed analytical description of the gravitational dynamics of two black holes and becomes the basis of the computation of many gravitational waveform templates [17, 18], which have been used in the data analysis of the gravitational wave signals detected by the LIGO and Virgo interferometers [19–22].

In 2016, Damour [23] developed another EOB theory with a post-Minkowskian (PM) approximation to release the assumption that v/c should be a small quantity in the EOB theory based on the PN approximation. Since then, the correlational research has attracted great attention [24–41]. Along this way, a self-consistent effective-one-body (SCEOB) theory [42] was set up for the real spinless two-body system in PM approximation, which can be applied to the study of the dynamics and the gravitational wave emission of coalescing nonspinning black holes with two mass parameters (m_1, m_2) . However, in general, a black hole has mass and spin. That is, a real spin black hole binary depends on the mass and spin parameters $(m_1, \mathbf{S}_1, m_2, \mathbf{S}_2)$. Therefore, the next important step is to extend the study of the SCEOB theory for the real spinless two-body system [42] to the case of a binary system of spinning black holes.

For the real spin two-body system, the basic idea of the EOB theory is to map the conservative dynamics of two compact objects with masses m_1 and m_2 and spins \mathbf{S}_1 and \mathbf{S}_2 into the dynamics of an effective particle with mass $m_0 = m_1 m_2 / (m_1 + m_2)$ and spin \mathbf{S}_* orbits around a massive black hole described by an effective rotating metric with mass parameter $M_0 = m_1 + m_2$ and rotational parameter a [43]. Thus, the first step in constructing the SCEOB theory for the binary system of spinning black holes is to find the effective rotating metric. Based on the effective metric [42] of a real spinless two-body system for the EOB theory in the PM approximation, an effective rotating metric for a real spin two-body system will be built in this study by means of the approach of constructing an effective rotating metric presented by Damour or Barausse et al. [43, 44].

The second step is to find an improved EOB Hamiltonian. The EOB Hamiltonian that includes spin effects has been studied extensively [16, 45–50]. Damour, Jaranowski, and Schafer [45] presented an EOB Hamiltonian to explore the calibration of numerical relativity waveforms of spinning black holes within the EOB formalism. Later, in the effective Kerr background spacetime, Barausse, Racine, and Buonanno [47] constructed an improved EOB Hamiltonian for a spinning test particle at a linear order of the particle spin. We use the Barausse–Racine–Buonanno proposal to construct an expression of an improved SCEOB Hamiltonian for a spinning test particle that orbits around a massive rotating black hole described by the effective rotating metric.

As a coalescing compact object binary system generates the GWs, the energy of the binary system will be lost, giving rise to the radiation reaction force (RRF). The RRF for these modes of the GWs must be determined to detect the “plus” and “cross” modes of the GWs. Therefore, the third step is to determine the RRF by means of the gravitational wave energy flux for the “plus” and “cross” modes of the GWs. In order to do that, the decoupled equation for the null tetrad component of the perturbed Weyl tensor ψ_4^B must be determined since it relates to the “plus” and “cross” modes of the GWs as $\psi_4^B = \frac{1}{2}(\ddot{h}_+ - i\ddot{h}_\times)$ at infinity. Before our studies, for the gravitational perturbation, the decoupled equation of ψ_4^B was only obtained in the Schwarzschild and Kerr spacetimes. In Ref. [51], by dividing the perturbation part of the metric into the odd and even parities, we found the decoupled equations of the tetrad component of the perturbed Weyl tensor ψ_4^B for the spinless binaries by taking the Regge–Wheeler gauge [52] in the effective metric. In Ref. [42], by means of the gauge transform property of the tetrad components of the perturbed Weyl tensors, we

then obtained a new decoupled equation with separable variables of ψ_4^B by taking a gauge in which ψ_1^B and ψ_3^B vanish in the effective metric. Compared with the two theories, we know that the computational effort for the RRF and waveform in the latter theory will be tremendously reduced. Especially, we know that the study for the spinless binaries in Ref. [42] to the spinning black holes can be extended by checking all conditions to obtain the decoupled equation of ψ_4^B . That is, we can find the decoupled equation of the tetrad component of the perturbed Weyl tensor ψ_4^B in the effective rotating metric and separate the decoupled equation in the radial and angular parts in a slowly rotating background and then obtain the RRF expression by means of the flux for the “plus” and “cross” modes of the GW energy emitted to infinity in the effective rotating spacetime.

The last step is to construct the waveform for the coalescing compact object binary systems. By comparing the tetrad component of the perturbed Weyl tensor ψ_4^B with the formula for the waveform [53] of the “plus” and “cross” modes of the GWs, we can read out the waveform h^{lm} easily, which is also based on the effective rotating spacetime. Furthermore, we can use Damour–Nagar–Pan’s proposals to improve the waveform [11, 48, 49, 54]. That is, based on the obtained waveform, the improved multipolar waveforms can be built as a product of the leading Minkowskian order term, the relativistic conserved energy or angular momentum of the effective moving source, an infinite number of leading logarithms entering the tail effect, an additional phase correction, the remaining PM effects, and a non-quasicircular effect.

The rest of the paper is organized as follows. Section II derives an effective rotating metric and an improved Hamiltonian of the EOB theory for a real spin two-body system. Section III presents a decoupled equation with a separation of variables and a formal solution of ψ_4^B in the effective rotating spacetime. The expressions of the RRF and the waveform for “plus” and “cross” modes of the GWs by means of ψ_4^B are then presented, and an SCEOB theory for the spin binaries based on the PM approximation is set up. The last section presents the final conclusions and discussions.

II. EFFECTIVE ROTATING METRIC AND HAMILTONIAN OF THE EOB THEORY FOR A REAL SPIN TWO-BODY SYSTEM

It is well known that the theoretical model of gravitational radiation is the basis of the gravitational waveform template, and the key point to constructing the theoretical model is to study the late dynamical evolution of a coalescing binary system of compact objects. The EOB theory is a novel approach to studying the two-body dynamics of compact objects with the goal of extending the analytical templates throughout the last stages of inspiral, plunge, merger, and ringdown. The dynamical evolution of the EOB theory for a real spin two-body system can be described by the Hamilton equations [16, 55]:

$$\frac{d\mathbf{r}}{d\hat{t}} = \{\mathbf{r}, \hat{H}_{\text{real}}\} = \frac{\partial \hat{H}_{\text{real}}}{\partial \mathbf{P}}, \quad (2.1a)$$

$$\frac{d\mathbf{P}}{d\hat{t}} = \{\mathbf{P}, \hat{H}_{\text{real}}\} + \hat{\mathcal{F}} = -\frac{\partial \hat{H}_{\text{real}}}{\partial \mathbf{r}} + \hat{\mathcal{F}}, \quad (2.1b)$$

$$\frac{d\mathbf{S}_1}{d\hat{t}} = \{\mathbf{S}_1, \nu \hat{H}_{\text{real}}\} = \nu \frac{\partial \hat{H}_{\text{real}}}{\partial \mathbf{S}_1} \times \mathbf{S}_1, \quad (2.1c)$$

$$\frac{d\mathbf{S}_2}{d\hat{t}} = \{\mathbf{S}_2, \nu \hat{H}_{\text{real}}\} = \nu \frac{\partial \hat{H}_{\text{real}}}{\partial \mathbf{S}_2} \times \mathbf{S}_2, \quad (2.1d)$$

where $\hat{t} \equiv t/M_0$, $\hat{H}_{\text{real}} = H_{\text{real}}/m_0$ is the reduced EOB Hamiltonian [43, 46, 47], and $\hat{\mathcal{F}} = \mathcal{F}/m_0$ is the reduced RRF [50], with $M_0 = m_1 + m_2$, $m_0 = m_1 m_2 / (m_1 + m_2)$, and $\nu = \frac{m_0}{M_0} = \frac{m_1 m_2}{(m_1 + m_2)^2}$.

To get the explicit expression of \hat{H}_{real} in the Hamilton equations (2.1), we should know the effective metric $g_{\mu\nu}^{\text{eff}}$ and the energy map between the EOB and real two-body systems. In this section, we first built up an effective rotating metric of the EOB theory for the real spin two-body system based on the effective metric of the EOB theory for the real spinless two-body system and then found an improved EOB Hamiltonian.

A. Effective rotating metric of the EOB theory for the real spin two-body system

The effective metric of a real spinless two-body system for the EOB theory, in the PM approximation, can be expressed as [42]

$$ds_{\text{eff}}^2 = \frac{\Delta_r^0}{r^2} dt^2 - \frac{r^2}{\Delta_r^0} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\varphi^2), \quad (2.2)$$

with

$$\Delta_r^0 = r^2 - 2GM_0r + \sum_{i=2}^{\infty} a_i \frac{(GM_0)^i}{r^{i-2}}, \quad (2.3)$$

where M_0 is the mass parameter of the black hole and

$$\begin{aligned} a_2 &= \frac{3(1-\Gamma)(1-5\gamma^2)}{\Gamma(3\gamma^2-1)}, \\ a_3 &= \frac{3}{2} \left[\frac{3-2\Gamma-3(15-8\Gamma)\gamma^2+6(25-16\Gamma)\gamma^4}{\Gamma(4\gamma^2-1)(3\gamma^2-1)} - \frac{2P_{30}}{(4\gamma^2-1)} \right], \\ &\dots \dots \end{aligned}$$

in which $\gamma = \frac{1}{2} \frac{\mathcal{E}^2 - m_1^2 - m_2^2}{m_1 m_2}$, $\Gamma = \frac{\mathcal{E}}{m_1 + m_2}$, \mathcal{E} is the energy of the real two-body system, and $P_{30} = \frac{18\gamma^2-1}{2\Gamma^2} + \frac{8\nu(3+12\gamma^2-4\gamma^4)}{\Gamma^2\sqrt{\gamma^2-1}} \operatorname{arcsinh} \sqrt{\frac{\gamma-1}{2}} + \frac{\nu}{\Gamma^2} \left(1 - \frac{103}{3}\gamma - 48\gamma^2 - \frac{2}{3}\gamma^3 + \frac{3\Gamma(1-2\gamma^2)(1-5\gamma^2)}{(1+\Gamma)(1+\gamma)} \right)$.

Based on the effective metric (2.2), using the approach of constructing an effective rotating metric presented by Damour or Barausse et al. [43, 44]¹, we find that the effective rotating metric for a real spin two-body system is described by

$$ds^2 = g_{\mu\nu}^{\text{eff}} dx^\mu dx^\nu = \frac{\Delta_r - a^2 \sin^2 \theta}{\Sigma} dt^2 - \frac{\Sigma}{\Delta_r} dr^2 - \Sigma d\theta^2 - \frac{\Lambda_t \sin^2 \theta}{\Sigma} d\phi^2 + \frac{2\omega_j \sin^2 \theta}{\Sigma} dt d\phi, \quad (2.4)$$

with

$$\begin{aligned} \Sigma &= \bar{\rho}\bar{\rho}^*, \quad \bar{\rho} = r + ia \cos \theta, \quad \bar{\rho}^* = r - ia \cos \theta, \quad \Delta_r = \Delta_r^0 + a^2, \\ \Lambda_t &= \varpi^4 - a^2 \Delta_r \sin^2 \theta, \quad \varpi = (r^2 + a^2)^{\frac{1}{2}}, \quad \omega_j = a(a^2 + r^2 - \Delta_r), \end{aligned}$$

where a is the rotational parameter. The metric can be applied to any post-Minkowskian orders. It is easy to show that the effective spacetime (2.4) is still type D .

B. Hamiltonian of the EOB theory for the real spin two-body system

The relationship between the energy of the EOB system \mathcal{E}_0 and that of a real spinless two-body system \mathcal{E} is given by [42]

$$\mathcal{E}_0 = \frac{\mathcal{E}^2 - m_1^2 - m_2^2}{2(m_1 + m_2)}. \quad (2.5)$$

¹ please note that, starting from the spinless metric (2.2), both the approaches presented by Damour or Barausse et al. will give the same effective rotating metric (2.4).

After Damour [16], we assume that Eq. (2.5) still holds for the real spin two-body system. By means of the effective rotating metric (2.4) and the energy map (2.5), and taking the approach of constructing the EOB Hamiltonian presented by Damour and Barausse et al., [44–47], we can construct an improved EOB Hamiltonian for a spinning black hole described by the effective rotating metric (2.4).

As shown in Ref [47], in curved spacetime, the total effective Hamiltonian of a spinning test particle at a linear order in the particle’s spin can be written as

$$\bar{H}_{\text{eff}}[g_{\mu\nu}^{\text{eff}}] = \bar{H}_{\text{NS}} + \bar{H}_{\text{S}}, \quad (2.6)$$

with

$$\bar{H}_{\text{NS}} = \beta^i P_i + \alpha \sqrt{m_0^2 - \gamma^{ij} P_i P_j + \mathcal{Q}_4(P)}, \quad (2.7)$$

$$\bar{H}_{\text{S}} = \left(\beta^i F_i^K + F_t^K - \frac{\alpha \gamma^{ij} P_i F_j^K}{\sqrt{m_0^2 + \gamma^{ij} P_i P_j}} \right) S_K, \quad (2.8)$$

where \bar{H}_{NS} is the Hamiltonian for a nonspinning particle in the effective rotating metric [13], \bar{H}_{S} is the first-order Hamiltonian in the particle’s spin approximation², $\mathcal{Q}_4(P)$ is a quartic term in the space momenta P_i which was introduced in Ref. [13], $\alpha = \frac{1}{\sqrt{g_{\text{eff}}^{tt}}}$, $\beta^i = \frac{g_{\text{eff}}^{ti}}{g_{\text{eff}}^{tt}}$, $\gamma^{ij} = \left(g_{\text{eff}}^{ij} - \frac{g_{\text{eff}}^{ti} g_{\text{eff}}^{tj}}{g_{\text{eff}}^{tt}} \right)$, S_K is the three-dimensional spin vector [46], and $F_\mu^K = \left(2E_{\mu TI} \frac{\bar{\omega}_J}{\bar{\omega}_T} + E_{\mu IJ} \right) \epsilon^{IJK}$, with $E_{\lambda\mu\nu} = \frac{1}{2} \eta_{AB} \tilde{e}_\mu^A \tilde{e}_{\nu;\lambda}^B$, $\bar{\omega}_\mu = \bar{P}_\mu - m_0 \tilde{e}_\mu^T$, $\bar{P}_i = P_i$, $\bar{P}_t = -\beta^i P_i - \alpha \sqrt{m_0^2 - \gamma^{ij} P_i P_j}$, $\bar{\omega}_T = \bar{\omega}_\mu \tilde{e}_T^\mu = \bar{P}_\mu \tilde{e}_T^\mu - m_0$, and $\bar{\omega}_I = \bar{\omega}_\mu \tilde{e}_I^\mu = \bar{P}_\mu \tilde{e}_I^\mu$.

For the effective rotating metric (2.4), we take the reference tetrad as

$$\begin{aligned} \tilde{e}_\mu^T &= \delta_\mu^t \sqrt{\frac{\Delta_r \Sigma}{\Lambda_t}}, & \tilde{e}_\mu^1 &= \delta_\mu^r \sqrt{\frac{\Sigma}{\Delta_r}}, \\ \tilde{e}_\mu^2 &= \delta_\mu^\theta \sqrt{\Sigma}, & \tilde{e}_\mu^3 &= -\frac{\omega_j \sin \theta}{\sqrt{\Lambda_t \Sigma}} \delta_\mu^t + \delta_\mu^\phi \sin \theta \sqrt{\frac{\Lambda_t}{\Sigma}}. \end{aligned} \quad (2.9)$$

After tedious calculations, we find that the Hamiltonian (2.6) for the effective spacetime (2.4) can be rewritten as

$$\bar{H}_{\text{eff}}[g_{\mu\nu}^{\text{eff}}] = \bar{H}_{\text{NS}} + \mathcal{K}^I S_I, \quad (2.10)$$

with

² We should note that the signature of the spacetime we used in this paper is different from that in Ref. [47].

$$\begin{aligned}
\mathcal{K}^1 &= -\frac{\sqrt{\Delta_r} \cos \theta}{\Lambda_t^2 \sqrt{\Sigma Q} (1 + \sqrt{Q}) \sin^2 \theta} \left[(1 + \sqrt{Q}) (\Delta_r \Sigma^2 + \varpi^4 (\varpi^2 - \Delta_r)) + a \omega_j \varpi^2 \sqrt{Q} \sin^2 \theta \right] \hat{P}_\phi \\
&+ \frac{\Delta_r (r \omega_j \Sigma + a \varpi^2 \xi) \sin \theta}{\Lambda_t^{3/2} \Sigma^2 \sqrt{Q} (1 + \sqrt{Q})} \hat{P}_r \hat{P}_\theta + \frac{a^2 \omega_j \Delta_r \cos \theta \sin^2 \theta}{\Lambda_t^{3/2} \Sigma \sqrt{Q} (1 + \sqrt{Q})} \left(1 + \sqrt{Q} + \frac{2 \Sigma \hat{P}_\phi^2}{\Lambda_t \sin^2 \theta} + \frac{\Delta_r \hat{P}_r^2}{\Sigma} \right), \\
\mathcal{K}^2 &= \left[\frac{\Delta_r (r \Sigma^2 - a^2 \xi \sin^2 \theta)}{\Lambda_t^2 \sqrt{\Sigma Q} \sin \theta} - \frac{(\varpi^4 \xi - r \omega_j^2 \sin^2 \theta)}{\Lambda_t^2 \sqrt{\Sigma} (1 + \sqrt{Q}) \sin \theta} \right] \hat{P}_\phi + \frac{a^2 \omega_j \Delta_r^{3/2} \cos \theta \sin^2 \theta}{\Lambda_t^{3/2} \Sigma^2 \sqrt{Q} (1 + \sqrt{Q})} \hat{P}_r \hat{P}_\theta \\
&+ \frac{\sqrt{\Delta_r} (r \omega_j \Sigma + a \varpi^2 \xi) \sin \theta}{\Lambda_t^{3/2} \Sigma \sqrt{Q} (1 + \sqrt{Q})} \left(1 + \sqrt{Q} + \frac{2 \Sigma}{\Lambda_t \sin^2 \theta} \hat{P}_\phi^2 + \frac{1}{\Sigma} \hat{P}_\theta^2 \right), \\
\mathcal{K}^3 &= -\frac{a^2 \Delta_r \cos \theta \sin \theta}{(\Lambda_t \Sigma)^{3/2} \sqrt{Q} (1 + \sqrt{Q})} (\Lambda_t + \sqrt{Q} \Delta_r \Sigma) \hat{P}_r \\
&- \frac{r \Lambda_t \Delta_r \Sigma + \sqrt{Q} (\Lambda_t (r \Delta_r \Sigma - \varpi^2 \xi) + \omega_j (a \varpi^2 \xi + r \omega_j \Sigma) \sin^2 \theta)}{\Lambda_t^{3/2} \Sigma^{5/2} \sqrt{Q} (1 + \sqrt{Q})} \hat{P}_\theta \\
&- \frac{\sqrt{\Delta_r}}{\Lambda_t^2 \Sigma \sqrt{Q} (1 + \sqrt{Q})} \left[a^2 \omega_j \Delta_r \cos \theta \sin \theta \hat{P}_r + (r \omega_j \Sigma + a \varpi^2 \xi) \hat{P}_\theta \right] \hat{P}_\phi, \tag{2.11}
\end{aligned}$$

where $\xi = a^2 r \sin^2 \theta - r \Delta_r + \frac{1}{2} \Sigma \Delta_r'$, $Q = 1 + \frac{\Delta_r}{\Sigma} \hat{P}_r^2 + \frac{1}{\Sigma} \hat{P}_\theta^2 + \frac{\Sigma}{\Lambda_t \sin^2 \theta} \hat{P}_\phi^2$, and $\hat{P}_i \equiv P_i/m_0$. Here and hereafter, a prime denotes derivation with respect to r .

Thus, by means of the energy map (2.5) and the total effective Hamiltonian (2.10), we find that the Hamiltonian can be expressed as $H_{\text{real}} = M_0 \sqrt{1 + 2\nu (\bar{H}_{\text{eff}}[g_{\mu\nu}^{\text{eff}}]/m_0 - 1)}$. Using the definitions of the reduced Hamiltonians $\hat{H}_{\text{real}} = H_{\text{real}}/m_0$ and $\hat{H}_{\text{eff}}[g_{\mu\nu}^{\text{eff}}] = \bar{H}_{\text{eff}}[g_{\mu\nu}^{\text{eff}}]/m_0$, we then find that an improved reduced EOB Hamiltonian which appeared in Eq (2.1) is described by

$$\hat{H}_{\text{real}} = \frac{1}{\nu} \sqrt{1 + 2\nu (\hat{H}_{\text{eff}}[g_{\mu\nu}^{\text{eff}}] - 1)}, \tag{2.12}$$

which shows that the improved reduced Hamiltonian \hat{H}_{real} is constructed in terms of the effective rotating metric.

III. RADIATION REACTION FORCE AND WAVEFORM FOR “PLUS” AND “CROSS” MODES OF GRAVITATIONAL WAVE

The key step to determine the RRF \mathcal{F} and the waveform h^{lm} for the “plus” and “cross” modes of the gravitational wave is to obtain the decoupled and variables separated equation

for the null tetrad component of the perturbed Weyl tensor ψ_4^B under the gravitational perturbation in the effective rotating spacetime. In this section, we first find the decoupled equation of ψ_4^B in the effective rotating spacetime. By separating the variables for the slowly rotating case, we then obtain the radial equation and its formal solution for ψ_4^B . At last, we present the formal expressions of the RRF and the waveform for the “plus” and “cross” modes of the gravitational wave in the effective rotating spacetime.

A. Decoupled equation of ψ_4^B for the gravitational perturbation in the effective rotating spacetime

Taking the null tetrad for the effective rotating metric (2.4) as

$$\begin{aligned}
l^\mu &= \left\{ \frac{r^2 + a^2}{\Delta_r}, 1, 0, \frac{a}{\Delta_r} \right\}, \\
n^\mu &= \left\{ \frac{r^2 + a^2}{2\rho\rho^*}, -\frac{\Delta_r}{2\rho\rho^*}, 0, \frac{a}{2\rho\rho^*} \right\}, \\
m^\mu &= \frac{1}{\sqrt{2\rho}} \left\{ ia \sin \theta, 0, 1, \frac{i}{\sin \theta} \right\}, \\
\bar{m}^\mu &= \frac{1}{\sqrt{2\rho^*}} \left\{ -ia \sin \theta, 0, 1, -\frac{i}{\sin \theta} \right\},
\end{aligned} \tag{3.1}$$

we find that all nonvanishing spin coefficients, the component of the trace-free Ricci tensor, and the component of the Weyl tensor in the effective spacetime (2.4) are given by

$$\begin{aligned}
\rho &= -\frac{1}{\bar{\rho}^*}, \quad \mu = -\frac{\Delta_r}{2\bar{\rho}\bar{\rho}^{*2}}, \quad \gamma = \frac{\Delta_r'}{4\bar{\rho}\bar{\rho}^*} + \mu, \quad \pi = \frac{ia \sin \theta}{\sqrt{2}\bar{\rho}^{*2}}, \\
\tau &= -\frac{ia \sin \theta}{\sqrt{2}\bar{\rho}\bar{\rho}^*}, \quad \alpha = \pi - \frac{\cot \theta}{2\sqrt{2}\bar{\rho}^*}, \quad \beta = \frac{\cot \theta}{2\sqrt{2}\bar{\rho}}, \\
\phi_{11} &= \frac{1}{8\bar{\rho}^2\bar{\rho}^{*2}} \left[4\Delta_r - 4r\Delta_r' + \bar{\rho}\bar{\rho}^*\Delta_r'' + 2r^2 - 5a^2 - a^2 \cos^2(2\theta) \right], \\
\psi_2 &= \frac{1}{12\bar{\rho}\bar{\rho}^{*3}} \left[12\Delta_r - 6\bar{\rho}^*\Delta_r' + \bar{\rho}^{*2}\Delta_r'' - 2(\bar{\rho}^{*2} + 6a(a + ir \cos \theta)) \right],
\end{aligned} \tag{3.2}$$

The equations in the Newman–Penrose formalism, which will be used to obtain the decoupled equation of ψ_4 , are [56, 57]

$$\Delta\lambda - \bar{\delta}\nu = -(\mu + \bar{\mu})\lambda - (3\gamma - \bar{\gamma})\lambda + (3\alpha + \bar{\beta} + \pi - \bar{\tau})\nu - \psi_4, \tag{3.3}$$

$$\begin{aligned}
\bar{\delta}\psi_3 - D\psi_4 + \bar{\delta}\phi_{21} - \Delta\phi_{20} &= 3\lambda\psi_2 - 2(\alpha + 2\pi)\psi_3 + (4\epsilon - \rho)\psi_4 - 2\nu\phi_{10} + 2\lambda\phi_{11} \\
&+ (2\gamma - 2\bar{\gamma} + \bar{\mu})\phi_{20} + 2(\bar{\tau} - \alpha)\phi_{21} - \bar{\sigma}\phi_{22},
\end{aligned} \tag{3.4}$$

$$\begin{aligned} \Delta\psi_3 - \delta\psi_4 + \bar{\delta}\phi_{22} - \Delta\phi_{21} &= 3\nu\psi_2 - 2(\gamma + 2\mu)\psi_3 + (4\beta - \tau)\psi_4 - 2\nu\phi_{11} - \bar{\nu}\phi_{20} \\ &+ 2\lambda\phi_{12} + 2(\gamma + \bar{\mu})\phi_{21} + (\bar{\tau} - 2\bar{\beta} - 2\alpha)\phi_{22}. \end{aligned} \quad (3.5)$$

The gravitational perturbation in the background spacetime can be described by

$$g_{\mu\nu} = g_{\mu\nu}^A + \varepsilon h_{\mu\nu}^B, \quad (3.6)$$

where the superscripts A and B denote the background and perturbation quantities, respectively, the background spacetime is taken as the effective metric $g_{\mu\nu}^A = g_{\mu\nu}^{\text{eff}}$, and ε is a small quantity. Describing the gravitational perturbation (3.6) by the Newman–Penrose formalism and keeping ε up to the first order as we did in Ref. [42], from Eqs. (3.3), (3.4), and (3.5) we obtain

$$\psi_4^B + (\Delta + 3\gamma - \bar{\gamma} + \mu + \bar{\mu})\lambda^B - (\bar{\delta} + 3\alpha + \bar{\beta} + \pi - \bar{\tau})\nu^B = 0, \quad (3.7)$$

$$\begin{aligned} \nu^B(3\psi_2 - 2\phi_{11}) - (\Delta + 2\gamma + 4\mu)\psi_3^B + (\delta + 4\beta - \tau)\psi_4^B \\ + (\Delta + 2\gamma + 2\bar{\mu})\phi_{21}^B - (\bar{\delta} + 2\alpha + 2\bar{\beta} - \bar{\tau})\phi_{22}^B = 0, \end{aligned} \quad (3.8)$$

$$\begin{aligned} \lambda^B(3\psi_2 + 2\phi_{11}) - (\delta + 2\alpha + 4\pi)\psi_3^B + (D - \rho)\psi_4^B \\ - (\bar{\delta} + 2\alpha - 2\bar{\tau})\phi_{21}^B + (\Delta + 2\gamma - 2\bar{\gamma} + \bar{\mu})\phi_{20}^B = 0. \end{aligned} \quad (3.9)$$

Here and hereafter, for simplicity and clarity, we omit the superscript A .

Chandrasekhar [58] pointed out that, for the linear perturbation described by Eq. (3.6), ψ_0^B and ψ_4^B are gauge invariant, while the ψ_1^B and ψ_3^B are not. Therefore, taking a gauge in which ψ_1^B and ψ_3^B vanish, we find that, in the effective rotating spacetime, the decoupled equation of ψ_4^B is described by

$$\begin{aligned} \left[\left((\Delta + 3\gamma - \bar{\gamma} + \mu + \bar{\mu}) - F_1 \right) (D - \rho) - F_2 \left((\bar{\delta} + 3\alpha + \bar{\beta} + \pi - \bar{\tau}) \right. \right. \\ \left. \left. - F_3 \right) (\delta + 4\beta - \tau) - F_4 \right] \psi_4^B = T_4^B, \end{aligned} \quad (3.10)$$

with

$$\begin{aligned} T_4^B = \left((\Delta + 3\gamma + \mu - \bar{\gamma} + \bar{\mu}) - F_1 \right) \left[(\bar{\delta} + 2\alpha - 2\bar{\tau})\phi_{21}^B - (\Delta + 2\gamma - 2\bar{\gamma} + \bar{\mu})\phi_{20}^B \right] \\ - F_2 \left((\bar{\delta} + 3\alpha + \bar{\beta} + \pi - \bar{\tau}) - F_3 \right) \left[(\bar{\delta} + 2\alpha + 2\bar{\beta} - \bar{\tau})\phi_{22}^B - (\Delta + 2\gamma + 2\bar{\mu})\phi_{21}^B \right], \end{aligned} \quad (3.11)$$

where

$$\begin{aligned}
F_1 &= -\frac{\Delta_r}{2\bar{\rho}\bar{\rho}^*} \frac{\partial \log[3\psi_2 + 2\phi_{11}]}{\partial r}, \\
F_2 &= \frac{3\psi_2 + 2\phi_{11}}{3\psi_2 - 2\phi_{11}}, \\
F_3 &= \frac{1}{\sqrt{2}\bar{\rho}^*} \frac{\partial \log[3\psi_2 - 2\phi_{11}]}{\partial \theta}, \\
F_4 &= 3\psi_2 + 2\phi_{11}.
\end{aligned} \tag{3.12}$$

To simplify the above equations, we introduce

$$\begin{aligned}
\mathcal{D}_n &= \partial_r + \frac{iK}{\Delta_r} + n \frac{\Delta'_r}{\Delta_r}, & \mathcal{D}_n^\dagger &= \partial_r - \frac{iK}{\Delta_r} + n \frac{\Delta'_r}{\Delta_r}, \\
\mathcal{L}_n &= \partial_\theta + Q + n \cot \theta, & \mathcal{L}_n^\dagger &= \partial_\theta - Q + n \cot \theta,
\end{aligned} \tag{3.13}$$

with

$$K = (r^2 + a^2)\sigma^\dagger + ma, \quad Q = \sigma^\dagger a \sin \theta + \frac{m}{\sin \theta}. \tag{3.14}$$

The general perturbation can be expressed as a superposition of different modes with a time and a φ dependence given by $e^{i(\sigma^\dagger t + m\varphi)}$. The intrinsic derivatives in Eqs. (3.10) and (3.11) then become

$$D = \mathcal{D}_0, \quad \Delta = -\frac{\Delta_r}{2\Sigma} \mathcal{D}_0^\dagger, \quad \delta = \frac{1}{\sqrt{2}\bar{\rho}} \mathcal{L}_0^\dagger, \quad \bar{\delta} = \frac{1}{\sqrt{2}\bar{\rho}^*} \mathcal{L}_0. \tag{3.15}$$

Taking $\psi_4^B = (\bar{\rho}^*)^{-4} \phi_4^B$, and using the spin coefficients given by (3.2), we find that the decoupled equation (3.10) can be rewritten as

$$\left[\Delta_r (\mathcal{D}_{-1}^\dagger + \frac{2\bar{\rho}\bar{\rho}^*}{\Delta_r} F_1) (\mathcal{D}_0 - \frac{3}{\bar{\rho}^*}) + F_2 (\mathcal{L}_{-1} - \sqrt{2}\bar{\rho}^* F_3) (\mathcal{L}_2^\dagger - \frac{3ia \sin \theta}{\bar{\rho}^*}) + 2\bar{\rho}\bar{\rho}^* F_4 \right] \phi_4^B = \mathcal{T}_4, \tag{3.16}$$

with

$$\begin{aligned}
\mathcal{T}_4 &= 4\pi G F_4 \left\{ \mathcal{L}_{-1} \left[\frac{\bar{\rho}^*}{3\psi_2 - 2\phi_{11}} \mathcal{L}_0 \left(\frac{1}{\bar{\rho}\bar{\rho}^*} T_{nn} \right) \right] + \frac{\Delta_r^2}{2} \mathcal{D}_0^\dagger \left[\frac{\bar{\rho}^*}{3\psi_2 + 2\phi_{11}} \mathcal{D}_0^\dagger \left(\frac{1}{\bar{\rho}^{-1}\bar{\rho}^*} T_{\bar{m}\bar{m}} \right) \right] \right. \\
&\quad \left. + \frac{\Delta_r^2}{\sqrt{2}} \left\{ \mathcal{D}_0^\dagger \left[\frac{\bar{\rho}^{-2}\bar{\rho}^*}{\Delta_r(3\psi_2 + 2\phi_{11})} \mathcal{L}_{-1} \left(\frac{1}{\bar{\rho}^2\bar{\rho}^*} T_{\bar{m}n} \right) \right] + \mathcal{L}_{-1} \left[\frac{\bar{\rho}^{-2}\bar{\rho}^*}{3\psi_2 - 2\phi_{11}} \mathcal{D}_0^\dagger \left(\frac{1}{\Delta_r} T_{\bar{m}n} \right) \right] \right\} \right\},
\end{aligned} \tag{3.17}$$

where $T_{nn} = \phi_{22}^B/(4\pi G)$, $T_{\bar{m}\bar{m}} = \phi_{21}^B/(4\pi G)$, and $T_{\bar{m}n} = \phi_{20}^B/(4\pi G)$.

It should be noted that although ψ_4^B is decoupled by taking the gauge in which ψ_1^B and ψ_3^B vanish, Eq. (3.16) with $a_i = 0$ ($i \geq 2$) is the same as that of the Kerr spacetime obtained in Ref. [56]. This clearly shows that ψ_4^B is gauge invariant for the linear perturbation described by (3.6).

B. Radial equation and solution of ψ_4^B for the gravitational perturbation in the effective rotating spacetime

To find the radial equation and the solution of ψ_4^B , we should separate the variables of Eq. (3.16) and work out the tetrad components of the energy momentum tensor of the system.

1. Separation of variables of ψ_4^B in the slowly rotating case

It is clear that Eq. (3.16) does not appear to be separable directly. However, if a slowly rotating background is close enough to spherical symmetry, we can separate the perturbation gravitational wave equation (3.16) in the radial and angular parts. In principle, the method can be extended to any order of the rotational parameter. Here, the perturbation equation is derived explicitly up to the first order.

For the slowly rotating case, we assume that ϕ_4^B can be decomposed into Fourier harmonic components according to

$$\phi_4^B = \sum_{\ell m} \frac{1}{\sqrt{2\pi}} \int d\omega e^{-i(\omega t - m\varphi)} {}_{-2}Y_{\ell m}(\theta) R_{\ell m}(r), \quad (3.18)$$

where we take $\sigma^\dagger = -\omega$, the angular function ${}_{-2}Y_{\ell m}(\theta)$ is called the spin-weighted spheroidal harmonic, which is normalized as

$$\int_0^\pi {}_{-2}Y_{\ell m}^*(\theta) {}_{-2}Y_{\ell m}(\theta) \sin\theta d\theta = 1. \quad (3.19)$$

We can then expand Eq. (3.16) to the first order in the rotation parameter a , which is expressed as

$$\mathcal{A}(\ell) \phi_4^B + a \mathcal{B} \phi_4^B - a \cos\theta \mathcal{C}(\ell) \phi_4^B = \mathcal{T}_4^{(0)} + a \mathcal{T}_4^{(1)}, \quad (3.20)$$

with

$$\begin{aligned}
\mathcal{A}(\ell) &= \Delta_r^0 \left[\mathcal{D}_{-1}^{0\dagger} \mathcal{D}_0^0 - \left(\frac{3}{r} \mathcal{D}_{-1}^{0\dagger} + F_{10} \mathcal{D}_0^0 \right) + \frac{3}{r^2} (1 + r F_{10}) \right] + 2r^2 F_{40} - F_{20} \lambda(\ell), \\
\mathcal{B} &= im \left[\left(\frac{3}{r} - F_{10} - \frac{2\Delta_r^{0'}}{\Delta_r^0} + \frac{2i r^2 \omega}{\Delta_r^0} \right) - 2F_{20} \left(i\omega + \frac{3}{r} \right) \right], \\
\mathcal{C}(\ell) &= i\Delta_r^0 \left[\left(\frac{3}{r^2} \mathcal{D}_{-1}^{0\dagger} + F_{11} \mathcal{D}_0^0 \right) - \frac{3}{r^3} (2 + r F_{10} + r^2 F_{11}) \right] \\
&\quad + 2F_{20} \left(\omega - \frac{3i}{r} \right) - 2ir^2 F_{41} - iF_{21} \lambda(\ell), \\
\lambda(\ell) &= (\ell + 2)(\ell - 1),
\end{aligned} \tag{3.21}$$

where we use the relation $(\mathcal{L}_{-1}^0 \mathcal{L}_2^{0\dagger})_{-2} Y_{\ell m}(\theta) = -\lambda(\ell)_{-2} Y_{\ell m}(\theta)$. Here and hereafter all quantities with the superscripts “0” mean that they are fixed at $a = 0$, and the functions

F_{ij} ($i = 1, 2, 3, 4$, and $j = 0, 1$) are defined as

$$\begin{aligned}
F_{10} &= -\frac{32\Delta_r^0 - 23r\Delta_r^{0'} + 7r^2\Delta_r^{0''} - r^3\Delta_r^{0(3)}}{r[8\Delta_r^0 - r(5\Delta_r^{0'} - r\Delta_r^{0''})]}, \\
F_{11} &= -\frac{3}{r^2[8\Delta_r^0 - r(5\Delta_r^{0'} - r\Delta_r^{0''})]^2} \left\{ 32(\Delta_r^0)^2 + 4r\Delta_r^0 [r(4 + r\Delta_r^{0(3)}) - 10\Delta_r^{0'}] \right. \\
&\quad \left. + r^2 [17(\Delta_r^{0'})^2 + r^2 (\Delta_r^{0''} (8 + \Delta_r^{0''}) - 2r\Delta_r^{0(3)}) - r\Delta_r^{0'} (16 + 6\Delta_r^{0''} + r\Delta_r^{0(3)})] \right\}, \\
F_{20} &= \frac{8\Delta_r^0 - r(5\Delta_r^{0'} - r\Delta_r^{0''})}{4\Delta_r^0 - r(2r + \Delta_r^{0'})}, \\
F_{21} &= -\frac{3}{r} \frac{4\Delta_r^0 - r[4\Delta_r^{0'} - r(2 + \Delta_r^{0''})]}{4\Delta_r^0 - r(2r + \Delta_r^{0'})}, \\
F_{40} &= \frac{1}{2r^4} [8\Delta_r^0 - r(5\Delta_r^{0'} - r\Delta_r^{0''})], \\
F_{41} &= \frac{3}{2r^5} [4\Delta_r^0 - r(2r + \Delta_r^{0'})].
\end{aligned} \tag{3.22}$$

By means of the identity [59]

$$\cos \theta_s Y_{\ell m} = \mathcal{Q}_{\ell+1s} Y_{\ell+1m} - \frac{ms}{\ell(\ell+1)}_s Y_{\ell m} + \mathcal{Q}_{\ell s} Y_{\ell-1m}, \tag{3.23}$$

with $\mathcal{Q}_\ell = \frac{1}{\ell} \sqrt{\frac{(\ell-s)(\ell+s)(\ell-m)(\ell+m)}{(2\ell-1)(2\ell+1)}}$, we find that the separation of variables of the gravitational wave equation Eq. (3.20) in the radial parts for the slowly rotating case is explicitly given by

$$\mathcal{A}(\ell) R_{\ell m} + a \left[\bar{\mathcal{B}} R_{\ell m} - \mathcal{C}(\ell-1, m) \mathcal{Q}_\ell R_{\ell-1m} - \mathcal{C}(\ell+1, m) \mathcal{Q}_{\ell+1} R_{\ell+1m} \right] = T_{\ell m \omega}(r), \tag{3.24}$$

where $\bar{\mathcal{B}} = \mathcal{B} + \frac{ms}{\ell(\ell+1)} \mathcal{C}(\ell, m)$, and

$$T_{\ell m \omega}(r) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} dt \int d\Omega \left(\mathcal{T}_4^{(0)} + a \mathcal{T}_4^{(1)} \right) e^{i(\omega t - m\varphi)} \frac{{}_{-2}Y_{\ell m}^*(\theta)}{\sqrt{2\pi}}. \tag{3.25}$$

2. Tetrad components of the energy-momentum tensor

By means of the definition in Ref. [60], we know that the energy-momentum tensor for the EOB theory, i.e., a particle that orbits around a massive black hole described by the effective metric, can be expressed as

$$T^{\mu\nu} = \frac{m_0}{\Sigma \sin \theta} \frac{dx^\mu}{dt/d\tau} \frac{dx^\nu}{d\tau} \delta(r - r(t)) \delta(\theta - \theta(t)) \delta(\varphi - \varphi(t)), \quad (3.26)$$

where $x^\mu = (t, r(t), \theta(t), \varphi(t))$ is a geodesic trajectory and τ is the proper time along the geodesic. The geodesic equations in the effective metric are

$$\begin{aligned} \Sigma \frac{d\theta}{d\tau} &= \pm \left[\mathcal{C} - \cos^2 \theta \left\{ a^2(1 - E^2) + \frac{L^2}{\sin^2 \theta} \right\} \right]^{1/2}, \\ \Sigma \frac{d\varphi}{d\tau} &= - \left(aE - \frac{L}{\sin^2 \theta} \right) + \frac{a}{\Delta_r} \left(E(r^2 + a^2) - aL \right), \\ \Sigma \frac{dt}{d\tau} &= - \left(aE - \frac{L}{\sin^2 \theta} \right) a \sin^2 \theta + \frac{r^2 + a^2}{\Delta_r} \left(E(r^2 + a^2) - aL \right), \\ \Sigma \frac{dr}{d\tau} &= \pm \left[[E(r^2 + a^2) - aL]^2 - \Delta_r [(Ea - L)^2 + r^2 + \mathcal{C}] \right]^{1/2}, \end{aligned} \quad (3.27)$$

where E , L , and \mathcal{C} are the energy, the z -component of the angular momentum [61], and the Carter constant of a test particle, respectively.

The tetrad components of the energy-momentum tensor can then be expressed as

$$\begin{aligned} T_{nn} &= m_0 \frac{C_{nn}}{\sin \theta} \delta(r - r(t)) \delta(\theta - \theta(t)) \delta(\varphi - \varphi(t)), \\ T_{\bar{m}n} &= m_0 \frac{C_{\bar{m}n}}{\sin \theta} \delta(r - r(t)) \delta(\theta - \theta(t)) \delta(\varphi - \varphi(t)), \\ T_{\bar{m}\bar{m}} &= m_0 \frac{C_{\bar{m}\bar{m}}}{\sin \theta} \delta(r - r(t)) \delta(\theta - \theta(t)) \delta(\varphi - \varphi(t)), \end{aligned} \quad (3.28)$$

with

$$\begin{aligned} C_{nn} &= \frac{1}{4\Sigma^3 \dot{t}} \left[E(r^2 + a^2) - aL + \Sigma \frac{dr}{d\tau} \right]^2, \\ C_{\bar{m}n} &= - \frac{1}{2\sqrt{2}\bar{\rho}^* \Sigma^2 \dot{t}} \left[E(r^2 + a^2) - aL + \Sigma \frac{dr}{d\tau} \right] \left[i \sin \theta \left(aE - \frac{L}{\sin^2 \theta} \right) + \Sigma \frac{d\theta}{d\tau} \right], \\ C_{\bar{m}\bar{m}} &= \frac{1}{2\bar{\rho}^{*2} \Sigma \dot{t}} \left[i \sin \theta \left(aE - \frac{L}{\sin^2 \theta} \right) + \Sigma \frac{d\theta}{d\tau} \right]^2, \end{aligned} \quad (3.29)$$

where $\dot{t} = dt/d\tau$.

Using Eqs. (3.17), (3.25), and (3.28), and performing integration by part, we obtain

$$\begin{aligned}
T_{\ell m \omega}(r) = & -\frac{4m_0 G}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dt \int d\theta e^{i\omega t - im\varphi(t)} \\
& \times \left\{ -\frac{1}{2} \mathcal{L}_1^\dagger \left[\frac{\bar{\rho}^*}{F_5} \mathcal{L}_2^\dagger ({}_{-2}Y_{\ell m} F_4) \right] C_{nn} \bar{\rho}^{*2} \bar{\rho} \delta(r - r(t)) \delta(\theta - \theta(t)) \right. \\
& + \frac{\Delta_r^2 \bar{\rho}^*}{2\sqrt{2}\bar{\rho}^2} \left((1 + F_2) \mathcal{L}_2^\dagger {}_{-2}Y_{\ell m} + \frac{{}_{-2}Y_{\ell m} \bar{\rho}^2}{\bar{\rho}^*} \frac{\partial}{\partial \theta} \frac{\bar{\rho}^*}{\bar{\rho}^2} + \frac{{}_{-2}Y_{\ell m}}{F_5} \frac{\partial F_4}{\partial \theta} \right) \\
& \times \mathcal{D}_0^\dagger \left[\frac{C_{\bar{m}n} \bar{\rho}^{*2} \bar{\rho}^2}{\Delta_r} \delta(r - r(t)) \delta(\theta - \theta(t)) \right] \\
& + \frac{1}{2\sqrt{2}} \mathcal{L}_2^\dagger \left[F_4 {}_{-2}Y_{\ell m} \frac{\partial}{\partial r} \left(\frac{\bar{\rho}^*}{\bar{\rho}^2 F_4} \right) \right] C_{\bar{m}n} \Delta_r \bar{\rho}^{*2} \bar{\rho}^2 \delta(r - r(t)) \delta(\theta - \theta(t)) \\
& \left. - \frac{1}{4} \Delta_r^2 {}_{-2}Y_{\ell m} F_4 \mathcal{D}_0^\dagger \left[\frac{\bar{\rho}^*}{F_4} \mathcal{D}_0^\dagger \left(\bar{\rho}^{-1} \bar{\rho}^{*2} C_{\bar{m}\bar{m}} \delta(r - r(t)) \delta(\theta - \theta(t)) \right) \right] \right\}, \quad (3.30)
\end{aligned}$$

where $F_5 = 3\psi_4^B - 2\phi_{11}$. We now expand Eq. (3.30) for $T_{\ell m \omega}(r)$ to the first order in the rotation parameter a , i.e.,

$$T_{\ell m \omega}(r) = T_{\ell m \omega}^{(0)}(r) + a T_{\ell m \omega}^{(1)}(r), \quad (3.31)$$

with

$$T_{\ell m \omega}^{(0)}(r) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} dt \int d\Omega \mathcal{T}_4^{(0)} e^{i(\omega t - m\varphi)} \frac{{}_{-2}Y_{\ell m}^*(\theta)}{\sqrt{2\pi}}, \quad (3.32)$$

$$T_{\ell m \omega}^{(1)}(r) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} dt \int d\Omega \mathcal{T}_4^{(1)} e^{i(\omega t - m\varphi)} \frac{{}_{-2}Y_{\ell m}^*(\theta)}{\sqrt{2\pi}}. \quad (3.33)$$

2-1. Zeroth-order source

For a source bounded in a finite range of r , it is convenient to rewrite the zeroth order of the source Eq. (3.32) as

$$\begin{aligned}
T_{\ell m \omega}^{(0)}(r) = & -m_0 G \int_{-\infty}^{\infty} dt e^{i\omega t - im\varphi(t)} (\Delta_r^0)^2 \left\{ (A_{nn}^{(0)} + A_{\bar{m}n}^{(0)} + A_{\bar{m}\bar{m}}^{(0)}) \delta(r - r(t)) \right. \\
& \left. + \left[(A_{\bar{m}n}^{(0)} + A_{\bar{m}\bar{m}}^{(0)}) \delta(r - r(t)) \right]' + \left[A_{\bar{m}\bar{m}}^{(0)} \delta(r - r(t)) \right]'' \right\}, \quad (3.34)
\end{aligned}$$

with

$$\begin{aligned}
A_{nn}^{(0)} &= -\frac{2r^4}{\sqrt{2\pi} (\Delta_r^0)^2} C_{nn}^{(0)} F_{20} \mathcal{L}_1^{0\dagger} \left[\mathcal{L}_2^{0\dagger} ({}_{-2}Y_{\ell m}(\theta)) \right], \\
A_{\bar{m}n}^{(0)} &= \frac{r^3}{\sqrt{\pi} \Delta_r^0} C_{\bar{m}n}^{(0)} \left[- (1 + F_{20}) \frac{iK^0}{\Delta_r^0} + \frac{F_{20}}{r} - F_{20,r} - \frac{F_{40,r}}{F_{40}} \right] \mathcal{L}_2^{0\dagger} ({}_{-2}Y_{\ell m}(\theta)),
\end{aligned}$$

$$\begin{aligned}
A_{\overline{mn}0}^{(0)} &= -\frac{r^2}{\sqrt{2\pi}} C_{\overline{mn}}^{(0)} {}_{-2}Y_{\ell m}(\theta) \left[i \left(\frac{K^0}{\Delta_r^0} \right)' - \frac{(K^0)^2}{(\Delta_r^0)^2} \right. \\
&\quad \left. + \frac{iK^0}{\Delta_r^0} \left(\frac{1}{r} + \frac{F_{40,r}}{F_{40}} \right) + \frac{F_{40,r}}{rF_{40}} + \left(\frac{F_{40,r}}{F_{40}} \right)' \right], \\
A_{\overline{mn}1}^{(0)} &= \frac{r^3}{\sqrt{\pi}\Delta_r^0} C_{\overline{mn}}^{(0)} (1 + F_{10}) \mathcal{L}_2^{0\dagger} \left({}_{-2}Y_{\ell m}(\theta) \right), \\
A_{\overline{mn}1}^{(0)} &= \frac{r^2}{\sqrt{2\pi}} C_{\overline{mn}}^{(0)} {}_{-2}Y_{\ell m}(\theta) \left(2i \frac{K^0}{\Delta_r^0} + \frac{1}{r} + \frac{F_{40,r}}{F_{40}} \right), \\
A_{\overline{mn}2}^{(0)} &= -\frac{r^2}{\sqrt{2\pi}} C_{\overline{mn}}^{(0)} {}_{-2}Y_{\ell m}(\theta), \tag{3.35}
\end{aligned}$$

where the functions $C_{ij}^{(0)}$ ($i, j = n, \overline{m}$) are given by

$$\begin{aligned}
C_{nn}^{(0)} &= \frac{\Delta_r^0}{4r^4 E} \left(E + \frac{dr}{d\tau} \right)^2, \\
C_{\overline{mn}}^{(0)} &= \frac{\Delta_r^0}{2\sqrt{2}r^3 E} \left(E + \frac{dr}{d\tau} \right) \left(\frac{iL}{r^2 \sin \theta} - \frac{d\theta}{d\tau} \right), \\
C_{\overline{mm}}^{(0)} &= \frac{\Delta_r^0}{2r^2 E} \left(\frac{d\theta}{d\tau} - \frac{iL}{r^2 \sin \theta} \right)^2. \tag{3.36}
\end{aligned}$$

2-2. First-order source

The first order of the source Eq. (3.33) can be expressed as

$$\begin{aligned}
T_{\ell m \omega}^{(1)}(r) &= -m_0 G \int_{-\infty}^{\infty} dt e^{i\omega t - im\varphi(t)} (\Delta_r^0)^2 \left\{ (A_{nn0}^{(1)} + A_{\overline{mn}0}^{(1)} + A_{\overline{mm}0}^{(1)}) \delta(r - r(t)) \right. \\
&\quad \left. + \left[(A_{\overline{mn}1}^{(1)} + A_{\overline{mm}1}^{(1)}) \delta(r - r(t)) \right]' + \left[A_{\overline{mm}2}^{(1)} \delta(r - r(t)) \right]'' \right\}, \tag{3.37}
\end{aligned}$$

with

$$\begin{aligned}
A_{nn0}^{(1)} &= -\frac{2r^4 C_{nn}^{(0)}}{\sqrt{2\pi}\Delta_r^0{}^2} \left\{ F_{20} \left[\left(\omega \sin \theta + \left(\frac{C_{nn}^{(1)}}{C_{nn}^{(0)}} - \frac{i \cos \theta}{r} \right) \mathcal{L}_1^{0+} \right) \mathcal{L}_2^{0+} + \omega \mathcal{L}_1^{0+} \sin \theta \right] {}_{-2}Y_{\ell m}(\theta) \right. \\
&\quad \left. - \frac{i F_{40}}{F_{50}} \left[\left(\frac{F_{51}}{F_{50}} + \frac{1}{r} \right) \mathcal{L}_1^{0+} \cos \theta \mathcal{L}_2^{0+} ({}_{-2}Y_{\ell m}(\theta)) - \frac{F_{41}}{F_{40}} \mathcal{L}_1^{0+} \mathcal{L}_2^{0+} (\cos \theta {}_{-2}Y_{\ell m}(\theta)) \right] \right\}, \\
A_{\overline{mn}0}^{(1)} &= -\frac{r^4}{\sqrt{\pi}\Delta_r^0} \left\{ \frac{1}{r^3} \left[\left(1 + r \frac{F'_{40}}{F_{40}} \right) r C_{\overline{mn}}^{(1)} \mathcal{L}_2^{0\dagger} - i \left(6 + 3r \frac{F'_{40}}{F_{40}} - r^2 \frac{F'_{41}}{F_{40}} \right) C_{\overline{mn}}^{(0)} \mathcal{L}_2^{0\dagger} \cos \theta \right. \right. \\
&\quad \left. \left. - ir^2 \frac{F'_{40}}{F_{40}} \frac{F_{41}}{F_{40}} C_{\overline{mn}}^{(0)} \mathcal{L}_2^{0\dagger} \cos \theta \right] + \frac{1}{r^2} \left(1 + r \frac{F'_{40}}{F_{40}} \right) C_{\overline{mn}}^{(0)} \omega \sin \theta \right. \\
&\quad \left. + \left\{ \left(C_{\overline{mn}}^{(1)} \mathcal{D}_0^0 + \frac{i m}{\Delta_r^0} C_{\overline{mn}}^{(0)} \right) \frac{1 + F_{20}}{r} - 3i \cos \theta \mathcal{D}_0^0 \frac{1 + F_{20}}{r^2} \right\} \mathcal{L}_2^{0\dagger} \right. \\
&\quad \left. + C_{\overline{mn}}^{(0)} \mathcal{D}_0^0 \left[\frac{1}{r} \left(i F_{21} \cos \theta \mathcal{L}_2^{0\dagger} + \left((1 + F_{20}) \omega + \frac{3i}{r} - \frac{i F_{41}}{F_{50}} \right) \sin \theta \right) \right] \right\} {}_{-2}Y_{\ell m}(\theta),
\end{aligned}$$

$$\begin{aligned}
A_{mm0}^{(1)} &= -\frac{r^2}{\sqrt{2\pi}} \left(C_{mm}^{(1)} \mathcal{H}_0 + C_{mm}^{(0)} \mathcal{H}_1 - \frac{4ia \cos \theta}{r} C_{mm}^{(0)} \mathcal{H}_0 \right) {}_{-2}Y_{\ell m}(\theta), \\
A_{mn1}^{(1)} &= \frac{r^3}{\sqrt{\pi} \Delta_r^0} \left\{ (1 + F_{20}) \left(C_{mn}^{(1)} - \frac{3i \cos \theta}{r} C_{mn}^{(0)} \right) \mathcal{L}_2^{0\dagger} \right. \\
&\quad \left. + C_{mn}^{(0)} \left[i F_{21} \cos \theta \mathcal{L}_2^{0\dagger} + \left((1 + F_{20}) \omega + \frac{3i}{r} + \frac{i F_{41}}{F_{50}} \right) \sin \theta \right] \right\} {}_{-2}Y_{\ell m}(\theta), \\
A_{mm1}^{(1)} &= \frac{r^2}{\sqrt{2\pi}} \left\{ C_{mm}^{(0)} \left(i \cos \theta \frac{F_{40} F'_{41} - F_{41} F'_{40}}{F_{40}^2} + \frac{2im}{\Delta_r^0} + \frac{i \cos \theta}{r^2} \right) \right. \\
&\quad \left. + \left(2i \frac{K^0}{\Delta_r^0} + \frac{1}{r} + \frac{F_{40,r}}{F_{40}} \right) \left(C_{mm}^{(1)} - \frac{4i \cos \theta}{r} C_{mm}^{(0)} \right) \right\} {}_{-2}Y_{\ell m}(\theta), \\
A_{mm2}^{(1)} &= \frac{r^2}{\sqrt{2\pi}} \left[\frac{4i \cos \theta}{r} C_{mm}^{(0)} - C_{mm}^{(1)} \right] {}_{-2}Y_{\ell m}(\theta), \tag{3.38}
\end{aligned}$$

where

$$\begin{aligned}
C_{nn}^{(1)} &= \frac{L r^2 - \Delta_r^0}{r^4 E} C_{nn}^{(0)} - \frac{L \Delta_r^0 (E + dr/d\tau)}{2Er^6}, \\
C_{mn}^{(1)} &= \left(\frac{L r^2 - \Delta_r^0}{r^4 E} + \frac{i \cos \theta}{r} \right) C_{mn}^{(0)} - \frac{\Delta_r^0}{2\sqrt{2}r^5 E} \left[iE \sin \theta (E + \frac{dr}{d\tau}) + L \left(\frac{iL}{r^2 \sin \theta} - \frac{d\theta}{d\tau} \right) \right], \\
C_{mm}^{(1)} &= \left(\frac{L r^2 - \Delta_r^0}{r^4 E} + \frac{2i \cos \theta}{r} \right) C_{mm}^{(0)} + \frac{i \sin \theta \Delta_r^0}{r^4} \left(\frac{d\theta}{d\tau} - \frac{iL}{r^2 \sin \theta} \right), \\
\mathcal{H}_0 &= i \left(\frac{K^0}{\Delta_r^0} \right)' - \frac{(K^0)^2}{(\Delta_r^0)^2} + \frac{iK^0}{\Delta_r^0} \left(\frac{1}{r} + \frac{F_{40,r}}{F_{40}} \right) + \frac{F_{40,r}}{rF_{40}} + \left(\frac{F_{40,r}}{F_{40}} \right)', \\
\mathcal{H}_1 &= \frac{m}{\Delta_r^0} \left[i \left(\frac{1}{r} + \frac{F'_{40}}{F_{40}} - \frac{\Delta_r^{0'}}{\Delta_r^0} \right) - \frac{2K^0}{\Delta_r^0} \right] - \frac{K^0 \cos \theta}{\Delta_r^0} \left[\frac{1}{r^2} + \frac{1}{F_{40}^2} (F_{40} F'_{41} - F_{41} F'_{40}) \right] \\
&\quad + \frac{i \cos \theta}{r^2 F_{40}^2} (F_{40} F'_{40} - r F_{41} F'_{40} + r F_{40} F'_{41}) - \frac{2i \cos \theta}{F_{40}^3} (F_{40} F'_{40} F'_{41} - F_{41} F'_{40}{}^2) \\
&\quad + \frac{i \cos \theta}{F_{40}^2} (F_{40} F_{41}'' - F_{41} F_{40}''). \tag{3.39}
\end{aligned}$$

3. Solution of ψ_4^B in the effective rotating spacetime

Using Eq. (3.31), we can rewrite Eq. (3.24) as

$$\begin{aligned}
\mathcal{A}(\ell) R_{\ell m} + a \left[\bar{\mathcal{B}} R_{\ell m} - \mathcal{C}(\ell - 1, m) \mathcal{Q}_\ell R_{\ell-1m} - \mathcal{C}(\ell + 1, m) \mathcal{Q}_{\ell+1} R_{\ell+1m} \right] \\
= T_{\ell m \omega}^{(0)}(r) + a T_{\ell m \omega}^{(1)}(r). \tag{3.40}
\end{aligned}$$

It is clear that Eq. (3.40) contains three modes, i.e., $R_{\ell m}$, $R_{\ell-1m}$, and $R_{\ell+1m}$. Therefore, for a calculation that is accurate to the first order in a , the $R_{\ell m}$ and $R_{\ell \pm 1m}$ terms in the square

brackets should be evaluated at the lower order. That is, we can solve Eq. (3.40) order by order by expanding the function $R_{\ell m}$ to the first order in the rotational parameter a as

$$R_{\ell m} = R_{\ell m}^{(0)} + a R_{\ell m}^{(1)}. \quad (3.41)$$

The equations for the zeroth and first orders of the rotational parameter a are respectively given by

$$\mathcal{A}(\ell) R_{\ell m}^{(0)} = T_{\ell m \omega}^{(0)}(r), \quad (3.42)$$

$$\mathcal{A}(\ell) R_{\ell m}^{(1)} + \left[\bar{\mathcal{B}} R_{\ell m}^{(0)} - \mathcal{C}(\ell - 1) \mathcal{Q}_\ell R_{\ell - 1 m}^{(0)} - \mathcal{C}(\ell + 1) \mathcal{Q}_{\ell + 1} R_{\ell + 1 m}^{(0)} \right] = T_{\ell m \omega}^{(1)}(r). \quad (3.43)$$

3-1. Formal solution of the zeroth-order equation

The zeroth-order equation can be obtained by substituting Eq. (3.21) into Eq. (3.42), which can be expressed explicitly as

$$\left[r^3 F_{40} (\Delta_r^0)^2 \frac{d}{dr} \left(\frac{1}{r^3 F_{40} \Delta_r^0} \frac{d}{dr} \right) + V \right] R_{\ell m \omega}^{(0)}(r) = T_{\ell m \omega}^{(0)}(r), \quad (3.44)$$

where

$$V = \frac{r^2 \omega (r^2 \omega + 2i \Delta_r^0)}{\Delta_r^0} - ir \omega (5 - r F_{10}) + \frac{3[\Delta_r^0 (1 + r F_{10}) + r \Delta_r^0]}{r^2} + 2r^2 F_{40} - \lambda(\ell) F_{20}.$$

It is interesting to note that if we take $a_i = 0$ ($i \geq 2$), Eq. (3.44) will reduce to the equation of the gravitational wave for ψ_4^B in the Schwarzschild spacetime [56].

The asymptotical homogeneous solutions for Eq. (3.44) are

$$R_{asy}^{in(0)} \rightarrow \begin{cases} B_{\ell m \omega}^{trans} (\Delta_r^0)^2 e^{-i\omega r^*}, & \text{for } r \rightarrow r_+, \\ r^3 B_{\ell m \omega}^{ref} e^{i\omega r^*} + r^{-1} B_{\ell m \omega}^{inc} e^{-i\omega r^*}, & \text{for } r \rightarrow +\infty, \end{cases} \quad (3.45)$$

$$R_{asy}^{up(0)} \rightarrow \begin{cases} C_{\ell m \omega}^{up} e^{i\omega r^*} + (\Delta_r^0)^2 C_{\ell m \omega}^{ref} e^{-i\omega r^*}, & \text{for } r \rightarrow r_+, \\ C_{\ell m \omega}^{trans} r^3 e^{i\omega r^*}, & \text{for } r \rightarrow +\infty, \end{cases} \quad (3.46)$$

where $r^* = \int \frac{r^2}{\Delta_r^0} dr$. Moreover, the inhomogeneous solution for Eq. (3.44) is

$$R_{\ell m \omega}^{(0)}(r) = \frac{1}{2i\omega C_{\ell m \omega}^{trans} B_{\ell m \omega}^{inc}} \left\{ R_{\ell m \omega}^{up(0)}(r) \int_{r_+}^r d\tilde{r} \frac{R_{\ell m \omega}^{in(0)}(\tilde{r}) T_{\ell m \omega}^{(0)}(\tilde{r})}{\tilde{r}^3 F_{40}(\tilde{r}) (\Delta_r^0)^2(\tilde{r})} \right. \\ \left. + R_{\ell m \omega}^{in(0)}(r) \int_r^\infty d\tilde{r} \frac{R_{\ell m \omega}^{up(0)}(\tilde{r}) T_{\ell m \omega}^{(0)}(\tilde{r})}{\tilde{r}^3 F_{40}(\tilde{r}) (\Delta_r^0)^2(\tilde{r})} \right\}, \quad (3.47)$$

where $R_{\ell m \omega}^{\text{up}(0)}(\tilde{r})$ and $R_{\ell m \omega}^{\text{in}(0)}(\tilde{r})$ are the homogeneous solutions of the radial equation (3.44). Therefore, the inhomogeneous solution for Eq. (3.44) at the infinity is

$$R_{\ell m \omega}^{(0)}(r \rightarrow \infty) \rightarrow \frac{r^3 e^{i\omega r^*}}{2i\omega B_{\ell m \omega}^{\text{inc}}} \int_{r_+}^{\infty} d\tilde{r} \frac{T_{\ell m \omega}^{(0)}(\tilde{r}) R_{\ell m \omega}^{\text{in}(0)}(\tilde{r})}{\tilde{r}^3 F_{40}(\tilde{r}) (\Delta_r^0)^2} \equiv \tilde{Z}_{\ell m \omega}^{(0)} r^3 e^{i\omega r^*}, \quad (3.48)$$

with

$$\tilde{Z}_{\ell m \omega}^{(0)} = \frac{\pi m_0 G}{i\omega B_{\ell m \omega}^{\text{inc}}} \left[A_0^{(0)} \frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} - A_1^{(0)} \left(\frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)' + A_2^{(0)} \left(\frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)'' \right]_{r_0, \theta_0} \delta(\omega - \omega_n), \quad (3.49)$$

where $\omega_n = m \Omega$, $A_0^{(0)} = A_{nn0}^{(0)} + A_{mn0}^{(0)} + A_{mm0}^{(0)}$, $A_1^{(0)} = A_{mn1}^{(0)} + A_{mm1}^{(0)}$, $A_2^{(0)} = A_{mm2}^{(0)}$, and (r_0, θ_0) are the values of (r, θ) on the geodesic trajectory. The zeroth-order formal solution of ψ_4^B is then shown by

$$\begin{aligned} \psi_4^{B(0)} &= \frac{1}{r} \sum_{\ell m n} \frac{\pi m_0 G}{i\omega_n B_{\ell m \omega_n}^{\text{inc}}} \left[A_0^{(0)} \frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} - A_1^{(0)} \left(\frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)' + A_2^{(0)} \left(\frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)'' \right]_{r_0, \theta_0} \\ &\times \frac{-2Y_{\ell m}}{\sqrt{2\pi}} e^{i\omega_n(r^* - t) + im\varphi}. \end{aligned} \quad (3.50)$$

3-2. Formal solution of the first-order equation

The first-order equation (3.43) can be written explicitly as

$$\left[r^3 F_{40} (\Delta_r^0)^2 \frac{d}{dr} \left(\frac{1}{r^3 F_{40} \Delta_r^0} \frac{d}{dr} \right) + V \right] R_{\ell m \omega}^{(1)}(r) = T_{\ell m \omega}^{(1)}(r) + T_{eff}^{(1)}(r), \quad (3.51)$$

with

$$\begin{aligned} T_{eff}^{(1)}(r) &= \mathcal{Q}_\ell \left[\frac{i\Delta_r^0}{r^2} (3 + r^2 F_{11}) \frac{d}{dr} + U(\ell - 1) \right] R_{\ell-1m}^{(0)}(r) \\ &+ \mathcal{Q}_{\ell+1} \left[\frac{i\Delta_r^0}{r^2} (3 + r^2 F_{11}) \frac{d}{dr} + U(\ell + 1) \right] R_{\ell+1m}^{(0)}(r) - \bar{\mathcal{B}} R_{\ell m}^{(0)}(r), \end{aligned} \quad (3.52)$$

where $R_{\ell m \omega}^{(0)}(r)$ is the inhomogeneous solution (3.47), and

$$\begin{aligned} U(\ell) &= (r^2 F_{11} - 3)\omega - \frac{3i}{r^3} \left[r \Delta_r^0 + \Delta_r^0 (2 + r F_{10} + r^2 F_{11}) \right] \\ &+ 2F_{20} \left(\omega - \frac{3i}{r} \right) - 2ir^2 F_{41} - iF_{21} \lambda(\ell). \end{aligned} \quad (3.53)$$

Now, we solve the first-order equation (3.51) by means of the Green function [62–64]. Since the homogeneous equations (3.44) and (3.51) take the same form, the asymptotic

solutions for the homogeneous solutions of Eq. (3.51) are

$$R_{asy}^{\text{in}(1)} \rightarrow \begin{cases} B_{\ell m \omega}^{\text{trans}} (\Delta_r^0)^2 e^{-i\omega r^*}, & \text{for } r \rightarrow r_+, \\ r^3 B_{\ell m \omega}^{\text{ref}} e^{i\omega r^*} + r^{-1} B_{\ell m \omega}^{\text{inc}} e^{-i\omega r^*}, & \text{for } r \rightarrow +\infty, \end{cases} \quad (3.54)$$

$$R_{asy}^{\text{up}(1)} \rightarrow \begin{cases} C_{\ell m \omega}^{\text{up}} e^{i\omega r^*} + (\Delta_r^0)^2 C_{\ell m \omega}^{\text{ref}} e^{-i\omega r^*}, & \text{for } r \rightarrow r_+, \\ C_{\ell m \omega}^{\text{trans}} r^3 e^{i\omega r^*}, & \text{for } r \rightarrow +\infty, \end{cases} \quad (3.55)$$

and the inhomogeneous solution for the radial equation (3.51) is

$$R_{\ell m \omega}^{(1)}(r) = \frac{1}{2i\omega C_{\ell m \omega}^{\text{trans}} B_{\ell m \omega}^{\text{inc}}} \left\{ R_{\ell m \omega}^{\text{up}(0)}(r) \int_{r_+}^r d\tilde{r} \frac{R_{\ell m \omega}^{\text{in}(0)}(\tilde{r}) (T_{\ell m \omega}^{(1)}(\tilde{r}) + T_{eff}^{(1)}(\tilde{r}))}{\tilde{r}^3 F_{40} (\Delta_r^0)^2} \right. \\ \left. + R_{\ell m \omega}^{\text{in}(0)}(r) \int_r^{\infty} d\tilde{r} \frac{R_{\ell m \omega}^{\text{up}(0)}(\tilde{r}) (T_{\ell m \omega}^{(1)}(\tilde{r}) + T_{eff}^{(1)}(\tilde{r}))}{\tilde{r}^3 F_{40} (\Delta_r^0)^2} \right\}, \quad (3.56)$$

where $R_{\ell m \omega}^{\text{up}(0)}(\tilde{r})$ and $R_{\ell m \omega}^{\text{in}(0)}(\tilde{r})$ are the homogeneous solutions of the radial equation (3.44).

Therefore, the inhomogeneous solution of Eq. (3.51) at infinity can be expressed as

$$R_{\ell m \omega}^{(1)}(r \rightarrow \infty) = \frac{r^3 e^{i\omega r^*}}{2i\omega B_{\ell m \omega}^{\text{inc}}} \int_{r_+}^{\infty} d\tilde{r} \frac{R_{\ell m \omega}^{\text{in}(0)}(\tilde{r}) (T_{\ell m \omega}^{(1)}(\tilde{r}) + T_{eff}^{(1)}(\tilde{r}))}{\tilde{r}^3 F_{40} (\Delta_r^0)^2} \equiv \tilde{Z}_{\ell m \omega}^{(1)} r^3 e^{i\omega r^*}. \quad (3.57)$$

Inserting Eqs. (3.37) and (3.52) into Eq. (3.57), we have

$$\tilde{Z}_{\ell m \omega}^{(1)} = \frac{m_0 G}{2i\omega B_{\ell m \omega}^{\text{inc}}} \int_{-\infty}^{\infty} dt e^{i\omega t - im\varphi(t)} \left[A_0^{(1)} \frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} - A_1^{(1)} \left(\frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)' + A_2^{(1)} \left(\frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)'' \right]_{r_0, \theta_0} \\ + \frac{1}{2i\omega B_{\ell m \omega}^{\text{inc}}} \int_{r_+}^{\infty} d\tilde{r} \frac{R_{\ell m \omega}^{\text{in}(0)}(\tilde{r})}{\tilde{r}^3 F_{40} (\Delta_r^0)^2} \left\{ \mathcal{Q}_{\ell} \left[\frac{i\Delta_{\tilde{r}}^0}{\tilde{r}^2} (3 + \tilde{r}^2 F_{11}(\tilde{r})) \frac{d}{d\tilde{r}} + U(\ell - 1) \right] R_{\ell-1m\omega}^{(0)}(\tilde{r}) \right. \\ \left. + \mathcal{Q}_{\ell+1} \left[\frac{i\Delta_{\tilde{r}}^0}{\tilde{r}^2} (3 + r^2 F_{11}(\tilde{r})) \frac{d}{d\tilde{r}} + U(\ell + 1) \right] R_{\ell+1m\omega}^{(0)}(\tilde{r}) - \bar{\mathcal{B}} R_{\ell m \omega}^{(0)}(\tilde{r}) \right\}, \quad (3.58)$$

where $A_0^{(1)} = A_{nn_0}^{(1)} + A_{\bar{m}n_0}^{(1)} + A_{\bar{m}\bar{m}0}^{(1)}$, $A_1^{(1)} = A_{\bar{m}n_1}^{(1)} + A_{\bar{m}\bar{m}1}^{(1)}$, and $A_2^{(1)} = A_{\bar{m}\bar{m}2}^{(1)}$. For the circular orbits, we can take $r(t) = r_0$, $\theta(t) = \theta_0$, and $\varphi(t) = \Omega t$, where Ω is the angular velocity. Noting that $\tilde{Z}_{\ell m \omega}^{(1)} = Z_{\ell m \omega}^{(1)} \delta(\omega - \omega_n)$ [42], we then find that Eq. (3.58) can be expressed as

$$\tilde{Z}_{\ell m \omega}^{(1)} = \frac{\pi m_0}{i\omega B_{\ell m \omega}^{\text{inc}}} \left\{ G \left[A_0^{(1)} \frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} - A_1^{(1)} \left(\frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)' + A_2^{(1)} \left(\frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)'' \right]_{r_0, \theta_0} \right. \\ \left. + B_{eff}^{(1)} \right\} \delta(\omega - \omega_n), \quad (3.59)$$

where

$$\begin{aligned}
B_{eff}^{(1)}\delta(\omega - \omega_n) = & \\
& \frac{1}{2\pi m_0} \int_{r_+}^{\infty} d\tilde{r} \frac{R_{\ell m \omega}^{\text{in}(0)}(\tilde{r})}{\tilde{r}^3 F_{40}(\Delta_r^0)^2} \left\{ \mathcal{Q}_\ell \left[\frac{i\Delta_{\tilde{r}}^0}{\tilde{r}^2} (3 + \tilde{r}^2 F_{11}(\tilde{r})) \frac{d}{d\tilde{r}} + U(\ell - 1) \right] R_{\ell-1 m \omega}^{(0)}(\tilde{r}) \right. \\
& \left. + \mathcal{Q}_{\ell+1} \left[\frac{i\Delta_{\tilde{r}}^0}{\tilde{r}^2} (3 + r^2 F_{11}(\tilde{r})) \frac{d}{d\tilde{r}} + U(\ell + 1) \right] R_{\ell+1 m \omega}^{(0)}(\tilde{r}) - \bar{\mathcal{B}} R_{\ell m \omega}^{(0)}(\tilde{r}) \right\}. \quad (3.60)
\end{aligned}$$

Therefore, the first-order formal solution of ψ_4^B is described by

$$\begin{aligned}
\psi_4^{B(1)} = & \frac{1}{r} \sum_{\ell m n} \frac{\pi m_0}{i\omega_n B_{\ell m \omega_n}^{\text{inc}}} \left\{ G \left[A_0^{(1)} \frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} - A_1^{(1)} \left(\frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)' + A_2^{(1)} \left(\frac{R_{\ell m \omega}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)'' \right]_{r_0, \theta_0} \right. \\
& \left. + B_{eff}^{(1)} \right\} \frac{-2Y_{\ell m}}{\sqrt{2\pi}} e^{i\omega_n(r^*-t) + im\varphi}. \quad (3.61)
\end{aligned}$$

3-3. Solution of ψ_4^B in the effective spacetime

Equations (3.50) and (3.61) show that the solution of ψ_4^B up to the first order of the rotational parameter a in the effective spacetime can be expressed as

$$\begin{aligned}
\psi_4^B = & \frac{1}{r} \sum_{\ell m n} \frac{\pi G m_0}{i\omega_n B_{\ell m \omega_n}^{\text{inc}}} \left\{ \left[(A_0^{(0)} + aA_0^{(1)}) \frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} - (A_1^{(0)} + aA_1^{(1)}) \left(\frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)' \right. \right. \\
& \left. \left. + (A_2^{(0)} + aA_2^{(1)}) \left(\frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)'' \right]_{r_0, \theta_0} + \frac{a}{G} B_{eff}^{(1)} \right\} \frac{-2Y_{\ell m}}{\sqrt{2\pi}} e^{i\omega_n(r^*-t) + im\varphi}. \quad (3.62)
\end{aligned}$$

C. Radiation reaction force and waveform for “plus” and “cross” modes of the gravitational wave

The reduced RRF is described by [50]

$$\hat{\mathcal{F}} = \frac{1}{\nu M_0 \Omega |\mathbf{r} \times \mathbf{P}|} \frac{dE}{dt} \mathbf{P}, \quad (3.63)$$

where $\Omega = |\mathbf{r} \times \dot{\mathbf{r}}|/r^2$ is the dimensionless orbital frequency and dE/dt is the energy flux of the gravitational radiation. For the quasicircular cases without precession, noting that $|\mathbf{r} \times \mathbf{P}| \approx p_\varphi$, we find that the reduced RRF that appeared in the Hamiltonian equation

(2.1) can be expressed explicitly as

$$\begin{aligned} \hat{\mathcal{F}} = & \frac{1}{\nu M_0 \Omega} \sum_{\ell=2}^{\infty} \sum_{m=1}^{\ell} \frac{2\pi m_0^2}{G \omega_n^4} \left| \frac{G}{B_{\ell m \omega_n}^{\text{inc}}} \left\{ \left[(A_0^{(0)} + aA_0^{(1)}) \frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} - (A_1^{(0)} + aA_1^{(1)}) \left(\frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)' \right. \right. \right. \\ & \left. \left. \left. + (A_2^{(0)} + aA_2^{(1)}) \left(\frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)'' \right]_{r_0, \theta_0} + \frac{a}{G} B_{eff}^{(1)} \right\} \right|^2 \frac{\mathbf{P}}{p_\varphi}, \end{aligned} \quad (3.64)$$

indicating that the reduced RRF is constructed in terms of the effective spacetime.

On the other hand, it is well known that the “plus” and “cross” modes of the gravitational wave can be expressed in terms of spin-weighted $s = -2$ spherical harmonics [53]:

$$h_+ - ih_\times = \sum_{l=2}^{\infty} \sum_{m=-l}^l h^{lm} \frac{{}_{-2}Y^{lm}(\theta, \varphi)}{\sqrt{2\pi}}. \quad (3.65)$$

Thus, by comparing Eq. (3.65) with the solution of ψ_4^B , we can read off the waveform easily, which is given by

$$\begin{aligned} h^{lm} = & \frac{1}{r} \frac{2\pi G m_0}{i\omega_n^3 B_{\ell m \omega_n}^{\text{inc}}} \left\{ \left[(A_0^{(0)} + aA_0^{(1)}) \frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} - (A_1^{(0)} + aA_1^{(1)}) \left(\frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)' \right. \right. \\ & \left. \left. + (A_2^{(0)} + aA_2^{(1)}) \left(\frac{R_{\ell m \omega_n}^{\text{in}(0)}(r)}{r^3 F_{40}} \right)'' \right]_{r_0, \theta_0} + \frac{a}{G} B_{eff}^{(1)} \right\} e^{i\omega_n(r^* - t)}. \end{aligned} \quad (3.66)$$

It is clear that the waveform is also constructed in terms of the effective rotating spacetime. Based on h^{lm} described by Eq. (3.66) and using Damour–Nagar’s proposals [11, 48, 49, 54], we can then construct an improved inspiral–plunge mode, i.e., an improved multipolar waveform can be built as a product of the leading Minkowskian order term $h_{\ell m}^{(M, \epsilon_p)}$, the relativistic conserved energy or angular momentum of the effective moving source $\hat{S}_{\text{eff}}^{(\epsilon_p)}$, an infinite number of leading logarithms that enter the tail effect $T_{\ell m}$, an additional phase correction $e^{i\delta_{\ell m}}$, the remaining PM effects $f_{\ell m}$, and a non-quasicircular effect $N_{\ell m}$.

The above discussions show that the improved reduced Hamiltonian (2.12), reduced RRF (3.64), and waveform (3.66) for the “plus” and “cross” modes of the gravitational wave, which are based on the same physical model, constitute a self-consistent EOB theory for the spin binaries based on the PM approximation.

IV. CONCLUSIONS AND DISCUSSIONS

From a practical point of view, it is well known that the general gravitational waveform template should be constructed for the spin binaries. Therefore, as a theory to construct

the gravitational waveform template, the self-consistent EOB theory should be set up for the spinning black hole binaries. In this paper, we extend the study on the SCEOB theory for the real spinless two-body system [42] to the case of the real spin two-body system.

Based on the effective metric (2.2) for a real spinless two-body system in the post-Minkowskian approximation obtained in Ref. [42], we used Damour’s or Barausse’s approach [43, 44] to find the effective rotating metric (2.4) for a real spin two-body system. The effective rotating metric is still type D . In principle, it can be applicable to any post-Minkowskian order. By means of the effective metric (2.4), the energy map (2.5), and the approach of constructing the EOB Hamiltonian for the spinning black hole binaries presented by Damour and Barausse et al., [44–47], we then obtain an improved reduced EOB Hamiltonian (2.12) of the EOB theory for the real spin two-body system at a linear order of the particle’s spin which appears in the Hamilton equations (2.1).

To work out the reduced RRF and the waveform for the “plus” and “cross” modes of the gravitational wave generated by a coalescing compact object binary system, we first used the gauge transform property of tetrad components of the perturbed Weyl tensors to find the decoupled equation (3.16) of the null tetrad component of the gravitational perturbed Weyl tensor ψ_4^B in the effective rotating spacetime. Next, noting that the decoupled equation (3.16) does not appear to be directly separable, we find that we can separate the equation for the radial and angular parts in the slowly rotating background, which is close enough to the spherical symmetry, and derive the explicitly radial equation (3.24) up to the first order of the rotational parameter a . However, it should be noted that, in principle, the method can be extended to any order of the rotational parameter a . Using the energy-momentum tensor (3.30) for a particle that orbits around a massive black hole described by the effective rotating metric, we then note that the radial equation can be solved order by order in the rotational parameter a and present a formal solution (3.62) of ψ_4^B up to the first order of the rotational parameter a by means of the Green function. Finally, with the help of the null tetrad component of the perturbed Weyl tensor ψ_4^B , we present the formal expressions of the reduced RRF (3.64) and the waveform (3.66) for “plus” and “cross” modes of the gravitational wave in the effective rotating spacetime.

It is interesting to note that the improved reduced EOB Hamiltonian (2.12), the reduced RRF (3.64), and the waveform (3.66) for the “plus” and “cross” modes of the gravitational wave, which are obtained in the same physical model, constitute a self-consistent EOB theory

for the spin black hole binaries based on the PM approximation.

Acknowledgments

We would like to thank professors S. Chen and Q. Pan for useful discussions on the manuscript. This work was supported by the Grant of NSFC Nos. 12035005 and 12122504, and National Key Research and Development Program of China No. 2020YFC2201400.

-
- [1] A. Einstein, *Über gravitationswellen*, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften Berlin. part 1: 154-167, (1918).
 - [2] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, *Gravitational waves in general relativity VII. Waves from axi-symmetric isolated systems*, Proc. R. Soc. London **A269**, 21 (1962).
 - [3] R. K. Sachs, *Gravitational waves in general relativity VIII. Waves in asymptotically flat space-time*, Proc. R. Soc. London **A270**, 103 (1962).
 - [4] S. Perlmutter, G. Aldering, M. Della Valle, S. Deustua, R. S. Ellis, S. Fabbro, A. Fruchter, G. Goldhaber, A. Goobar, D. E. Groom, I. M. Hook et al., *Discovery of a supernova explosion at half the age of the universe and its cosmological implications*, Nature **391**, 51 (1998).
 - [5] Wentao Liu, Xiongjun Fang, Jiliang Jing, Anzhong Wang, *Gauge Invariant Perturbations of General Spherically Symmetric Spacetimes*, Sci. China, Phys. Mech. Astron. **66**, 210411 (2023).
 - [6] Y. H. Zou, M. J. Wang and J. L. Jing, *Test of a model coupling of electromagnetic and gravitational fields by using high-frequency gravitational waves*, Sci. China, Phys. Mech. Astron. **64**, 250411 (2021).
 - [7] B. C. Barish and R. Weiss, *LIGO and the Detection of Gravitational Waves*, Phys. Today **52**, 44 (1999).
 - [8] S. J. Waldman (LIGO Scientific Collaboration), *Status of LIGO at the start of the fifth science run*, Class. Quantum Grav. **23**, S653 (2006).
 - [9] F. Acernese et al. (Virgo Collaboration), *The Virgo status*, Class. Quantum Grav. **23**, S635 (2006).

- [10] B. F. Schutz, Fundamental physics with LISA, *Class. Quantum Grav.* **26**, 094020 (2009).
- [11] Y. Pan, A. Buonanno, L. T. Buchman, T. Chu, L. E. Kidder, H. P. Pfeiffer, M. A. Scheel, *Effective-one-body waveforms calibrated to numerical relativity simulations: Coalescence of nonprecessing, spinning, equal-mass black holes*, *Phys. Rev. D* **81**, 084041 (2010).
- [12] A. Buonanno and T. Damour, *Effective one-body approach to general relativistic two-body dynamics*, *Phys. Rev. D* **59**, 084006 (1999).
- [13] T. Damour, P. Jaranowski and G. Schafer, *On the determination of the last stable orbit for circular general relativistic binaries at the third post-Newtonian approximation*, *Phys. Rev. D* **62**, 084011 (2000).
- [14] T. Damour, B. R. Iyer, and A. Nagar, *Improved resummation of post-Newtonian multipolar waveforms from circularized compact binaries*, *Phys. Rev. D* **79**, 064004 (2009).
- [15] A. Buonanno and T. Damour, *Transition from inspiral to plunge in binary black hole coalescences*, *Phys. Rev. D* **62**, 064015 (2000).
- [16] T. Damour, *Coalescence of Two Spinning Black Holes: An effective one-body approach*, *Phys. Rev. D* **64**, 124013 (2001).
- [17] A. Taracchini et al., *Effective-one-body model for black-hole binaries with generic mass ratios and spins*, *Phys. Rev. D* **89**, 061502 (2014).
- [18] A. Bohé et al., *Improved effective-one-body model of spinning, nonprecessing binary black holes for the era of gravitational-wave astrophysics with advanced detectors*, *Phys. Rev. D* **95**, no. 4, 044028 (2017).
- [19] B. P. Abbott et al. (LIGO Scientific and Virgo Collaborations), *Phys. Rev. Lett.* **116**, 061102 (2016).
- [20] B. P. Abbott et al. (LIGO Scientific and Virgo Collaborations), *Phys. Rev. Lett.* **116**, 241103 (2016).
- [21] B. P. Abbott et al. (LIGO Scientific and Virgo Collaborations), *Phys. Rev. Lett.* **118**, 221101 (2017).
- [22] B. P. Abbott et al. (LIGO Scientific and Virgo Collaborations), *Phys. Rev. Lett.* **119**, 141101 (2017).
- [23] T. Damour, *Gravitational scattering, post-Minkowskian approximation and Effective One-Body theory*, *Phys. Rev. D* **94**, 104015 (2016).
- [24] D. Bini and T. Damour, *Gravitational spin-orbit coupling in binary systems, post-*

- Minkowskian approximation, and effective-one-body theory*, Phys. Rev. D **96**, 104038 (2017).
- [25] D. Bini and T. Damour, *Gravitational spin-orbit coupling in binary systems at the second post-Minkowskian approximation*, Phys. Rev. D **98**, 044036 (2018).
- [26] T. Damour, *High-energy gravitational scattering and the general relativistic two-body problem*, Phys. Rev. D **97**, 044038 (2018).
- [27] A. Antonelli, A. Buonanno, J. Steinhoff, M. van de Meent and J. Vines, *Energetics of two-body Hamiltonians in post-Minkowskian gravity*, Phys. Rev. D **99**, 104004 (2019)
- [28] T. Damour, *Classical and quantum scattering in post-Minkowskian gravity*, Phys. Rev. D **102**, 024060 (2020)
- [29] D. Bini, T. Damour and A. Gericco, *Scattering of tidally interacting bodies in post-Minkowskian gravity*, Phys. Rev. D **101**, 044039 (2020)
- [30] G. He and W. Lin, *Second-order time delay by a radially moving Kerr-Newman black hole*, Phys. Rev. D **94**, 063011(2016)
- [31] X. K. He, J. L. Jing and Z. J. Cao, *Generalized gravitomagnetic field and gravitational waves*, Sci. China, Phys. Mech. Astron. **62**, 110422 (2019).
- [32] L. Blanchet and A. Fokas, *Equations of motion of self-gravitating N-body systems in the first post-Minkowskian approximation*, Phys. Rev. D **98**, 084005 (2018)
- [33] C. Cheung, I. Rothstein and M. Solon, *From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion*, Phys. Rev. Lett. **121** , 251101 (2018)
- [34] J. Vines, J. Steinhoff and A. Buonanno, *Spinning-black-hole scattering and the test-black-hole limit at second post-Minkowskian order*, Phys. Rev. D **99**, 064054 (2019)
- [35] A. Cristofoli, N. Bjerrum-Bohr, P. Damgaard and P. Vanhove, *Post-Minkowskian Hamiltonians in general relativity*, Phys. Rev. D **100**, 084040 (2019)
- [36] A. Collado, P. Vecchia and R. Russo, *Revisiting the second post-Minkowskian eikonal and the dynamics of binary black holes*, Phys. Rev. D **100**, 066028 (2019)
- [37] J. Plefka, C. Shi, J. Steinhoff and T. Wang, *Breakdown of the classical double copy for the effective action of dilaton-gravity at NNLO*, Phys. Rev. D **100**, 086006 (2019)
- [38] D. Bini, T. Damour and A. Gericco, *Scattering of tidally interacting bodies in post-Minkowskian gravity*, Phys. Rev. D **101**, 044039 (2020)
- [39] C. Cheung and M. Solon, *Tidal Effects in the Post-Minkowskian Expansion*, Phys. Rev. Lett. **125** , 191601 (2020)

- [40] Z. Bern, C. Cheung, R. Roiban, C. H. Shen, M. P. Solon and M. Zeng, *Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order*, Phys.Rev. Lett. **122**, 201603 (2019).
- [41] Z. Bern, C. Cheung, R. Robin, C. H. Shen, M. P. Solon and M. Zeng, *Black Hole Binary Dynamics from the Double Copy and Effective Theory*, JHEP,**10**,206(2019).
- [42] Jiliang Jing, Sheng Long, WeiKe Deng, Mengjie Wang, and Jieci Wang, *New self-consistent effective one-body theory for spinless binaries based on the post-Minkowskian approximation*, Sci. China, Phys. Mech. Astron. **65**, 100411 (2022), ArXiv: 2208.02420.
- [43] E. Barausse, A. Buonanno, *Extending the effective-one-body Hamiltonian of black-hole binaries to include next-to-next-to-leading spin-orbit couplings*, Phys. Rev. D **84**, 104027 (2011).
- [44] T. Damour, *Coalescence of two spinning black holes: An effective one-body approach*, Phys. Rev., **64**, 124013 (2001).
- [45] T. Damour, P. Jaranowski, and G. Schafer, *Effective one body approach to the dynamics of two spinning black holes with next-to-leading order spin-orbit coupling*, Phys. Rev. D, **78**, 024009 (2008).
- [46] E. Barausse, E. Racine, and A. Buonanno, *Hamiltonian of a spinning test particle in curved spacetime*, Phys. Rev. D **80**, 104025 (2009).
- [47] E. Barausse, and A. Buonanno, *Improved effective-one-body Hamiltonian for spinning black-hole binaries*, Phys. Rev. D **81**, 084024 (2010).
- [48] A. Buonanno, Y. Pan, H. P. Pfeiffer, M. A. Scheel, L. T. Buchman, L. E. Kidder, *Effective-one-body waveforms calibrated to numerical relativity simulations: Coalescence of nonspinning, equal-mass black holes*, Phys. Rev. D **79**, 124028 (2009).
- [49] T. Damour and A. Nagar, *Improved analytical description of inspiralling and coalescing black-hole binaries*, Phys. Rev. D **79**, 081503(R) (2009).
- [50] A. Buonanno, Y. Chen, and T. Damour, *Transition from inspiral to plunge in precessing binaries of spinning black holes*, Phys. Rev. D **74**, 104005 (2006).
- [51] Jiliang Jing, Shuai Chen, Manman Sun, Xiaokai He, Mengjie Wang, and Jieci Wang, *Self-consistent Effective-one-body theory for spinless binaries based on post-Minkowskian approximation I: Hamiltonian and decoupled equation for ψ_4^B* , Sci. China, Phys. Mech. Astron. **65**, 260411 (2022), ArXiv: 2112.09838.
- [52] J. E. Thompson, H. Chen and B. F. Whiting, *Gauge invariant perturbations of the*

- Schwarzschild spacetime, *Class. Quantum Grav.* **34** (2017) 174001.
- [53] L. E. Kidder, *Using full information computing modes of post-Newtonian waveforms from inspiralling compact binaries in circular orbit*, *Phys. Rev. D* **77**, 044016 (2008).
- [54] T. Damour and A. Nagar, *Faithful effective-one-body waveforms of small-mass-ratio coalescing black hole binaries*, *Phys. Rev.* **D76**, 064028 (2007).
- [55] A. Taracchini, Y. Pan, A. Buonanno, E. Barausse, M. Boyle, et al., *Prototype effective-one-body model for nonprecessing spinning inspiral-merger-ringdown waveforms*, *Phys.Rev.* **D86**, 024011 (2012).
- [56] S. A. Teukolsky, *Perturbations of a rotating black hole. I. Fundamental equations for gravitational, electromagnetic, and neutrino field perturbations*, *Astrophys. J.* **185**, 635 (1973).
- [57] M. Carmeli, *Classical fields: general relativity and gauge theory*, A Wiley-interscience Publication, New York, (1982).
- [58] S. Chandrasekhar, *The mathematical theory of black hole*, Oxford University Press, New York, Oxford, (1993).
- [59] Dario Nunez, Juan Carlos Degollado, and Carlos Palenzuela Y. Kojima, *One dimensional description of the gravitational perturbation in a Kerr background*, *Phys. Rev. D* **81**, 064011 (2010); Y. Kojima, *Equations governing the nonradial oscillations of a slowly rotating relativistic star*, *Phys. Rev.* **D46**, 4289 (1992).
- [60] M. Sasaki and T. Nakamura, *The Regge-Wheeler equation with sources for both even and odd parity perturbations of the Schwarzschild geometry*, *Phys. Lett. A* **87**, 85 (1981).
- [61] Xuan Zhou, Songbai Chen, Jiliang Jing, *Effect of noncircularity on the dynamic behaviors of particles in a disformal rotating black-hole spacetime*, *Sci. China, Phys. Mech. Astron.* **65**, 250411 (2022).
- [62] M. Sasaki and H. Tagoshi, *Analytic black hole perturbation approach to gravitational radiation*, *Living Rev. Relativity*, **6**, 6 (2003).
- [63] E. Poisson, *Gravitational radiation from a particle in circular orbit around a black hole. I. Analytical results for the nonrotating case*, *Phys. Rev.* **D47**, 1497 (1993).
- [64] H. Tagoshi and M. Sasaki, *Post-Newtonian expansion of gravitational waves from a particle in circular orbit around a Schwarzschild black hole*, *Prog. Theor. Phys.* **92**, 745 (1994).