

Constraint on Lorentz invariance violation from Vela pulsar

Hao Li^a, Jie Zhu^a, Bo-Qiang Ma^{a,b,c,*}

^a*School of Physics, Peking University, Beijing 100871, China*

^b*Center for High Energy Physics, Peking University, Beijing 100871, China*

^c*Collaborative Innovation Center of Quantum Matter, Beijing, China*

Abstract

The High Energy Stereoscopic System (H.E.S.S.) Collaboration reported the discovery of a novel radiation component from the Vela pulsar by their Cherenkov telescopes. It is of great importance that gamma rays with energies of at least 20 TeV are recorded unexpectedly. The H.E.S.S. Collaboration argued that such results may challenge the state-of-the-art models for the high-energy emission of pulsars. We point out in this work that these results also provide a unique opportunity to constrain certain Lorentz invariance violation parameters, leading to the realization of studying Lorentz invariance violation by using gamma-ray pulsars. The Lorentz invariance violation scale is constrained at the level of $E_{LV,1} > 1.66 \times 10^{17}$ GeV for the linear scenario, and $E_{LV,2} > 3.53 \times 10^{10}$ GeV for the quadratic scenario. We anticipate that digging into the detailed features of the data of the Vela pulsar and analyzing potentially more very-high-energy photon data from pulsars in the future would improve the constraints on Lorentz invariance violation.

Keywords: Lorentz invariance violation, the Vela pulsar, very-high-energy gamma rays

Utilizing high-energy photons from distant objects in the Universe is one of the most important approaches to searching Lorentz invariance violation (LIV) [1, 2]. In general, these sources are required to be at a very large distance from the Earth and the emissions of energetic photons are highly variable, with a time-structure of milliseconds [1, 2]. Therefore, gamma-ray bursts (GRBs) have long been the central focus for the study of LIV following this method (see Refs. [3, 4] and references therein). There are also proposals that very-high-energy (VHE) photons from pulsars can be used to study LIV [5–7]. Just recently, results from the High Energy Stereoscopic System (H.E.S.S.) provide a unique opportunity to searching Lorentz invariance violation from VHE cosmic photons. The Cherenkov telescopes of H.E.S.S. detected multi-TeV gamma-rays from the Vela pulsar, PSR B0833045, of which the radiation component extends up to energies of at least 20 TeV [8]. These results are already analyzed by the H.E.S.S. Collaboration, to show that the state-of-the-art models for the high-energy emission of pulsars may be challenged [8]. However, besides using these results to limit the models for pulsars, it is noteworthy that the photons from the Vela pulsar are very energetic, and the phasograms of the Vela pulsar have time structures of milliseconds. Although this object is quite close to the earth, at a distance of 287 pc [8, 9], we suggest that the Vela pulsar can be used to constrain parameters of LIV, with the same method as that utilized in the case of GRBs, because of the photons of energies of at least 20 TeV and the high temporal variability of this source, which has a spin period of 89 ms [8]. In the following, we perform a quick and rough analysis without

more detailed data and constrain the leading order and next-to-leading order LIV parameters for photons.

It is speculated from quantum gravity that the Lorentz invariance might be broken at the Planck scale ($E_{Pl} \approx 1.22 \times 10^{19}$ GeV), and that the light speed may have a variation with the energy of the photon. For energy $E \ll E_{Pl}$, the modified dispersion relation (MDR) of the photon can be expressed to the leading order as

$$E^2 = p^2 c^2 \left[1 - s_n \left(\frac{pc}{E_{LV,n}} \right)^n \right]. \quad (1)$$

Assuming that the traditional relation $v = \partial E / \partial p$ holds, we have the following speed relation

$$v(E) = c \left[1 - s_n \frac{n+1}{2} \left(\frac{pc}{E_{LV,n}} \right)^n \right], \quad (2)$$

where $n = 1$ or $n = 2$ as usually assumed, $s_n = \pm 1$ indicates whether high-energy photons travel faster ($s_n = -1$) or slower ($s_n = +1$) than low-energy photons, and $E_{LV,n}$ represents the n th-order Lorentz violation scale. Usually, we should consider the expansion of the Universe [10, 11]. However, since the Vela pulsar is located nearby at the distance of $L = 287$ pc, we can ignore the expansion of the Universe and assume that the traveling time of cosmic particles can be written as $T = L/v(E)$. Thus the arrival time difference between a high-energy photon with energy E and a low-energy light is

$$\Delta T = \frac{L}{v(E)} - \frac{L}{c} = s_n \frac{(n+1)L}{2c} \left(\frac{E}{E_{LV,n}} \right)^n, \quad (3)$$

and the n th-order Lorentz violation scale can be written as

$$E_{LV,n} = \left(s_n \frac{(1+n)L}{2c\Delta T} \right)^{\frac{1}{n}} E. \quad (4)$$

*Corresponding author

Email address: mabq@pku.edu.cn (Bo-Qiang Ma)

One aspect we need to notice is where are these ultra-high-energy photons produced, since the strong gravitational field close to the pulsar may influence the traveling time difference. From Ref.[8], these ultra-high-energy photons are most likely created by the inverse-Compton (IC) scattering of high-energy electrons with low-energy photons. To produce such high-energy photons, the energies of the electrons must be higher. These electrons were accelerated for a long time in the electromagnetic field of the pulsar before the IC scattering, which means that these ultra-high-energy photons are produced far away from the pulsar. In other words, we can dismiss the influence of the gravitational field. During the propagation of the photons, the interstellar medium (ISM) may slow down the photons. But for ultra-high-energy photons, we can dismiss this effect because if ultra-high-energy photons interact with ISM, ultra-high-energy photons will be absorbed and create new particles. From Fig. 1 of Ref. [8], the phasograms of Vela as measured with H.E.S.S. CT1–4 for energies above 5 TeV and with H.E.S.S. CT5 in the 10 to 80 GeV range, we can see a possible time difference of peak P2 between energy above 5 TeV and 10 – 80 GeV. We can see that the high-energy photons travel slower than low-energy photons, and that suggests $s_n = +1$. The time difference between the two peaks is one-half of the bin width of the phasogram for energies above 5 TeV ($0.01 \times$ spin period). Considering that the time difference read from the phasograms can be also from bin division, which means that the actual time difference can be smaller, we can obtain that $\Delta T < 0.01 T_{\text{period}} = 0.89$ ms. Since $5 \text{ TeV} \gg 80 \text{ GeV}$, we can ignore the contribution of the time difference from low-energy photons, and the energy of high-energy photons has a relation $E > 5 \text{ TeV}$. From Eq. (4), for $n = 1$, we have

$$E_{LV,1} = \frac{L}{c\Delta T} E > 1.66 \times 10^{17} \text{ GeV}. \quad (5)$$

For $n = 2$, we have

$$E_{LV,2} = \left(\frac{3L}{2c\Delta T} \right)^{\frac{1}{2}} E > 3.53 \times 10^{10} \text{ GeV}. \quad (6)$$

It is worth noting that, the results we obtained in Eq. (5) and Eq. (6) are somewhat weaker compared to some existing constraints in the literature [3, 4] (see, also, a conservative bound $E_{LV,1} \geq 3.6 \times 10^{17} \text{ GeV}$ in Refs. [12–21]). However these results can be expected to be refined by analyzing the data of the Vela pulsar more thoroughly and utilizing potentially more observations of very-high-energy photons from pulsars in the future. We anticipate that more stringent constraints on Lorentz invariance violation parameters can be obtained. On the other hand, the constraints obtained in this work, albeit weak, reveal the feasibility of utilizing pulsar photons to constrain Lorentz invariance violation parameters. Therefore in this work, we show that the Vela pulsar expands the range of sources that can be chosen to effectively study Lorentz invariance violation.

Our approach can be also extended to analyze the traveling time difference for photons with very big energy differences from the Crab pulsar, which is the most powerful pulsar in our Galaxy. The distance of the Crab pulsar is 6.5 times that of

the Vela pulsar, while the Crab pulsar spins 30 times a second, with a spin period around 33 ms, which is comparable with 89 ms of the Vela pulsar. Therefore we can estimate that the Crab pulsar may provide stronger or comparable constraints on the Lorentz violation parameter of photons if there are data of around 10 TeV photons from Crab pulsar. Unfortunately at the moment, we have no information about 10-TeV scale photons from the Crab pulsar for our analysis, what we know is that the pulsed emission up to 1.5 TeV has been detected from the Crab pulsar [22]. With the constraints of Eqs. (5) and (6), we can provide a rough formula that the predicted time delay between high-energy photons and low-energy photons satisfies $\Delta T < 5.8 \text{ ms} \times (E/5 \text{ TeV})^n$, where E is the energy of high-energy photons and $n = 1, 2$ is the order of Lorentz violation.

Acknowledgements

This work is supported by National Natural Science Foundation of China under Grants No. 12335006 and No. 12075003.

References

- [1] G. Amelino-Camelia, J. Ellis, N. E. Mavromatos, D. V. Nanopoulos, *Distance measurement and wave dispersion in a Liouville-String approach to quantum gravity*, International Journal of Modern Physics A 12 (03) (1997) 607–623. [arXiv:9605211](https://arxiv.org/abs/9605211), [doi:10.1142/S0217751X97000566](https://doi.org/10.1142/S0217751X97000566). URL <https://www.worldscientific.com/doi/abs/10.1142/S0217751X97000566>.
- [2] G. Amelino-Camelia, J. Ellis, N. E. Mavromatos, D. V. Nanopoulos, S. Sarkar, *Tests of quantum gravity from observations of γ -ray bursts*, Nature 393 (6687) (1998) 763–765. [doi:10.1038/31647](https://doi.org/10.1038/31647). URL <http://www.nature.com/articles/31647>.
- [3] J.-J. Wei, X.-F. Wu, *Tests of Lorentz invariance* (2022). [arXiv:2111.02029](https://arxiv.org/abs/2111.02029), [doi:10.1007/978-981-16-4544-0_132-1](https://doi.org/10.1007/978-981-16-4544-0_132-1). URL https://doi.org/10.1007/978-981-16-4544-0_132-1.
- [4] P. He, B.-Q. Ma, *Lorentz Symmetry Violation of Cosmic Photons*, Universe 8 (6) (2022) 323. [arXiv:2206.08180](https://arxiv.org/abs/2206.08180), [doi:10.3390/universe8060323](https://doi.org/10.3390/universe8060323).
- [5] A. N. Otte, *Prospects of performing Lorentz invariance tests with VHE emission from pulsars*, in: 32nd International Cosmic Ray Conference, Vol. 7, 2012, pp. 256–259. [arXiv:1208.2033](https://arxiv.org/abs/1208.2033), [doi:10.7529/ICRC2011/V07/1302](https://doi.org/10.7529/ICRC2011/V07/1302).
- [6] B. Zitzer, *Lorentz Invariance Violation Limits from the Crab Pulsar using VERITAS*, in: 33rd International Cosmic Ray Conference, 2013, p. 1147. [arXiv:1307.8382](https://arxiv.org/abs/1307.8382).
- [7] M. L. Ahnen, et al., *Constraining Lorentz invariance violation using the Crab Pulsar emission observed up to TeV energies by MAGIC*, Astrophys. J. Suppl. 232 (1) (2017) 9. [arXiv:1709.00346](https://arxiv.org/abs/1709.00346), [doi:10.3847/1538-4365/aa8404](https://doi.org/10.3847/1538-4365/aa8404).
- [8] The H.E.S.S. Collaboration et al. and R. Zanin, M. Kerr et al., *Discovery of a radiation component from the Vela pulsar reaching 20 teraelectronvolts*, Nat Astron (2023). [doi:10.1038/s41550-023-02052-3](https://doi.org/10.1038/s41550-023-02052-3).
- [9] R. Dodson, D. Legge, J. E. Reynolds, P. M. McCulloch, *The Vela pulsar’s proper motion and parallax derived from VLBI observations*, Astrophys. J. 596 (2003) 1137–1141. [arXiv:astro-ph/0302374](https://arxiv.org/abs/astro-ph/0302374), [doi:10.1086/378089](https://doi.org/10.1086/378089).
- [10] U. Jacob, T. Piran, *Lorentz-violation-induced arrival delays of cosmological particles*, Journal of Cosmology and Astroparticle Physics 2008 (01) (2008) 031. [arXiv:0712.2170](https://arxiv.org/abs/0712.2170), [doi:10.1088/1475-7516/2008/01/031](https://doi.org/10.1088/1475-7516/2008/01/031). URL <https://iopscience.iop.org/article/10.1088/1475-7516/2008/01/031>.
- [11] J. Zhu, B.-Q. Ma, *Lorentz-violation-induced arrival time delay of astroparticles in Finsler spacetime*, Phys. Rev. D 105 (12) (2022) 124069. [arXiv:2206.07616](https://arxiv.org/abs/2206.07616), [doi:10.1103/PhysRevD.105.124069](https://doi.org/10.1103/PhysRevD.105.124069).
- [12] L. Shao, Z. Xiao, B.-Q. Ma, *Lorentz violation from cosmological objects with very high energy photons*, Astroparticle Physics 33 (5-6) (2010) 312–315. [arXiv:0911.2276](https://arxiv.org/abs/0911.2276), [doi:10.1016/j.astropartphys.2010.03.003](https://doi.org/10.1016/j.astropartphys.2010.03.003). URL <https://linkinghub.elsevier.com/retrieve/pii/S09276505100006>.

- [13] S. Zhang, B.-Q. Ma, Lorentz violation from gamma-ray bursts, *Astroparticle Physics* 61 (2015) 108–112. [arXiv:1406.4568](#), [doi:10.1016/j.astropartphys.2014.04.008](#).
- [14] H. Xu, B.-Q. Ma, Light speed variation from gamma-ray bursts, *Astroparticle Physics* 82 (2016) 72–76. [arXiv:1607.03203](#), [doi:10.1016/j.astropartphys.2016.05.008](#).
URL <http://linkinghub.elsevier.com/retrieve/pii/S0927650516300792>
- [15] H. Xu, B.-Q. Ma, Light speed variation from gamma ray burst GRB 160509A, *Physics Letters B* 760 (2016) 602–604. [arXiv:1607.08043](#), [doi:10.1016/j.physletb.2016.07.044](#).
URL <https://linkinghub.elsevier.com/retrieve/pii/S0370269316303859>
- [16] G. Amelino-Camelia, G. D’Amico, G. Rosati, N. Loret, In vacuo dispersion features for gamma-ray-burst neutrinos and photons, *Nature Astronomy* 1 (2017) 1–12. [arXiv:1612.02765v1](#), [doi:10.1038/s41550-017-0139](#).
- [17] H. Xu, B.-Q. Ma, Regularity of high energy photon events from gamma ray bursts, *Journal of Cosmology and Astroparticle Physics* 2018 (01) (2018) 050–050. [arXiv:1801.08084](#), [doi:10.1088/1475-7516/2018/01/050](#).
URL <https://iopscience.iop.org/article/10.1088/1475-7516/2018/01/050>
- [18] Y. Liu, B.-Q. Ma, Light speed variation from gamma ray bursts: criteria for low energy photons, *The European Physical Journal C* 78 (10) (2018) 825. [arXiv:1810.00636](#), [doi:10.1140/epjc/s10052-018-6294-y](#).
URL <http://link.springer.com/10.1140/epjc/s10052-018-6294-y>
- [19] H. Li, B.-Q. Ma, Light speed variation from active galactic nuclei, *Science Bulletin* 65 (4) (2020) 262–266. [doi:10.1016/j.scib.2019.11.024](#).
URL <https://linkinghub.elsevier.com/retrieve/pii/S2095927319306693>
- [20] J. Zhu, B.-Q. Ma, Pre-burst events of gamma-ray bursts with light speed variation, *Physics Letters B* 820 (2021) 136518. [doi:10.1016/j.physletb.2021.136518](#).
- [21] Y. Chen, B.-Q. Ma, Novel pre-burst stage of gamma-ray bursts from machine learning, *Journal of High Energy Astrophysics* 32 (2021) 78–86. [doi:https://doi.org/10.1016/j.jheap.2021.09.002](#).
URL <https://www.sciencedirect.com/science/article/pii/S2214404821000380>
- [22] S. Ansoldi, et al., Teraelectronvolt pulsed emission from the Crab pulsar detected by MAGIC, *Astron. Astrophys.* 585 (2016) A133. [arXiv:1510.07048](#), [doi:10.1051/0004-6361/201526853](#).