

# Edge-connectivity keeping trees in $k$ -edge-connected graphs\*

Qing Yang, Yingzhi Tian<sup>†</sup>

College of Mathematics and System Sciences, Xinjiang University, Urumqi, Xinjiang, 830046, PR China

**Abstract** Mader [J. Combin. Theory Ser. B 40 (1986) 152-158] proved that every  $k$ -edge-connected graph  $G$  with minimum degree at least  $k + 1$  contains a vertex  $u$  such that  $G - \{u\}$  is still  $k$ -edge-connected. In this paper, we prove that every  $k$ -edge-connected graph  $G$  with minimum degree at least  $k + 2$  contains an edge  $uv$  such that  $G - \{u, v\}$  is  $k$ -edge-connected for any positive integer  $k$ . In addition, we show that for any tree  $T$  of order  $m$ , every  $k$ -edge-connected graph  $G$  with minimum degree greater than  $4(k+m)^2$  contains a subtree  $T'$  isomorphic to  $T$  such that  $G - V(T')$  is  $k$ -edge-connected.

**Keywords:** Edge-connectivity;  $k$ -edge-connected graphs; Trees

## 1 Introduction

Throughout this paper, graph always means a finite, undirected graph without multiple edges and loops. Let  $G$  be a graph with *vertex set*  $V(G)$ , *edge set*  $E(G)$  and *order*  $|V(G)|$ . For a vertex  $v \in V(G)$ , the *neighborhood* of  $v$  is  $N_G(v)$ . For a vertex set  $S \subseteq V(G)$ , the *neighborhood*  $N_G(S)$  of  $S$  in  $G$  is  $(\cup_{v \in S} N_G(v)) \setminus S$ . The *degree*  $d_G(v)$  of  $v$  in  $G$  is  $|N_G(v)|$ . The *minimum degree*  $\delta(G)$  of  $G$  is  $\min_{v \in V(G)} d_G(v)$ . The *induced subgraph*  $G[S]$ , is the graph with vertex set  $S$ , where two vertices in  $S$  are adjacent if and only if they are adjacent in  $G$ . And  $G - S$  is the induced graph  $G[V(G) \setminus S]$ . For an edge set  $U \subseteq E(G)$ , let  $G - U$  be the subgraph of  $G$  with vertex set  $V(G)$  and edge set  $E(G) \setminus U$ . For graph-theoretical terminologies and notation not defined here, we follow [1].

The *connectivity* of  $G$ , denoted by  $\kappa(G)$ , is the minimum size of a vertex set  $S \subseteq V(G)$  such that  $G - S$  is disconnected or has only one vertex. The *edge-connectivity* of  $G$ , denoted by  $\kappa'(G)$ , is the minimum size of an edge set  $U \subseteq E(G)$  such that  $G - U$  is disconnected. The graph  $G$  is said to be  *$k$ -connected* (respectively,  *$k$ -edge-connected*) if  $\kappa(G) \geq k$  (respectively,  $\kappa'(G) \geq k$ ). Note that a trivial graph is both 1-connected and 1-edge-connected, but is not  $k$ -connected and  $k$ -edge-connected for any  $k > 1$ .

In 1972, Chartrand, Kaigars and Lick [2] proved that there is a redundant vertex in a  $k$ -connected graph with minimum degree at least  $\lfloor \frac{3k}{2} \rfloor$ .

**Theorem 1.1.** (Chartrand et al.[2]) *Every  $k$ -connected graph  $G$  with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor$  contains a vertex  $u$  such that  $\kappa(G - \{u\}) \geq k$ .*

\*The research is supported by National Natural Science Foundation of China (11861066).

<sup>†</sup>Corresponding author. E-mail: tianyzhxj@163.com (Y. Tian), qingyangmath@sina.com (Q. Yang).

In 2008, Fujita and Kawarabayashi [4] showed that every  $k$ -connected graph  $G$  with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + 2$  contains an edge  $uv$  such that  $\kappa(G - \{u, v\}) \geq k$ . Furthermore, they conjectured that every  $k$ -connected graph with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + f_k(m) - 1$  contains a connected subgraph  $H$  of order  $m$  such that  $\kappa(G - V(H)) \geq k$ , where  $f_k(m)$  is nonnegative. Mader [11] confirmed the conjecture and proved that  $f_k(m) = m$ . In addition, the connected subgraph  $H$  can be chosen as a path. Meanwhile, Mader conjectured that the result would hold even if we replace the path by any tree with the same order.

**Conjecture 1.** (Mader [11]) *For any tree  $T$  with order  $m$ , every  $k$ -connected graph  $G$  with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + m - 1$  contains a tree  $T' \cong T$  such that  $\kappa(G - V(T')) \geq k$ .*

Actually, the theorem in [3] implies Conjecture 1 holds for  $k = 1$ . When  $k = 2$ , Conjecture 1 holds for star, path-star, spider and caterpillar et al., see [5-7, 9, 14-15] for references. In 2022, Hong and Liu [8] confirmed Conjecture 1 for  $k = 2$  and  $k = 3$ . Conjecture 1 remains open for  $k \geq 4$ . However, Mader [12] showed that Conjecture 1 holds if  $\delta(G) \geq 2(k - 1 + m)^2 + m - 1$ .

In this paper, we consider a similar problem in the  $k$ -edge-connected graphs. Mader [10] showed that there is a redundant vertex in a  $k$ -edge-connected graph with minimum degree at least  $k + 1$ .

**Theorem 1.2.** (Mader [10]) *Every  $k$ -edge-connected graph  $G$  with  $\delta(G) \geq k + 1$  contains a vertex  $u$  such that  $\kappa'(G - \{u\}) \geq k$ .*

In this paper, we prove that there exists a pair of redundant adjacent vertices in a  $k$ -edge-connected graph  $G$  with  $\delta(G) \geq k + 2$  for any positive integer  $k$ . Meanwhile, we also obtain that for any tree  $T$  of order  $m$ , every  $k$ -edge-connected graph  $G$  with  $\delta(G) > 4(k + m)^2$  contains a subtree  $T' \cong T$  such that  $\kappa'(G - V(T')) \geq k$ . In the next section, we will introduce some definitions and lemmas that will be used to prove the main results. In Section 3, we present the main results. Concluding remarks will be given in the last section.

## 2 Preliminaries

Let  $U$  be an edge-cut of  $G$  and  $F$  be the union of at least one component, but not all components of  $G - U$ . Then  $F$  is called a *semifragment* of  $G$  to  $U$ . And  $F' = G - V(F)$  is called a *complementary semifragment* of  $F$  in  $G$  to  $U$ . In fact,  $F'$  is also a semifragment of  $G$  to  $U$ . In addition, if  $U$  is a minimum edge-cut of  $G$ , then  $G - U$  has exactly two components  $F$  and  $F'$ . Each of  $F$  and  $F'$  is a *fragment* of  $G$  to  $U$ , and  $F'$  is a complementary fragment of  $F$  in  $G$  to  $U$ . Sometimes, we simply say  $F$  is a fragment of  $G$ .

For two vertex set  $V_1, V_2 \subseteq G$ , the edge set between  $V_1$  and  $V_2$  in  $G$  is denoted by  $E_G[V_1, V_2]$  and the number of all edges between  $V_1$  and  $V_2$  in  $G$  is denoted by  $d_G(V_1, V_2)$ , which is  $|E_G[V_1, V_2]|$  or  $\sum_{u \in V_1} |N_G(u) \cap V_2|$ . For convenience, this paper may use  $G[V_1]$  and  $G[V_2]$  for  $V_1$  and  $V_2$ , respectively. For examples, we use  $G[V_1] \cap G[V_2]$ ,  $G[V_1] \cup G[V_2]$  and  $d_G(G[V_1], G[V_2])$  for  $V_1 \cap V_2$ ,  $V_1 \cup V_2$  and  $d_G(V_1, V_2)$ , respectively.

**Lemma 2.1.** *Let  $e$  and  $e_1$  be two nonadjacent edges of  $G$ , and, let  $U$  and  $U_1$  be minimum edge-cuts of  $G - V(e)$  and  $G - V(e_1)$ , respectively. Assume that  $F$  is a fragment of  $G - V(e)$*

to  $U$  and  $F_1$  is a fragment of  $G - V(e_1)$  to  $U_1$ . And we denote  $F' = G - V(e) - V(F)$  and  $F'_1 = G - V(e_1) - V(F_1)$ . Furthermore, we assume  $e \in E(F_1)$ ,  $e_1 \in E(F')$  and  $F \cap F_1 \neq \emptyset$  (see Figure 1). Then we have the following properties.

- ( $\alpha$ ) If  $F' \cap F'_1 \neq \emptyset$  holds, then  $F \cap F_1$  is a fragment of  $G - V(e)$ .  
 ( $\beta$ ) If  $F' \cap F'_1 = \emptyset$  holds, then  $|V(F')| < |V(F_1)|$ .




Figure 1: Illustration for Lemma 2.1

**Proof.** ( $\alpha$ ) Since  $F \cap F_1 \neq \emptyset$ , we have

$$\begin{aligned} d_{G-V(e)}(F \cap F_1, F' \cup F'_1) &\geq \kappa'(G - V(e)) \\ &= d_G(F, F') \\ &= d_G(F \cap F_1, F') + d_G(F \cap F'_1, F'). \end{aligned} \quad (1)$$

Thus

$$d_G(F \cap F_1, F \cap F'_1) \geq d_G(F \cap F'_1, F') \quad (2)$$

by  $d_{G-V(e)}(F \cap F_1, F' \cup F'_1) = d_G(F \cap F_1, F') + d_G(F \cap F_1, F \cap F'_1)$ .

If  $F' \cap F'_1 \neq \emptyset$  holds, then

$$\begin{aligned} d_{G-V(e_1)}(F' \cap F'_1, F \cup F_1) &\geq \kappa'(G - V(e_1)) \\ &= d_G(F_1, F'_1) \\ &= d_G(F' \cap F'_1, F_1) + d_G(F \cap F'_1, F_1). \end{aligned} \quad (3)$$

Thus

$$d_G(F \cap F'_1, F' \cap F'_1) \geq d_G(F \cap F'_1, F_1) \quad (4)$$

by  $d_{G-V(e_1)}(F' \cap F'_1, F \cup F_1) = d_G(F' \cap F'_1, F_1) + d_G(F' \cap F'_1, F \cap F'_1)$ . By inequalities (2) and (4), we get

$$\begin{aligned} d_G(F \cap F_1, F \cap F'_1) &\geq d_G(F \cap F'_1, F') \\ &\geq d_G(F \cap F'_1, F' \cap F'_1) \\ &\geq d_G(F \cap F'_1, F_1) \\ &\geq d_G(F \cap F_1, F \cap F'_1), \end{aligned} \quad (5)$$

that is,  $d_G(F \cap F_1, F \cap F'_1) = d_G(F \cap F'_1, F')$ . Moreover, we have

$$\begin{aligned}
d_{G-V(e)}(F \cap F_1, G - V(e) - (F \cap F_1)) &= d_{G-V(e)}(F \cap F_1, F' \cup F'_1) \\
&= d_G(F \cap F_1, F') + d_G(F \cap F_1, F \cap F'_1) \\
&= d_G(F \cap F_1, F') + d_G(F \cap F'_1, F') \quad (6) \\
&= d_G(F, F') \\
&= \kappa'(G - V(e)).
\end{aligned}$$

Thus,  $U' = E_{G-V(e)}[F \cap F_1, G - V(e) - (F \cap F_1)]$  is a minimum edge-cut of  $G - V(e)$  and  $F \cap F_1$  is a fragment of  $G - V(e)$  to  $U'$ .

( $\beta$ ) If  $F' \cap F'_1 = \emptyset$  holds, then  $F' - V(e_1) \subseteq F_1$ . Since  $V(e) \in V(F_1)$ ,  $V(e) \notin V(F')$  and  $F \cap F_1 \neq \emptyset$ , we have  $|V(F')| < |V(F_1)|$ .  $\square$

**Lemma 2.2.** *Let  $G$  be a  $k$ -edge-connected graph with  $\delta(G) \geq k + 2$  and let  $e_0$  be an edge of graph  $G$  such that  $\kappa'(G - V(e_0)) < k$ . Assume that  $U_0$  is a minimum edge-cut of  $G - V(e_0)$  and  $F_0$  is a fragment of  $G - V(e_0)$  to  $U_0$ . Then there exist an edge  $e_1 \in E(G[V(F_0) \cup V(e_0)])$  with  $\kappa'(G - V(e_1)) < k$  and a fragment  $F_1$  of  $G - V(e_1)$  with  $F_1 \subseteq G[V(F_0) \cup V(e_0)]$  such that the following property holds.*

*For any  $e \in E(F_1)$ , if  $\kappa'(G - V(e)) < k$  holds, then any fragment  $F$  of  $G - V(e)$  with  $e_1 \notin E(F)$  satisfies  $F \cap F_1 = \emptyset$ . See Figure 2 for an illustration.*




Figure 2: Illustration for Lemma 2.2

**Proof.** Let  $E_0 = \{e \in E(G[V(F_0) \cup V(e_0)]) : \kappa'(G - V(e)) < k\}$  and for any  $e \in E_0$ , let

$$\mathcal{U}_e = \{U : U \text{ is a minimum edge-cut of } G - V(e)\}.$$

For any  $U_e \in \mathcal{U}_e$ , we denote

$$\mathcal{F}_{U_e} = \{F : F \text{ is a fragment of } G - V(e) \text{ to } U_e \text{ with } F \subseteq G[V(F_0) \cup V(e_0)]\}.$$

Furthermore, let  $\mathcal{F} = \cup_{e \in E_0} (\cup_{U_e \in \mathcal{U}_e} \mathcal{F}_{U_e})$ . Then  $e_0 \in E_0$ ,  $U_0 \in \mathcal{U}_{e_0}$  and  $F_0 \in \mathcal{F}_{U_0}$ . Then

Then  $e \in E_0$  by  $F_1 \subseteq G[V(F_0) \cup V(e_0)]$ . Let  $U$  be a minimum edge-cut of  $G - V(e)$ ,  $F$  be a fragment of  $G - V(e)$  to  $U$  with  $e_1 \notin E(F)$  and  $F' = G - V(e) - V(F)$  be the complementary fragment of  $F$  in  $G - V(e)$  to  $U$ . Then  $e_1 \in E(F')$ .

Suppose  $F \cap F_1 \neq \emptyset$ . By Lemma 2.1 ( $\alpha$ ), if  $F' \cap F'_1 \neq \emptyset$  holds, then  $F \cap F_1$  is a fragment of  $G - V(e)$ . Since  $F_1 \subseteq G[V(F_0) \cup V(e_0)]$ ,  $\kappa'(G - V(e)) < k$  and  $e \in E(F_1)$ , we get  $F \cap F_1 \in \mathcal{F}$ . By  $e \in E(F_1)$  and  $e \notin E(F)$ , we have  $|F \cap F_1| < |V(F_1)|$ , which contradicts the choice of  $F_1$ . By Lemma 2.1 ( $\beta$ ), if  $F' \cap F'_1 = \emptyset$  holds, then  $|V(F')| < |V(F_1)|$ . Since  $F' \cap F'_1 = \emptyset$  and  $e_1 \in E(F')$ , we obtain

$$F' \subseteq G[V(F_1) \cup V(e_1)] \subseteq G[V(F_0) \cup V(e_0)].$$

Thus  $F' \in \mathcal{F}$ , also a contradiction.  $\square$

### 3 Main results

**Theorem 3.1.** *For any positive integer  $k$ , every graph  $G$  with  $\kappa'(G) \geq k$  and  $\delta(G) \geq k + 2$  contains an edge  $e$  such that  $\kappa'(G - V(e)) \geq k$ .*

**Proof.** By contrary, assume that  $\kappa'(G - V(e)) < k$  for any  $e \in E(G)$ . Then we have the following claims.

**Claim 1.** There exist an edge  $e_1 \in E(G)$  and a fragment  $F_1$  of  $G - V(e_1)$  such that

$$2 \leq |V(F_1)| < k.$$

Let  $e_0 \in E(G)$ . Assume that  $U_0$  is a minimum edge-cut of  $G - V(e_0)$  and  $F_0$  is a fragment of  $G - V(e_0)$  to  $U_0$ . Since  $\kappa'(G - V(e_0)) < k$ , by Lemma 2.2, there exist  $e_1 \in E(G[V(F_0) \cup V(e_0)])$  and a fragment  $F_1 \subseteq G[V(F_0) \cup V(e_0)]$  of  $G - V(e_1)$  such that for any  $e \in E(F_1)$  and any fragment  $F$  of  $G - V(e)$  with  $e_1 \notin E(F)$ , we have  $F \cap F_1 = \emptyset$ . Let  $F' = G - V(e) - V(F)$  and  $F'_1 = G - V(e_1) - V(F_1)$ . Thus  $F \subseteq F'_1$ . Since  $\kappa'(G - V(e)) < k$ , we have  $d_G(V(e), F) > 0$ . By  $F \subseteq F'_1$ , we get  $d_G(V(e), F'_1) > 0$ . If there exists a vertex  $u \in V(F_1)$  such that  $N_G(u) \cap V(F'_1) = \emptyset$ , then by  $d_G(V(e), F'_1) > 0$  for any  $e \in E(F_1)$ , we have  $N_G(v) \cap V(F'_1) \neq \emptyset$  for any  $v \in N_G(u) \cap V(F_1)$ . Since

$$\begin{aligned} |N_G(u) \cap V(F_1)| &= d_G(u) - |N_G(u) \cap V(F'_1)| - |N_G(u) \cap V(e_1)| \\ &\geq k + 2 - 0 - 2 \\ &= k, \end{aligned} \tag{7}$$

we have  $k \leq |N_G(u) \cap V(F_1)| \leq d_G(F_1, F'_1) = \kappa'(G - V(e_1)) < k$ , a contradiction. Thus, for any  $w \in V(F_1)$ , we obtain  $N_G(w) \cap V(F'_1) \neq \emptyset$ . Since  $d_G(F_1, F'_1) < k$ , we have  $|V(F_1)| < k$ . Thus Claim 1 holds.

By claim 1, the following claim is established.

**Claim 2.**  $k \geq 3$ .

Since  $G$  is a simple graph with  $\delta(G) \geq k + 2$ , we have

$$d_G(F_1, F'_1) \geq |V(F_1)|(k - |V(F_1)| + 1)$$

by

$$\begin{aligned}
|N_G(z) \cap V(F'_1)| &= d_G(z) - |N_G(z) \cap V(F_1)| - |N_G(z) \cap V(e_1)| \\
&\geq k + 2 - (|V(F_1)| - 1) - 2 \\
&= k - |V(F_1)| + 1
\end{aligned} \tag{8}$$

for any  $z \in V(F_1)$ . Thus  $d_G(F_1, F'_1) \geq |V(F_1)|(k - |V(F_1)| + 1) \geq 2(k - 1)$  by  $2 \leq |V(F_1)| < k$ . This, together with  $d_G(F_1, F'_1) < k$ , we obtain  $k < 2$ , which contradicts to  $k \geq 3$  (by Claim 2).  $\square$

The following lemma will be used to prove that there exists a redundant subtree in  $k$ -edge-connected graphs with minimum degree greater than  $4(k + m)^2$ .

**Lemma 3.2.** (Thomassen [13]) *Any graph  $G$  with  $\delta(G) > 4k^2$  contains a  $k$ -connected subgraph  $H$  with  $|V(H)| > 4k^2$  such that  $|N_G(G - V(H))| \leq 2k^2$ .*

**Theorem 3.3.** *For any tree  $T$  with order  $m$ , every  $k$ -edge-connected graph  $G$  with  $\delta(G) > 4(k + m)^2$  contains a tree  $T' \cong T$  such that  $G - V(T')$  is still  $k$ -edge-connected.*

**Proof.** By Lemma 3.2, there exists a  $(k + m)$ -connected subgraph  $H$  with  $|V(H)| > 4(k + m)^2$  in  $G$  such that  $|N_G(G - V(H))| \leq 2(k + m)^2$ . Let  $\bar{H} = G - V(H)$ . Since  $\delta(G) > 4(k + m)^2$ , we obtain  $\delta(H - N_G(\bar{H})) \geq m$ , thus there is a subtree  $T' \cong T$  in  $H - N_G(\bar{H})$ . If  $G - V(T')$  is  $k$ -edge-connected, then the result holds. So, we assume  $\kappa'(G - V(T')) \leq k - 1$ .

Let  $U$  be a minimum edge-cut of  $G - V(T')$ . Then  $|U| \leq k - 1$ . Let  $F$  be a fragment of  $G - V(T')$  to  $U$  and  $F' = G - V(T') - V(F)$ . Note that  $F'$  is also a fragment of  $G - V(T')$  to  $U$ . We give some notations as follows (see Figure 3 for an illustration).

$$\begin{aligned}
H_1 &= F \cap H, & H_2 &= F' \cap H; \\
\bar{H}_1 &= F \cap \bar{H}, & \bar{H}_2 &= F' \cap \bar{H}; \\
D_1 &= V(U) \cap F, & D_2 &= V(U) \cap F'.
\end{aligned}$$




Figure 3: Illustration for the proof of Theorem 3.3

By  $|U| \leq k - 1$ , we get  $|D_1| \leq k - 1$  and  $|D_2| \leq k - 1$ . Since  $H$  is  $(k + m)$ -connected,  $H - D_1 \cup V(T')$  is connected, thus

$$H_1 \setminus D_1 = \emptyset \text{ or } H_2 = \emptyset. \tag{9}$$

And by  $H - D_2 \cup V(T')$  is connected, we have

$$H_2 \setminus D_2 = \emptyset \text{ or } H_1 = \emptyset. \quad (10)$$

If both  $H_1 \setminus D_1 = \emptyset$  and  $H_2 \setminus D_2 = \emptyset$ , then  $|V(H) \setminus V(T')| \leq |D_1 \cup D_2| \leq 2(k-1)$ , which is a contradiction to  $|V(H)| > 4(k+m)^2$ . Thus

$$H_1 \setminus D_1 \neq \emptyset \text{ or } H_2 \setminus D_2 \neq \emptyset. \quad (11)$$

If  $H_1 \neq \emptyset$  holds, then  $H_2 \setminus D_2 = \emptyset$  by (10). Suppose that  $H_2 \neq \emptyset$  holds. Then  $H_1 \setminus D_1 = \emptyset$  by (9), a contradiction to (11). Thus  $H_2 = \emptyset$ . Since  $F'$  is a fragment of  $G - V(T')$  to  $U$ , we have  $F' \neq \emptyset$ , thus  $\bar{H}_2 \neq \emptyset$ . So,

$$k \leq \kappa'(G) \leq d_G(\bar{H}_2, V(G) \setminus \bar{H}_2) \leq |U| \leq k-1,$$

a contradiction.

If  $H_1 = \emptyset$  holds, then  $\bar{H}_1 \neq \emptyset$  by  $F \neq \emptyset$ . Thus

$$k \leq \kappa'(G) \leq d_G(\bar{H}_1, V(G) \setminus \bar{H}_1) \leq |U| \leq k-1,$$

also a contradiction.  $\square$

## 4 Concluding remarks

**Theorem 4.1.** (Diwan and Tholiya [3]) *For any tree  $T$  with order  $m$ , every 1-connected graph  $G$  with  $\delta(G) \geq m$  contains a tree  $T' \cong T$  such that  $G - V(T')$  is 1-connected.*

Since the definitions of 1-edge-connected graph and 1-connected graph are equivalent, there exists a redundant subtree in a 1-edge-connected graph with minimum degree of at least  $m$  by Theorem 4.1.

**Theorem 4.2.** *For any tree  $T$  with order  $m$ , every 1-edge-connected graph  $G$  with  $\delta(G) \geq m$  contains a tree  $T' \cong T$  such that  $G - V(T')$  is 1-edge-connected.*

**Theorem 4.3.** (Hong et al. [8]) *For any tree  $T$  with order  $m$ , every 2-connected (respectively, 3-connected) graph  $G$  with  $\delta(G) \geq m+2$  (respectively,  $\delta(G) \geq m+3$ ) contains a subtree  $T' \cong T$  such that  $\kappa(G - V(T')) \geq 2$  (respectively,  $\kappa(G - V(T')) \geq 3$ ).*

Using almost the same proof as Theorem 4.3, we can prove that there is a redundant subtree in a 2-edge-connected (respectively, 3-edge-connected) graph with minimum degree of at least  $m+2$  (respectively,  $m+3$ ). Specifically, we only need to substitute “2-connected” and “3-connected” in the proof of Theorem 4.3 with “2-edge-connected” and “3-edge-connected”, respectively.

**Theorem 4.4.** *For any tree  $T$  with order  $m$ , every 2-edge-connected (respectively, 3-edge-connected) graph  $G$  with  $\delta(G) \geq m+2$  (respectively,  $\delta(G) \geq m+3$ ) contains a subtree  $T' \cong T$  such that  $\kappa'(G - V(T')) \geq 2$  (respectively,  $\kappa'(G - V(T')) \geq 3$ ).*

Motivated by the results above, we would like to propose the following conjecture, which relates to redundant subtree in a generally  $k$ -edge-connected graphs. As the case  $k = 1$  solved completely by Theorem 4.2, we assume  $k \geq 2$  in the following conjecture.

**Conjecture 2.** *For any tree  $T$  of order  $m$  and any integer  $k \geq 2$ , every  $k$ -edge-connected graph  $G$  with minimum degree at least  $k + m$  contains a subtree  $T' \cong T$  such that  $G - V(T')$  is still  $k$ -edge-connected.*

Since  $K_{k+m} - V(T)$  is not  $k$ -edge-connected for any tree  $T$  with order  $m$ , where  $K_{k+m}$  denotes the complete graph of order  $k + m$ , the minimum degree bound would be best possible when Conjecture 2 were true. Theorem 1.2 and Theorem 3.1 confirm Conjecture 2 for  $m = 1$  and  $m = 2$ , respectively. For  $k = 2$  and  $k = 3$ , by Theorem 4.4, Conjecture 2 is also true.

### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### References

- [1] J. A. Bondy, U. S. R. Murty, Graph Theory, Graduate Texts in Mathematics 244, Springer, Berlin, 2008.
- [2] G. Chartrand, A. Kaigars, D.R. Lick, Critically  $n$ -connected graphs, Proc. Amer. Math. Soc. 32 (1972) 63-68.
- [3] A. A. Diwan, N. P. Tholiya, Non-separating trees in connected graphs, Discrete Math. 309 (16) (2009) 5235-5237.
- [4] S. Fujita, K. Kawarabayashi, Connectivity keeping edges in graphs with large minimum degree, J. Combin. Theory Ser. B 98 (2008) 805-811.
- [5] T. Hasunuma, K. Ono, Connectivity keeping trees in 2-connected graphs, J. Graph Theory 94 (2020) 20-29.
- [6] T. Hasunuma, Connectivity Keeping Trees in 2-Connected Graphs with Girth Conditions, Algorithmica 83 (2021) 2697-2718.
- [7] Y. Hong, Q. Liu, C. Lu, Q. Ye, Connectivity keeping caterpillars and spiders in 2-connected graphs, Discrete Math. 344 (3) (2021) 112236.
- [8] Y. Hong, Q. Liu, Mader's conjecture for graphs with small connectivity, J. Graph Theory 101 (3) (2022) 379-388.
- [9] C. Lu, P. Zhang, Connectivity keeping trees in 2-connected graphs, Discrete Math. 343 (2) (2020) 1-4.
- [10] W. Mader, Kritisch  $n$ -fach kantenzusammenhängende Graphen, J. Combin. Theory Ser. B 40 (1986), 152-158.



- [11] W. Mader, Connectivity keeping paths in  $k$ -connected graphs, *J. Graph Theory* 65 (2010) 61-69.
- [12] W. Mader, Connectivity keeping trees in  $k$ -connected graphs, *J. Graph Theory* 69 (2012) 324-329.
- [13] C. Thomassen, The Erdős-Pósa property for odd cycles in graphs of large connectivity, *Combinatorica* 21 (2001) 321-333.
- [14] Y. Tian, J. Meng, H. Lai, L. Xu, Connectivity keeping stars or double-stars in 2-connected graphs, *Discrete Math.* 341 (4) (2018) 1120-1124.
- [15] Y. Tian, H. Lai, L. Xu, J. Meng, Nonseparating trees in 2-connected graphs and oriented trees in strongly connected digraphs, *Discrete Math.* 342 (2) (2019) 344-351.