

On the monotonicity of discrete entropy for log-concave random vectors on \mathbb{Z}^d

Matthieu Fradelizi¹, Lampros Gavalakis^{*1}, and Martin Rapaport¹

¹Univ Gustave Eiffel, Univ Paris Est Creteil, CNRS, LAMA UMR8050 F-77447 Marne-la-Vallée, France

July 9, 2024

Abstract

We prove the following type of discrete entropy monotonicity for sums of isotropic, log-concave, independent and identically distributed random vectors X_1, \dots, X_{n+1} on \mathbb{Z}^d :

$$H(X_1 + \dots + X_{n+1}) \geq H(X_1 + \dots + X_n) + \frac{d}{2} \log \left(\frac{n+1}{n} \right) + o(1),$$

where $o(1)$ vanishes as $H(X_1) \rightarrow \infty$. Moreover, for the $o(1)$ -term, we obtain a rate of convergence $O\left(H(X_1)e^{-\frac{1}{4}H(X_1)}\right)$, where the implied constants depend on d and n . This generalizes to \mathbb{Z}^d the one-dimensional result of the second named author (2023). As in dimension one, our strategy is to establish that the discrete entropy $H(X_1 + \dots + X_n)$ is close to the differential (continuous) entropy $h(X_1 + U_1 + \dots + X_n + U_n)$, where U_1, \dots, U_n are independent and identically distributed uniform random vectors on $[0, 1]^d$ and to apply the theorem of Artstein, Ball, Barthe and Naor (2004) on the monotonicity of differential entropy. In fact, we show this result under more general assumptions than log-concavity, which are preserved up to constants under convolution. Namely, we consider families of random variables for which, as the determinant of the covariance matrix increases, the probability mass function **i**) is bounded above in terms of the determinant of the covariance matrix, **ii**) has subexponential tails, **iii**) has (discrete) bounded variation. In order to show that log-concave distributions satisfy our assumptions in dimension $d \geq 2$, more involved tools from convex geometry are needed because a suitable position is required. We show that, for a log-concave function on \mathbb{R}^d in isotropic position, its integral, barycenter and covariance matrix are close to their discrete counterparts. Moreover, in the log-concave case, we weaken the isotropicity assumption to what we call *almost isotropicity*. One of our technical tools is a discrete analogue to the upper bound on the isotropic constant of a log-concave function, which extends to dimensions $d \geq 1$ a result of Bobkov, Marsiglietti and Melbourne (2022) in dimension one and may be of independent interest.

Keywords — entropy, discrete log-concavity, entropy power inequality, central limit theorem, convex bodies, isotropic constant


2020 Mathematics subject classification — 94A17; 52C07; 39B62

1 Introduction

1.1 Monotonicity of entropy

The *differential entropy* of an \mathbb{R}^d -valued random vector X with density f is defined as

$$h(X) = - \int_{\mathbb{R}^d} f(x) \log f(x) dx,$$

^{*}L.G. has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 101034255  and by the Bézout Labex, funded by ANR, reference ANR-10-LABX-58.

if the integral exists. When X is supported on a strictly lower-dimensional set than \mathbb{R}^d (and hence does not have a density with respect to the d -dimensional Lebesgue measure), one sets $h(X) = -\infty$. The *entropy power* of X is defined by $N(X) = e^{2h(X)/d}$.

Let X_1, \dots, X_n be i.i.d. continuous random variables. A generalization of the classical Entropy Power Inequality (EPI) of Shannon [22] and Stam [24], due to Artstein, Ball, Barthe and Naor, [1] states that

$$h\left(\frac{1}{\sqrt{n+1}} \sum_{i=1}^{n+1} X_i\right) \geq h\left(\frac{1}{\sqrt{n}} \sum_{i=1}^n X_i\right),$$

which by the scaling property of differential entropy is equivalent to

$$h\left(\sum_{i=1}^{n+1} X_i\right) \geq h\left(\sum_{i=1}^n X_i\right) + \frac{d}{2} \log\left(\frac{n+1}{n}\right). \quad (1)$$

It can also be interpreted as the monotonic increase of entropy along the Central Limit Theorem (CLT) [3]. Later, other proofs and generalizations were given by Shlyakhtenko [23], Tulino and Verdú [26] and Madiman and Barron [17].

Let A be a discrete (finite or countable) set and let X be a random variable supported on A with probability mass function (p.m.f.) p on A . The discrete (Shannon) entropy of X is

$$H(X) = - \sum_{x \in A} p(x) \log p(x).$$

An exact analogue of (1) cannot be true for discrete random variables as can be seen by taking $n = 1, d = 1$ and considering deterministic or even close to deterministic random variables. Nevertheless, Tao [25], using ideas from additive combinatorics, proved that for any independent and identically distributed random variables X_1, X_2 taking values in a finite subset of a torsion-free (abelian) group

$$H(X_1 + X_2) \geq H(X_1) + \frac{1}{2} \log 2 - o(1), \quad (2)$$

where $o(1) \rightarrow 0$ as $H(X_1) \rightarrow \infty$. This can be seen as a discrete analogue of (1) for $d = n = 1$. Besides (2), other discrete versions of the EPI have been studied by several authors. In [11, 12], a discrete EPI similar to (2) was proved, with a worse constant than $\frac{1}{2} \log 2$, but with sharp quantitative bounds for the $o(1)$ -term (see also [9] for a detailed discussion on connections with that work). An EPI for the binomial family was proved in [13, Theorem 1.1]; in [20, Theorem 4.6] it was shown that $e^{2H(X)} - 1$ is superadditive with respect to convolution on the class of uniform distributions on finite subsets of the integers (see also [18] for extensions to Rényi entropy and [19] for dimensional extensions). Finally the constant $\frac{1}{2} \log 2$ was shown to be improvable for the class of uniform distributions in [27].

In addition to (2), Tao conjectured that for any independent and identically distributed random variables X_1, \dots, X_{n+1} taking values in a finite subset of a torsion-free group

$$H\left(X_1 + \dots + X_{n+1}\right) \geq H\left(X_1 + \dots + X_n\right) + \frac{1}{2} \log\left(\frac{n+1}{n}\right) - o(1), \quad (3)$$

as $H(X_1) \rightarrow \infty$ depending on n . A special case of this conjecture was recently proven in [8], where it is shown that (3) is satisfied by log-concave random variables on the integers with explicit rate for the $o(1)$ that is exponential in $H(X_1)$:

Theorem 1 ([8]). *Let X_1, \dots, X_n be i.i.d. log-concave random variables on \mathbb{Z} . Then*

$$H\left(\sum_{i=1}^{n+1} X_i\right) \geq H\left(\sum_{i=1}^n X_i\right) + \frac{1}{2} \log\left(\frac{n+1}{n}\right) - O\left(H(X_1)e^{-H(X_1)}\right),$$

as $H(X_1) \rightarrow \infty$.

This statement can be interpreted as a type of “approximate” monotonicity in the *discrete* entropic CLT [10]. We recall here that an integer valued random variable X with p.m.f. p is said to be *log-concave* if

$$p(k)^2 \geq p(k-1)p(k+1), \quad (4)$$

for every $k \in \mathbb{Z}$.

1.2 Main results and paper outline

Our goal is to extend Theorem 1 to higher dimensions and to the following larger class of probability distributions.

Definition 2. Let $\mathcal{C}(c_1, c_2, c_3)$ be the class of p.m.f.s p on \mathbb{Z}^d for which, denoting $\sigma := \det(\text{Cov}(p))^{\frac{1}{2d}}$,

1.

$$\max_{k \in \mathbb{Z}^d} p(k) \leq \frac{c_1}{\sigma^d},$$

2. for all $k \in \mathbb{Z}^d$ with $\|k\|_\infty \geq c_2\sigma$,

$$p(k) \leq \frac{c_2}{\sigma^d} e^{-\frac{\|k\|_1}{c_2\sigma}} \quad \text{and}$$

3.

$$\sum_{i=1}^d \sum_{k \in \mathbb{Z}^d} |p(k) - p(k - e_i)| \leq \frac{c_3}{\sigma}.$$

Our first main result is the following one:

Theorem 3. For any i.i.d. random vectors X_1, \dots, X_{n+1} on \mathbb{Z}^d such that their common p.m.f. belongs to the class $\mathcal{C}(c_1, c_2, c_3)$, for some absolute constants c_1, c_2, c_3 that depend on the dimension only,

$$H(X_1 + \dots + X_{n+1}) \geq H(X_1 + \dots + X_n) + \frac{d}{2} \log\left(\frac{n+1}{n}\right) - o(1) \quad (5)$$

as $H(X_1) \rightarrow \infty$. Moreover, we have the rate of convergence $o(1) = O_{d,n}\left(H(X_1)e^{-\frac{1}{d}H(X_1)}\right)$. In particular, (5) holds for any i.i.d. log-concave random vectors X_1, \dots, X_{n+1} on \mathbb{Z}^d with almost isotropic extension.

Note that regarding Tao's conjecture 3, our method for this class of random variables improves the result by a constant d , where d is the dimension. In the following subsection, we will explicitly detail the definitions we use for *discrete log-concavity in \mathbb{Z}^d* as well as the definition of *almost isotropic extension*. Theorem 3 readily follows from the next Theorem together with the generalized EPI in \mathbb{R}^d and can be interpreted as an approximation result of discrete and continuous entropies in higher dimensions:

Theorem 4. For any i.i.d. random vectors X_1, \dots, X_{n+1} on \mathbb{Z}^d such that their common p.m.f. belongs to the class $\mathcal{C}(c_1, c_2, c_3)$, for some absolute constants c_1, c_2, c_3 that depend on the dimension only,

$$h(X_1 + \dots + X_n + U_1 + \dots + U_n) = H(X_1 + \dots + X_n) + o(1), \quad (6)$$

where $o(1) \rightarrow 0$ as $H(X_1) \rightarrow \infty$, h stands for the differential entropy, H denotes the discrete Shannon entropy and U_1, \dots, U_n are independent continuous uniforms on $[0, 1]^d$. Moreover, we have the rate of convergence $o(1) = O_{d,n}\left(H(X_1)e^{-\frac{1}{d}H(X_1)}\right)$. In particular, (6) holds for any i.i.d. log-concave random vectors X_1, \dots, X_{n+1} on \mathbb{Z}^d with almost isotropic extension.

Remark 5. Let X be a log-concave random vector on \mathbb{Z}^d with almost isotropic extension. Denote the covariance matrix of X by K_X , $\sigma^2 := \det(K_X)^{\frac{1}{d}}$ and let U be a uniform on the open box $[0, 1]^d$. Then

$$H(X) = h(X + U) \leq \frac{d}{2} \log\left(\det\left(K_X + \frac{1}{12}I_d\right)^{\frac{1}{d}} 2\pi e\right) \simeq \frac{d}{2} \log\left(\sigma^2 2\pi e\right), \quad (7)$$

by the fact that the Gaussian maximizes the entropy under fixed covariance matrix, which implies that for any random variable X on \mathbb{R}^d one has

$$h(X) \leq \frac{d}{2} \log(2\pi e (\det \text{Cov}(X))^{1/d}). \quad (8)$$

Thus, $H(X) \rightarrow \infty$ implies $\sigma \rightarrow \infty$ and therefore it suffices to prove (6) with $o(1) \rightarrow 0$ as $\sigma \rightarrow \infty$.

We prove Theorem 4 in Section 2. First we use the second property of Definition 2, which is an exponential tail bound, in order to control the *entropy tails*. We use the third property, which is a type of smoothness, to control the error term arising from the approximation of the density of $\sum_i X_i + U_i$ by the p.m.f. of $\sum_i X_i$. Combined with the Taylor-type estimate of Lemma 17, which is analogous to an estimate used in [25], this approximation is the main technical step in the proof of Theorem 4. The first property of Definition 2 ensures that Lemma 17 is applicable.

In Section 3 we will show that the assumption that the p.m.f. of X_1 satisfies these three properties is weaker than almost isotropic log-concavity in the sense that any almost isotropic log-concave p.m.f. on \mathbb{Z}^d satisfies these properties with appropriate constants. We remark here that, since our definition of log-concavity reduces to the usual one for the one-dimensional case, Theorems 3 and 4 are also relaxed versions of the corresponding results obtained in [8]. We prove the following theorem, which allows us to establish that if p is log-concave with almost isotropic extension then p belongs to the class $\mathcal{C}(c_1(d), c_2(d), c_3(d))$ for some absolute constants $c_1(d), c_2(d), c_3(d)$ that depend on the dimension only (in particular, this theorem is essential for proving item 1 in Theorem 18):

Theorem 6. *Suppose p is a log-concave p.m.f. on \mathbb{Z}^d with almost isotropic extension and covariance matrix $\text{Cov}(p)$. Then there exists a constant C'_d depending on the dimension only, such that*

$$\max_{k \in \mathbb{Z}^d} p(k) \leq \frac{C'_d}{\det(\text{Cov}_{\mathbb{Z}^d}(p))^{\frac{1}{2}}}$$

provided that $\det(\text{Cov}_{\mathbb{Z}^d}(p))$ is large enough depending on d .

The above Theorem can be seen as d -dimensional analogue of the following result, due to Bobkov, Marsiglietti and Melbourne [4], who also studied discrete versions of the EPI (up to multiplicative constants) for Rényi entropies of log-concave distributions and which is an important tool in the one-dimensional case:

Theorem 7. [4, Theorem 1.1] *If a random variable X follows a discrete log-concave p.m.f. f on \mathbb{Z} , then*

$$\max_{k \in \mathbb{Z}} f(k) \leq \frac{1}{\sqrt{1 + 4\sigma^2}}, \quad (9)$$

where $\sigma^2 = \text{Var}(X)$.

Theorem 6 may also be seen as a discrete analogue of a dimensional upper bound on the isotropic constant L_f . The corresponding (continuous) lower bound on L_f can be deduced from (8) and the fact that $e^{-h(X)} \leq \max(f)$:

$$L_f = \left(\max_{\mathbb{R}^d} f \right)^{\frac{1}{d}} \det(\text{Cov}(f))^{\frac{1}{2d}} \geq e^{-\frac{h(X)}{d}} \det(\text{Cov}(f))^{\frac{1}{2d}} \geq \frac{1}{\sqrt{2\pi e}}.$$

It is well known that a dimensional upper bound also exists and we denote by L_d the maximum of the isotropic constants L_f among log-concave functions f in \mathbb{R}^d . The latter is related to the famous hyperplane conjecture (or slicing problem) from convex geometry, one of the central questions in the field. The hyperplane conjecture is equivalent to the isotropic constant conjecture [5, Theorem 3.1.2].

Hyperplane conjecture: there exists a universal constant $c > 0$ such that, for any dimension d and for any convex body K in isotropic position and of volume 1 in \mathbb{R}^d and any direction $\theta \in \mathbb{S}^{d-1}$, one has $|K \cap \theta^\perp|_{d-1} \geq c$.

Isotropic constant conjecture: there exists a universal constant C such that for any dimension d , one has $L_d \leq C$.

A recent result of Klartag [16] gives the best known constant depending on the dimension d for the slicing problem and thus also the best upper bound known for L_d :

$$L_d \leq C \sqrt{\log d}, \quad (10)$$

for some absolute constant C .

Our method for proving Theorem 6 is to use the corresponding continuous result. To this end, we obtain the approximations

$$\begin{aligned} \left| \int_{\mathbb{R}^d} f - \sum_{\mathbb{Z}^d} f \right| &= o_d(1), \quad \text{as } \sigma \rightarrow \infty, \\ \left| \int_{\mathbb{R}^d} xf - \sum_{k \in \mathbb{Z}^d} kf(k) \right| &= O_d(1), \quad \text{as } \sigma \rightarrow \infty, \\ \left| \det(\text{Cov}_{\mathbb{Z}^d}(f)) - \det(\text{Cov}_{\mathbb{R}^d}(f)) \right| &= O_d(\sigma^{2d-1}), \quad \text{as } \sigma \rightarrow \infty, \end{aligned}$$

for any isotropic log-concave density f . This is done in Section 3.4. Before that, after recalling some known facts about Ball's bodies in Section 3.1, we prove the exponential tails bound in Section 3.2. Although our results hold under the more general almost isotropicity assumption, for better illustration of the ideas, we assume first in Sections 3.1–3.4 that the continuous extension f is isotropic. In Section 3.5 we show how to relax this assumption (see Remark 36).

Finally, in Section 4 we conclude with a brief discussion of our assumptions as well as open questions regarding fundamental issues of log-concavity in \mathbb{Z}^d .

1.3 Notations and definitions

In this subsection, we will detail the notations as well as key definitions.

Big- and small- O notation. Let f be a real-valued function and g another strictly positive function. We write $f = O(g)$ if there exist positive absolute constants N, C such that $|f(x)| \leq Cg(x)$ for every $x \geq N$. If N, C are absolute up to a parameter d , we write $f = O_d(g)$. Analogously, we write $f = \Omega(g)$, if $|f(x)| \geq Cg(x)$ for every $x \geq N$. If $f = \Omega(g)$ and $f = O(g)$, we write $f = \Theta(g)$ (with the analogous definitions for Ω_d and Θ_d). When it is more convenient, we will write $f \lesssim_d g$ for $f = O_d(g)$ and $f \simeq_d g$ for $f = \Theta(g)$. We write $f(x) = o(g(x))$ if $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0$.

Convex bodies. A *convex body* is a convex set that is compact and has a non-empty interior. For a convex body $K \in \mathbb{R}^d$, one denotes by $|K|$ its volume. The *convex hull* of a set A , denoted by $\text{conv}(A)$, is the smallest convex set containing the set A .

Log-concavity and convexity in \mathbb{Z}^d . For $d > 1$, there are more than one definitions of log-concavity that have been used in different contexts. We will use a quite general definition that implies several other notions of log-concavity, the one of *extensible log-concave functions*. The reader is referred to Murota [21] for an extensive discussion of discrete convexity in higher dimensions. A function $f : \mathbb{Z}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ is said to be *convex-extensible* if there exists a convex function $\bar{f} : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ such that

$$\bar{f}(z) = f(z) \quad (\forall z \in \mathbb{Z}^d).$$

We refer to \bar{f} as the *extension* of f . Based on the above, we define

Definition 8 (Log-concave extensible functions). A function $f : \mathbb{Z}^d \rightarrow \mathbb{R}_+$ is said to be log-concave extensible if there exists $V : \mathbb{Z}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ convex-extensible such that

$$f(z) = e^{-V(z)}.$$

Throughout when we say that f is log-concave, we mean that f is log-concave extensible. We say that a random vector X , taking values in \mathbb{Z}^d and p.m.f. $p : \mathbb{Z}^d \rightarrow [0, 1]$ is log-concave, if p is log-concave.

Similarly, we define a \mathbb{Z}^d -convex set (commonly referred in the literature as *hole-free* set [21]):

Definition 9 (\mathbb{Z}^d -convexity). A set $A \subset \mathbb{Z}^d$ is said to be \mathbb{Z}^d -convex if and only if its indicator function χ_A , defined by $\chi_A(x) = 1$ if $x \in A$ and 0 otherwise, is log-concave-extensible or equivalently if

$$A = \text{conv}(A) \cap \mathbb{Z}^d.$$

Notice that the support of a log-concave function f on \mathbb{Z}^d is \mathbb{Z}^d -convex.

Remark 10. For $d = 1$ our definition of log-concavity is equivalent to the usual definition $p(k)^2 \geq p(k-1)p(k+1)$, $k \in \mathbb{Z}$, which is preserved under convolution (e.g. [14]). However, for $d > 1$ log-concavity may not be preserved in general, as pointed out by Murota [21, Example 3.15] by considering two log-concave distributions supported on $S_1 = \{(0, 0), (1, 1)\}$ and $S_2 = \{(0, 1), (1, 0)\}$ respectively. This is because the Minkowski sum $S_1 + S_2$ is not \mathbb{Z}^2 -convex. On the other hand, in the i.i.d. case, we shall see in our Proposition 38 below that $\sum_{i=1}^n A$ is \mathbb{Z}^d -convex for any \mathbb{Z}^d -convex set A . We do not know whether our definition of log-concavity is preserved under self-convolution.

Notion of isotropicity. The isotropic constant of a function $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is defined by

$$L_f := \left(\frac{\max_{\mathbb{R}^d} f}{\int_{\mathbb{R}^d} f} \right)^{\frac{1}{d}} \det(\text{Cov}(f))^{\frac{1}{2d}}, \quad (11)$$

where $\text{Cov}(f)$ is the *inertia* or *covariance* matrix defined, for $1 \leq i, j \leq d$, by

$$[\text{Cov}(f)]_{ij} := \frac{\int_{\mathbb{R}^d} x_i x_j f(x) dx}{\int_{\mathbb{R}^d} f(x) dx} - \frac{\int_{\mathbb{R}^d} x_i f(x) dx \int_{\mathbb{R}^d} x_j f(x) dx}{\left(\int_{\mathbb{R}^d} f(x) dx\right)^2}.$$

We say that f is *isotropic* if $\text{Cov}(f) = \sigma^2 \mathbf{I}_d$, for some $\sigma > 0$, where \mathbf{I}_d is the $d \times d$ identity matrix. Let K be a convex body in \mathbb{R}^d . Then its covariance matrix is $\text{Cov}(K) := \text{Cov}(\mathbf{1}_K)$ and its isotropic constant is $L_K := L_{\mathbf{1}_K}$. The convex body K is called *isotropic* if $\mathbf{1}_K$ is isotropic.

Similarly, in the discrete case $p : \mathbb{Z}^d \rightarrow \mathbb{R}_+$, we define the covariance matrix $\text{Cov}(p)$ by

$$[\text{Cov}(p)]_{ij} := \frac{\sum_{k \in \mathbb{Z}^d} k_i k_j p(k)}{\sum_k p(k)} - \frac{\sum_{k \in \mathbb{Z}^d} k_i p(k) \sum_{k \in \mathbb{Z}^d} k_j p(k)}{\left(\sum_{k \in \mathbb{Z}^d} p(k)\right)^2}.$$

Definition 11. A family $\{f\}_{\sigma \in \mathbb{R}_+}$ of non-negative functions on \mathbb{R}^d is *almost isotropic* if, as $\sigma \rightarrow \infty$,

$$\begin{aligned} \text{Cov}(f)_{ij} &= \sigma^2 + O(\sigma) \quad \text{for } i = j, \\ &= O(\sigma) \quad \text{for } i \neq j. \end{aligned}$$

Remark 12. It is straightforward to check that the family $\{f\}_{\sigma \in \mathbb{R}_+}$ is almost isotropic if and only if

$$\|\text{Cov}(f) - \sigma^2 \mathbf{I}_d\|_{\text{op}} = O(\sigma),$$

where $\|A\|_{\text{op}}$ is the operator norm.

We are interested in log-concave densities f , for which $\det \text{Cov}(f) \rightarrow \infty$. Thus, when we write that f is almost isotropic, it is meant that f represents a family of densities with the underlying parameter $\sigma := \det \text{Cov}(f)^{\frac{1}{2d}}$ growing large. When its not clear from the context, we write $\text{Cov}_{\mathbb{R}}(f)$ and $\text{Cov}_{\mathbb{Z}}(f)$ to distinguish between the continuous and discrete covariance matrix of f .

2 Proofs of main results

Our first observation is that the class $\mathcal{C}(c_1, c_2, c_3)$ defined in Definition 2 is closed under self-convolution up to constants, as shown by the following proposition:

Proposition 13. Suppose $p \in \mathcal{C}(c_1(d), c_2(d), c_3(d))$ for some constants $c_1(d), c_2(d) \geq 1, c_3(d)$ depending on the dimension only. Then the convolution of p with itself n times, $p_n := p * \dots * p$, satisfies

$$p_n \in \mathcal{C}(n^{\frac{d}{2}} c_1(d), n^{\frac{d}{2}} c'_2(d, n), \sqrt{n} c_3(d))$$

for some possibly different constant $c'_2(d, n)$ that depends on the dimension and n .

That is, denoting $\sigma := \det(\text{Cov}(p))^{\frac{1}{2d}}$, we have

1.

$$\max_{k \in \mathbb{Z}^d} p_n(k) \leq \frac{c_1(d)}{\sigma^d}.$$

2. For all $k \in \mathbb{Z}^d$ with $\|k\|_{\infty} \geq c'_2(d, n)\sigma$,

$$p_n(k) \leq \frac{c'_2(d, n)}{\sigma^d} e^{-\frac{\|k\|_1}{c'_2(d, n)\sigma}},$$

where in fact we may take

$$c'_2(d, n) = (3 \cdot 4^{d+1} c_2(d))^{2^n - 1}.$$

3.

$$\sum_{i=1}^d \sum_{k \in \mathbb{Z}^d} |p_n(k) - p_n(k - e_i)| \leq \frac{c_3(d)}{\sigma}.$$

Proof. First we note that, since $\text{Cov}(p_n) = n\text{Cov}(p)$, one has $\det(\text{Cov}(p_n))^{\frac{1}{2d}} = \sqrt{n} \det(\text{Cov}(p))^{\frac{1}{2d}}$, so $p_n \in \mathcal{C}(n^{\frac{d}{2}}c_1(d), n^{\frac{d}{2}}c'_2(d, n), \sqrt{nc_3}(d))$ is indeed equivalent to the three statements.

For 1 we have

$$\max_k p_n(k) = \max_{j \in \mathbb{Z}^d} p_{n-1}(j)p(k-j) \leq \max_k p(k) \sum_{j \in \mathbb{Z}^d} p_{n-1}(j) = \max_k p(k) \leq \frac{c_1(d)}{\sigma^d}.$$

To prove that property 2 is satisfied, we proceed by induction. Assume that p , as well as p_n , satisfy condition 2 of Definition 2 with the constant $c'_2(d, n)$. Let $k \in \mathbb{Z}$ such that $\|k\|_\infty \geq c'_2(d, n+1)\sigma$. We have

$$\begin{aligned} p_{n+1}(k) &= \sum_j p(j)p_n(k-j) \\ &= \sum_{\|j\|_\infty \leq c_2(d)} p(j)p_n(k-j) + \sum_{\substack{\|j\|_\infty < c_2(d) \\ \|k-j\|_\infty < c'_2(d, n)}} p(j)p_n(k-j) + \sum_{\substack{\|j\|_\infty < c_2(d) \\ \|k-j\|_\infty \leq c'_2(d, n)}} p(j)p_n(k-j). \end{aligned} \quad (12)$$

For the first sum in (12), we note that for $\|j\|_\infty \leq c_2(d)\sigma$, $\|k-j\|_1 \geq \|k\|_1 - \|j\|_1 \geq \|k\|_1 - \sqrt{d}\|j\|_\infty \geq \|k\|_1 - \sqrt{d}c_2(d)\sigma \geq \|k\|_1 - c'_2(d, n)\sigma$ and $\|k-j\|_\infty \geq \|k\|_\infty - \|j\|_\infty \geq c'_2(d, n+1)\sigma - c_2(d)\sigma \geq c'_2(d, n)\sigma$, by the elementary bound $\|x\|_\infty \leq \|x\|_1 \leq \sqrt{d}\|x\|_\infty, x \in \mathbb{R}^d$. Therefore we can bound that sum as

$$\sum_{\|j\|_\infty \leq c_2(d)} p(j)p_n(k-j) \leq \frac{c'_2(d, n)}{\sigma^d} \sum_{\|j\|_\infty \leq c_2(d)} p(j)e^{-\frac{\|k-j\|_1}{c'_2(d, n)\sigma}} \leq \frac{c'_2(d, n)}{\sigma^d} e^{-\frac{\|k\|_1}{c'_2(d, n)\sigma}}. \quad (13)$$

For the second sum in (12), using $c_2(d) \leq c'_2(d, n)$, we have

$$\begin{aligned} &\sum_{\|j\|_\infty < c_2(d)} \sum_{\|k-j\|_\infty < c'_2(d, n)} p(j)p_n(k-j) \\ &\leq \frac{c_2(d)c'_2(d, n)}{\sigma^{2d}} \sum_{\substack{\|j\|_\infty < c_2(d) \\ \|k-j\|_\infty < c'_2(d, n)}} e^{-\frac{-\|j\|_1 - \|k-j\|_1}{c'_2(d, n)\sigma}} \\ &\leq \frac{c_2(d)c'_2(d, n)}{\sigma^{2d}} \sum_{j \in \mathbb{Z}^d} e^{-\frac{-\|j\|_1 - \|k\|_1 - \|j\|_1}{c'_2(d, n)\sigma}} \\ &\leq \frac{c_2(d)c'_2(d, n)}{\sigma^{2d}} \left[\sum_{0 \leq m \leq \|k\|_1} \sum_{\|j\|_1 = m} e^{-\frac{-m + \|k\|_1 + m}{c'_2(d, n)\sigma}} + \sum_{m \geq \|k\|_1} \sum_{\|j\|_1 = m} e^{-\frac{-2m + \|k\|_1}{c'_2(d, n)\sigma}} \right]. \end{aligned} \quad (14)$$

The first double sum in (14) is

$$\begin{aligned} \sum_{0 \leq m \leq \|k\|_1} \sum_{\|j\|_1 = m} e^{-\frac{-m + \|k\|_1 + m}{c'_2(d, n)\sigma}} &\leq e^{-\frac{\|k\|_1}{c'_2(d, n)\sigma}} |\{j : \|j\|_1 \leq \|k\|_1\}| \\ &\leq e^{-\frac{\|k\|_1}{c'_2(d, n)\sigma}} |\{j : \|j\|_\infty \leq \|k\|_1\}| \\ &\leq e^{-\frac{\|k\|_1}{c'_2(d, n)\sigma}} (2\|k\|_1 + 1)^d \leq 4^d \|k\|_1^d e^{-\frac{\|k\|_1}{c'_2(d, n)\sigma}} \leq 4^d \sigma^d e^{-\frac{\|k\|_1}{2c'_2(d, n)\sigma}}, \end{aligned} \quad (15)$$

where the last inequality holds at least as long as

$$\|k\|_1 \geq 2c'_2(d, n)d\sigma \left(\log(2c'_2(d, n)d) \right)^2, \quad (16)$$

since $x^d \leq e^{\frac{x}{2c}}$, for $x \geq 2cd(\log(2cd))^2, d \geq 1, c \geq 2$. On the other hand, the second double sum in (14) is

$$\sum_{m \geq \|k\|_1} \sum_{\|j\|_1 = m} e^{-\frac{-2m + \|k\|_1}{c'_2(d, n)\sigma}} \leq e^{\frac{\|k\|_1}{c'_2(d, n)\sigma}} \sum_{m \geq \|k\|_1} |\{j : \|j\|_\infty \leq m\}| e^{-\frac{2m}{\sigma c'_2(d, n)}} \quad (17)$$

$$\leq 4^d e^{\frac{\|k\|_1}{c_2'(d,n)\sigma}} \sum_{m \geq \|k\|_1} m^d e^{-\frac{2m}{\sigma c_2'(d,n)}} \quad (18)$$

$$\leq 4^d \sigma^d e^{\frac{\|k\|_1}{c_2'(d,n)\sigma}} \sum_{m \geq \|k\|_1} e^{-\frac{3m}{2\sigma c_2'(d,n)}} \quad (19)$$

$$= 4^d \sigma^d e^{\frac{\|k\|_1}{c_2'(d,n)\sigma}} \frac{e^{-\frac{3\|k\|_1}{2\sigma c_2'(d,n)}}}{1 - e^{-\frac{3\|k\|_1}{2\sigma c_2'(d,n)}}} \leq 2 \cdot 4^d \sigma^d e^{-\frac{\|k\|_1}{2\sigma c_2'(d,n)}}, \quad (20)$$

where (19) holds at least as long as (16) is satisfied since $m \geq \|k\|_1$ and similarly for the last inequality. Plugging the bounds (15), (20) in (14) we obtain

$$\sum_{\|j\|_\infty < c_2(d) \quad \|k-j\|_\infty < c_2'(d,n)} p(j)p_n(k-j) \leq 4^{d+1} \frac{c_2(d)c_2'(d,n)^2}{\sigma^d} e^{-\frac{\|k\|_1}{2\sigma c_2'(d,n)}}, \quad (21)$$

provided that (16) holds.

For the third sum in (12), we observe that $\|k-j\|_\infty \leq c_2'(d,n)\sigma$ implies $\|j\|_1 \geq \|k\|_1 - \|k-j\|_1 \geq \|k\|_1 - c_2'(d,n)\sqrt{d}\sigma$, and therefore

$$\begin{aligned} \sum_{\|j\|_\infty < c_2(d) \quad \|k-j\|_\infty \leq c_2'(d,n)} p(j)p_n(k-j) &\leq \sum_{\|j\|_\infty < c_2(d) \quad \|k-j\|_\infty \leq c_2'(d,n)} \frac{c_2(d)}{\sigma^d} e^{-\frac{\|j\|_1}{c_2'(d,n)\sigma}} p_n(k-j) \\ &\leq \frac{c_2'(d,n)}{\sigma^d} e^{\frac{-\|k\|_1}{c_2'(d,n)\sigma} + \sqrt{d}} \sum_{j \in \mathbb{Z}} p(k-j) \\ &\leq \frac{e^{\sqrt{d}} c_2'(d,n)}{\sigma^d} e^{\frac{-\|k\|_1}{c_2'(d,n)\sigma}}. \end{aligned} \quad (22)$$

Clearly, the worst bound out of (13), (21) and (22) is (21) and the dominating condition for k is (16). By (12), we can therefore bound

$$p_{n+1}(k) \leq 3 \cdot 4^{d+1} \frac{c_2(d)c_2'(d,n)^2}{\sigma^d} e^{-\frac{\|k\|_1}{2\sigma c_2'(d,n)}} \leq \frac{c_2'(d,n+1)}{\sigma^d} e^{-\frac{\|k\|_1}{\sigma c_2'(d,n+1)}}, \quad (23)$$

with $c_2'(d,n+1) = 3 \cdot 4^{d+1} c_2(d)c_2'(d,n)^2 = (3 \cdot 4^{d+1} c_2(d))^{2^{n-1}}$ by the inductive hypothesis. Note that this choice of $c_2'(d,n+1)$ also ensures that (16) is satisfied and therefore the proof of 2. is complete.

For property 3, let X be a random vector with p.m.f. $p_X = p$. Then $S_n := \sum_{i=1}^n X_i$, where X_i are i.i.d. copies of X , has p.m.f. p_n . We observe that

$$\sum_{i=1}^d \sum_{k \in \mathbb{Z}^d} |p_n(k) - p_n(k - e_i)| = \sum_{i=1}^d \|p_{S_n} - p_{S_n + e_i}\|_{\text{TV}}.$$

For any probability measures μ_1, μ_2, ν_1 and ν_2 , the total variation distance satisfies the convolution inequality

$$\|\mu_1 * \mu_2 - \nu_1 * \nu_2\|_{\text{TV}} \leq \|\mu_1 - \nu_1\|_{\text{TV}} + \|\mu_2 - \nu_2\|_{\text{TV}}.$$

This can be proved either directly or by the fact that total variation is an f -divergence and therefore satisfies a data processing inequality. Choosing $\mu_1 = \nu_1 = p_{S_{n-1}}$, $\mu_2 = p_X$ and $\nu_2 = p_{X+e_i}$, we get

$$\|p_{S_n} - p_{S_n + e_i}\|_{\text{TV}} \leq \|p_X - p_{X+e_i}\|_{\text{TV}}$$

Using the assumption on X , the last inequality yields

$$\sum_{i=1}^d \|p_{S_n} - p_{S_n + e_i}\|_{\text{TV}} \leq \sum_{i=1}^d \|p_X - p_{X+e_i}\|_{\text{TV}} \leq \frac{c_3(d)}{\sigma}.$$

□

Lemma 14. *Let X be a random vector with p.m.f. p on \mathbb{Z}^d satisfying conditions 2, 3 of Definition 2. Let $U^{(n)} = \sum_{i=1}^n U_i$ where U_i are i.i.d. uniform random vectors on the cube $[0, 1]^d$. Denote by $h_n^{(d)}(x) = \prod_{i=1}^d h_n(x_i)$ the density of U_n and by $f_n^{(d)}$ the density of $U^{(n)} + X$. Then,*

$$1. \quad \forall z \in [0, 1]^d, \forall n \geq 1, \sum_{j \in \{0, \dots, n-1\}^d} h_n^{(d)}(j+z) = 1.$$

2. For $x = k + z$ where $z \in [0, 1]^d$ and $k \in \mathbb{Z}^d$, $|f_n^{(d)}(x) - p(k)| \leq \frac{2^{d-1} c_3}{d}$ with c_3 the constant that appears in condition 3 of Definition 2.

3. For $x = k + z$ where $z \in [0, 1]^d$ and $k \in \mathbb{Z}^d$, with $\|k\|_\infty \geq c_2 \sigma + n$,

$$f_n^{(d)}(x) \leq \left(n^d (e^{\frac{n d}{c_2 \sigma}} + 1) + 1 \right) \frac{c_2}{\sigma^d} e^{-\frac{\|k\|_1}{c_2 \sigma}}.$$

Proof. First, we prove 1. Let us prove that the one-dimensional case holds i.e. $\sum_{j=0}^{n-1} h_n(j+z) = 1$, for any $z \in [0, 1]$. For this, one does an induction on n . For $n = 1$, since $S_1 = U_1$, $h_1(z) = 1_{[0,1]}$ so the base case is immediate. Now let us suppose by induction that the claim holds for n . Noticing that the support of h_n is $[0, n]$ and using the induction hypothesis, we get

$$\begin{aligned} \sum_{j=0}^n h_{n+1}(j+z) &= \sum_{j=0}^n \int_0^1 h_n(j+z-u) du = \sum_{j=0}^n \left(\int_0^z h_n(j+z-u) du + \int_z^1 h_n(j+z-u) du \right) \\ &= z + \int_0^z h_n(n+z-u) du + \int_z^1 \left(\sum_{k=0}^{n-1} h_n(k+1+z-u) \right) du = z + 1 - z = 1. \end{aligned}$$

For the d -dimensional case, observe that since $h_n^{(d)}$ is a product, for all $z \in [0, 1]^d$,

$$\sum_{j \in \{0, \dots, n-1\}^d} h_n^{(d)}(j+z) = \sum_{j \in \{0, \dots, n-1\}^d} \left(\prod_{i=1}^d h_n(z_i + j_i) \right) = \prod_{i=1}^d \left(\sum_{j_i=0}^{n-1} h_n(z_i + j_i) \right) = 1,$$

where, for the last equality, we have used the above one-dimensional argument.

Let us prove the second item. Using item 1, one has

$$f_n^{(d)}(x) - p(k) = f_n^{(d)}(k+z) - p(k) = \sum_{i \in \mathbb{Z}^d} p(k-i) h_n^{(d)}(i+z) - p(k) = \sum_{i \in \mathbb{Z}^d} u(i) v(i), \quad (24)$$

where $u(i) = p(k-i) - p(k)$, $v(i) = h_n^{(d)}(z+i)$. Note that the sum actually ranges over $i \in \{0, \dots, n-1\}^d$, since $h_n^{(d)}(i+z) = 0$ outside this discrete cube. Also, for $i = (i_1, \dots, i_d) \in \mathbb{Z}^d$, denote

$$W(i) := (T(v))_i = \sum_{j \geq i} v_j,$$

where $j \geq i$ means that $j_l \geq i_l$ for all $1 \leq l \leq d$. By a straightforward induction,

$$v(i) = (T^{-1}(W))_i = \sum_{\epsilon \in \{-1, 1\}^d} \left(\prod_{j=1}^d (-\epsilon_j) \right) W \left(i + \sum_{k=1}^d \frac{\epsilon_k + 1}{2} e_k \right).$$

Let us introduce the standard dot product in $l^2(\mathbb{Z})$ as $\langle z, h \rangle_{l^2(\mathbb{Z})} := \sum_{i \in \mathbb{Z}^d} z(i) h(i)$ for any $z, h \in \mathbb{R}^{\mathbb{Z}^d}$. Then the sum $\sum_{i \in \mathbb{Z}^d} u(i) v(i)$ can be rewritten as $\langle u, T^{-1}W \rangle_{l^2(\mathbb{Z})}$ with

$$T^{-1} := \sum_{\epsilon \in \{-1, 1\}^d} \left(\prod_{j=1}^d (-\epsilon_j) \right) S,$$

where S is the left-shift operator, that is, $(S W)_i = W \left(i + \sum_{k=1}^d \frac{\epsilon_k + 1}{2} e_k \right)$. It is well known that the adjoint S° of S , is the right-shift operator or in another words $(S^\circ(W))_i = W \left(i - \sum_{k=1}^d \frac{\epsilon_k + 1}{2} e_k \right)$. Thus

$$\begin{aligned} \sum_{i \in \mathbb{Z}^d} u(i) v(i) &= \sum_{i \in \mathbb{Z}^d} \sum_{\epsilon \in \{-1, 1\}^d} u_i \left(\prod_{j=1}^d (-\epsilon_j) \right) W \left(i + \sum_{k=1}^d \frac{\epsilon_k + 1}{2} e_k \right) \\ &= \langle u, T^{-1}(W) \rangle_{l^2(\mathbb{Z})} = \langle (T^{-1})^\circ u, W \rangle_{l^2(\mathbb{Z})} \\ &= \sum_{i \in \mathbb{Z}^d} W(i) \sum_{\epsilon \in \{-1, 1\}^d} \left(\prod_{j=1}^d (-\epsilon_j) \right) u \left(i - \sum_{k=1}^d \frac{\epsilon_k + 1}{2} e_k \right) \end{aligned}$$

Thus, by the triangular inequality, summing up to the last coordinate which can be either 1 or -1 , we obtain

$$\begin{aligned} \left| \sum_{i \in \mathbb{Z}^d} u(i)v(i) \right| &\leq \sum_{i \in \mathbb{Z}^d} \sum_{\epsilon \in \{-1, 1\}^{d-1}} \left| u \left(i - \sum_{k=1}^d \frac{\epsilon_k + 1}{2} e_k \right) - u \left(i - \sum_{k=1}^{d-1} \frac{\epsilon_k + 1}{2} e_k - e_d \right) \right| \\ &\leq 2^{d-1} \sum_{i \in \mathbb{Z}^d} |u(i) - u(i - e_d)|. \end{aligned}$$

Now, using the same argument for $j \in \{1, \dots, d\}$, instead of $j = d$ and summing over j one gets

$$|f_n(x) - p(k)| = \left| \sum_{i \in \mathbb{Z}^d} u(i)v(i) \right| \leq \frac{2^{d-1}}{d} \sum_{i \in \mathbb{Z}^d} \sum_{j=1}^d |p(i) - p(i + e_j)| \leq \frac{2^{d-1}}{d} \frac{c_3}{\sigma},$$

where, in the last inequality, we have used that X satisfies condition 3 of Definition 2.

Finally, for item 3, we have by (24), since $|h_n| \leq 1$,

$$\begin{aligned} |f_n^{(d)}(x) - p(k)| &= \left| \sum_{i \in \{0, \dots, n-1\}^d} (p(k-i) - p(k))h_n(z+i) \right| \\ &\leq \sum_{i \in \{0, \dots, n-1\}^d} p(k-i) + n^d p(k). \end{aligned}$$

But, using hypothesis 2, the assumption $\|k\|_\infty \geq c_2\sigma + n$ and the observations that $\|k-i\|_\infty \geq \|k\|_\infty - n \geq c_2\sigma$ and $\|k-i\|_1 \geq \|k\|_1 - nd$, we have

$$|f_n^{(d)}(x) - p(k)| \leq n^d \frac{c_2}{\sigma^d} e^{-\frac{\|k\|_1 - nd}{c_2\sigma}} + n^d \frac{c_2}{\sigma^d} e^{-\frac{\|k\|_1}{c_2\sigma}}$$

and the claimed bound follows since $f_n^{(d)}(x) \leq |f_n^{(d)}(x) - p(k)| + p(k)$. \square

Remark 15. The upper bound of Lemma 14, item 2 depends on d , but not on n . Hence, for $d = 1$ our new method of proof improves the bound [8, Eq. (21)] by removing the dependence on n , i.e. the number of uniforms being added to the discrete random variable.

We are now ready to prove Theorem 4. Theorem 3 will readily follow from this, together with the generalized EPI in \mathbb{R}^d .

Proof of Theorem 4. Let $F(x) = x \log \frac{1}{x}$, $x > 0$ and note that $F(x)$ is increasing for $x \leq 1/e$. Denote $S_n = \sum_{i=1}^n X_i$, $U^{(n)} = \sum_{i=1}^n U_i$, and let $f_{S_n+U^{(n)}}$ be the density of $S_n + U^{(n)}$ on the \mathbb{R}^d and p_{S_n} the p.m.f. of S_n . Note that p_{S_n} satisfies the three parts of Proposition 13. We have

$$\begin{aligned} &h(X_1 + \dots + X_n + U_1 + \dots + U_n) \\ &= \sum_{k: \|k\|_\infty \leq 2} \int_{k+[0, 1)^d} F(f_{S_n+U^{(n)}}(x)) dx + \sum_{k: \|k\|_\infty < 2} \int_{k+[0, 1)^d} F(f_{S_n+U^{(n)}}(x)) dx. \end{aligned} \quad (25)$$

First, we bound the second term. From part 3 of Lemma 14, in that range of k and for any $x \in k+[0, 1)^d$,

$$f_{S_n+U^{(n)}}(x) \leq \frac{C'_d n}{\sigma^d} e^{-\frac{\|k\|_1}{c'_2(d,n)\sigma}} \quad (26)$$

for some constants $C'_d n, c'_2(d,n) > 0$ and σ large enough. Therefore, for σ large enough $f_{S_n+U^{(n)}} \leq 1/e$ so that F is increasing. Moreover, using (26) and that $\|x\|_\infty \leq \|x\|_1 \leq \sqrt{d}\|x\|_\infty$, $x \in \mathbb{R}^d$, we have, for σ large enough,

$$\sum_{k: \|k\|_\infty < 2} \int_{k+[0, 1)^d} F(f_{S_n+U^{(n)}}(x)) dx \lesssim d n \sum_{k: \|k\|_\infty < 2} \int_{k+[0, 1)^d} F\left(\frac{C'_d n}{\sigma^d} e^{-\frac{\|k\|_1}{c'_2(d,n)\sigma}}\right) dx \quad (27)$$

$$\begin{aligned} &\lesssim d n \sum_{k: \|k\|_\infty < 2} \frac{1}{\sigma^d} e^{-\frac{\|k\|_\infty}{c'_2(d,n)\sigma}} \frac{\|k\|_\infty}{\sigma} \\ &+ \sum_{k: \|k\|_\infty < 2} \frac{1}{\sigma^d} e^{-\frac{\|k\|_\infty}{c'_2(d,n)\sigma}} \log \sigma. \end{aligned} \quad (28)$$

Observing that the first term in (28) dominates and using a similar calculation as the one leading to (20), we obtain

$$\begin{aligned} & \sum_{k: \|k\|_\infty < 2} \int_{k+[0,1]^d} F(f_{S_n+U^{(n)}}(x)) dx \\ & \lesssim_{d,n} \frac{1}{\sigma^{d+1}} \sum_{k: \|k\|_\infty < 2} e^{-\frac{\|k\|_\infty}{c_2'(d,n)\sigma}} \|k\|_\infty \end{aligned} \quad (29)$$

$$\lesssim_{d,n} \frac{1}{\sigma^{d+1}} \sum_{m < 2} m^{d+1} e^{-\frac{m}{c_2'(d,n)\sigma}} \quad (30)$$

$$\lesssim_{d,n} \sum_{m < 2} e^{-\frac{m}{2c_2'(d,n)\sigma}} \lesssim_{d,n} \frac{e^{-\frac{\sigma}{2c_2'(d,n)}}}{1 - e^{-\frac{1}{2c_2'(d,n)\sigma}}} \lesssim_{d,n} \sigma e^{-\frac{\sigma}{2c_2'(d,n)}} = O(\sigma e^{-\Theta(\cdot)}). \quad (31)$$

Now we show that the first term in (25) is approximately $H(S_n)$. To that end we apply the estimate (38) of Lemma 17 to the integrand with $\mu = \Theta_{d,n}(\frac{1}{\sigma})$, $D = \Theta_{d,n}(\sigma^{2d-1})$, $M = \sigma^{2d}$, $a = f_{S_n+U^{(n)}}(x)$ and $b = p_{S_n}(k)$. The assumption of the lemma is satisfied for an appropriate choice of the implied constants since by Lemma 14 and part 1 of Proposition 13,

$$a, b \lesssim_{d,n} \max_{k \in \mathbb{Z}^d} p_{S_n}(k) + O_{d,n}\left(\frac{1}{\sigma}\right) \lesssim_{d,n} \frac{1}{\sigma} \simeq_{d,n} \frac{D}{M}.$$

Thus, since there are at most $\Theta(\sigma^{2d})$ elements in the set $\{k \in \mathbb{Z}^d : \|k\|_\infty \leq \sigma^2\}$,

$$\begin{aligned} & \sum_{k: \|k\|_\infty \leq 2} \left| \int_{k+[0,1]^d} F(f_{S_n+U^{(n)}}(x)) dx - F(p_{S_n}(k)) \right| \\ & \lesssim_{d,n} \sigma^{2d} \frac{\log \sigma}{\sigma^{2d+1}} + \log \sigma \sum_{k \in \mathbb{Z}^d} \int_{k+[0,1]^d} |f_{S_n+U^{(n)}}(x) - p_{S_n}(k)| dx \end{aligned} \quad (32)$$

$$\lesssim_{d,n} \frac{\log \sigma}{\sigma} + \log \sigma \sum_{k \in \mathbb{Z}^d} \sup_{x \in k+[0,1]^d} |g_n(k, x)| \quad (33)$$

$$\lesssim_{d,n} \frac{\log \sigma}{\sigma}, \quad (34)$$

where $g_n(k, x)$ is given by Lemma 14 applied to the random variable S_n , which satisfies the assumption $p_{S_n} \in \mathcal{C}(c_1'(d, n), c_2'(d, n), c_3'(d, n))$ for some constants that depend on d, n , and therefore

$$\sum_k \sup_{x \in k+[0,1]^d} |g_n(k, x)| \lesssim \det(\text{Cov}(S_n))^{-\frac{1}{2d}} \simeq_{d,n} \frac{1}{\sigma}.$$

Therefore, by (34),

$$\begin{aligned} \left| H(S_n) - \sum_{k: \|k\|_\infty \leq 2} \int_{k+[0,1]^d} F(f_{S_n+U^{(n)}}(x)) dx \right| & \lesssim_{d,n} \left| H(S_n) - \sum_{k: \|k\|_\infty \leq 2} F(p_{S_n}(k)) \right| + \frac{\log \sigma}{\sigma} \\ & \lesssim_{d,n} \sum_{k: \|k\|_\infty < 2} F(p_{S_n}(k)) + \frac{\log \sigma}{\sigma}. \end{aligned} \quad (35)$$

But, in view of (26) and (27), we can bound the discrete tails in the same way:

$$\sum_{k: \|k\|_\infty < 2} F(p_{S_n}(k)) \lesssim_{d,n} \frac{\sigma}{e^{\Theta_{d,n}(\cdot)}}. \quad (36)$$

Plugging the bound (36) into (35), we get

$$\begin{aligned} & \left| H(S_n) - \sum_{k: \|k\|_\infty \leq 2} \int_{k+[0,1]^d} F(f_{S_n+U^{(n)}}(x)) dx \right| \\ & \leq O_{d,n}\left(\frac{\sigma}{e^{\Theta_{d,n}(\cdot)}}\right) + O_{d,n}\left(\frac{\log \sigma}{\sigma}\right) \\ & = O_{d,n}\left(\frac{\log \sigma}{\sigma}\right). \end{aligned}$$

Finally, by (25) and the exponential bounds on the entropy tails (34), (36), we obtain

$$\begin{aligned} |h(S_n + U^{(n)}) - H(S_n)| &= O_{d,n} \left(\frac{\log \sigma}{\sigma} \right) + O_{d,n} \left(\frac{\sigma}{e^{\Theta_{d,n}(\cdot)}} \right) \\ &= O_{d,n} \left(\frac{\log \sigma}{\sigma} \right), \end{aligned} \quad (37)$$

completing the proof. \square

Proof of Theorem 3. By the continuous EPI (1),

$$h \left(\sum_{i=1}^{n+1} X_i + U_i \right) \geq h \left(\sum_{i=1}^n X_i + U_i \right) + \frac{d}{2} \log \left(\frac{n+1}{n} \right)$$

and by Theorem 4 applied to the differential entropies on both sides

$$H \left(\sum_{i=1}^{n+1} X_i \right) + O_{d,n} \left(\frac{\log \sigma}{\sigma} \right) \geq H \left(\sum_{i=1}^n X_i \right) + O_{d,n} \left(\frac{\log \sigma}{\sigma} \right) + \frac{d}{2} \log \left(\frac{n+1}{n} \right).$$

But by (7), $O_{d,n} \left(\frac{\log \sigma}{\sigma} \right) = O_{d,n} \left(H(X_1) e^{-\frac{1}{d} H(X_1)} \right)$ and the result follows. \square

Remark 16. *The rate can be improved to superpolynomial by splitting the sum according to σ^{1+} and optimising over ϵ .*

Lemma 17 ([8]). *Let $D, M \geq 1$ and, for $x > 0$, consider $G(x) = F(x) - x \log M$, where $F(x) = -x \log x$. Then, for $0 \leq a, b \leq \frac{D}{M}$ and any $0 < \mu < \frac{1}{e}$, we have the estimate*

$$|G(b) - G(a)| \leq \frac{2\mu}{M} \log \frac{1}{\mu} + |b - a| \log \frac{eD}{\mu},$$

whence

$$|F(b) - F(a)| \leq \frac{2\mu}{M} \log \frac{1}{\mu} + |b - a| \log \frac{eDM}{\mu}. \quad (38)$$

Proof. See [8]. \square

3 Discrete isotropic log-concave random vectors

This section is devoted to the proof of the following result, which shows that discrete log-concave extensible random vectors with isotropic extension satisfy the properties of Definition 2. In Section 3.5 we show that this can be generalized to discrete log-concave extensible random vectors with almost isotropic extension.

Theorem 18. *Suppose p is a centered log-concave p.m.f. on \mathbb{Z}^d with isotropic extension and covariance matrix $\text{Cov}(p) = \sigma^2 \text{I}_d$. Then there are constants $c_1(d), c_2(d), c_3(d)$ such that $p \in \mathcal{C}(c_1(d), c_2(d), c_3(d))$, provided that $\sigma = \det(\text{Cov}_{\mathbb{Z}^d}(p))^{\frac{1}{2d}}$ is large enough depending on d , where $\mathcal{C}(c_1(d), c_2(d), c_3(d))$ is defined in Definition 2. That is, if σ is large enough depending on d*

1.

$$\max_{k \in \mathbb{Z}^d} p(k) \leq \frac{c_1(d)}{\sigma^d}.$$

2. For every k with $\|k\|_\infty \geq c_2(d)\sigma$

$$p(k) \leq \frac{C_2(d)}{\sigma^d} e^{-\frac{\|k\|_1}{c_2(d)\sigma}}.$$

3.

$$\sum_{i=1}^d \sum_{k \in \mathbb{Z}^d} |p(k) - p(k - e_i)| \leq \frac{c_3(d)}{\sigma}.$$

Proof. Part 1 follows directly from Corollary 32. Part 2 follows from Lemma 22, the well-known fact that for a centered log-concave function f on \mathbb{R}^d

$$f(0) \simeq_d \frac{1}{\det(\text{Cov}(f))^{\frac{1}{2}}}$$

and Corollary 31. Finally, Part 3 follows directly from Proposition 33. \square

3.1 Ball's bodies

In what follows, we set ω_d to be the volume of the unit Euclidean ball denoted by B_2^d . For a given convex body $K \in \mathbb{R}^d$ with 0 in its interior, its *inradius* $r(K)$ is defined as the largest $r > 0$ for which $rB_2^d \subset K$ and its *circumradius* $R(K)$ is defined as the smallest $R > 0$ such that $K \subset RB_2^d$. The support function of K is defined as $h_K(x) = \max_{y \in K} \langle x, y \rangle$ and the radial function of K is $\rho_K(x) = \inf\{t > 0; tx \in K\} = 1/\|x\|_K$, where $\|x\|_K$ denotes the gauge associated to K .

For most results in this section we will assume that $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a centered, isotropic, log-concave density, that is $\int_{\mathbb{R}^d} f = 1$, $\int_{\mathbb{R}^d} xf = 0$ and $\int_{\mathbb{R}^d} x^T x f = \sigma^2 \mathbf{I}_d$. To this function f , we attach its Ball's bodies. This important family of bodies was introduced by Ball [2]. We refer to the book [5] for the properties of these.

Definition 19. Let $f : \mathbb{R}^d \rightarrow [0, \infty)$ be an integrable, centered log-concave function. For any $p > 0$, the set $K_p(f)$ is defined as follows

$$K_p(f) := \left\{ x \in \mathbb{R}^d : \int_0^\infty pr^{p-1} f(rx) dr \geq f(0) \right\}.$$

Ball [2] established that the set $K_p(f)$ is a convex body. Moreover, its radial function is

$$\rho_{K_p(f)}(x) = \left(\frac{1}{f(0)} \int_0^\infty pr^{p-1} f(rx) dr \right)^{\frac{1}{p}} \text{ for } x \neq 0.$$

From integration in polar coordinates (see [5, Proposition 2.5.3]), it follows that $K_{d+1}(f)$ is centered and, for any $p \geq 0$ and $u \in \mathbb{S}^{d-1}$,

$$\int_{K_{d+p}(f)} |\langle x, u \rangle|^p dx = \frac{1}{f(0)} \int_{\mathbb{R}^d} |\langle x, u \rangle|^p f(x) dx. \quad (39)$$

Finally, the following inclusion relations between $K_p(f)$ and $K_q(f)$ hold [5, Proposition 2.5.7], for $p < q$,

$$\frac{\Gamma(p+1)^{\frac{1}{p}}}{\Gamma(q+1)^{\frac{1}{q}}} K_q(f) \subset K_p(f) \subset e^{\frac{d}{p} - \frac{d}{q}} K_q(f).$$

Applying these inclusions first for $p = d$ and $q = d+1$ then for $p = d+1$ and $q = d+2$ and using classical inequalities on Gamma functions, it follows that there exist universal constants $0 < c_1 < c_2$ such that for any dimension $d \geq 1$, one has

$$c_1 K_{d+1}(f) \subset K_d(f) \subset c_2 K_{d+1}(f) \quad \text{and} \quad c_1 K_{d+2}(f) \subset K_{d+1}(f) \subset c_2 K_{d+2}(f). \quad (40)$$

We shall apply to $K_{d+1}(f)$ the following theorem due to Kannan, Lovász and Simonovits [15, Theorem 4.1] to which we give a new and simple proof.

Theorem 20. Let K be a centered convex body in \mathbb{R}^d and $u \in \mathbb{S}^{d-1}$. Then

$$\frac{h_K(u)^2}{d(d+2)} \leq \frac{1}{|K|} \int_K \langle x, u \rangle^2 dx = \langle \text{Cov}(K)u, u \rangle \leq \frac{d}{d+2} h_K(u)^2. \quad (41)$$

Proof. Let $f(t) = |\{x \in K; \langle x, u \rangle = t\}|$. Denote by $[-b, a]$ its support. Then one has $a = h_K(u)$,

$$\int_{\mathbb{R}} f(t) dt = |K|, \quad \int_{\mathbb{R}} tf(t) dt = \int_K \langle x, u \rangle dx \quad \text{and} \quad \int_{\mathbb{R}} t^2 f(t) dt = \int_K \langle x, u \rangle^2 dx.$$

Now we define $g : \mathbb{R} \rightarrow \mathbb{R}_+$ by

$$g(t) = \alpha \left(1 + \frac{t}{da} \right)^{d-1} \mathbf{1}_{[-da, a]}(t), \quad \text{where } \alpha = \frac{1}{a \left(1 + \frac{1}{d} \right)^d} \int_{\mathbb{R}} f(t) dt.$$

And let $h : \mathbb{R} \rightarrow \mathbb{R}_+$ be defined by $h(t) = dh(-dt)$. Then, it is not difficult to see that

$$\int_{\mathbb{R}} g(t) dt = \int_{\mathbb{R}} h(t) dt = \int_{\mathbb{R}} f(t) dt \quad \text{and} \quad \int_{\mathbb{R}} tg(t) dt = \int_{\mathbb{R}} th(t) dt = \int_{\mathbb{R}} tf(t) dt = 0. \quad (42)$$

We shall prove that

$$\int_{\mathbb{R}} t^2 g(t) dt \leq \int_{\mathbb{R}} t^2 f(t) dt \leq \int_{\mathbb{R}} t^2 h(t) dt,$$

from which the inequality (41) follows immediately by a simple calculation. If $f \neq g$, it follows from (42) that there exist at least two points $t_1(g) < t_2(g)$ such that $g - f$ changes sign at these points and, in the same way, if $f \neq h$ there exists at least two points $t_1(h) < t_2(h)$ such that $f - h$ changes sign at these points. Since the function $f^{1/(d-1)}$ is concave on its support $[-b, a]$ and the functions $g^{1/(d-1)}$ and $h^{1/(d-1)}$ are affine on their respective supports $[-da, a]$ and $[-\frac{a}{d}, a]$, it follows that the functions $g - f$ and $f - h$ have exactly two sign changes and that they are non positive on $[t_1, t_2]$ and non negative outside $[t_1, t_2]$. Let us define v to be either $g - f$ or $f - h$. Then, for any $t \in \mathbb{R}$, one has $(t - t_1)(t - t_2)v(t) \geq 0$. Integrating this inequality, we deduce that $\int t^2 v(t) dt \geq 0$, which is the result. \square

We shall deduce from the preceding theorem the following important technical lemma. We note that Lemma 21 (and its consequences) is the only part, where the isotropicity assumption is used.

Lemma 21. *Let $d \geq 1$ be an integer. There exist two constants $0 < C'_d < C_d$ such that for any $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ centered, isotropic, log-concave density and for every $\theta \in \mathbb{S}^{d-1}$,*

$$C'_d \leq \left(\int_0^\infty dr^{d-1} f(r\theta) dr \right)^{\frac{1}{d}} \leq C_d, \quad (43)$$

where C_d and C'_d are constants depending only on the dimension d . In fact, we may take

$$C'_d = \frac{c_1^{d+2}}{\sqrt{2\pi}e^{\frac{3}{2}}} \quad \text{and} \quad C_d = (d+1)c_2^{d+2}L_d, \quad (44)$$

where c_1 and c_2 are the absolute constants appearing in (40).

Proof. The function f being isotropic, we have $\text{Cov}(f) = \sigma^2 I_d$, for some $\sigma > 0$ and $\int f = 1$, thus $L_f = \max(f)^{\frac{1}{d}}\sigma$. Note that proving (43) is equivalent to proving that

$$C'_d B_2^d \subset f(0)^{\frac{1}{d}} K_d(f) \subset C_d B_2^d.$$

Applying Theorem 20 to the centered body $K_{d+1}(f)$, we get that for any $u \in \mathbb{S}^{d-1}$,

$$\frac{h_{K_{d+1}(f)}(u)^2}{d(d+2)} \leq \frac{1}{|K_{d+1}(f)|} \int_{K_{d+1}(f)} \langle x, u \rangle^2 dx \leq \frac{d}{d+2} h_{K_{d+1}(f)}(u)^2. \quad (45)$$

Then, we use (39) for $p = 2$ to get that

$$\int_{K_{d+2}(f)} \langle x, u \rangle^2 dx = \frac{1}{f(0)} \int_{\mathbb{R}^d} \langle x, u \rangle^2 f(x) dx = \frac{\sigma^2}{f(0)}.$$

This gives

$$c_1^{d+2} \frac{\sigma^2}{f(0)} \leq \int_{K_{d+1}(f)} \langle x, u \rangle^2 dx \leq c_2^{d+2} \frac{\sigma^2}{f(0)}.$$

Using again the inclusion relations (40), we get upper and lower bound of $|K_{d+1}(f)|$ by $|K_d(f)| = 1/f(0)$, by (39) for $p = 0$. Thus, altogether we have

$$c_1^{2d+2} \sigma^2 \leq \frac{1}{|K_{d+1}(f)|} \int_{K_{d+1}(f)} \langle x, u \rangle^2 dx \leq c_2^{2d+2} \sigma^2.$$

Using these inequalities, (45) and the inclusion relations (40), we get

$$\sqrt{\frac{d+2}{d}} c_1^{d+2} \sigma B_2^d \subset K_d(f) \subset \sqrt{d(d+2)} c_2^{d+2} \sigma B_2^d.$$

Now, from [6, Theorem 4], we have $f(0) \geq e^{-d} \max(f)$ hence

$$\frac{L_f}{e} \leq f(0)^{\frac{1}{d}} \sigma = \left(\frac{f(0)}{\max(f)} \right)^{\frac{1}{d}} L_f \leq L_d.$$

Using that $L_f \geq 1/\sqrt{2\pi}e$, we conclude. \square

3.2 A concentration lemma

The following concentration lemma will be required to establish the concentration property 2.

Lemma 22 (Concentration Lemma). *Let $c_d := 3^{\frac{1}{d}} C_d$, where C_d is the constant from (44). Then, for every log-concave, isotropic, centered density function f and for every $x \in \mathbb{R}^d$ such that $\|x\|_2 \geq c_d/f(0)^{\frac{1}{d}}$,*

$$f(x) \leq f(0)2^{-\|x\|_2 \frac{f(0)^{\frac{1}{d}}}{c_d}}. \quad (46)$$

Proof. Let $x_{\max} \in \mathbb{R}^d$ be a value where the maximum of f is attained and for every $\theta \in \mathbb{S}^{d-1}$, let $r_{\max}(\theta) \in \mathbb{R}^d$ be a value where the maximum of $r \mapsto f(r\theta)$ is attained. Since f is log-concave, for every $\theta \in \mathbb{S}^{d-1}$, the function $r \mapsto f(r\theta)$ is non-increasing for $r \geq r_{\max}(\theta)$. Therefore, for every $r_0(\theta) \geq r_{\max}(\theta)$,

$$\int_0^\infty dr^{d-1} f(r\theta) dr \geq \int_{r_{\max}(\theta)}^{r_0(\theta)} dr^{d-1} f(r\theta) dr \geq f(r_0(\theta))(r_0(\theta)^d - r_{\max}(\theta)^d). \quad (47)$$

We choose $r_0 = r_0(\theta)$ as

$$r_0(\theta) = \left(2d \int_0^\infty \frac{r^{d-1} f(r\theta)}{f(0)} dr + r_{\max}^d \right)^{\frac{1}{d}}, \quad (48)$$

so that, by (47), one has $f(r_0(\theta)) \leq \frac{f(0)}{2}$. Since $r \mapsto f(r\theta)$ is non-decreasing on $[0, r_{\max}(\theta)]$, one has

$$\int_0^{+\infty} dr^{d-1} f(r\theta) dr \geq \int_0^{r_{\max}(\theta)} dr^{d-1} f(0) dr = f(0)r_{\max}(\theta)^d,$$

and therefore, by Lemma 21,

$$r_{\max}(\theta) \leq \frac{(d \int_0^{+\infty} r^{d-1} f(r\theta) dr)^{\frac{1}{d}}}{f(0)^{\frac{1}{d}}} \leq \frac{C_d}{f(0)^{\frac{1}{d}}},$$

where C_d is given by (44). Thus, by (48) and the definition of c_d ,

$$r_0(\theta) \leq \left(\frac{3}{f(0)} \right)^{\frac{1}{d}} C_d = \frac{c_d}{f(0)^{\frac{1}{d}}}.$$

Since for every $r \geq r_0(\theta)$, one has $r_0(\theta) = \frac{r_0(\theta)}{r} \cdot r + (1 - \frac{r_0(\theta)}{r}) \cdot 0$, by log-concavity, we deduce

$$f(r) \leq f(0) \left(\frac{f(r_0(\theta))}{f(0)} \right)^{\frac{r}{r_0(\theta)}} \leq f(0)2^{-\frac{r}{r_0(\theta)}} \leq f(0)2^{-\frac{rf(0)^{\frac{1}{d}}}{c_d}}.$$

Hence, if $\|x\|_2 \geq c_d/f(0)^{\frac{1}{d}}$, then

$$f(x) \leq f(0)2^{-\|x\|_2 \frac{f(0)^{\frac{1}{d}}}{c_d}}.$$

□

3.3 Sum of maxima of isotropic log-concave functions

The following lemma bounds the sum of the maxima of a log-concave density using our previous concentration result.

Lemma 23. *Let f be a centered, isotropic, log-concave density on \mathbb{R}^d with covariance $\sigma^2 \mathbf{I}_d$. Let $0 \leq i \leq 2$ and $0 \leq j \leq d-1$. Then, as $\sigma \rightarrow \infty$,*

$$\sum_{l \in \mathbb{Z}^{d-j}} |l_1|^i \max_{k \in \mathbb{Z}^j} f(k_1, \dots, k_j, l_1, \dots, l_{d-j}) = O_d \left(\frac{1}{\sigma^{j-i}} \right). \quad (49)$$

Proof. Set $\lambda = c_d f(0)^{-1/d}$, with c_d is given in Lemma 22. Then $f(x) \leq f(0)2^{-\|x\|_2}$, for $\|x\|_2 \geq \lambda$. Moreover, from the definition of L_f , its bounds and the inequality $f(0) \geq e^{-d} \max(f)$, we have

$$\frac{c_d \sigma}{L_d} \leq \lambda = c_d f(0)^{-1/d} \leq \frac{c_d e \sigma}{L_f} \leq c_d \sqrt{2\pi e^3} \sigma.$$

Note that

$$\sum_{l \in \mathbb{Z}^{d-j}} |l_1|^i \max_{k \in \mathbb{Z}^j} f(k, l) = \sum_{l \in \mathbb{Z}^{d-j} \|\cdot\|_\infty \leq} |l_1|^i \max_{k \in \mathbb{Z}^j} f(k, l) + \sum_{l \in \mathbb{Z}^{d-j} \|\cdot\|_\infty <} |l_1|^i \max_{k \in \mathbb{Z}^j} f(k, l),$$

The first sum is easily upper bounded by $O_d(\frac{1}{j-i})$. Indeed,

$$\sum_{l \in \mathbb{Z}^{d-j} \|\cdot\|_\infty \leq} |l_1|^i \max_{k \in \mathbb{Z}^j} f(k, l) \leq (\max f)(2\lambda)^{d+i-j} = \frac{L_f^d}{\sigma^d} (2\lambda)^{d+i-j} \leq \frac{L_d^d (2c_d \sqrt{2\pi e^3})^{d+i-j}}{\sigma^{j-i}} = O_d\left(\frac{1}{\sigma^{j-i}}\right).$$

Using the tails estimates of Lemma 22 and the fact that for $l \in \mathbb{Z}^{d-j}$, one has $\|l\|_2 \geq \|l\|_\infty$ and $\|l\|_2 \geq \frac{\|l\|_1}{\sqrt{d-j}} \geq \frac{\|l\|_1}{\sqrt{d}}$ the second sum can be expressed as follows

$$\begin{aligned} \sum_{l \in \mathbb{Z}^{d-j} \|\cdot\|_\infty <} |l_1|^i \max_{k \in \mathbb{Z}^j} f(k, l) &\leq f(0) \sum_{l \in \mathbb{Z}^{d-j} \|\cdot\|_\infty <} |l_1|^i 2^{-\frac{\|l\|_1}{\lambda\sqrt{d}}} \\ &= f(0) \left(\sum_{|l_1| <} |l_1|^i 2^{-\frac{|l_1|}{\lambda\sqrt{d}}} \right) \left(\sum_{|n| <} 2^{-\frac{|n|}{\lambda\sqrt{d}}} \right)^{d-j-1}. \end{aligned}$$

Putting $x = 2^{-1/\sqrt{d}}$ and assuming that $\lambda\sqrt{d} \geq 1$, we have $x \in [\frac{1}{2}, 1)$ and $\frac{1}{2} \leq \lambda\sqrt{d}(1-x) \leq \ln(2)$. Thus

$$\sum_{|n| <} 2^{-\frac{|n|}{\lambda\sqrt{d}}} = 2 \sum_{n <} x^n \leq \frac{2}{1-x} \leq 4\lambda\sqrt{d}.$$

Hence, for $i = 0$, replacing λ by its value, for $\sigma \geq \frac{L_d}{\sqrt{dc_d}}$, we have $\lambda \geq 1/\sqrt{d}$ and thus

$$\sum_{l \in \mathbb{Z}^{d-j} \|\cdot\|_\infty <} \max_{k \in \mathbb{Z}^j} f(k, l) \leq f(0) (4\lambda\sqrt{d})^{d-j} = (4\sqrt{dc_d})^{d-j} f(0)^{\frac{1}{d}} \leq \frac{(4\sqrt{dc_d})^{d-j} L_d^j}{\sigma^j} = O_d\left(\frac{1}{\sigma^j}\right).$$

And, for $i = 1$, it is not difficult to see that, for $\lambda \geq 1/\sqrt{d}$, we have

$$\sum_{|l_1| <} |l_1| 2^{-\frac{|l_1|}{\lambda\sqrt{d}}} = 2 \sum_{n <} n x^n = 2x \left(\sum_{n <} x^n \right)' \leq \frac{6}{(1-x)^2} \leq 24d\lambda^2.$$

Thus, again for $\sigma \geq \frac{L_d}{\sqrt{dc_d}}$, which ensures that $\lambda \geq 1/\sqrt{d}$, we deduce

$$\sum_{l \in \mathbb{Z}^{d-j} \|\cdot\|_\infty <} |l_1| \max_{k \in \mathbb{Z}^j} f(k, l) \leq f(0) \times 24d\lambda^2 (4\lambda\sqrt{d})^{d-j-1} \leq 2f(0) (4\lambda\sqrt{d})^{d-j+1} = O_d\left(\frac{1}{\sigma^{j-1}}\right).$$

In the same way, for $i = 2$, we easily get that, for $\lambda \geq 1/\sqrt{d}$,

$$\sum_{|l_1| <} l_1^2 2^{-\frac{|l_1|}{\lambda\sqrt{d}}} = 2 \sum_{n <} n^2 x^n = 2x^2 \left(\sum_{n <} x^n \right)'' + 2 \sum_{n <} n x^n \leq \frac{14}{(1-x)^3} + \frac{12}{(1-x)^2} \leq 26(2\lambda\sqrt{d})^3.$$

Thus, again for $\sigma \geq \frac{L_d}{\sqrt{dc_d}}$, which ensures that $\lambda \geq 1/\sqrt{d}$, we deduce

$$\sum_{l \in \mathbb{Z}^{d-j} \|\cdot\|_\infty <} l_1^2 \max_{k \in \mathbb{Z}^j} f(k, l) \leq f(0) \times 26(2\lambda\sqrt{d})^3 (4\lambda\sqrt{d})^{d-j-1} \leq 4f(0) (4\lambda\sqrt{d})^{d-j+2} = O_d\left(\frac{1}{\sigma^{j-2}}\right).$$

□

3.4 Approximating the integral, mean and covariance discretely

In this section, we will approximate the isotropic constant of an isotropic, log-concave density $f \in \mathbb{R}^d$ in a discrete way. To do so, we will first approximate the integral as a sum and subsequently approximate the continuous covariance of f by its discrete covariance.

Proposition 24. *Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a log-concave isotropic density function with covariance matrix of the form $\sigma^2 I_d$. Then*

$$\left| \int_{\mathbb{R}^d} f dx - \sum_{k \in \mathbb{Z}^d} f(k) \right| = O_d\left(\frac{1}{\sigma}\right).$$

Proof. Let us proceed by induction. Let us first consider the one dimensional base case. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a log-concave function and let suppose that the maximum of f is attained at $x_0 \in \mathbb{R}$. Let $k_0 \in \mathbb{Z}$ such that $k_0 \leq x_0 < k_0 + 1$. Then, one has

$$\begin{aligned} \int_{\mathbb{R}} f dx &= \sum_{k \in \mathbb{Z}} \int_k^{k+1} f(x) dx \\ &\geq \sum_{k \leq k_0} f(k) + \sum_{k \geq k_0+1} f(k+1) + \min\{f(k_0), f(k_0+1)\} \\ &= \sum_{k \in \mathbb{Z}} f(k) - f(k_0) - f(k_0+1) + \min\{f(k_0), f(k_0+1)\} \\ &= \sum_{k \in \mathbb{Z}} f(k) - \max\{f(k_0), f(k_0+1)\} \\ &= \sum_{k \in \mathbb{Z}} f(k) - \max_{\mathbb{Z}} f. \end{aligned}$$

The reverse inequality is analogous and thus one gets

$$\left| \int_{\mathbb{R}} f dx - \sum_{k \in \mathbb{Z}} f(k) \right| \leq \max_{\mathbb{Z}} f \leq \max_{\mathbb{R}} f. \quad (50)$$

For $d = 1$, we get $\max_{\mathbb{R}}(f) = \frac{L_f}{\sigma} \leq \frac{1}{\sigma}$, by [7], which proves the base case. Notice that the inequality (50) holds as soon as f is quasi-concave on \mathbb{R} , i.e. if f is first non-decreasing then non-increasing and not necessarily a density.

Now, let $x \in \mathbb{R}$, $y \in \mathbb{R}^{d-1}$ and let $F(y) := \int_{\mathbb{R}} f(x, y) dx$. Note that F is log-concave as the marginal of a log-concave function. Moreover F is an isotropic density and $\text{Cov}_{\mathbb{R}^{d-1}}(F) = \sigma^2 I_{d-1}$. One has

$$\left| \int_{\mathbb{R}^d} f(x, y) dx dy - \sum_{k \in \mathbb{Z}^d} f(k) \right| \leq \left| \int_{\mathbb{R}^{d-1}} F(y) dy - \sum_{y \in \mathbb{Z}^{d-1}} F(y) \right| + \sum_{y \in \mathbb{Z}^{d-1}} \left| \int_{\mathbb{R}} f(x, y) dx - \sum_{k \in \mathbb{Z}} f(k, y) \right|.$$

By the inductive hypothesis the first term is upper bounded by C_{d-1}/σ for some constant $C_{d-1} > 0$. Since for every $y \in \mathbb{Z}^{d-1}$, the function $x \mapsto f(x, y)$ is quasi-concave, one may use (50) to get that the second term is upper bounded by

$$\sum_{y \in \mathbb{Z}^{d-1}} \max_{x \in \mathbb{Z}} f(x, y).$$

We conclude by Lemma 23. □

A remark from convex geometry: the case of isotropic convex bodies. Applying the previous proposition to $f = \mathbb{1}_K/|K|_d$, where K is an isotropic convex body in \mathbb{R}^d and using that, in this case, $\sigma = L_K|K|^{1/d}$, we deduce the following proposition.

Proposition 25. *For any isotropic convex body K in \mathbb{R}^d ,*

$$\left| \frac{\#(K \cap \mathbb{Z}^d)}{|K|_d} - 1 \right| \leq \frac{M_d}{|K|^{1/d}},$$

for some constant $M_d > 0$, depending only on the dimension, where $\#K$ denotes the cardinality of a discrete set K .

Remark 26. The hypothesis of isotropicity is necessary, since in \mathbb{Z}^d , for $d \geq 2$, it is easy to construct convex bodies that do not contain integer points, but whose volumes are increasingly large.

Proposition 27. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a centered, isotropic, log-concave density. Then, for $1 \leq i \leq d$,

$$\sum_{k \in \mathbb{Z}^d} k_i f(k) = \int_{\mathbb{R}^d} x_i f(x) dx + O_d(1) = O_d(1).$$

In particular, for $d = 1$, the following finite bound holds for any log-concave centered integrable function

$$\left| \int_{\mathbb{R}} x f(x) dx - \sum_{k \in \mathbb{Z}} k f(k) \right| \leq (e + 1) \sum_{k \in \mathbb{Z}} f(k).$$

Proof. The proof is by induction again. Let us make a one dimensional remark that will be used on the base case of the induction. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a centered log-concave function and let suppose that the maximum of f is attained at $x_0 \in \mathbb{R}$ and let us suppose without loss of generality that $x_0 > 0$. Then since f is log-concave and by the inequality [6, Theorem 4] one has

$$\int_{\mathbb{R}} f(x) dx \geq \int_0^{x_0} f(x) dx \geq f(0)x_0 \geq \frac{\max f}{e} x_0. \quad (51)$$

Hence, we get $x_0 \max f \leq e \int_{\mathbb{R}} f(x) dx$. The case of the summation is analogous. Let $k_0 \in \mathbb{Z}$ be such that $k_0 \leq x_0 < k_0 + 1$. Then, we deduce

$$\sum_{k \in \mathbb{Z}} f(k) \geq \sum_{k=0}^{k_0} f(k) \geq (k_0 + 1)f(0) \geq \frac{\max f}{e} (k_0 + 1). \quad (52)$$

Let us start the induction. Let f be a centered, isotropic, log-concave density and let suppose that the maximum of f is attained at x_0 . We decompose the integral $\int_{\mathbb{R}} x f(x) dx$ as follows

$$\begin{aligned} \int_{\mathbb{R}} x f(x) dx &= \sum_{k \tau 0} \int_k^{k+1} x f(x) dx + \sum_{k=0}^{k_0-1} \int_k^{k+1} x f(x) dx + \int_{k_0}^{k_0+1} x f(x) dx + \sum_{k \geq k_0+1} \int_k^{k+1} x f(x) dx \\ &\geq \sum_{k \tau 0} k f(k+1) + \sum_{k=0}^{k_0-1} k f(k) + k_0 \min\{f(k_0), f(k_0+1)\} + \sum_{k \geq k_0+1} k f(k+1) \\ &\geq \sum_{k \in \mathbb{Z}} k f(k) - k_0 f(k_0) - k_0 f(k_0+1) + k_0 \min\{f(k_0), f(k_0+1)\} - \sum_{k \in \mathbb{Z}} f(k). \end{aligned}$$

Therefore,

$$\begin{aligned} \int_{\mathbb{R}} x f(x) dx &\geq \sum_{k \in \mathbb{Z}} k f(k) - \sum_{k \in \mathbb{Z}} f(k) - k_0 \max f \\ &\geq \sum_{k \in \mathbb{Z}} k f(k) - \sum_{k \in \mathbb{Z}} f(k) - e \sum_{k \in \mathbb{Z}} f(k) = \sum_{k \in \mathbb{Z}} k f(k) - (e + 1) \sum_{k \in \mathbb{Z}} f(k), \end{aligned}$$

where in the last line we have used the inequality (52). Bounding from the above $\int_{\mathbb{R}} x f(x) dx$ we obtain a similar inequality. Thus, we get

$$\left| \int_{\mathbb{R}} x f(x) dx - \sum_{k \in \mathbb{Z}} k f(k) \right| \leq (e + 1) \sum_{k \in \mathbb{Z}} f(k),$$

and the one-dimensional case follows by Proposition 24.

For the inductive step, let $F : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$ be defined, for $x \in \mathbb{R}, y \in \mathbb{R}^{d-1}$, by

$$F(y) := \int_{\mathbb{R}} f(x, y) dx.$$

Then F is log-concave, centered and isotropic as the marginal of a log-concave, centered and isotropic density and, for $i = 1, \dots, d-1$

$$\int_{\mathbb{R}^d} y_i f(x, y) dx dy = \int_{\mathbb{R}^{d-1}} y_i F(y) dy.$$

We have

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} y_i f(x, y) dx dy - \sum_{k_1 \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^{d-1}} k_i f(k_1, k) \right| \\ & \leq \left| \int_{\mathbb{R}^d} y_i f(x, y) dx dy - \sum_{k \in \mathbb{Z}^{d-1}} k_i F(k) \right| + \left| \sum_{k \in \mathbb{Z}^{d-1}} k_i F(k) - \sum_{k \in \mathbb{Z}^d} k_i f(k) \right|. \end{aligned}$$

But the first term is $O_d(1)$ by inductive hypothesis and for the second we note that $f(\cdot, k)$ is log-concave for every $k \in \mathbb{Z}^{d-1}$, so we may apply (50) to get, for any $k \in \mathbb{Z}^{d-1}$,

$$\left| F(k) - \sum_{k_1 \in \mathbb{Z}} f(k_1, k) \right| = \left| \int_{\mathbb{R}} f(x, k) dx - \sum_{k_1 \in \mathbb{Z}} f(k_1, k) \right| \leq \max_{x \in \mathbb{R}} f(x, k),$$

which gives

$$\left| \int_{\mathbb{R}^d} y_i f(x, y) dx dy - \sum_{k_1 \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^{d-1}} k_i f(k_1, k) \right| \leq O_d(1) + \sum_{k \in \mathbb{Z}^{d-1}} k_i \max_{x \in \mathbb{R}} f(x, k) = O_d(1),$$

where the last equality follows by Lemma 23. \square

Now we wish to show that a continuous isotropic, log-concave density f in \mathbb{R}^d is almost isotropic in the discrete sense, meaning that its discrete covariance matrix $\text{Cov}_{\mathbb{Z}}(f)$ has $O(\sigma)$ off-diagonal elements and $\sigma^2 + O(\sigma)$ diagonal elements.

Proposition 28. *Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a centered, isotropic, log-concave density with $\text{Cov}(f) = \sigma^2 \mathbf{I}_d$. Then*

$$\sum_{k \in \mathbb{Z}^d} f(k) k_i^2 = \sigma^2 + O_d(\sigma), \text{ for every } 1 \leq i \leq d.$$

Proof. We proceed by induction on the dimension. For the one-dimensional case let $x_0 = \inf\{x \in \mathbb{R} : f(x) = \max f\}$. Then

$$\int_{\mathbb{R}} f(x) x^2 dx = \int_{x \leq x_0} f(x) x^2 dx + \int_{x < x_0} f(x) x^2 dx.$$

Noting that f is non-decreasing on $(-\infty, x_0]$ and non-increasing on $[x_0, +\infty)$ by log-concavity, we get

$$\begin{aligned} \int_{\mathbb{R}} f(x) x^2 dx &= \sum_{k \tau \lfloor x_0 \rfloor} \int_{[k, k+1)} f(x) x^2 dx + \sum_{k < \lfloor x_0 \rfloor} \int_{[k, k+1)} f(x) x^2 dx + \int_{\lfloor x_0 \rfloor}^{\lfloor x_0 \rfloor + 1} f(x) x^2 dx \\ &\leq \sum_{k \tau \lfloor x_0 \rfloor} f(k+1) \left(k^2 + k + \frac{1}{3} \right) + \sum_{k < \lfloor x_0 \rfloor} f(k) \left(k^2 + k + \frac{1}{3} \right) + \int_{\lfloor x_0 \rfloor}^{\lfloor x_0 \rfloor + 1} f(x) x^2 dx \\ &\leq \sum_{k \in \mathbb{Z}} f(k) k^2 - \sum_{k \tau \lfloor x_0 \rfloor} f(k+1) k - \sum_{k < \lfloor x_0 \rfloor} f(k) k + f(x_0) \left(\lfloor x_0 \rfloor^2 + \lfloor x_0 \rfloor + \frac{1}{3} \right) \\ &\leq \sum_{k \in \mathbb{Z}} f(k) k^2 - \sum_{k \in \mathbb{Z}} f(k) k + f(x_0) \left(\lfloor x_0 \rfloor^2 + \lfloor x_0 \rfloor + \frac{1}{3} \right) \end{aligned} \tag{53}$$

Now we observe that $f(x_0) = \max f \leq \frac{1}{\sigma}$ and, by (51), $x_0 = O(\sigma)$. Hence, the last term in (53) is $O(\sigma)$. Furthermore, by Proposition 24, the second term in (53) is $O(1)$. Thus,

$$\int_{\mathbb{R}} f(x) x^2 dx \leq \sum_{k \in \mathbb{Z}} f(k) k^2 + O(\sigma).$$

Similarly, using the piecewise monotonicity of f in the reverse way, we obtain

$$\int_{\mathbb{R}} f(x) x^2 dx \geq \sum_{k \in \mathbb{Z}} f(k) k^2 + O(\sigma).$$

This proves the base case.

Let $x \in \mathbb{R}$, $y \in \mathbb{R}^{d-1}$ and let $F(y) := \int_{\mathbb{R}} f(x, y) dx$. Then, by the inductive hypothesis applied to F , for $i = 1, \dots, d-1$,

$$\int_{\mathbb{R}^d} f(x, y) y_i^2 dx dy = \int_{\mathbb{R}^{d-1}} F(y) y_i^2 dy = \sum_{k \in \mathbb{Z}^{d-1}} F(k) k_i^2 + O_d(\sigma). \quad (54)$$

Since, for every $k \in \mathbb{Z}^{d-1}$, the function $f(\cdot, k)$ is log-concave in \mathbb{R} , we have, by (50),

$$\left| F(k) - \sum_{k' \in \mathbb{Z}} f(k', k) \right| = \left| \int_{\mathbb{R}} f(x, k) dx - \sum_{k' \in \mathbb{Z}} f(k', k) \right| \leq \max_x f(x, k). \quad (55)$$

So by (54) and (55) we get, for $i = 1, \dots, d-1$

$$\begin{aligned} \int_{\mathbb{R}^d} f(x, y) y_i^2 dx dy &= \sum_{k' \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^{d-1}} f(k', k) k_i^2 + \sum_{k \in \mathbb{Z}^{d-1}} O(\max_x f(x, k)) k_i^2 + O_d(\sigma) \\ &= \sum_{k' \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^{d-1}} f(k', k) k_i^2 + O_d(\sigma), \end{aligned}$$

by Lemma 23, and the result follows. \square

We will need the following lemma for the base case of the induction of Proposition 30.

Lemma 29. *Let f be a centered, isotropic, log-concave density in \mathbb{R}^2 , with $\text{Cov}(f) = \sigma^2 \mathbf{I}_2$. For $x \in \mathbb{R}$, let*

$$N_{\max}(x) := \inf\{y \in \mathbb{R} : f(x, y) = \max_y f(x, y)\}.$$

Then

$$\int_{\mathbb{R}} x N_{\max}(x) \max_y f(x, y) dx = O(\sigma) \quad (56)$$

and

$$\sum_{k \in \mathbb{Z}} k N_{\max}(k) \max_y f(k, y) = O(\sigma). \quad (57)$$

Proof. Let c be an absolute constant to be determined later and denote the set $A := \{x \in \mathbb{R} : |x|, |N_{\max}(x)| \leq c\sigma\}$. We have

$$\int_{\mathbb{R}} x N_{\max}(x) \max_y f(x, y) dx = \int_{A_\sigma} x N_{\max}(x) \max_y f(x, y) dx + \int_{A_\sigma^c} x N_{\max}(x) \max_y f(x, y) dx$$

The first term can be trivially bounded as

$$\left| \int_{A_\sigma} x N_{\max}(x) \max_y f(x, y) dx \right| \leq O(\sigma^3) \max f = O(\sigma).$$

For the second term we have, for some absolute constant C'_2 and for c large enough,

$$\begin{aligned} \left| \int_{A_\sigma^c} |x| N_{\max}(x) \max_y f(x, y) dx \right| &\leq \max(f) \int_{A_\sigma^c} x N_{\max}(x) e^{-\frac{|x| + N_{\max}(x)}{\sigma}} C'_2 dx \\ &\leq O\left(\frac{1}{\sigma^2}\right) \sigma \int_{x < c} x e^{-\frac{x}{\sigma} C'_2} dx \\ &= O(\sigma). \end{aligned} \quad (58)$$

$$= O(\sigma). \quad (59)$$

Here (58) follows by using the fact that $ye^{-y} \leq 1$. This gives (56), (57) follows in a similar way. \square

Proposition 30. *Let f be a centered, continuous, isotropic, log-concave density in \mathbb{R}^d , with $\text{Cov}(f) = \sigma^2 \mathbf{I}_d$. Then, for all $i \neq j$*

$$\sum_{k \in \mathbb{Z}^d} f(k) k_i k_j = O_d(\sigma).$$

Proof. We proceed by induction on d . Consider first the base case $d = 2$. Then, letting $N_{\max}(x) := \inf\{y \in \mathbb{R} : f(x, y) = \max_y f(x, y)\}$ and $M_{\max}(y) := \inf\{x \in \mathbb{R} : f(x, y) = \max_x f(x, y)\}$ and assuming without loss of generality that the infima are achieved,

$$\int_{\mathbb{R}_+^2} f(x, y)xydx = \int_{\mathbb{R}_+} x \int_{0 \leq y \leq N_{\max}(x)} f(x, y)dy + \int_{y < N_{\max}(x)} f(x, y)dydx.$$

Now

$$\begin{aligned} \int_{0 \leq y \leq N_{\max}(x)} f(x, y)dy &\leq \sum_{0 \leq k \leq N_{\max}(x)} \int_{[k, k+1)} f(x, y)dy \\ &\leq \sum_{0 \leq k \leq N_{\max}(x)-1} (k+1)f(x, k+1) + \max_y f(x, y)(N_{\max}(x) + 1) \end{aligned} \quad (60)$$

and

$$\int_{y < N_{\max}(x)} f(x, y)dy \leq \sum_{k \geq N_{\max}(x)} (k+1)f(x, k). \quad (61)$$

By (60) and (61)

$$\int_{\mathbb{R}_+^2} f(x, y)xydx \leq \int_{\mathbb{R}} x \left(\sum_{k \geq 0} kf(x, k) + \sum_{k \geq N_{\max}(x)} f(x, k) + \max_y f(x, y)(N_{\max}(x) + 1) \right) dx \quad (62)$$

For the first term in (62) we have

$$\begin{aligned} \int_{\mathbb{R}} x \sum_{k \geq 0} kf(x, k)dx &\leq \sum_{k \geq 0} \sum_{0 \leq m \leq M_{\max}(k)-1} k(m+1)f(m+1, k) + \sum_{k \geq 0} \sum_{m < M_{\max}(k)} k(m+1)f(m, k) \\ &\quad + \sum_{k \geq 0} k(M_{\max}(k) + 1) \max_x f(x, k) \\ &\leq \sum_{k \geq 0} \sum_{m \geq 0} kmf(m, k) + \sum_{k \geq 0} \sum_{m \geq 0} kf(m, k) + \sum_{k \geq 0} k(M_{\max}(k) + 1) \max_x f(x, k) \end{aligned} \quad (63)$$

$$\leq \sum_{k \geq 0} \sum_{m \geq 0} kmf(m, k) + \sum_{k \geq 0} \sum_{m \geq 0} kf(m, k) + O(\sigma), \quad (64)$$

where the last inequality follows by Lemma 29. On the other hand, for the second term in (62) we have

$$\int_{\mathbb{R}} \sum_{k \geq N_{\max}} xf(x, k)dx \leq \left(\int_{\mathbb{R}} \sum_{k \in \mathbb{Z}} x^2 f(x, k)dx \right)^{\frac{1}{2}} = O(\sigma) \quad (65)$$

by the Cauchy-Schwarz inequality and Proposition 28. Finally, the third term in (62) is

$$\int_{\mathbb{R}} \max_y f(x, y)(N_{\max}(x) + 1)dx = O(\sigma) \quad (66)$$

by Lemma 29. Plugging (64), (65) and (66) into (62) we obtain

$$\int_{\mathbb{R}_+^2} f(x, y)xydx \leq \sum_{k \geq 0} \sum_{m \geq 0} kmf(m, k) + \sum_{k \geq 0} \sum_{m \geq 0} kf(m, k) + O(\sigma). \quad (67)$$

Using the reverse bounds to those in (60) and (61) obtained by the monotonicity of f , we get the lower bound

$$\int_{\mathbb{R}_+^2} f(x, y)xydx \geq \sum_{k \geq 0} \sum_{m \geq 0} kmf(m, k) + \sum_{k \geq 0} \sum_{m \geq 0} kf(m, k) + O(\sigma).$$

In a completely analogous way, we may bound the integral on the rest of the quadrants and obtain the approximations

$$\int_{\mathbb{R}_-^2} f(x, y)xydx = \sum_{k \leq 0} \sum_{m \leq 0} kmf(m, k) + \sum_{k \leq 0} \sum_{m \leq 0} kf(m, k) + O(\sigma), \quad (68)$$

$$\int_{\mathbb{R}_- \times \mathbb{R}_+} f(x, y)xydx = \sum_{k \geq 0} \sum_{m \leq 0} kmf(m, k) + \sum_{k \geq 0} \sum_{m \leq 0} kf(m, k) + O(\sigma) \quad \text{and} \quad (69)$$

$$\int_{\mathbb{R}_+ \times \mathbb{R}_-} f(x, y)xydx = \sum_{k \leq 0} \sum_{m \geq 0} kmf(m, k) + \sum_{k \leq 0} \sum_{m \geq 0} kf(m, k) + O(\sigma). \quad (70)$$

Since, by Proposition 27, $\sum_k \sum_m k f(m, k) = O(1)$, by adding (67)–(70) we obtain

$$0 = \int_{\mathbb{R}^2} f(x, y) x y dx = \sum_{k \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} k m f(m, k) + O(\sigma).$$

This completes the base case $d = 2$.

The inductive step is similar to that in the proof of Proposition 28. Let $x \in \mathbb{R}$, $y = (y_1, \dots, y_{d-1}) \in \mathbb{R}^{d-1}$ and let $F(y) := \int_{\mathbb{R}} f(x, y) dx$. Then, by the inductive hypothesis, for $1 \leq i, j \leq d-1$,

$$\int_{\mathbb{R}^d} f(x, y) y_i y_j dy = \int_{\mathbb{R}^{d-1}} F(y) y_i y_j dy = \sum_{k \in \mathbb{Z}^{d-1}} F(k) k_i k_j + O_d(\sigma). \quad (71)$$

Since for every $k = (k_1, \dots, k_{d-1}) \in \mathbb{Z}^{d-1}$, the function $f(\cdot, k)$ is log-concave on \mathbb{R} , we have by Proposition 24

$$F(k) - \sum_{k' \in \mathbb{Z}} f(k', k) = \int_{\mathbb{R}} f(x, k) dx - \sum_{k' \in \mathbb{Z}} f(k', k) = O_d\left(\max_x f(x, k)\right). \quad (72)$$

So by (71) and (72) we get, for $1 \leq i, j \leq d-1$

$$\int_{\mathbb{R}^d} f(x, y) y_i y_j dx dy = \sum_{k' \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^{d-1}} f(k', k) k_i k_j + \sum_{k \in \mathbb{Z}^{d-1}} O\left(\max_x f(x, k)\right) k_i k_j + O_d(\sigma). \quad (73)$$

But by the Cauchy-Schwarz inequality and Lemma 23, $\sum_{k \in \mathbb{Z}^{d-1}} O\left(\max_x f(x, k)\right) k_i k_j = O_d(\sigma)$ and the result follows. \square

Corollary 31. *Let f be a centered, isotropic, log-concave density in \mathbb{R}^d , with $\text{Cov}(f) = \sigma^2 \text{I}_d$. Then*

$$\det\left(\text{Cov}_{\mathbb{Z}^d}(f)\right) = \sigma^{2d} + O_d(\sigma^{2d-1}).$$

Proof. By Proposition 30

$$[\text{Cov}_{\mathbb{Z}^d}(f)]_{ij} = O_d(\sigma), \quad \text{for } i \neq j$$

and

$$[\text{Cov}_{\mathbb{Z}^d}(f)]_{ij} = \sigma^2 + O_d(\sigma), \quad \text{for } i = j.$$

Thus,

$$\begin{aligned} \det\left(\text{Cov}_{\mathbb{Z}^d}(f)\right) &= \sum_{\tau} \text{sgn}(\tau) \prod_{i=1}^d \text{Cov}_{\mathbb{Z}^d}(f)_{i\tau(i)} \\ &= \sigma^{2d} + O(\sigma^{2d-1}), \end{aligned}$$

since for τ being the identity permutation $\text{sgn}(\tau) \prod_{i=1}^d \text{Cov}_{\mathbb{Z}^d}(f)_{i\tau(i)} = (\sigma^2 + O(\sigma))^d = \sigma^{2d} + O(\sigma^{2d-1})$ and for any other permutation τ'

$$\prod_{i=1}^d \text{Cov}_{\mathbb{Z}^d}(f)_{i\tau'(i)} = O((\sigma^2)^{d-2} \sigma^2) = O_d(\sigma^{2d-2}).$$

\square

Finally, Corollary 32 below is a version of Theorem 6 with the assumption that the continuous function f is isotropic. It can be observed from the proof that the dependence of C'_d on d can be taken to be the same as in the continuous case (cf. (10)). It can also be seen that due to use of Corollary 31, σ needs to be taken at least $\Omega(d!)$. We do not know whether this is the best (lowest) rate.

Corollary 32 (Theorem 6 for isotropic f). *Let p be log-concave p.m.f. on \mathbb{Z}^d , whose continuous log-concave extension, say f , is isotropic. Then there exists a constant C'_d that depends on the dimension only, such that*

$$\max_{k \in \mathbb{Z}^d} p(k) \leq \frac{C'_d}{\det\left(\text{Cov}(p)\right)^{\frac{1}{2}}}$$

provided that $\sigma := \det\left(\text{Cov}_{\mathbb{R}}(f)\right)$ is large enough depending on d .

Proof. Since p is extensible log-concave, there exists a continuous log-concave function f (not necessarily a density) such that $f(k) = p(k)$ for all $k \in \mathbb{Z}^d$ and by assumption f is isotropic. Thus

$$\max_{k \in \mathbb{Z}^d} p(k) \leq \max_{x \in \mathbb{R}^d} f(x) = \frac{L_f^d \int_{\mathbb{R}^d} f}{\sigma^d} \leq \frac{C_d \int_{\mathbb{R}^d} f}{\sigma^d}, \quad (74)$$

where C_d is an upper bound of L_f^d . Noting that by definition $\text{Cov}(Zp) = \text{Cov}(p)$ for any normalising constant $Z \in \mathbb{R}_+$ and applying Proposition 24 and Corollary 31 to $\frac{f(\cdot)}{\int f}$, we obtain

$$\max_{k \in \mathbb{Z}^d} p(k) \leq \frac{C_d(1 + O_d(\frac{1}{\sigma}))}{\left(\det(\text{Cov}(p)) + O_d(\sigma^{2d-1})\right)^{\frac{1}{2}}} \leq \frac{2C_d}{\det(\text{Cov}(p))^{\frac{1}{2}}}$$

provided that σ is large enough depending on d , using that $O_d(\sigma^{2d-1}) \geq -\frac{1}{2} \det(\text{Cov}(p)) \simeq -\frac{1}{2} \sigma^{2d}$ as $\sigma \rightarrow \infty$. \square

Proposition 33. *Fix $d \geq 1$ and let p be a log-concave p.m.f. on \mathbb{Z}^d with almost isotropic extension. Then, for every $1 \leq i \leq d$,*

$$\sum_{k \in \mathbb{Z}^d} |p(k) - p(k - e_i)| = O_d \left(\frac{1}{\det(\text{Cov}(p))^{\frac{1}{2d}}} \right),$$

where $e_i \in \mathbb{Z}^d$ is defined as the vector with the i -th coordinate 1 and all the other coordinates 0.

Proof. For any one-dimensional quasi-concave non-negative real sequence $(a_n)_{n \in \mathbb{Z}}$, $\sum_{n \in \mathbb{Z}} |a_n - a_{n-1}| \leq 2 \max_{n \in \mathbb{Z}} a_n$. Thus, by log-concavity, denoting $\bar{k}_i = \sum_{j \neq i} k_j e_j \in \mathbb{Z}^{d-1}$, one has

$$\sum_{k \in \mathbb{Z}^d} |p(k) - p(k - e_i)| \leq 2 \sum_{\bar{k}_i \in \mathbb{Z}^{d-1}} \max_{k_i \in \mathbb{Z}} p(k_1, \dots, k_{i-1}, k_i, k_{i+1}, \dots, k_d),$$

Hence, as in (49), we have

$$\sum_{\bar{k}_i \in \mathbb{Z}^{d-1}} \max_{k_i \in \mathbb{Z}} p(k_1, \dots, k_{i-1}, k_i, k_{i+1}, \dots, k_d) = O_d \left(\frac{1}{\det(\text{Cov}(p))^{\frac{1}{2d}}} \right)$$

and the result follows. \square

3.5 Relaxing the isotropicity assumption on f

In this section, we relax the assumptions of Lemma 21, so that we obtain conclusions of the same kind under some weaker assumptions on the eigenvalues of the continuous covariance matrix $\text{Cov}_{\mathbb{R}}(f)$. Specifically, we will prove:

Proposition 34. *For every centered and log-concave density $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and for every $\theta \in \mathbb{S}^{d-1}$ one has*

$$c_1^{d+2} f(0)^{\frac{1}{d}} \sqrt{\lambda_{\min}(\text{Cov}(f))} \leq \left(\int_0^\infty r^{d-1} f(r\theta) dr \right)^{\frac{1}{d}} \leq (d+1) c_2^{d+2} f(0)^{\frac{1}{d}} \sqrt{\lambda_{\max}(\text{Cov}(f))}.$$

where c_1 and c_2 are the constants appearing in (40).

Before giving the proof, we mention the consequences of Proposition 34.

Corollary 35. *Let f be a centered, almost isotropic log-concave density. Then for every $\theta \in \mathbb{S}^{d-1}$*

$$\int_0^\infty r^{d-1} f(r\theta) dr = \Theta_d(1),$$

as $\sigma := \det(\text{Cov}(f)) \rightarrow \infty$.

Proof. Since f is almost isotropic, we have

$$\lambda_{\min}(\text{Cov}(f)) \simeq_d \lambda_{\max}(\text{Cov}(f)) \simeq_d \sigma^2$$

as $\sigma \rightarrow \infty$. The result follows from Proposition 34 since $f(0) \simeq_d \max f \simeq_d \frac{1}{\sigma^d}$. \square

Remark 36. The isotropicity assumption was only used in Lemma 21, which in turn allowed us to prove the concentration Lemma 22 which was repeatedly evoked in Section 3 to bound the sums of maxima of log-concave functions and thus also the error terms. By Corollary 35, a version of Lemma 22 holds true for almost isotropic densities, although with possibly different constants and for large enough σ . Therefore, all the previous results of Section 3 hold true and thus also the statements of Theorem 3 and 4 under the almost isotropic log-concave assumption.

Remark 37. Note that if the function f is in addition isotropic i.e. $\text{Cov}(f) = \sigma^2 \mathbf{I}_d$, one has that $f(0) \simeq_d \frac{1}{\sigma^d}$ and so we recover the result of Lemma 21.

For the above-mentioned proposition, the following lemma which gives bounds for the inradius and circumradius of convex bodies in not necessarily isotropic position will be essential .

Proof of Proposition 34. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a centered log-concave density. For simplicity, we denote $\lambda_{\min} = \lambda_{\min}(\text{Cov}(f))$ and analogously for λ_{\max} . Following the same steps as in the proof of Lemma 21, we get that, for any $u \in \mathbb{S}^{d-1}$,

$$\frac{\lambda_{\min}}{f(0)} \leq \int_{K_{d+2}(f)} \langle x, u \rangle^2 dx = \frac{1}{f(0)} \int_{\mathbb{R}^d} \langle x, u \rangle^2 f(x) dx \leq \frac{\lambda_{\max}}{f(0)}.$$

Using the inclusion relations (40), this gives

$$c_1^{2d+2} \lambda_{\min} \leq \frac{1}{|K_{d+1}(f)|} \int_{K_{d+1}(f)} \langle x, u \rangle^2 dx \leq c_2^{2d+2} \lambda_{\max}.$$

Using these inequalities, (45) and again the inclusion relations (40), we get

$$\sqrt{\frac{d+2}{d}} c_1^{d+2} \sqrt{\lambda_{\min}} B_2^d \subset K_d(f) \subset \sqrt{d(d+2)} c_2^{d+2} \sqrt{\lambda_{\max}} B_2^d,$$

which gives inequality (34). □

4 Concluding Remarks and Open Questions

1. As noted in the Introduction, for $d = 1$, our definition of log-concavity is equivalent to the usual definition $p(k)^2 \geq p(k-1)p(k+1), k \in \mathbb{Z}$, which is preserved under convolution (e.g. [14]). However, for $d > 1$ log-concavity may not be preserved in general as we have already observed in Remark 10. On the other hand in the i.i.d. case, Proposition 38 below shows that $\sum_{i=1}^n A$ is convex for any discrete convex set A , meaning that $\text{conv}(A) \cap \mathbb{Z}^d = A$. We do not know whether our definition of log-concavity is preserved under self-convolution.

Proposition 38. *Let $A \subset \mathbb{Z}^d$ be \mathbb{Z}^d convex. Then, for any integer $n \geq 2$, $A_n := \sum_{i=1}^n A$ is \mathbb{Z}^d -convex.*

Proof. It is clear that

$$A_n \subset \text{conv}(A_n) \cap \mathbb{Z}^d.$$

For the inverse inclusion, we first note that $\text{conv}(A_n) = n \text{conv}(A)$. Thus, if $a \in \text{conv}(A_n) \cap \mathbb{Z}^d$, then there are $a_1, a_2 \in A$ and $\lambda \in [0, 1]$ such that $a = n((1-\lambda)a_1 + \lambda a_2) \in \mathbb{Z}^d$.

We distinguish n cases depending on the value of λ . If $\frac{k}{n} \leq \lambda \leq \frac{k+1}{n}$ for k integer such that $0 \leq k \leq n-1$, we express a conveniently as follows

$$a = (n\lambda - k)a_2 + (1 - (n\lambda - k))a_1 + (n - k - 1)a_1 + ka_2.$$

Thus, we get that $a \in \text{conv}(A) \cap \mathbb{Z}^d + A_{n-k-1} + A_k$. By \mathbb{Z}^d -convexity, $\text{conv}(A) \cap \mathbb{Z}^d = A$ and $A_{n-k-1} + A_k = A_{n-1}$. Therefore, $a \in A + A_{n-1} = A_n$. □

It is therefore natural to ask the following question:

Question 39. *Let X_1, X_2 be i.i.d. log-concave random vectors on \mathbb{Z}^d . Is $X_1 + X_2$ log-concave? Furthermore, if X_1, \dots, X_n are i.i.d. log-concave, is $\sum_{i=1}^n X_i$ log-concave for every n ?*

We note here that our assumptions of Definition 2 are preserved up to constants under convolution as shown in Proposition 13 and therefore, assuming that the p.m.f. of X_1 is log-concave ensures that the p.m.f. of the sum also satisfies these assumptions.

Answering Question 39 even for quantized multivariate Gaussians seems to be non-trivial. Nevertheless, the following simple example shows that the answer is positive for quantized isotropic multivariate Gaussians.

Example 40. Let X_1 be a random variable with p.m.f. on \mathbb{Z}^d proportional to $e^{-\frac{\|k\|_2^2}{2\sigma^2}}$; that is X_1 is a multivariate centered isotropic Gaussian quantized on \mathbb{Z}^d . Then X_1 has the distribution of (Z_1, \dots, Z_d) , where Z_i are i.i.d. one-dimensional quantized centered Gaussians with variance σ^2 . Note that if p is a p.m.f. on \mathbb{Z}^d which is a product of log-concave p.m.f.s $\{p_i\}_{i=1}^d$ on \mathbb{Z}

$$p(k) = \prod_{i=1}^d p^{(i)}(k_i), \quad k = (k_1, \dots, k_n) \in \mathbb{Z}^d,$$

then p is clearly log-concave on \mathbb{Z}^d .

Therefore, since log-concavity is preserved under convolution in dimension one [4] and the coordinates of $\sum_{i=1}^n X_i$ are independent, it follows that $\sum_{i=1}^n X_i$ is log-concave for every n .

2. For Theorem 18 and therefore for Theorems 3 and 4 as well, we have assumed that there exists a continuous extension f , which is almost isotropic. We have then shown that f is also almost isotropic in the discrete sense. It would be more natural to start with the assumption that the discrete p.m.f. is isotropic. However, in this case we would not be able to use the continuous toolkit to prove our concentration lemma in Section 3.2. Nevertheless, we suspect that if the discrete p.m.f. is isotropic, then there exists a continuous extension that is almost isotropic and therefore our results for discrete log-concave distributions would still hold under this assumption:

Question 41. Let X be a log-concave random vector with p.m.f. p on \mathbb{Z}^d . Assume that p is almost isotropic. Is there a continuous log-concave extension of p on \mathbb{R}^d which is almost isotropic?

References

- [1] S. Artstein, K. Ball, F. Barthe, and A. Naor. Solution of Shannon’s problem on the monotonicity of entropy. *Journal of the American Mathematical Society*, 17(4):975–982, 2004.
- [2] K. Ball. Logarithmically concave functions and sections of convex sets in \mathbf{R}^n . *Studia Math*, 88(1):69–84, 1988.
- [3] A. R. Barron. Entropy and the central limit theorem. *The Annals of Probability*, pages 336–342, 1986.
- [4] S. G. Bobkov, A. Marsiglietti, and J. Melbourne. Concentration functions and entropy bounds for discrete log-concave distributions. *Combinatorics, Probability and Computing*, 31(1):54–72, 2022.
- [5] S. Brazitikos, A. Giannopoulos, P. Valettas, and B.-H. Vritsiou. *Geometry of isotropic convex bodies*, volume 196. American Mathematical Soc., 2014.
- [6] M. Fradelizi. Sections of convex bodies through their centroid. *Archiv der Mathematik*, 69(6):515–522, 1997.
- [7] M. Fradelizi. Hyperplane sections of convex bodies in isotropic position. *Beiträge Algebra Geom.*, 40(1):163–183, 1999.
- [8] L. Gavalakis. Approximate discrete entropy monotonicity for log-concave sums. *Combinatorics, Probability and Computing*, page 1–14, 2023.
- [9] L. Gavalakis. Discrete generalised entropy power inequalities for log-concave random variables. In *2023 IEEE International Symposium on Information Theory (ISIT)*, pages 42–47, 2023.
- [10] L. Gavalakis and I. Kontoyiannis. Entropy and the discrete central limit theorem. *Stochastic Processes and their Applications*, page 104294, 2024.
- [11] S. Haghshatshoar, E. Abbe, and E. Telatar. Adaptive sensing using deterministic partial hadamard matrices. In *2012 IEEE International Symposium on Information Theory Proceedings*, pages 1842–1846. Ieee, 2012.

- [12] S. Haghhighatshoar, E. Abbe, and I. E. Telatar. A new entropy power inequality for integer-valued random variables. *IEEE transactions on information theory*, 60(7):3787–3796, 2014.
- [13] P. Harremoés, C. Vignat, et al. An entropy power inequality for the binomial family. *JIPAM. J. Inequal. Pure Appl. Math*, 4(5):93, 2003.
- [14] S. Hoggar. Chromatic polynomials and logarithmic concavity. *Journal of Combinatorial Theory, Series B*, 16(3):248–254, 1974.
- [15] R. Kannan, L. Lovász, and M. Simonovits. Isoperimetric problems for convex bodies and a localization lemma. *Discrete & Computational Geometry*, 13:541–559, 1995.
- [16] B. Klartag. Logarithmic bounds for isoperimetry and slices of convex sets. *Ars Inveniendi Analytica*, 6 2023.
- [17] M. Madiman and A. Barron. Generalized entropy power inequalities and monotonicity properties of information. *IEEE Transactions on Information Theory*, 53(7):2317–2329, 2007.
- [18] M. Madiman, J. Melbourne, and C. Roberto. Bernoulli sums and Rényi entropy inequalities. *Bernoulli*, 29(2):1578–1599, 2023.
- [19] M. Madiman, L. Wang, and J. O. Woo. Majorization and Rényi entropy inequalities via Sperner theory. *Discrete Mathematics*, 342(10):2911–2923, 2019.
- [20] M. Madiman, L. Wang, and J. O. Woo. Entropy inequalities for sums in prime cyclic groups. *SIAM Journal on Discrete Mathematics*, 35(3):1628–1649, 2021.
- [21] K. Murota. *Discrete convex analysis*. SIAM Monographs on Discrete Mathematics and Applications. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2003.
- [22] C. E. Shannon. A mathematical theory of communication. *The Bell system technical journal*, 27(3):379–423, 1948.
- [23] D. Shlyakhtenko. A free analogue of Shannon’s problem on monotonicity of entropy. *Adv. Math.*, 208(2):824–833, 2007.
- [24] A. J. Stam. Some inequalities satisfied by the quantities of information of Fisher and Shannon. *Information and Control*, 2(2):101–112, 1959.
- [25] T. Tao. Sumset and inverse sumset theory for Shannon entropy. *Combinatorics, Probability and Computing*, 19:603–639, 07 2010.
- [26] A. M. Tulino and S. Verdú. Monotonic decrease of the non-Gaussianness of the sum of independent random variables: a simple proof. *IEEE Trans. Inform. Theory*, 52(9):4295–4297, 2006.
- [27] J. O. Woo and M. Madiman. A discrete entropy power inequality for uniform distributions. In *2015 IEEE International Symposium on Information Theory (ISIT)*, pages 1625–1629. IEEE, 2015.