

VORTICITY BUDGET OF WEAK THERMAL CONVECTION IN KEPLERIAN DISKS

PIN-GAO GU

Department of Physics, University of Texas at Austin, Austin, TX 78712; gu@physics.utexas.edu

Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218

Accepted to ApJ 2000 April 19

ABSTRACT

By employing the equations of mean-square vorticity (enstrophy) fluctuations in strong shear flows, we demonstrate that unlike energy production of turbulent vorticity in non-rotating shear flows, the turbulent vorticity of weak convection in Keplerian disks cannot gain energy from vortex stretching/tilting by background shear unless the associated Reynolds stresses are negative. This is because the epicyclic motion is an energy sink of the radial component of mean-square turbulent vorticity in Keplerian disks when Reynolds stresses are positive. Consequently, weak convection cannot be self-sustained in Keplerian flows. This agrees with the results implied from the equations of mean-square velocity fluctuations in strong shear flows. Our analysis also sheds light on the explanation of the simulation result in which positive kinetic helicity is produced by the Balbus-Hawley instability in a vertically stratified Keplerian disk. We also comment on the possibility of outward angular momentum transport by strong convection based on azimuthal pressure perturbations and directions of energy cascade.

Subject headings: accretion, accretion disks — convection — hydrodynamics — MHD — turbulence

1. INTRODUCTION

Turbulence is usually driven by waves or instabilities in fluids, taping accessible free energy into fluctuating velocity fields. For astronomical objects, accessible free energy can come from body forces (*e.g.*, gravity or Lorentz force) or from large-scale fluid motions (*e.g.*, rotation or shear).

In Keplerian disks, thermal convection is an important ingredient in the thermal-viscous instability model which explains the semi-periodic changes of light curves of dwarf novae (Meyer & Meyer-Hofmeister 1981) and soft X-ray transients (Mineshige & Wheeler 1989). However, it has been argued that convective instability cannot access free thermal energy (*i.e.*, higher entropy near the mid-plane) to transport energy vertically outward if it can access free rotational energy (*i.e.*, higher angular velocities at small radii) to transport angular momentum radially outward. Recently a more general argument has been established to include the turbulence which is not thermally driven and whose azimuthal pressure perturbation is small. The equations of averaged turbulent kinetic energy due to local hydrodynamic mixing can be expressed as (Balbus & Hawley 1998)

$$\begin{aligned} & \frac{\partial}{\partial t} \left\langle \frac{\rho v_r^2}{2} \right\rangle + \nabla \cdot \left\langle \frac{1}{2} \rho v_r^2 \mathbf{v} \right\rangle \\ & = 2\Omega \langle \rho v_r v_\theta \rangle - \left\langle v_r \frac{\partial \delta P}{\partial r} \right\rangle - \text{losses}, \end{aligned} \quad (1)$$

$$\begin{aligned} & \frac{\partial}{\partial t} \left\langle \frac{\rho v_\theta^2}{2} \right\rangle + \nabla \cdot \left\langle \frac{1}{2} \rho v_\theta^2 \mathbf{v} \right\rangle \\ & = -\frac{\kappa^2}{2\Omega} \langle \rho v_r v_\theta \rangle - \left\langle v_\theta \frac{\partial \delta P}{r \partial \theta} \right\rangle - \text{losses}, \end{aligned} \quad (2)$$

where “losses” represent the energy sink due to viscosity,

v_i is the turbulent velocity field, δP is the pressure perturbation, ρ is the mass density, Ω is the Keplerian angular velocity, and the epicyclic frequency $\kappa^2 = (2\Omega/r)d(r^2\Omega)/dr$. The notation $\langle \rangle$ in equations (1) and (2) denotes the averaging carried out over the ensemble of turbulent cells, indicating the time/space correlation between two fluctuating quantities. In the case of accretion disks, the time average is taken over several eddy turnover time (but still smaller than the viscous time scale), and the space average is performed over 2π in the azimuthal direction, over the whole disk vertical scale height¹, and over several eddy sizes in the radial direction (but still much smaller than the disk radius).

As indicated by equation (2), the epicyclic term is an energy sink of $\langle v_\theta^2 \rangle$ if $\langle v_r v_\theta \rangle > 0$. Therefore any local hydrodynamic instabilities cannot grow when azimuthal pressure perturbation is small. If a disk, however, is heated by other sources, the energy of weak convection can be maintained and then transport angular momentum inward. When thermal convection is weak, the associated negative Reynolds stress is a result of conservation of angular momentum of turbulent elements during the process of local mixing in Rayleigh-stable disks such as Keplerian flows (Balbus 2000).

Besides considering the epicyclic effect which damps fluctuating velocity fields, the same damping effect should apply to fluctuating vorticity fields which characterize strong turbulence. It is also worth investigating how fluctuating vorticity fields interact with background vorticity when convection is maintained by other heating sources such as MHD turbulence driven by the Balbus-Hawley instability (Balbus & Hawley 1998), in order to determine the typical linear modes for convection in nonlinear regime (Gu, Vishniac, & Cannizzo 2000, hereafter GVC).

¹Averaging over the disk height is not performed when one would like to investigate the detailed vertical structures (*e.g.*, see Stone *et al.* 1996; Miller & Stone 2000.)

As high levels of fluctuating vorticities are observed in non-rotating shear flows at high Reynolds numbers, we must be wondering why the generation of fluctuating vorticities in Rayleigh-stable disks cannot be achieved by the usual vortex dynamics such as vortex stretching or vortex tilting, as noted and numerically simulated by Hawley, Balbus, & Winters 1999.

Perhaps one of the most difficult tasks in dealing with equations of turbulence is to determine the signs of correlations between fluctuating quantities. Positive, negative, or no correlations are usually related to the properties of background flows such as stratification, rotation, background shear, or body forces. Positive Reynolds stresses in non-rotating shear flows, for example, are always related to the mean flow moving in $+x$ direction with negative gradient in $+y$ direction. This situation is reversed in Keplerian flows for weak convection (Ryu & Goodman 1992; Kley *et al.* 1993; Kumer *et al.* 1995; Cabot 1996; Stone & Balbus 1996; Balbus 2000). Positive kinetic helicity occurring in magnetized disks shown in the numerical simulation by Brandenburg & Donner 1997 probably means that the Coriolis force is of less importance in Keplerian disks where MHD turbulence is driven by the Balbus-Hawley instability. Similar to these examples, we would expect that background shear and rotation can determine the signs of correlations in the equations of fluctuating vorticities.

Beginning with the perturbed vorticity equations in a non-rotating flow, we determine the signs of turbulent correlation based on positive Reynolds stresses without worrying about nonlinear shear instabilities in §2. In §3, we apply the analysis presented in §2 to Keplerian flows, showing that mean-square vorticity perturbations of weak convection cannot grow with positive Reynolds stresses. Positive kinetic helicity driven by the Balbus-Hawley instability in a stratified Keplerian disk is analyzed through vortex equations in §4. In the last section, we comment on the possibility of outward angular momentum by strong convection in accretion disks based on the recent development of theories concerning about azimuthal pressure perturbations and directions of energy cascade.

2. MOMENTUM CONSERVATION AND VORTICITY EQUATIONS IN NON-ROTATING SHEAR FLOWS

The vorticity equation reads

$$\frac{D\mathbf{w}}{Dt} = (\mathbf{w} \cdot \nabla) \mathbf{u} - (\nabla \cdot \mathbf{u}) \mathbf{w} + \frac{\nabla \rho \times \nabla P}{\rho^2} + \nabla \times \nu \nabla^2 \mathbf{u}, \quad (3)$$

where $\mathbf{w} = \nabla \times \mathbf{u}$, $D/Dt = \partial_t + (\mathbf{u} \cdot \nabla)$, P is the pressure, ρ is the mass density, and ν is the kinematic viscosity. Consider a 3-D turbulent flow with a background shear dV/dx without rotation, where $V(x)$ is the background flow in $+y$ direction. Let the vertical scale height be much smaller than the scale heights in x and y directions. If the eddy growth rate is smaller than the shearing rate (*i.e.*, weak convection), linearizing the vorticity equation ($\mathbf{u} = V\hat{j} + \mathbf{v}$, and $\boldsymbol{\omega} = \nabla \times \mathbf{v}$) gives the equations of mean-square vorticity (enstrophy) perturbation:

$$\frac{D}{Dt} \left\langle \frac{\omega_x^2}{2} \right\rangle \approx \left\langle \omega_x \frac{\partial v_x}{\partial z} \right\rangle \frac{dV}{dx}$$

²This constraint is called the “weak-shear limit” in Shu 1974. In fact, we are dealing with convection with the growth rates smaller than the shearing rate in this paper. In order to avoid confusion, we do not use the term “weak-shear” here.

$$- \frac{1}{\rho^2} \frac{\partial \rho}{\partial z} \left\langle \omega_x \frac{\partial \delta P}{\partial y} \right\rangle + \frac{1}{\rho^2} \frac{\partial P}{\partial z} \left\langle \omega_x \frac{\partial \delta \rho}{\partial y} \right\rangle - \text{losses}, \quad (4)$$

$$\begin{aligned} \frac{D}{Dt} \left\langle \frac{\omega_y^2}{2} \right\rangle &\approx \langle \omega_x \omega_y \rangle \frac{dV}{dx} + \left\langle \omega_y \frac{\partial v_y}{\partial z} \right\rangle \frac{dV}{dx} \\ &+ \frac{1}{\rho^2} \frac{\partial \rho}{\partial z} \left\langle \omega_y \frac{\partial \delta P}{\partial x} \right\rangle - \frac{1}{\rho^2} \frac{\partial P}{\partial z} \left\langle \omega_y \frac{\partial \delta \rho}{\partial x} \right\rangle - \text{losses}, \end{aligned} \quad (5)$$

$$\begin{aligned} \frac{D}{Dt} \left\langle \frac{\omega_z^2}{2} \right\rangle &\approx - \langle v_x \omega_z \rangle \frac{d^2 V}{dx^2} \\ &- \left\langle \omega_z \left(\frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} \right) \right\rangle \frac{dV}{dx} - \text{losses}, \end{aligned} \quad (6)$$

where the terms “losses” represent the energy sink due to viscosity, v_i and ω_i are fluctuating velocity and vorticity respectively, and D/Dt denotes $\partial_t + V\partial_y$. We have ignored the term $\langle \omega_i \omega_j \partial_j v_i \rangle$ which represents stochastic stretching of vortex by turbulent shear. This effect is smaller than stretching by background shear as long as background shear rate is larger than turbulent growth rate. Similar to the role played by $\partial \delta P / \partial r$ in equation (1), the terms associated with $\partial \delta P / \partial x$ and $\partial \delta \rho / \partial x$ are energy sources of $\langle \omega_y^2 \rangle$ in equation (5). Since these terms come from the baroclinic term in equation (3), they appear as a result of vertical stratification.

The production or annihilation of turbulent vorticity relies on pressure perturbations, density perturbations, and the signs of correlations between fluctuating vorticities and fluctuating velocity shears as shown in the above equations. However, the term associated with ∂_y is usually small for weak convection as a result of strong shear. Assuming that all perturbation quantities are proportional to $\exp(ik_x x + ik_y y + ik_z z + i\omega t)$, we have the linear perturbation equations for adiabatic convection under Boussinesq approximation:

$$i\bar{\omega} v_x + ik_x \Psi = 0, \quad (7)$$

$$i\bar{\omega} v_y + \frac{dV}{dx} v_x + ik_y \Psi = 0, \quad (8)$$

$$i\bar{\omega} v_z + ik_z \Psi + \delta g_z = 0, \quad (9)$$

$$i\bar{\omega} \delta - v_z \partial_z \ln \left(\frac{P^{1/\Gamma}}{\rho} \right) = 0, \quad (10)$$

$$k_x v_x + k_y v_y + k_z v_z = 0, \quad (11)$$

with the shearing constraint (Shu 1974)²

$$k_y < k_x \frac{i\bar{\omega}}{|dV/dx|}, \quad (12)$$

where $\bar{\omega}$ is the frequency measured by a local observer moving with the mean flow, $\delta \equiv \delta \rho / \rho$ is the local fractional density perturbation, $\Psi \equiv \delta P / \rho$ is the pressure perturbation divided by the density, v_x , v_y , and v_z are velocity perturbations, k_x , k_y and k_z are the wavenumbers in x , y and z directions. The validity of the plane-wave approximation for k_x is assured by the shearing constraint for

thermal convection in the presence of a shear flow. The plane-wave approximation for k_z in the presence of a vertical gradient is not a bad assumption for weak convection since first of all, the dispersion relation (19) inferred from the above Boussinesq equations in the case of accretion disks is the same as the one derived by Ruden, Papaloizou, & Lin 1988, who consider the vertical wavelength of axisymmetrical modes for convection as a function of z owing to vertical stratification. Second of all, as we shall see in the following text, the equations (7)–(12) serve only as an example to illustrate how the phenomenological interpretation works for explaining the directions of angular momentum transport done by Balbus 2000, and how convective eddies approximately look like in a shear flow, all of which are not very sensitive to the detailed vertical structures of disks. We are going to adopt the same phenomenological strategy to evaluate the signs of correlations in the equations of perturbed enstrophy, and consequently to investigate if turbulent vorticities can be generated by vortex stretching/tilting in the regime where the terms associated with ∂_y are small.

By virtue of equations (7) through (12) in the weak-convection limit (*i.e.*, the convective growth rate $i\bar{\omega} < |dV/dx|$), one can show that the terms associated with ∂_y (or say k_θ) in the above Boussinesq equations and in the equations of perturbed enstrophy are smaller than the terms associated with dV/dx , owing to the shearing constraint. Without the pressure perturbation in the y direction, equation (8) states that a turbulent element can conserve momentum in the y direction when it moves.

In the case $dV/dx < 0$, conservation of momentum of turbulent elements in the y direction described by equation (8) implies that $\langle v_x v_y \rangle > 0$, *i.e.*, as observed in the comoving frame, positive (negative) values of v_x should occur more frequently than negative (positive) ones when v_y is positive (negative), or vice versa. By virtue of equation (8), $\langle v_y^2 \rangle$ can grow when $\langle v_x v_y \rangle > 0$ (Balbus & Hawley 1998). The major concern for a successful growth of turbulent vorticity is that the first term of the right hand side in equation (4) must be positive because the other terms associated with ∂_y are small. Since it is the equation (8) which determines the sign of $\langle v_x v_y \rangle$ and the growth of turbulence, we would expect that the same equation plays the same role in equation (4). By virtue of equation (8) again, we have

$$\frac{1}{2} \left\langle \frac{\partial \gamma}{\partial z} \frac{\partial v_y^2}{\partial z} \right\rangle + \gamma \left\langle \left(\frac{\partial v_y}{\partial z} \right)^2 \right\rangle = -\frac{dV}{dx} \left\langle \frac{\partial v_y}{\partial z} \frac{\partial v_x}{\partial z} \right\rangle, \quad (13)$$

where the growth rate $\gamma \equiv i\bar{\omega} > 0$. The first term on the left hand is positive since turbulent velocity fields usually increase with convective growth rate. As a result, the correlation $\langle \partial_z v_y \partial_z v_x \rangle$ is positive. The first term on the right hand side of equation (4) is $\approx -\langle \partial_z v_y \partial_z v_x \rangle dV/dx$ which is therefore positive, leading to successful growth of $\langle \omega_x^2 \rangle$. In fact, this positive sign can be understood in another way. The equation of vorticity $D\omega_x/Dt \approx \partial_z v_x (dV/dx)$ suggests that ω_x tends to be negative (positive) when $\partial_z v_x > 0$ (< 0) and $dV/dx < 0$ (> 0). Hence ω_x and $\partial_z v_x$ are negatively correlated in a non-rotating shear flow. In other words, $\langle \omega_x^2 \rangle$ can grow from the interaction between background (dV/dx) and turbulent ($\partial_z v_x$) shear.

Although the signs of correlations associated with back-

ground shear in equation (5) are not important because there are energy source terms due to pressure/density perturbations, we can possibly determine the sign of $\langle \omega_x \omega_y \rangle$ phenomenologically. In the case of $dV/dx < 0$, if we consider a turbulent element moving toward $+x$ direction with positive sense of ω_y , the element will tend to move toward $+y$ direction more frequently so that the negative sense of ω_x is created as seen in the comoving frame. This effect of vortex rotation is therefore described by the correlation $\langle \omega_x \omega_y \rangle < 0$. The signs is reversed when $dV/dx > 0$. After all, the term $\langle \omega_x \omega_y \rangle dV/dx$ is an energy source of $\langle \omega_z^2 \rangle$.

The term $\langle v_x \omega_z \rangle$ in equation (6) describes the vorticity transport in the flows which have nonuniform background vorticity (*i.e.*, $d^2V/dx^2 \neq 0$). Assume that the angular momentum of vertical vortex tubes are nearly conserved; *i.e.*, the loss term in equation (6) is small. If the gradient of background vorticity d^2V/dx^2 is positive (negative), the local turbulent mixing will transport the vorticity toward $-x$ ($+x$) direction, or vice versa. This means that the term $-\langle v_x \omega_z \rangle d^2V/dx^2 > 0$. However, the production provided by vorticity transport is supposed to be small since v_x and the correlation length across the mean shear flows are severely shortened by strong shear. This point can be also understood in terms of the fact that the term $\langle v_x \omega_z \rangle$ is zero in a shear flow without a gradient of a background state in the x direction. Introducing a gradient of background state gives rise to a factor about one over the scale height along the x direction which is much smaller than one over the vertical scale height associated with other terms in equation (6).

The second term in equation (6) describes vortex generation due to the interaction between background vorticity dV/dx and variation of cross sections of vortices. The equation of motion $D\omega_z/Dt \approx -(\partial_x v_x + \partial_y v_y)(dV/dx)$ suggests that ω_z tends to be negative (positive) when $\partial_x v_x + \partial_y v_y < 0$ (> 0) and $dV/dx < 0$ (> 0). This means that the second term in equation (6) is the energy source of $\langle \omega_z^2 \rangle$. For a nearly incompressible disturbance, $\partial_x v_x + \partial_y v_y < 0$ means vortex stretching (*i.e.*, $\partial_z v_z > 0$), and $\partial_x v_x + \partial_y v_y > 0$ means vortex squeezing (*i.e.*, $\partial_z v_z < 0$) in the vertical direction.

After all, the turbulent vorticities ω_x and ω_y can grow simultaneously via vortex tilting by background shear (dV/dx) and by turbulent shear ($\partial_j v_i$). The turbulent vorticity ω_z can grow via vortex stretching/squeezing. The different vortex dynamics associated with ω_z comes from the fact that ω_z does not couple with ω_x and ω_y as shown in equations (4), (5), and (6). We note that vortex stretching/tilting is a 3-D effect and cannot exist in 2-D fluids ($\omega_x \approx \omega_y \approx 0$) which are usually employed to model vortex generation in larger scales \gg vertical scale height h (*i.e.*, the shallow water approximation). Large-scale vortex production in 2-D usually relies on the baroclinic term in the first place, and subsequently vortexes can evolve by mutual interactions such as merging into large ones which have long lifetimes (Adams & Watkins 1995; Godon & Livio 1999). In this paper, we do not consider this evolution.

Before we apply the same approach to Keplerian flows, we need to emphasize that the above analysis just serves as preliminary calculations for Keplerian flows. The turbulent fields driven by thermal convection in a planar

shear flow are more complicated than those in a Keplerian flow. In the former case, thermal convection excites a non-linear shear instability which overwhelms the convective motion (Stone & Balbus 1996), and therefore the above analysis for thermal convection in a planar shear flow is quite questionable; for example, the term $\langle \omega_i \omega_j \partial_j v_i \rangle$ which we ignored becomes most important (Tennekes & Lumley 1972). In the case of Keplerian flows, however, numerical simulations thus far have shown that strong epicyclic motion suppresses non-linear shear instabilities (Hawley, Balbus, & Winters 1999, however see Richard & Zahn 1999, and Klahr 2000). Hence the above analysis should be reasonable for weak convection in a Keplerian flow. We shall see in the next section that the result implied from the equations of mean-square vorticity perturbations are indeed consistent with the result inferred from the equations of mean-square velocity fluctuations.

3. ANGULAR MOMENTUM CONSERVATION AND VORTICITY EQUATIONS IN A KEPLERIAN FLOW

In Keplerian flows, the vorticity equations in cylindrical coordinates become

$$\begin{aligned} \frac{D}{Dt} \left\langle \frac{\omega_r^2}{2} \right\rangle &\approx \left\langle \omega_r \frac{\partial v_r}{\partial z} \right\rangle \frac{\kappa^2}{2\Omega} \\ &- \frac{1}{\rho^2} \frac{\partial \rho}{\partial z} \left\langle \omega_r \frac{\partial \delta P}{r \partial \theta} \right\rangle - \text{losses}, \end{aligned} \quad (14)$$

$$\begin{aligned} \frac{D}{Dt} \left\langle \frac{\omega_\theta^2}{2} \right\rangle &\approx \langle \omega_r \omega_\theta \rangle r \frac{d\Omega}{dr} \\ &+ \left\langle \omega_\theta \frac{\partial v_\theta}{\partial z} \right\rangle \frac{\kappa^2}{2\Omega} + \frac{1}{\rho^2} \frac{\partial \rho}{\partial z} \left\langle \omega_\theta \frac{\partial \delta P}{\partial r} \right\rangle - \text{losses}, \end{aligned} \quad (15)$$

$$\begin{aligned} \frac{D}{Dt} \left\langle \frac{\omega_z^2}{2} \right\rangle &\approx - \langle v_r \omega_z \rangle \frac{d}{dr} \left(\frac{\kappa^2}{2\Omega} \right) \\ &- \left\langle \omega_z \left(\frac{\partial v_r}{\partial r} + \frac{\partial v_\theta}{r \partial \theta} \right) \right\rangle \frac{\kappa^2}{2\Omega} - \text{losses}, \end{aligned} \quad (16)$$

where D/Dt denotes $\partial_t + \Omega \partial_\theta$. In a Keplerian disk threaded with sub-thermal magnetic fields, the ‘‘loss’’ terms in above equations include the turbulent damping due to radial mixing driven by the Balbus-Hawley instability. We do not show the terms associated with density perturbations explicitly because unlike $\delta P/P \lesssim \delta \rho/\rho$ in non-rotating shear flows owing to equation (7), $\delta P/P \sim \delta \rho/\rho$ in rotating shear flows due to the Coriolis force (see equation (18) below). Once again ω_z does not couple with ω_r and ω_θ .

Although perturbation equations (7) through (12) should be modified by adding the Coriolis force in a rotating flow, the Coriolis term 2Ω is comparable to the shear term $r d\Omega/dr$. Therefore the terms associated with ∂_θ are small compared to the first term on the right hand side in equation (14), analogous to equation (4). In a Keplerian flow, neglecting azimuthal pressure perturbations means that turbulent elements conserve angular momenta before they mix with background fluid. This leads to a transport of angular momentum down to the angular momentum gradient, giving rise to negative Reynolds stresses in a Keplerian disk (Balbus 2000). This picture is described by

the azimuthal perturbed equation of motion which is similar to equation (8) except dV/dx is replaced by $\kappa^2/2\Omega$; *i.e.*,

$$\gamma v_\theta = - \frac{\kappa^2}{2\Omega} v_r = - \left(r \frac{d\Omega}{dr} + 2\Omega \right) v_r, \quad (17)$$

where we have dropped the term $ik_\theta \Psi$. The above equation indicates that $\langle v_\theta^2 \rangle$ cannot grow when $\langle v_r v_\theta \rangle > 0$ (Balbus & Hawley 1998), which is different from the situation in a non-rotating shear flow owing to the appearance of the Coriolis force.

In fact, negative Reynolds stresses can be realized as follows. Consider a turbulent element is moving with positive v_r . While shear ($r d\Omega/dr$) tries to move the element to the $+\hat{\theta}$, as observed in a corotating frame, the Coriolis force (2Ω) tried to move the element to the $-\hat{\theta}$. Since $2\Omega > r |d\Omega/dr|$ in Keplerian disks (*i.e.*, $\kappa^2 > 0$), the Coriolis force is the winner. Therefore the turbulent element tends to have $-v_\theta$, leading to $\langle v_r v_\theta \rangle < 0$. Without rotation (Coriolis effect), $\langle v_x v_y \rangle > 0$ in a non-rotating shear flow with negative dV/dx . The contribution to $\langle v_r v_\theta \rangle$ from the radial deviation of turbulent azimuthal motion due to the Coriolis force is small since for weak convection in a Keplerian disk, equation (7) becomes

$$ik_r \Psi \sim 2\Omega v_\theta, \quad (18)$$

where the term $i\bar{\omega} v_r$ is small and has been ignored. This means that the strong radial gradient of pressure perturbation is roughly balanced by the Coriolis force, resulting in negligible contribution to $\langle v_r v_\theta \rangle$. In other words, $\langle v_r v_\theta \rangle$ is mostly determined by equation (17). We note that in contrast to equation (7), the extra term due to the Coriolis force in equation (18) indicates that the radial gradient of pressure perturbation should be large enough to make convection grow in a rotating flow. Namely, the Coriolis force is a stabilized factor to convection. Owing to equations (17) and (18), the linear theory without radiative and turbulent damping indicates that γ^2 is smaller than the Brunt-Väisälä frequency $-N^2$ (GVC; Ruden *et al.* 1988):

$$\gamma^2 = \frac{-N^2 - A\Omega^2}{A + 1}, \quad (19)$$

where $A \equiv (k_z/k_r)^2$. It has been known in terrestrial experiments that the critical Rayleigh numbers of rotating Rayleigh-Benard convection are increased above their non-rotating values (*e.g.*, Zhong *et al.* 1993).

The above analysis means that for weak convection, the signs of correlations in Keplerian flows in equations (14) (15), and (16) are reversed compared with those in the non-rotating flows with $dV/dx < 0$ when we replace $x - y$ with $r - \theta$. In other words, we have $\langle v_r v_\theta \rangle < 0$, $\langle \omega_r \omega_\theta \rangle > 0$, $\langle \omega_r \partial_z v_r \rangle > 0$, and $\langle \omega_z (\partial_r v_r + \partial_\theta v_\theta/r) \rangle < 0$ when $\kappa^2 > 0$. The right hand side of equation (14) becomes positive and $\langle \omega_r^2 \rangle$ can grow. Although the first term on the right hand side of equation (15) turns out to be negative, $\langle \omega_\theta^2 \rangle$ can be maintained by the radial gradient of perturbed pressure owing to strong epicyclic effect.

If we assume that weak convection transports angular momentum down to the angular velocity gradient, then the signs of correlations in equation (14) and equation (15) are

the same as those in the non-rotating flows with $dV/dx < 0$ when we replace $x - y$ with $r - \theta$. This means that the first term on the right hand of equation (14) is negative because $\kappa^2 > 0$. Hence $\langle \omega_r^2 \rangle$ cannot grow and weak convection dies away.

If the disks are Rayleigh-unstable (*i.e.*, $\kappa^2 < 0$), the signs of all terms in equations (14), (15), and (16) remain unchanged compared to those with positive κ^2 except that the first term on the right hand side in equation (15) becomes positive because $\langle \omega_r \omega_\theta \rangle < 0$. In other words, a Rayleigh-unstable disk behaves as a non-rotating shear flow in the sense that $\langle v_r v_\theta \rangle > 0$. We note that our analysis for Rayleigh-unstable disks is oversimplified because a non-linear shear instability should be excited.

4. POSITIVE KINETIC HELICITY AS A RESULT OF THE BALBUS-HAWLEY INSTABILITY

A weakly magnetized Keplerian disk is linearly unstable to the Balbus-Hawley instability. In nonlinear regime, this instability can drive MHD turbulence which in turn leads to a dynamo process unless the magnetic Reynolds numbers are low (Balbus & Hawley 1998). Brandenburg & Donner 1997 shows that this dynamo process observed in simulations can be imitated by an $\alpha - \Omega$ dynamo in a vertically stratified disk, with a negative $\alpha_{\theta\theta}$ in the upper disk plane (where $\alpha_{\theta\theta}$ is the θ - θ component of kinetic helicity tensor α). A negative $\alpha_{\theta\theta}$ means that the kinetic helicity $\langle v_z \omega_z \rangle$ is positive for nearly isotropic turbulence. This contradicts with the usual notion that the kinetic helicity of convection driven by the Coriolis force is usually negative. Brandenburg 1998 suggests that the right-handed helical turbulence results from the combined effect of the Balbus-Hawley instability (*i.e.*, $\langle \delta B_r \delta B_\theta \rangle < 0$) and magnetic buoyancy.

Although we have concentrated on convection-like eddies and have not taken into account the effect of subthermal magnetic fields in this paper, equations (6) and (16) could shed light on preferred directions of helical turbulence if we realize that a Coriolis-dominated disk characterized by equation (16) becomes a shear-dominated disk symbolized by equation (6) due to the Balbus-Hawley instability. As explained in the preceding section, $\langle \omega_z (\partial_r v_r + \partial_\theta v_\theta / r) \rangle < 0$ in a nonmagnetized Keplerian disk. Assume that a turbulent element expands (contracts) as it goes upward (downward) in the upper disk plane. The above negative correlation means that a turbulent element tends to have a negative (positive) ω_z due to expansion (contraction). Consequently, $\langle v_z \omega_z \rangle < 0$ (see the vortex plot associated with positive κ^2 in Figure 1). This result is consistent with the picture in which the Coriolis force is the winner over shear so that turbulent eddies are mostly left-handed; *i.e.*, as observed in a corotating frame, left-handed helical turbulent motion results from a positive vorticity ($\kappa^2 > 0$) of the mean flow. However, introducing the Balbus-Hawley instability changes the sign of Reynolds stresses from negative to positive in Keplerian disks because angular momentum of a turbulent element is changed by fluctuating magnetic torques (Balbus 2000), leading to the condition for the instability changed from the Rayleigh criterion $\kappa^2 < 0$ to the Chandrasekhar-Velikov criterion $d\Omega^2/dr < 0$ (Velikhov 1959; Chandrasekhar 1960; Fricke 1969) in a Keplerian flow. Equivalently speaking, the Velikov-Chandrasekhar

criterion is a manifestation of conservation of angular velocity of turbulent elements due to the fact that the fluctuating magnetic torque counteracts the torque of the Coriolis force in a weak field limit (Fricke 1969). Without the torque by the Coriolis force, azimuthal dynamics of turbulence in Keplerian disks becomes that in a non-rotating shear flow. Furthermore, if the contribution to $\langle v_r v_\theta \rangle$ due to the radial deviation of azimuthal motion is small (like the situation described by equation (18); radial deviation of azimuthal motion is usually small owing to shear), then the signs of correlation quantities in a weakly magnetized disk should behave as the one described by negative vorticity of the mean flow (such as the correlation quantities in equations (4), (5), and (6)), giving rise to a positive kinetic helicity (see the vortex picture associated with negative dV/dx in Figure 1). The only physical difference of vortex dynamics between a non-rotating shear flow and a weakly magnetized disk is that in the later case, a positive kinetic helicity coupled with a positive vorticity of the mean flow becomes an energy sink of $\langle \omega_z^2 \rangle$ as shown in equation (16). This is analogous to the result that a positive Reynolds stress is an energy sink of v_θ^2 in Keplerian disks (Hawley, Balbus, & Winters 1999).




Fig. 1. The expansion of a rising turbulent element described by equation (16) or by equation (6) when $\kappa^2 > 0$ or $dV/dx > 0$ (left), and when $\kappa^2 < 0$ or $dV/dx < 0$ (right). As observed in a corotating frame, $\langle \omega_z (\partial_r v_r + \partial_\theta v_\theta / r) \rangle < 0$ and $\langle v_z \omega_z \rangle < 0$ when $\kappa^2 > 0$. These signs are reversed when $\kappa^2 < 0$. A negative (positive) ω_z , as a result of a positive (negative) vorticity of the mean flow, can give rise to a negative (positive) $\langle v_r v_\theta \rangle$ when radial deviation of azimuthal motion is small. As argued in the text, the picture on the right could also apply to the ω_z produced by the Balbus-Hawley instability which generates positive Reynolds stresses in Keplerian disks. In the magnetic case, however, the vortex motion illustrated in the figure becomes an energy sink of $\langle \omega_z^2 \rangle$. $k_\theta < k_r$ is a result of shear.

Although our analysis is related to convection-like eddies (*i.e.*, $k_z \sim 1/h$), the picture we present here for estimating the sign of kinetic helicity does not contradict the result that the Parker instability, a magnetohydrodynamic instability with $k_z \sim 1/h$, is suppressed by MHD turbulence (Vishniac & Diamond 1992). The main point presented here depends solely on the sign of Reynolds stresses which is assumed to be largely caused by the azimuthal deviation of radial motion. Therefore, the picture that we sketch here applies to any turbulent eddies which expand (contract) as they rise (descent). We note that rising turbulent eddies can expand statistically even without any types of buoyancy. As a matter of fact, the vertical structures of large-scale magnetic fields generated by the Balbus-

Hawley instability in vertically stratified accretion disks are stable against magnetic buoyancy (Stone *et al.* 1996; Miller & Stone 2000). The detailed information, such as the magnitude of this kinetic helicity and the expansion of turbulent eddies in a stably stratified disk, should depend strongly on the detailed structure of the flow, such as azimuthal perturbations (*i.e.*, anisotropic turbulence) and correct vertical dynamics.

5. DISCUSSION AND OUTLOOK: COMMENTS ON ANGULAR MOMENTUM TRANSPORT BY STRONG CONVECTION

In this paper, we demonstrate that weak convection cannot be self-sustained by vortex stretching and tilting in a Keplerian flow, suggested by the equations of enstrophy fluctuations. This is because epicyclic term is an energy sink of $\langle \omega_r^2 \rangle$ when $\langle v_r v_\theta \rangle > 0$. The situation is reversed when weak convection has a negative Reynolds stress, leading to a successful growth of thermal convection. Our determination of the signs of correlation quantities in the equations of enstrophy is based on the point of view in which the signs of Reynolds stresses are determined by conservation of angular momentum for weak convection (Balbus 2000). Although our determination of signs of correlation quantities is not a rigorous proof but quite suggestive, the results are consistent with those suggested by the equations of velocity fluctuations. Full three-dimensional numerical simulations are needed to verify our thought.

Positive kinetic helicity in a weakly magnetized, vertically stratified Keplerian disk implied from our analysis agrees with the simulation result by Brandenburg & Donner 1997. However, the physical picture shown in Figure 1 is totally different from the plot sketched in Figure 7 in their paper: in a corotating frame of a Keplerian disk, the direction of Coriolis effect (2Ω) should be opposite to the direction of shear effect ($d\Omega/dr$), resulting in a positive vorticity ($\kappa^2 > 0$). In our analysis for turbulence, $\langle w_z(\partial_r v_r + \partial_\theta v_\theta/r) \rangle < 0$ (> 0) when vorticity is positive (negative), giving rise to negative (positive) $\langle v_z \omega_z \rangle$ as long as $\langle v_z(\partial_r v_r + \partial_\theta v_\theta/r) \rangle$ is positive. Since the Balbus-Hawley instability generates a positive $\langle v_r v_\theta \rangle$ which resembles the hydrodynamic turbulence with a negative vorticity, MHD turbulence driven by the Balbus-Hawley instability should display right-handed helical motion as observed in numerical simulations. However, if rising eddies contract statistically, our analysis will give rise to the conventional sign of kinetic helicities in a rotating fluid influenced mostly by the Coriolis force. The expansion of rising turbulent elements in a stably stratified disk could be just a result of a special profile of underlying background states, or might result from the possible scenario that turbulent mixing, an effect definitely not included in usual laminar analyses, alters the condition of flux freezing in some degree. On the contrary, the simple explanation by Brandenburg 1998 relying on magnetic buoyancy is not consistent with the simulations for vertically stratified accretion disks stirred by the Balbus-Hawley instability.

GVC studied convective energy transport in Keplerian disks stirred by the Balbus-Hawley instability. The typical modes of convection are determined from linear perturbation theory and nonlinear saturation which is caused by the secondary Kelvin-Helmholtz instability (*i.e.*, interaction between convective shear and background vorticity)

and by MHD turbulent damping. From equations (14), (15), and (16), we have seen how convective vorticity/shear possibly interacts with background vorticity/shear and how this interaction becomes an energy sink of turbulent vorticity. This means that the quasi-linear approach for the mixing-length theory by GVC based upon the secondary Kelvin-Helmholtz instability is qualitatively correct.

However, the issue of angular momentum transport by weak convection is probably not of great importance in reality. According to the thermal-viscous instability model, weak convection occurs in Keplerian disks when the Shakura-Sunyaev viscosity α_{SS} (Shakura & Sunyaev 1973) is large ~ 0.1 , as a result of strong radiative losses (GVC; Cabot 1996). Furthermore, based on GVC, large α_{SS} means strong MHD turbulent mixing driven by the Balbus-Hawley instability, which suppresses weak convection by smoothing away momentum and entropy anisotropies associated with convective bubbles mostly in the radial direction (see equation (19) below): $A \ll 1$ when $-N^2 \ll \Omega$. Consequently, only the convection with larger Rossby numbers $Ro \sim 1$ can survive in a magnetized Keplerian disk. Weak convection can occur at smaller radii in protostellar disks (*e.g.*, see D'Alessio *et al.* 1998). In such a cold disk, a significant ‘dead zone’ implies that the Balbus-Hawley instability may not function properly, but accretion might occur near the surfaces of disks where cosmic rays (Gammie 1996), irradiation (Glassgold *et al.* 1997), or rigorous heating beyond the bottom of photosphere as manifested by protostellar winds and flares might increase the ionization level beyond the critical values (Gammie and Menou 1998). However, the heat due to layer accretion should reduce the temperature gradient of the ‘dead zone’ in the vertical direction, possibly leading to the suppression of weak convection as a result of enhanced radiative losses in the radial direction (GVC). Therefore, the question becomes, does strong convection transport angular momentum inward or outward? Low Reynolds number simulation by Cabot & Pollock 1992 found strong convection transports angular momentum outward.

Azimuthal pressure perturbations have drawn large attention in the literature recently, as suggested by the equations of mean-square velocity perturbation and by, in this paper, the equations of enstrophy. Axisymmetry/nonaxisymmetry of convection is actually an intimate issue of the damping of convection due to MHD turbulent mixing (Klahr, Henning, & Kley 1999). When MHD turbulent mixing is large (*i.e.*, large α_{SS}), modes with large k_r are all suppressed owing to strong radial mixing. Similarly, modes with large k_r should also succumb to strong radiative losses in radial direction. Modes with small k_r can survive in a shearing environment only if k_θ is also very small, suggesting nearly axisymmetrical convection. As mentioned in the preceding paragraph, weak convection occurs when α_{SS} is large. Nearly axisymmetrical convection is therefore a manifestation of weak convection which struggles with turbulent and radiative damping in a differentially rotating flow. In other words, nonaxisymmetry of strong ($Ro \sim 1$) convection means that azimuthal pressure perturbations could be important to violation of conservation of angular momentum. If this happens, the equations of velocity and vorticity fluctuations could allow strong

convection to transport angular momentum radially outward. Without a doubt, the quasi-linear analysis based on equations of velocity and vorticity fluctuations is very suggestive.

The other point of view regarding the nature of positive/negative viscosity is the direction of turbulent energy cascade. Negative eddy viscosity manifests the process in which turbulent fields do not only extract energy from the mean flow, but the extracted energy is also passed up to larger scales and finally goes back to the mean flow (Starr 1968). The connection between negative viscosity and inverse energy cascade in two-dimensional flows has been studied in fluid society (Kraichnan 1976; Pouquet 1978; Chechkin *et al.* 1999). Cabot 1996 and Klahr *et al.* suggested that negative viscosity is a result of an inverse energy cascade of nearly axisymmetrical convection and that positive viscosity results from nonaxisymmetrical convection in which 3D hydrodynamical turbulence is resumed. In fact, the collapse of 3D hydrodynamical turbulence to 2D is not necessarily related only to axisymmetrical patterns, but is a natural result of turbulent anisotropy due to strong body forces, which has been manifested by ample examples in geophysics. Hossain 1994 found that when a strong rigid rotation ($Ro < 1$) is turned on, the turbulent velocity fields perpendicular to the direction of rotation (*i.e.* $x - y$ plane) are strongly correlated along the direction of rotation (*i.e.* z direction). Therefore 3D turbulence (direct cascade) reduces to an approximate 2D state (inverse cascade). In the case of accretion disks, epicyclic effect should enforce turbulent fields of weak convection to have an approximate 2D state, giving rise to negative viscosity due to an inverse energy cascade. If convection is strong, however, the collapse to 2D breaks down and this probably results in positive/zero eddy viscosity due to a direct energy cascade.

The numerical simulation by Klahr 2000 found that the gradient of fluctuating angular momentum generated by weak convection could flatten background gradient of angular momentum. Consequently, a secondary instability occurs at a fast rate of some fraction of Ω . This induced instability overwhelms weak convection and drives angular momentum outward. In terms of the quasi-linear analysis based on velocity and vorticity fluctuations, outward angular momentum transport by this secondary instability is a result of strong azimuthal pressure perturbations and stochastic stretching of vortex by turbulent shear. By virtue of nonlinear phenomena of turbulent cascade, this fast secondary instability diminishes epicyclic effect and thereby forces fluid back to an approximate 3-D hydrodynamical state.

If thermal convection generates positive eddy viscosity in Keplerian disks, it does not necessarily mean that it can be self-sustained. It has been a concern that a self-sustained convection would violate the second law of thermodynamics: in the case of a system in which the eddies are thermally driven, the heat dissipated from the mean flow cannot again be used to drive the eddies (Starr 1968). If thermal convection can be self-maintained in Keplerian disks, this could mean that convection is not *totally* thermally driven, in the sense that it can access the available rotational energy at smaller radii via secondary or nonlinear instabilities (Kumer *et al.* 1995; Richard & Zahn 1999). However, the reverse of the above statement is not

necessarily true. Klahr 2000 shows that circumstellar disks with outward transport of angular momentum by convection cool continuously throughout the simulation.

Direction of angular momentum transport by convection is also an important issue in advection-dominated accretion flows (ADAFs). When α_{SS} (not due to convection) is small, strong convection ($Ro \sim 1$) occurs when convection transports angular momentum inward (or outward less efficiently than it transports energy), and accretion is suppressed in ADAFs (Narayan, Igumenshchev, & Abramowicz 2000; Quataert & Gruzinov 2000). However, as noted in this section, strong convection is unlikely to be nearly axisymmetrical and is doubtful to be able to collapse to a 2D state especially in a thick disk where the direction of strong entropy gradient has a component along angular momentum gradient. Numerical observations of a transition of direction of angular momentum (if it exists), a transition of direction of energy cascade, and a transition of pattern formation for convection in both Keplerian accretion disks and ADAFs is worthwhile when Ro and Ra are the control parameters, where Ra should be the ‘‘turbulent’’ Rayleigh number characterized by radiative/advection losses and MHD turbulent damping.

While the *perturbed* baroclinic terms in the equations of perturbed enstrophy act as the energy source of fluctuating vorticities, we do not consider the terms associated with the *background* baroclinic term in this paper, due to the thin-disk approximation. That is, Ω does not vary with z dramatically in thin disks, resulting in a barotropic and Keplerian disk (*e.g.*, see Frank, King, & Raine 1992). The background baroclinic effect is usually of less importance on local faster turbulence in disks; for instance, the Biermann battery owing to the same mechanism is usually ignored when a mean-field dynamo is able to operate via mean kinetic helicities driven by local turbulent events such as thermal convection in stars or supernovae/stellar winds in galactic disks. In spite of their appearance in thick disks which behave as fast rotating stars, it is also quite doubtful that the background baroclinic term can have important influence on the onset of convective instability in these cases. As long as the convective growth rate is not too small, the effect of large scale circulation due to the baroclinic term to smooth away entropy inhomogeneity associated with small-scale mixing such as thermal convection in this case is usually negligible, compared with the same damping mechanism achieved by MHD turbulence. On the other hand, as we mention in the end of §2, a large-scale vortex can be initiated from the background baroclinic effect. Large-scale eddies could be also formed via merging convective vortexes (Zhong *et al.* 1993; Godon & Livio 1999) or through helicity fluctuations of convection (Branover *et al.* 1999). The nonlinear interaction between large-scale vortexes and convective vorticities is worth investigating.

Finally we note that our analysis in this paper is applied to local turbulent mixing in accretion disks, such as thermal convection and the Balbus-Hawley instability. Turbulent transport mediated by global waves, such as internal, Rossby, or spiral shock waves, is not subject to our analysis. Moreover, convective turbulence might behave differently in the environment of MHD turbulence since ideal invariants during turbulent cascades are different between hydrodynamical and MHD turbulence (*e.g.*, see Biskamp

1997).

We are deeply indebted to Ethan T. Vishniac for many useful discussions, and for providing the information about Klahr 2000. We are also grateful to the anonymous referee

for insightful comments and suggestions. This work was completed in the High Energy Physics Laboratory at The University of Texas at Austin, and we would like to thank Roy Schwitters for his generous hospitality.

REFERENCES

- Adams, F. C. & Watkins, R. 1995, *ApJ*, 451, 314
 Balbus, S. A. 2000, *ApJ*, in press, (astro-ph/9906315)
 Balbus, S. A. & Hawley, J. F. 1998, *Rev. Mod. Phys.*, 70, 1
 Biskamp, D. 1997, *Nonlinear Magnetohydrodynamics*, Cambridge Univ. Press
 Brandenburg, A. 1998, *Theory of Black Hole Accretion Disks (Cambridge Contemporary Astrophysics)* ed. M. A. Abramowicz, G. Bjornsson, & J. Pringle, pp. 61-90. Cambridge Univ. Press
 Brandenburg, A. and Donner, K. J. 1997, *MNRAS*, 288, L29
 Branover, H., Eidelman, A., Golbraikh, E., & Moiseev, S. 1999, *Turbulence and Structures*, Academic Press
 Cabot, W. 1996, *ApJ*, 465, 874
 Cabot, W., & Pollack, J. B. 1992, *Geophys. Astrophys. Fluid Dyn.*, 64, 97
 Chandrasekhar, S. 1960, *Proc. Nat. Acad. Sci.*, 46, 253
 Chechkin, A. V., Kopp, M. I., & Yanovsky, V. V. 1999, *JETP*, 86, 357
 D'Alessio, P., Cantó, J., Calvet, N., & Lizano, S. 1998, *ApJ*, 500, 411
 Frank, J., King, A., & Raine D. 1992, *Accretion Power in Astrophysics*, Cambridge Univ. Press
 Fricke, K. 1969, *A&A*, 1, 388
 Gammie, C. F. 1996, *ApJ*, 457, 355
 Gammie, C.F., and Menou, K. 1998, *ApJ*, 492, L75
 Glassgold, A. E., Najita, J., & Igea, J. 1997, *ApJ*, 480, 344 (Erratum: *ApJ*, 485, 920)
 Godon, P. & Livio, M. 1999 *ApJ*, 523, 350
 Gu, P.-G., Vishniac, E. T., & Cannizzo, J. K. 2000, *ApJ*, 534, 380 (GVC)
 Hawley, J. F., Balbus, S. A., & Winters, W. F. 1999, *ApJ*, 518, 394
 Hossain, M. 1994, *Phys. Fluids*, 6, 1077
 Klahr, H. H. 2000, a talk in the meeting "Physics of Accretion and Associated Outflows" in Copenhagen
 Klahr, H. H., Henning, Th., & Kley, W. 1999, *ApJ*, 514, 325
 Kley, W., Papaloizou, J. C. B., & Lin, D. N. C. 1993, *ApJ*, 416, 679
 Kraichnan, R. H. 1976, *J. Atmos. Sci.*, 33, 1521
 Kumar, P., Narayan, R., & Loeb, A. 1995, *ApJ*, 453, 480
 Meyer, F. & Meyer-Hofmeister, E. 1981, *A&A*, 104L, 10
 Miller, K. A., & Stone, J. M. 2000, *ApJ*, in press, (astro-ph/9912135)
 Mineshige, S. & Wheeler, J. C. 1989, *ApJ*, 343, 241
 Narayan, R., Igumenshchev, I. V., & Abramowicz, M. A. 2000, *ApJ*, in press, (astro-ph/9912449)
 Pouquet, A. 1978, *J. Fluid Mech.*, 88, 1
 Quataert, E. & Gruzinov, A., 2000, *ApJ*, in press, (astro-ph/9912440)
 Richard, D., & Zahn, J. 1999, *A&A*, 347, 734
 Ruden, S. P., Papaloizou, J. C. B., & Lin, D. N. C. 1988, *ApJ*, 329, 739
 Ryu, D., & Goodman, J. 1992, *ApJ*, 388, 438
 Starr, V. P. 1968, *Physics of Negative Viscosity Phenomena*, McGraw-Hill, New York
 Shakura, N. I., and Sunyaev, R. A. 1973, *A&A*, 24, 337
 Shu, F. H. 1974, *A&A*, 33, 55
 Stone, J. M. & Balbus S. A. 1996, *ApJ*, 464, 364
 Stone, J., Hawley, J.F., Gammie, C.F., & Balbus, S.A. 1996, *ApJ*, 463, 656
 Tennekes, H., & Lumley, J. L. 1972, *A First Course in Turbulence*, MIT Press, Cambridge, Mass.
 Velikhov, E. P. 1959, *Soviet Phys. - JETP*, 36, 1398
 Vishniac, E. T., & Diamond, P. H. 1992, *ApJ*, 398, 561
 Zhong, F., Ecke, R. E., & Steinberg, V. 1993, *J. Fluid Mech.*, 249, 135