

A New Association Of Post-T Tauri Stars Near The Sun ¹

Carlos A. O. Torres², Licio da Silva³, Germano R. Quast², Ramiro de la Reza³ and Evgueni Jilinski^{3,4}

beto@lna.br, ldasilva@eso.org, germano@lna.br, delareza@on.br, jilinski@on.br

ABSTRACT

Observing ROSAT sources in an area $20^\circ \times 25^\circ$ centered at the high latitude ($b = -59^\circ$) active star ER Eri, we found evidences for a nearby association, that we call the Horologium Association (HorA), formed by at least 10 very young stars, some of them being bona fide Post-T Tauri stars. We suggest other six stars as possible members of this proposed association. We examine several requirements that characterize a young stellar association. Although no one of them, isolated, gives an undisputed prove of the existence of the HorA, all together practically create a strong evidence for it. In fact, the Li line intensities are between those of the older classical T Tauri stars and the ones of the Local Association stars.

The space velocity components, of the HorA relative to the Sun ($U = -9.5 \pm 1.0$, $V = -20.9 \pm 1.1$, $W = -2.1 \pm 1.9$) are not far from those of the Local Association, so that it could be one of its last episodes of star formation. In this region of the sky there are some hotter and non-X-ray active stars, with similar space velocities, that could be the massive members of the HorA, among them, the nearby Be star Achernar. The maximum of the mass distribution function of its probable members is around $0.7 - 0.9 M_\odot$. We estimate its distance as ~ 60 pc and its size as ~ 50 pc. If spherical, this size would be larger than the surveyed area and many other members could have been missed.

We also observed 3 control regions, two at northern and southern galactic latitudes and a third one in the known TW Hya Association (TWA), and the properties and distribution of their young stars strengthen the reality of the HorA. Contrary to the TWA, the only known binaries in the HorA are 2 very wide systems. The HorA is much more isolated from clouds and older (~ 30 Myr) than the TWA and could give some clues about the lifetime of the disks around T Tauri stars. Actually, none of the proposed members is an IRAS source indicating an advanced stage of the evolution of their primitive accreting disks.

ER Eri itself was found to be a RSCVn-like system.

Subject headings: (Galaxy:) open clusters and associations: individual (Horologium Association) — (Galaxy:) solar neighborhood — stars: pre-main sequence — X-rays

1. INTRODUCTION

Surveys of young pre-main-sequence (PMS) stars based on IRAS colors detect mainly the classical T Tauri stars (CTT), with ages less than 10 Myr, due to their important dusty accretion disks. Among them, one of the most comprehensive was the “Pico dos Dias Survey” (PDS) (Gregorio-Hetem et al. 1992; Torres et al. 1995; Torres 1998). The weak T Tauri stars (WTT) are mainly detected by the X-rays produced in

¹Based on observations made under the Observatório Nacional-ESO agreement for the joint operation of the 1.52 m ESO telescope and at the Observatório do Pico dos Dias, operated by MCT/Laboratório Nacional de Astrofísica, Brazil

²Laboratório Nacional de Astrofísica/MCT, CP 21, 37504-360 Itajubá, MG, Brazil

³Observatório Nacional/MCT, Rua General José Cristiano 77, 20921-030, Rio de Janeiro, Brazil

⁴Pulkovo Observatory, St. Petersburg, Russia

their active coronae, since their disks in general contain no more sufficient dust. Finally, there should exist even more evolved young stars, with masses smaller or comparable to the Sun and ages between about 10 to 70 Myr - the so-called Post-T Tauri stars (PTT). If star formation were constant during the last 100 Myr, a large number of uniformly distributed PTT should exist in the solar neighborhood. Nevertheless, the first searches for PTT resulted in very few objects (Herbig 1978). This lack of discoveries may indicate that star formation was not constant in time. However, as the disks around PTT are even more evolved than those around WTT, they should have been dissipated or agglutinated into planetesimals and they hardly would be detected in surveys based on the far infrared or even on H_α emission, as it can be very weak in PTT.

The detection of X-ray sources by the ROSAT All-Sky Survey (RASS) (Neuhäuser 1997) outside important star formation regions gave some hope that these are the expected population of PTT. Nevertheless, several considerations cast doubts about the PTT nature of those dispersed objects (Briceño et al. 1997; Palla & Galli 1997), indicating that most of them could be active main sequence (MS) stars. Jeffries (1995) suggests that most of them are connected with the Local Association. On the other hand, Frinck (1999) found that some of those objects could be PMS stars ejected from forming clouds. She also notes that the WTT around the Chamaleon region appear to be formed in small short-lived cloudlets (Feigelson 1996). In any case, this large population of X-ray sources around star forming clouds seems to be represented by a mixture of PMS and young MS objects.

Do genuine PTT exist and where are they? The main point of this work is that PTT can be found in physically characterized dispersed groups, with ages larger than the mean lifetime of the original clouds. In such case they may be located far from any cloud.

In fact, we discovered the first of this kind of association when we were searching for new T Tauri stars among IRAS point sources in a five degree radius around TW Hya, an already known isolated CTT at high galactic latitude ($b = 23^\circ$) (de la Reza et al. 1989; Gregorio-Hetem et al. 1992). Hipparcos parallaxes show that this is a

nearby association, the distance of TW Hya being 56.4 ± 7 pc (Wichmann et al. 1998). Several research groups became interested in this association and other members were found [see. e.g., Hoff et al. (1998); Jensen et al. (1998); Webb et al. (1999); Jayawardhana et al. (1999b)]. Today nearly 13 young systems totalling 21 stars and a brown dwarf are known to belong to the TW Hya association (TWA) (Webb et al. 1999; Sterzik et al. 1999) and this constitutes a sufficiently large density of PMS stars in a localized region of the sky to consider it as a real cluster. Furthermore, the stars appear to have a common origin (Kastner et al. 1997; Webb et al. 1999) and the best age estimated for the association is ~ 10 Myr (Soderblom et al. 1998; Webb et al. 1999).

Other groups of WTT, apparently isolated from clouds, have been detected at larger distances, as the η Cha cluster at 100 pc (Mamajek et al. 1999) or the one in front of the translucent clouds MBM7 and MBM55 (Hearty et al. 1999). The former appears to be more clearly defined, being nearly 40 times more compact than the TWA and seems to be related to the Sco-Cen OB association. Studies of this kind of clusters are important to understand the beginning of dispersion of regions containing also massive stars. We propose in this paper the existence of a new nearby association around ER Eri (PDS 1 = Hen 1), older and more isolated from clouds than the TWA.

2. SEARCH STRATEGY AND OBSERVATIONS

The high galactic latitude ($b = -59^\circ$) active star ER Eri, despite not being an IRAS source, was observed during the PDS and classified as a WTT (Gregorio-Hetem et al. 1992; Torres 1998). Distinctly from our earlier search for the TWA region, where candidates were selected by their infrared emission, around ER Eri we used RASS sources.

We began with an area of $10^\circ \times 10^\circ$ centered on ER Eri:

$$01:30 < \alpha < 02:50 \text{ and } -60^\circ < \delta < -50^\circ$$

and later we enlarged it to about $20^\circ \times 25^\circ$:

$$01:00 < \alpha < 03:30 \text{ and } -70^\circ < \delta < -45^\circ$$

in order to find the limits of the possible association. We also chose three control areas, two of

them of $10^\circ \times 10^\circ$ where no isolated PMS star was known to exist *a priori*.

The first control area was chosen in the southern galactic hemisphere:

$$06:20 < \alpha < 07:40 \text{ and } -65^\circ < \delta < -55^\circ$$

The second area was in the northern galactic hemisphere:

$$09:40 < \alpha < 10:20 \text{ and } -15^\circ < \delta < -05^\circ$$

And the third one was in the direction of the known TWA, measuring $13^\circ \times 14^\circ$:

$$10:45 < \alpha < 11:45 \text{ and } -38^\circ < \delta < -24^\circ$$

As possible optical counterparts, we selected stars brighter than ~ 13 mag within the error boxes of the RASS, using the GSC and Tycho catalogues. We found 19, 46, 23, 33 and 23 X-ray sources with observable counterparts, respectively in the inner and outer areas around ER Eri, in the northern, in the southern and in the TWA regions. In the outer area around ER Eri we discarded, for observation, 10 bright stars, 9 being of the Bright Star Catalogue. These stars are, in general, well known and, in fact, none is a good candidate for their spectral type and space velocities. The only dwarf later than F is ζ^1 Ret, that has been studied recently by del Peloso et al. (2000). These ten stars are included at the end of Table 1 and we take them into account when doing the statistics in this paper. We observed then 36 X-ray sources in this outer region. Figure 3 summarizes graphically our strategy around ER Eri.

In September 1997 and January 1998 we observed the candidates, taking low resolution spectra using the B&C spectrograph installed at the Cassegrain focus of the 1.52 m ESO telescope at La Silla, Chile. The configuration used (grating of 1200 lmm^{-1} and the CCD #39) gives a final dispersion of $0.98 \text{ \AA pixel}^{-1}$. In 1999, using the same telescope, we obtained high resolution spectra for the selected candidates, with the FEROS echelle spectrograph (Kaufer et al. 1999), which gives a total coverage of 4000 \AA and a resolving power of 50000. To complete the extended area around ER Eri we took also some spectra using the coudé spectrograph (grating of 600 lmm^{-1} ; resolution of 9000; spectral coverage of 450 \AA centered

at 6500 \AA , using the CCD #101) of the 1.60 m telescope of the Observatório do Pico dos Dias (OPD), Brazil.

$UBV(RI)_C$ photometry for most of the selected stars (except in the southern control region) was obtained using FOTRAP (Jablonski et al. 1994) at the 0.60 m telescope of the OPD.

3. RESULTS

The main results of these surveys are given in Tables 1 to 4. In these tables we present the possible optical counterparts, the photometric data, our spectral types and the equivalent widths of the Li line $\lambda 6707.8 \text{ \AA}$ (W_{Li}).

The Li I line, when present in late type stars, can provide a first estimate for the age (Jeffries 1995), selecting possible PTT, moreover if the H_α line is in emission or filled-in. In such way we consider young stars those with spectral types between G0 and K2 with $W_{Li} > 0.05 \text{ \AA}$, but for later types just the presence of the Li line will be a sufficient condition. For hotter stars the Li line is not a good discriminator for age (see section 8) and, in general, we will limit our discussion to stars later than F in this paper.

In Tables 5 and 6 we present the kinematical and physical data of the possible young stars of the ER Eri and of the control regions, respectively. The data in italic mean deduced values, taking into account duplicity or estimated ones, for parallaxes and radial velocities, as we will see later. To obtain the Li abundances presented in column 13, we used our modified version of a LTE code kindly made available to us by Monique Spite, of Paris-Meudon Observatory. These Li abundances (and the $v \sin i$, column 14) were determined by fitting synthetic spectra to the observed ones. In the computation of the synthetic spectra, all the known atomic lines in the range $\lambda 6702 - \lambda 6712$ were taken into account. As the molecular bands are not taken into account, the uncertainties are higher for M type stars. The used stellar atmospheric models are from Gustafsson and collaborators (Edvardsson et al. 1993). For all stars we used $\log g = 4.5$, microturbulence velocity of 2 km s^{-1} and solar abundances. The Li abundance is not very sensitive to gravity and the main source of errors is the inaccuracy of the used effective temperatures (column 11).

In the area around ER Eri there are 19 X-ray sources having possible young stars as counterparts (8 in the inner area). The 4 visual binaries (ERX 22, ERX 33, ERX 37 and ERX 47)⁵ were counted as one entry each and we excluded in this count ER Eri = ERX 24 for reasons we will see below. We included in Table 5 three F stars kinematically similar to the later stars, presenting weak Li line (ERX 19, ERX 35 and ERX 38). Nevertheless, they will be excluded in any statistical consideration.

There is an indication that a young stellar association may exist in the direction of the inner area, as the number of X-ray sources associated with PTT candidates relative to all selected sources with optical counterparts, as explained in section 2, is greater than in all other regions. Actually, it is $\sim 40\%$ in the inner area, decreasing to $\sim 25\%$ in the additional region. This latter frequency is similar to the one of the southern control region, decreasing even more in the northern region ($\sim 10\%$), as can be seen in Table 8.

The known TWA region may furnish another useful comparison when searching for PTT associations far from clouds. The limits in right ascension and declination for this region were established somewhat larger than the other control regions to include the five known members of the association at the moment of the survey. We observed this region in the same way as the others and detected 9 possible young stars among 23 X-ray selected optical counterparts. Five of them (TRX 9, TRX 11, TRX 12, TRX 18 and TRX 20) are the already known members (de la Reza et al. 1989; Gregorio-Hetem et al. 1992), two (TRX 16 and TRX 21) were independently discovered by Sterzik et al. (1999) and Webb et al. (1999).

The frequency of possible young stars in the known TWA is very similar to the one we found in the inner ER Eri region. Is there a new young association in that part of the sky? To answer this question we used several criteria in order to characterize the young stars of this possible association and to define its true members. These involve their kinematical properties, relative proximity in the sky, ages, X-ray emission, the intensity of Li lines, the behavior of the H_α line and

stellar rotation. In the following we will discuss these different criteria.

4. KINEMATICAL PROPERTIES

In Tables 5 and 6 we present in columns 2 to 5 the proper motions, radial velocities and parallaxes of the young stars. Proper motions and parallaxes measured by the Hipparcos satellite are available for 11 of the 23 candidate members of the association around ER Eri and for 6 stars of the control areas. For the other stars the proper motions are from the TYCHO-2 catalogue (Høg et al. 2000) (exceptions: for TRX 12, we used Webb et al. (1999); for ERX 16 and SRX 11, we calculated the proper motions using AC2000 and GSC). Both catalogues are equally accurate (Høg et al. 2000) and the errors are usually less than 4 mas yr^{-1} . These translate to tangential velocity errors of $\sim 1 \text{ km s}^{-1}$ if the stars are at 50 pc . As proper motions of stars ERX 32 and ERX 53 are not available, they will not be taken into account in kinematical considerations.

We obtained the radial velocities using FEROS or OPD coudé spectra and their errors are similar to those in tangential velocities. We obtained more than one radial velocity for some stars: ERX 4 (3 measurements), ERX 16 (2), ERX 21 (6), ERX 22N (3), ERX 26 (2), ERX 32 (2), ERX 33N (2), ERX 33S (2) and ERX 45 (2). None of these stars presents noticeable variations and we assume also that the others are not spectroscopic binaries (SB). But we found that ER Eri (ERX 24) is a double line SB with an orbital period of 5.9255 days and the radial velocity used in Table 5 is the γ velocity (9.0 km s^{-1}). The analysis of this system showed that ER Eri is a RS CVn type binary at $\sim 330 \text{ pc}$ (Torres et al. 2000) and, although being the bull's-eye for our search, it has nothing to do with the proposed association. This indicates that some of the other objects classified as “young stars” in this work may be found to be other kinds of active stars when examined in more detail. For ERX 35, a known SB, we used the γ velocity from Sahade & Hernández (1964). The radial velocities in Table 6 for SRX 8S, a faint optical companion of SRX 8N, and for SRX 28, not measured by us, have been assumed to be 30 km s^{-1} , about the mean value for the region. Idem for SRX 18, a double line SB, for which we

⁵For convenience, the stars will be designated in this paper by their entries in Tables 1 to 4.

have only one spectrum. For TRX 10N we used the value of its brighter companion.

To estimate the distances for the stars in the ER Eri region not measured by Hipparcos, we began supposing that they all are in the Zero-Age MS (ZAMS). With the distances thus obtained we computed the space velocity components relative to the Sun, U , V (in the direction of galactic rotation) and W . These initial results are plotted, in the U - V plane, in Figure 1a. As can be visualized in the figure there is a systematic difference between the stars with the reliable parallaxes from Hipparcos (full circles) and the others. The latter are shifted in the direction towards $(U, V) = (0, 0)$. This distribution is highly improbable and it can be explained only if the distances are underestimated. There are two ways to produce this underestimation:

a) Almost all stars not measured by Hipparcos are binaries with similar components. This will increase the luminosities and decrease the derived distances. Even though possible, this is very unlikely as the only difference between the group with higher velocities and that with lower ones lies in the fact that the stars of the former are brighter and, therefore, measured by Hipparcos. This is an obvious contradiction with the hypothesis.

b) The stars not measured by Hipparcos are more luminous than predicted by ZAMS and, of course, younger.

If we assume this much more probable hypothesis (but keeping in mind that for some stars hypothesis (a) may be true) we can obtain kinematical parallaxes by adjusting them so as to minimize the moduli of the velocity vectors centered on the mean vector of the Hipparcos member stars. The space velocity components of the 6 stars with measured parallaxes (and in this particular context we included the F stars to increase the statistical significance) are:

$$(U, V, W) = (-8.8 \pm 1.3, -20.8 \pm 1.1, -1.4 \pm 1.5)$$

These errors are compatible with the very low observational errors for these stars ($\sim 0.3 \text{ km s}^{-1}$), plus an additional scattering of $\sim 1 \text{ km s}^{-1}$, expected for the original velocity dispersion during star formation. After obtaining a first set of kinematical parallaxes, we redefined the central vector including the stars within 2σ of the distribution, but excluding now the F stars. We minimized

again the moduli, but the changes were small and we stopped the process at this point. The distances and space velocity components obtained in this way are presented in columns 5 to 8 of Table 5 and the new U - V diagram is shown in Figure 1b. There is now a compact core formed by 12 stars, two of them being F stars, ERX 22N and ERX 19. One is a binary, ERX 37, and for it we will take the mean values. The remaining 10 stars have the following mean space velocity components:

$$(U, V, W) = (-9.5 \pm 1.0, -20.9 \pm 1.1, -2.1 \pm 1.9)$$

The rather large σ in the W velocity is mainly due to the radial velocity of ERX 14. To reduce it to the mean W velocity we would need to reduce the radial velocity by $\sim 4 \text{ km s}^{-1}$. This could be achieved by a large, but not impossible, measurement error or ERX 14 may be a single line SB. Without it we would have $W = -1.6 \pm 1.2$.

These ten stars form the “probable members” of an association which henceforth we will call the “*Horologium Association*” (HorA), since the majority of the members are in the constellation of the Horologium and we will try to prove that this is a real association.

The next star that might be included in the HorA is ERX 4, at 3.8σ of the vector defined by the 10 stars (or 5.9σ excluding ERX 14) and we can consider this star, that lies in a corner of the surveyed area, a limit case of possible member. Only a very improbable conjugation of errors could bring its velocity near the mean of the 10 stars. The next one, the F star ERX 35, will be at 10.6σ (or 15.5σ excluding ERX 14). Even if we include ERX 4, this star will be at 6.8σ . It is almost impossible that observational errors could bring this or any other selected star to the core of Figure 1b. We “rejected” these stars as members of the association and deduced their parallaxes as if they were on the ZAMS (Table 5). We call “possible members”, in addition to ERX 4, the F stars in the core and the stars ERX 32 and ERX 53, with unknown proper motions but physically not rejectable. For reasons presented in section 6, we classified the star ERX 37N as possible member. (The star ERX 24, or ER Eri, is retained in this table as a “rejected young star”, in spite of being older, for comparison to other possible mismatched cases).

The space velocity components obtained for the

HorA are similar to those of the complex for the Local Association

$$(U, V, W) = (-11.6, -20.7, -10.5)$$

represented by the Pleiades and the α Per groups, with ages between 20 and 150 Myr (Montes et al. 1999). Jeffries (1995) has previously noted such similarity between the kinematics of Li rich active late type stars and the Local Association. But for the HorA the scattering of space velocities and the modulus of W are lower than the values of the Local Association. In this sense, these values are distinct from any of the young Pleiades subgroups found by Asiain et al. (1999).

In columns 6 to 8 of Table 6 we give the space velocity components relative to the Sun for the control regions. To estimate parallaxes of the stars not in the Hipparcos catalogue of the northern and southern control areas and of the non-members of the TWA, we suppose they are on the ZAMS. For the TWA, we obtain kinematical parallaxes for the three stars with known proper motions in the same way as for the HorA, using as a mean vector the velocities of TRX 9 and TRX 18. With these five members the distance of the TWA is 45 pc. Of course, a better estimation would be obtained using also the proposed members outside our surveyed area, but there is no published complete set of data for none of them. Anyway, the distances obtained here are in agreement with Kastner et al. (1997). (TWA 9 has a Hipparcos parallax of 19.9 mas, near our mean distance for the TWA.) The space velocity components obtained from these five stars are:

$$(U, V, W) = (-11.7 \pm 1.0, -19.5 \pm 1.5, -4.7 \pm 0.7)$$

They are close to those of the HorA.

For the control regions (Table 6), there is a concentration in the U-V plane at about $(-8, -28)$ for five stars in the southern region. Do these young stars form another distinct physical association? Only SRX 1 and SRX 6 have trigonometric parallaxes - they seem mutually related - and we can use as a mean vector their velocities. But the obtained kinematical parallaxes for the stars SRX 22, SRX 18 and SRX 28 would place them *below* the MS. Actually, the last two stars would fall far below the MS. Put in another way, their distances and the scattering in space velocities would increase if they were younger. The result for SRX 11

is almost the same as if it were on the MS. Thus, the entries in Table 6 represent the best we can do for this region. Therefore, this concentration has discrepant parallaxes, a scattering in W velocities of 4 km s^{-1} and an age near the ZAMS, rejecting it as a possible association.

Two possibilities arise: either the concentration in the U-V plane is produced by a statistical fluctuation, as it seems the case for the southern control region, or it is the result of a real association, as for the TWA region. A fluctuation is not as improbable as it might appear at first glance since, all stars being young, they have space velocity components resembling those of the Local Association. Nevertheless, a real association must have a set of other characteristics alike and that is the case for the HorA, as we will show below.

Another way to test if the found concentration is a distinct one is to confront it with the more homogeneous sample formed by the Hipparcos catalogue. In Figure 2 we have plotted all stars measured by Hipparcos (excluding those that are probable or possible members of Table 5), with distances smaller than 90 pc in the same area of the sky of the HorA. To compute space velocity components, we adopted for all stars an artificial radial velocity of 12 km s^{-1} (close to the mean of the proposed association). In this way, the dispersion of space velocities is somewhat reduced. There is no concentration anywhere in Figure 2, even at the position of HorA velocity components. This is what may be expected if the HorA is real, formed by young active objects with constrained kinematics, and not a general behavior of the stars in the solar neighborhood. There are six other stars near the core of the HorA velocity distribution and, as they could be its non-active members, we got them together in Table 7. All have spectral types earlier than F6, as expected if they were the more massive and non X-ray emitting stars of the HorA. The hotter one is Achernar (HD 10144), a well known Be star, already leaving the MS, whose evolutive age is similar to that of the proposed association. If we use the radial velocity given in SIMBAD, its space velocity components are near those of the proposed association $(-6.1, -24.2, -6.3)$ (A in Fig. 1).

5. APPEARANCE IN THE SKY

In Figure 3 we show a map of the selected X-ray sources in the $20^\circ \times 25^\circ$ ER Eri region. The sources not associated with young objects are evenly distributed. The young stars not belonging to the HorA are also almost evenly distributed (we attributed the lack of stars at lower right ascensions to their small number). The probable and possible members of the HorA seem more assembled to the center.

It is yet difficult to be sure about the boundaries of the HorA, but towards the North there is a lack of members for at least 7° . This northern strip may be converted into one more control region. In this region, of the 20 observed plus 4 bright stars, less than 10% may be young. (Now and hereafter in this section, we will consider only stars later than F, binaries taken as one star.) The stars in this northern strip in Table 5 are marked with asterisks. Defining *a posteriori* the declination limits of the HorA region in the same way as we did for the TWA region (that is, $-66^\circ < \delta < -52^\circ$), it contains 33 selected X-ray sources and 5 bright stars not observed, 16 being young stars (42%), 13 of them probable or possible members of the HorA. This is a 270 square degrees area and the density of selected X-ray sources is 0.14 stars per square degree, being 0.06 for all young stars and 0.05 for the proposed members of the HorA (that is, possible and probable non-F stars).

Similarly, in the selected 182 square degrees TWA region, the densities of selected X-ray sources, young stars and members of the association are, respectively, 0.13, 0.05 and 0.04. In the northern strip (174 square degrees) and in the northern and southern control regions the densities of X-ray selected sources and young stars are, respectively, 0.14 and 0.01; 0.23 and 0.02; 0.33 and 0.08. The density of X-ray sources in the southern region is much higher than in all other regions and this explains its higher density of young stars. There is no obvious explanation for such excess as the young stars of this region seems not to be physically related. It is possible that this larger density in the southern region reflects only a preferential exploration of the ecliptic poles by the ROSAT satellite. Taking this higher X-ray density into account, the HorA is the densest region, slightly denser than the inner TWA. (The

statistics of the several classes of selected X-ray sources of the observed regions - or sub-regions - are summarized in Table 8.)

Considering only the members of both associations, the density of the HorA is higher than that of the TWA (0.05 to 0.038). It should be noted that two of the stars found by Webb et al. (1999) in the TWA, that are not in our surveyed region, are very near our boundaries. But three others are far from the center of the TWA, two of them being $\sim 10^\circ$ from its apparent border. This shows how difficult it is to be sure about the limits of the HorA. The list of the proposed candidates of the TWA outside our surveyed area (Webb et al. 1999; Sterzik et al. 1999) is presented in Table 9 with our measured photometric data (except for the bright star HR 4796).

The membership in the TWA is easier to establish due to its younger age. Even if its boundaries are not yet known, it is very improbable that it extends all over the sky as there is no star so young in any other of our surveyed areas. For the HorA this is more difficult to settle on as the stars may be rather similar to Local Association stars. Nevertheless, looking at the space motions in the control regions, only NRX 1 could be considered as a possible, but not probable, member of the HorA. (The members of the TWA, as noted before, have space motions not far from those of the HorA).

Using only the stars with known trigonometric parallaxes, the distance of the HorA would be 46 ± 5 pc. The mean distance, using also the kinematical parallaxes, is ~ 60 pc. The distance, based on Hipparcos, is an underestimation, as it measures only the brighter, that means nearer, members. ERX 8, the faintest member directly measured by Hipparcos, and also the reddest one, is of visual magnitude 9. The kinematical parallaxes give a dispersed association, ~ 50 pc in diameter. If we suppose that the original velocity dispersion during star formation is equal to the average modulus of the velocity vectors ($\sim 1.8 \text{ km s}^{-1}$), obtained as explained in section 4, the size of the HorA would be, again, ~ 50 pc after 30 Myr. This size, at ~ 60 pc, implies that the true angular extent of the HorA could be $\sim 50^\circ$, much larger than our observed region. In this case, many other young objects may be missing in our survey. A possible one is AB Dor, whose space velocity components (-8.5 , -25 and -14.6) are not far from

those of the HorA. Even though the W velocity is nearer to that of the Local Association, the ratio $\log(F_x/F_b) = -3.0$ and the age of 30 Myr (Collier Cameron & Foing 1997) are similar to those of the HorA. A more extended survey is necessary to decide about such kind of possibility.

The parallaxes for the stars of the TWA give a diameter of ~ 20 pc. Taking into account an age of ~ 10 Myr (Soderblom et al. 1998; Webb et al. 1999), this would imply an original velocity dispersion similar to the one we found for the HorA. If spherical, this diameter results in an angular extent of $\sim 25^\circ$, which is the size needed to embrace all the members found by Webb et al. (1999).

6. AGES AND STELLAR MASSES

An approach to the evolutionary state of the HorA can be made using a diagram of absolute magnitude M_V versus $(B-V)$. When calculating absolute magnitudes, no correction for interstellar extinction was applied as these stars are nearby and are not IRAS sources. Actually, the possible member ERX 19, a F type star, is a very weak IRAS source, detected only in $12 \mu\text{m}$, 0.24J , near the detection limit. Two other stars in Table 5 are associated with weak IRAS sources, ERX 35 and ERX 38, but they do not belong to the HorA. In all these cases the IRAS fluxes come from the stellar photospheres and from here on we will consider that there are no IRAS sources in the association.

In Figure 4 we present the M_V versus $(B-V)$ diagram, its respective isochrones and evolutionary paths for some stellar masses calculated by Siess et al. (1997). The proposed members not measured by Hipparcos had their parallaxes estimated kinematically as explained in section 4. As almost all probable members are near the isochrone of 30 Myr, the two stars without proper motions, ERX 32 and ERX 53, had their distances estimated supposing they are on this isochrone.

Although the distances of the majority of the stars of the HorA were obtained only from kinematical considerations, they have a small scattering in the evolutive diagram as almost all are on the isochrone of 30 Myr, as expected for a young association. This is very unlikely to result by chance alone and we propose this as a first approximation for the age of the HorA.

Approximate stellar masses can be obtained by

direct comparison with the positions of the stars in Figure 4 with respect to the stellar evolution paths. The mass distribution function has a maximum around $0.7 - 0.9 M_\odot$. There are some differences between this mass distribution and that of the other young stars in the control regions. The HorA stars are nearer and less massive objects than the other young field stars and more massive than those of the TWA. That is, most of the HorA members are K type stars at distances from 40 to 80 pc whereas most of the young field stars are G type stars at 50 to 100 pc and the TWA members are late K or M type stars. Where are the M stars of the HorA and the young K-M stars of the control regions? Probably beyond our observational limit of 13 mag. As we established this limit using GSC (that is, B magnitude ~ 13), our limit for redder stars is $V \sim 12$. For $B-V = 1.0$ on the ZAMS, this will represent stars beyond 100 pc and for $B-V = 1.5$, with an age of 30 Myr, stars beyond 30 pc. The few young K type field stars detected indicate that they are more Li depleted than those of the HorA. Depletion makes the Li line more difficult to be detected and, in fact, some stars with a dubious Li line in the B&C spectra present W_{Li} values between 0.05 and 0.1 \AA when observed with FEROS. As the stars of the TWA are younger, the M stars are intrinsically brighter and less Li depleted, being then easier to detect. We interpret these mass distributions as another indication that the age of the HorA is between that of the TWA and the young field stars. In any way, there seems to be a lack of G and early K type stars in the TWA.

This may be not the only difference between the TWA and the HorA. In the TWA most of the stars are binaries (Webb et al. 1999) and they are very few in the HorA. In fact, with the instruments used, we are unable to detect close visual binaries in the HorA - ERX 22 and ERX 37 being very wide. The primary of ERX 37, lying above the 30 Myr isochrone, is a candidate to be investigated for duplicity. As the other stars are on the isochrone (except for ERX 16, see below), any eventual secondary must have a very large luminosity difference. We have very few radial velocity measurements to detect single line SB. Nevertheless, at the sky position of the HorA, the radial velocity is mainly in the W direction. Even if, in general, we have not measured the stars more

than once, the slightly larger scattering in the W velocity, obtained in section 4, shows there is only a small possibility that some of these stars are SB. The best candidate is ERX 14. For the TWA we can compare the observations of this survey to those of the PDS and, at least, there is one SB - TRX 20 (PDS 55). Double line SB might be detected in only one exposure and we found five previously unknown SB (see Tables 1, 2 and 3), but none is a proposed member of the HorA.

Even considering this limited information we have an indication that the frequency of binaries in the HorA is less than in the TWA. The existence of some yet undetected binaries may decrease the scattering in the evolutive diagram even more.

6.1. Comments on some objects

Some of the objects in Figure 4 deserve detailed comments:

ERX 4 (CPD -64 120): This star is in the lower right corner of Figure 3 and has discordant space motions. It lies on the ZAMS and this may be another reason to exclude it as proposed member of the HorA. But, as its parallax was determined kinematically, the accidental errors that would conspire against suitable space velocity components could also underestimate the distance. We have observed thrice for radial velocity with similar values. It presents large variations in H_α - from absorption to a small emission. Thus, we keep ERX 4 as a limit case of membership.

ERX 14 (GSC 8047-0232): The probable member with the large discordant W . Its position in the evolutive diagram is compatible with the supposition that it may be a single line SB. We have only one radial velocity observation of this star.

ERX 16 (CD -53 386): It is the only probable member having a discrepant position in the evolutive diagram, and, besides, this is the only star in Table 5 without TYCHO-2 proper motions. Nevertheless, to check our proper motions, we used also the PPM and the USNO-A2, without any appreciable difference. For its absolute magnitude to be compatible with the age of 30 Myr, the distance should be ~ 80 pc. This would result in $(U, V, W) = -6.2, -16.8, -5.1$. Even not giving a

perfect fit to the kinematics or age test to belong to the HorA, it is still close enough to keep it as probable member.

ERX 22 (HD 13246): The stars of this system are separated by $52.76''$ but whereas star ERX 22N was not directly measured by Hipparcos, TYCHO-2 gives very similar proper motions and Torres (1986) has not found changes in their relative position since 1919, indicating that both stars constitute a physical pair separated by 2380 AU. Therefore, the parallax of ERX 22N was taken as being equal to that of the primary.

ERX 33 (HD 16699): This visual binary has been detected as such by Hipparcos, with a separation of $8.74''$, and our identical measured values of the radial velocities confirm it as a physical system. ERX 33S seems younger than the HorA by its position in the evolutive diagram. The spectrum of the primary (ERX 33N) has a cooler system of lines superimposed (like a late G or early K star) and Cutispoto et al. (1999) explain it by a close visual binary. (Actually, the USNO-A2 has two stars - the same? - in the field, of 11.5 mag., but they hardly could be the faint component.) Cutispoto et al. (1999) measured also the $v \sin i$ of this system and found almost the reverse of our values. One possibility is that ERX 33S is a double line SB too, not resolved in both works. This could explain the positions in the evolutive diagram not in good agreement for coeval stars and, in this case, more compatible with an age similar to that of the HorA. Anyway, these stars should not be considered members of the HorA due to their space velocities.

ERX 37 (CD -53 544): This system had already been proposed by Jeffries et al. (1996) as an isolated pair of PTT. The separation is about $22''$ and both stars are X-ray sources, the hotter one (ERX 37S) having a stronger flux, compatible with its larger $v \sin i$ (80 km s^{-1}) - one of the highest for this association. Our radial velocities are similar for both stars, as also found by Jeffries et al. (1996), indicating that they could form a physical pair. The main problem is the large difference in Li abundance between the components. One possible explanation is that ERX 37N may be a field

dMe star and the stars would not form a physical pair, only ERX 37S being member of the HorA. Nevertheless, if ERX 37N has an age of 30 Myrs, it would be at the same distance as the kinematical one obtained for ERX 37S, and this can hardly be accepted as being due to chance. Another possibility, adopted by Jeffries et al. (1996), takes into account a proposed dependence of Li depletion with rotation. Our measured low Li abundance (-1.2) for the slow rotator ERX 37N is in the direction of such proposal. Cautiously, we will consider ERX 37N as only a possible member of this association. ERX 37S is above the 30 Myr isochrone and this discrepant position could be an indication for duplicity.

ERX 32 and ERX 53 (GSC 8056-0432 and GSC 8499-0304): The distances for these stars are estimated supposing they are on the isochrone of 30 Myr, as they have no measured proper motions. But they appear to be members of the HorA for their large X-ray fluxes and large W_{Li} , as we will see in the following sections. Actually, the W_{Li} for ERX 32 seems very high compared to the depletion expected for a M3 star, but its $v\sin i$ is also high for this spectral type.

7. X-RAY FLUXES AND ROTATION

In column 10 of Tables 5 and 6 we present the ratios of X-ray fluxes to total luminosities, following the procedure of Jensen et al. (1998) to obtain the actual X-ray fluxes. In double systems we suppose that the X-ray fluxes are equally contributed by both components (except for ERX 37, for which we used the individual X-ray fluxes from Jeffries et al. (1996)). The values for the probable and possible members are high and compatible with those expected for PMS stars (Damiani et al. 1995) - except for the F stars ERX 19 and ERX 22S - and they are near the empirical saturation level for the Pleiades age (Kastner et al. 1997). In fact, the mean ratio $\log(F_x/F_b)$ for the probable members (-3.2 ± 0.2) is compatible with the evolutionary age (Kastner et al. 1997) and similar to that we obtained for the TWA (-3.0 ± 0.3). Only half of the young field stars have such high values.

Among the probable or possible members of the HorA, there are five stars with $v\sin i > 30 \text{ km s}^{-1}$ and four among young field stars: one in the north-

ern strip, ERX 26, two in the border of the HorA region, ERX 50 and ERX 54, and one SB, SRX 18. There is also one star in the TWA, a SB too, TRX 20.

The $\log(F_x/F_b)$ values for the probable and possible members of the HorA, excluding F stars, are weakly correlated to $v\sin i$ ($R = 0.64$). Although this is expected if rotation is one of the main physical causes of the X-ray emission in late type stars, even this weak correlation is not easily obtained since differences in stellar inclinations and evolutionary stages smear out the correlations for non-coeval stars. In fact, there is nothing like that for the young field stars. As almost all measured stars of the TWA present small $v\sin i$ it is impossible to draw any conclusion for this association.

The rotation of ER Eri shows that, if this star is synchronised, $R\sin i = 4.4 R_\odot$, which is incompatible with a WTT. Actually we found that this star is a RS CVn type system (Torres et al. 2000) at 330 pc and it was only by a very good luck that we found the HorA taking it as the bull's eye of our search.

8. LITHIUM AND H_α LINES

Because PTT are in an intermediate evolutionary stage between CTT or young WTT and active MS stars, genuine PTT are expected to have a peculiar H_α behavior (Pallavicini et al. 1992). Actually, the $W_{H\alpha}$ of the members of the HorA behave in a monotonic mode with respect to temperature, with fill-in at $B-V \sim 0.9$ and in emission for later type stars. For the young field stars there is more scattering, but the different distributions of stellar temperatures between both samples makes a detailed comparison difficult.

One of the best confirmations of the PTT nature of the stars, whose membership to the HorA was established by their space motions, comes from the Li lines. Martín (1997), comparing the behavior of the W_{Li} with temperature for a sample of CTT and stars of some young clusters, including the Pleiades, finds a gap (the PTT-gap) for stars cooler than 5250 K, between the high W_{Li} formed by CTT and the lower W_{Li} formed by young MS stars belonging to the Local Association. A similar conclusion is obtained by Jeffries (1995). The Li abundance is not a good criterium to discriminate PTT above the cutoff at $T_{eff} = 5250 \text{ K}$, as

there is almost no Li depletion in hotter stars.

The W_{Li} of the stars of the HorA are mainly in the lower edge of the PTT-gap (Fig. 5). The figure shows that the HorA is younger than the youngest open clusters of the Local Association plotted by Martín (1997) and Jeffries (1995). The two stars of the HorA with W_{Li} similar to the ones of the young field stars have the smallest $v\sin i$.

Ten young field stars are cooler than the cutoff in temperature, but only two of them may be younger than the Local Association: SRX 8S and TRX 10N. The latter seems a physical binary whose primary has a W_{Li} normal for a young star of its temperature, whereas the weak value of the secondary was obtained from a low resolution spectrum and is of low quality, making it a poor candidate for PTT. SRX 8S, the optical companion of the brighter SRX 8N, had its W_{Li} measured also on a low resolution spectrum. Its $W_{H\alpha}$ indicates some filling-in, but as it is very near the temperature cutoff, it is not a good PTT candidate too.

The stars of the TWA are on the upper edge of the PTT-gap, and some of its members are similar to CTT or WTT. The gap between the TWA and the HorA stars in Figure 5 indicates the amount of Li depletion in ~ 20 Myr.

9. DISCUSSION

In view of all the above arguments, we can conclude that there is, in the direction of the star ER Eri, an association of very young stars, younger than the majority of the groups of the Local Association and older than the TWA, characterized by the space motions and with properties that we have not found in other parts of the sky. We named it the ‘‘Horologium Association’’. Its isolation from clouds is quite remarkable. Even if there are some dispersed infrared cirrus at a few degrees from its center (Désert et al. 1988), the nearest clouds are at 25° from it. Moreover, these are CO translucent clouds, where there is no evidence for star formation (Magnani et al. 1996; Caillault et al. 1995). In any case, considering the proposed age for this association (~ 30 Myr), it is hopeless trying to find its birth place.

It is also interesting to note that contrary to the TWA, which is formed by a mixture of a few CTT as TWHya itself and a larger number of

PTT, the HorA is formed only by PTT where none of them is an IRAS source, indicating the absence of dusty accretion disks. This suggests that, at their advanced evolutionary stage, the circumstellar disks have been dissipated or agglomerated into planetesimals and, therefore, the HorA can give an upper limit for the lifetime of the disks. Jayawardhana et al. (1999b) discussed the problem raised by the existence of some isolated CTT having active disks in the TWA, as TWHya itself. There is also the case of the visual binary PDS 50 in the same cluster, where the component B could be evolving from a CTT to WTT stage (Jayawardhana et al. 1999a). We suppose that the TWA will evolve into the HorA stage and that the disks around their CTT will lose the IRAS properties in a time scale of less than 30 Myr. Considering that the majority of objects in the TWA are WTT, the few CTT would be in their last stages of activity. The transformation of the disks would consist in converting their dust into pre-planetary material. Quite recently Kürster et al. (1999) have detected a planet of 2.2 Jupiter masses in an Earth-like orbit (320 days) around the star ι Hor (ERX 38), one of the rejected members in Table 5. As this star seems not to belong to the HorA, we could not use the age of this association to get a better idea of the time scale of the formation of a planet around a young star. On the other hand, Barrado y Navascués et al. (1999) propose that the protoplanetary disk of β Pic has only 20 Myr, less than the age of the HorA. Probably there is no universal time scale for the apparition of planets, that should depend on the properties of the accreting disk.

10. CONCLUSIONS

Exploring a region of about $20^\circ \times 25^\circ$ around the high galactic latitude ($b = -59^\circ$) active star ER Eri, previously classified as a WTT (Gregorio-Hetem et al. 1992; Torres 1998), we found evidences of a new association, the ‘‘Horologium Association’’, formed mainly by bona fide low mass ($\sim 0.8 M_\odot$) PTT. Its probable and possible members are marked with asterisks in Table 1 and are discriminated in Table 5. Possible hot members are presented in Table 7. ER Eri itself was found to be a background RSCVn-like system. Since we found no low mass PTT in the two control areas, as can be seen in Figure 5, we believe that

HorA-like stellar groups are not numerous in the solar vicinity, indicating a non constant rate of star formation in the last 100 Myr. This new association is presently represented by at least 10 members (Table 5), having an age of ~ 30 Myr and is older than the isolated TWA, with an age of approximately 10 Myr. Until now there are no detected binaries in the HorA (stars ERX 22N and ERX 22S have a separation of 2380 AU and should be considered as common proper motion stars). If confirmed, this marked difference with the TWA must be due to different intrinsic conditions of star formation of these two associations. The HorA is even more isolated from clouds than the TWA and, in any case, we can expect that its original cloud could have dissipated in less than 30 Myr.

The distances of the stars in the HorA cover an interval from nearly 40 to about 90 pc (at a mean distance of ~ 60 pc) giving a diameter of ~ 50 pc, compatible with the size produced after 30 Myr by an initial velocity dispersion of ~ 1.8 km s $^{-1}$. If this size is the same in angular extent, it surpasses the surveyed region and could contain many other members, such as the interesting young star AB Dor. In the surveyed region there may be hotter members, that are not X-ray sources, (for example, the nearby Be star Achernar) having similar ages, distances and space motions. If this is the case, the HorA could be the remnant of an old OB association, but, due to the kinematical errors, there is little hope to localize its birth place.

The space velocity components of the HorA are near but distinct from those of the Local Association. In fact, in all surveyed regions we found young field stars, possible members of the Local Association, with a compatible Li depletion. But, for its well defined kinematics, physical properties, Li abundances and restricted location in the sky, the HorA can be distinguished from these stars, resembling more the general behavior of the TWA.

Considering the observational limitations, as that of the magnitude limit of the Hipparcos measurements, we are aware of the challenge that represents the detection of a coeval moving group of stars with an age around 30 Myrs. To arrive at this, we examine several requirements that characterize a young stellar association. If no one of them, isolated, is completely sufficient, none of them are mutually contradictory and all together practically create the necessary condition for the

establishment of the HorA as a real nearby association. Following Chereul et al. (1998) and considering that large moving groups, more commonly known as Eggen's superclusters, are not real clusters but are formed by a chance coincidence of smaller coeval streams, we can conceive the HorA as one small, coeval structure in the Pleiades supercluster. But its overall characteristics are distinct from those of the young Pleiades subgroups found by Asiain et al. (1999). In any case, the HorA may be one of the last episodes of star formation of the Local Association and could be useful to understand better its fine structure.

We thank L. Siess for providing the theoretical evolutive diagram used in this work. Drs. S. Frinck, B. Reipurth, G. Cutispoto, A. Andrei and N. Drake for instructive discussions. L. da Silva thanks ESO for the data reduction facilities. The suggestions and careful analysis of an anonymous referee contributed substantially in the improvement of this paper. We thank also the Centre de Données Astronomiques de Strasbourg (CDS) and NASA for the use of their electronic facilities, specially SIMBAD and ADS. This work was partially supported by the CNPq, grants to L da Silva and R. de la Reza, processes 200580/97 and 301375/86, respectively.

REFERENCES

- Asiain, R., Figueras, F., & Torra, J. 1999, *A&A*, 350, 434
- Barrado y Navascués, D., Stauffer, J. R., Hartmann, L., & Balachandran, S. C. 1997, *ApJ*, 475, 313
- Barrado y Navascués, D., Stauffer, J. R., Song, I., & Caillault, J. P. 1999, *ApJ*, 520, L123
- Briceño, C., Hartmann, L. W., Stauffer, J. R., Gagné, M., Stern, R. A., & Caillault, J. P. 1997, *AJ*, 113, 740
- Caillault, J. P., Magnani, L., & Fryer, C. 1995, *ApJ*, 441, 261
- Chereul, E., Crézé, M., & Bienaymé, O. 1998, *A&A*, 340, 384
- Collier Cameron, A., & Foing, B. H. 1997, *Observatory*, 117, 218
- Cutispoto, G., Pastori, L., Tagliaferri, G., Messina, S., Pallavicini, R. 1999, *A&AS*, 138, 87
- Damiani, F., Micela, G., Sciortino, S., & Harnden, F. R., Jr. 1995, *ApJ*, 446, 331
- de la Reza, R., Torres, C. A. O., Quast, G., Castilho, B. V., & Vieira, G. L. 1989, *ApJ*, 343, L61
- del Peloso, E. F., da Silva, L. & Porto de Mello, G. F. 2000, *A&A*, in press
- Désert, F. X., Bazell, D., & Boulanger, F. 1988, *ApJ*, 334, 815
- Edvardsson, B., Andersen, J., Gustafsson, G., Lambert, D. L., Nissen, P. E., & Tomkin, J. 1993, *A&A*, 275, 101
- Frinck, S. 1999, thesis, University of Heidelberg
- Feigelson, E. D. 1996, *ApJ*, 468, 306
- Gregorio-Hetem, J., Lépine, J. R. D., Quast, G. R., Torres, C. A. O., & de la Reza, R. 1992, *AJ*, 103, 549
- Hearty, T., Magnani, L., Caillault, J. P., Neuhäuser, R., Schmitt, J. H. M. M., & Stauffer, J. 1999, *A&A*, 341, 163
- Herbig, G. H. 1978, in *Problems of Physics and Evolution of the Universe*, edited by L. V. Mirzoyan, Publishing House of the Armenian Acad. of Sciences, Yerevan, p. 171
- Hoff, W., Henning, T., & Pfau, W. 1998, *A&A*, 336, 242
- Høg, E., Fabricius, C., Makarov, V. V., Urban, S., Corbin, T., Wycoff, G., Bastian, U., Schwkendiek, P., & Wicenc, A. 2000, *A&A*, 355, L27
- Jablonski, F., Baptista, R., Barroso, J., Jr., Gneiding, C. D., Rodrigues, F., & Campos, R. P. 1994, *PASP*, 106, 1172
- Jayawardhana, R., Hartmann, L., Fazio, G., Scott Fisher, R., Telesco, C. M., & Piña, R. K. 1999a, *ApJ*, 520, L41
- Jayawardhana, R., Hartmann, L., Fazio, G., Scott Fisher, R., Telesco, C. M., & Piña, R. K. 1999b, *ApJ*, 521, L129
- Jensen, E. L. N., Cohen, D. H., & Neuhäuser, R. 1998, *AJ*, 116, 414
- Jeffries, R. D. 1995, *MNRAS*, 273, 559
- Jeffries, R. D., Buckley, D. A. H., James, D. J., & Stauffer, J. R. 1996, *MNRAS*, 281, 1001
- Kaufer, A., Stahl, S., Tubbesing, S., Nørregaard, P., Avila, G., Francois, P., Pasquini, L., & Pizzella, A. 1999, *The Messenger*, 95, 8
- Kastner, J. H., Zuckerman, B., Weintraub, D. A., & Forveille, T. 1997, *Science*, 277, 67
- Kenyon, S. J., & Hartmann, L. 1995, *ApJS*, 101, 117
- Kürster, M., Endl, M., Els, S., Hatzes, A. P., Cochran, W. D., Döbereiner, S., & Dennerl, K. 1999, *A&A*, 353, L33
- Magnani, L., Hartmann, D., & Speck, B. G. 1996, *ApJS*, 106, 447
- Mamajek, E. E., Lawson, W. A., & Feigelson, E. D. 1999, *ApJ*, 516, L77
- Martín, E. L. 1997, *A&A*, 321, 492
- Montes, D., Latorre, A., & Fernández-Figueroa 1999, *astro-ph9906069*

- Neuhäuser, R. 1997, *Science*, 276, 1363
- Pallavicini, R., Pasquini, L., & Randich, S. 1992, *A&A*, 261, 245
- Palla, F., & Galli, D. 1997, *ApJ*, 476, L35
- Sahade, J., & Hernández, C. A. 1964, *Ann. d'Ap.*, 27, 11
- Siess, L., Forestini, M., & Dougados, C. 1997, *A&A*, 324, 556
- Soderblom, D. R., King, J. R., Siess, L., Noll, K. S., Gilmore, D. M., Henry, T. J., Nelan, E., Burrows, C. J., Brown, R. A., Perryman, M. A. C., Benedict, G. F., McArthur, B. J., Franz, O. G., Wasserman, L. H., Jones, B. F., Latham, D. W., Torres, G., & Stefanik, R. P. 1998, *ApJ*, 498, 385
- Sterzik, M. F., Alcalá, J. M., Covino, E., & Petr, M. G. 1999, *A&A*, 346, L41
- Torres, C. A. O. 1998, *Publicação Especial do Observatório Nacional*, 10/99
- Torres, C. A. O., Quast, G., da Silva, L., & de la Reza, R., 2000, in *Poster proceedings of IAU Symp 200, Birth and Evolution of Binary Stars*, edited by Bo Reipurth & Hans Zinnecker, p. 118.
- Torres, C. A. O., Quast, G., de la Reza, R., Gregorio-Hetem, J., & Lépine, J. R. D., 1995, *AJ*, 109, 2146
- Torres, G., 1986, *A&AS*, 64, 105
- Webb, R. A., Zuckerman, B., Platais, I., Patience, J., White, R. J., Schwartz, M. J., & McCarthy, C. 1999, *ApJ*, 512, L63
- Wichmann, R., Bastian, U., Krautter, J., Jankovics, I., & Ruciński, S. M. 1998, *MNRAS*, 301, L39

Fig. 1.— (U, V) space velocity components for stars in the ER Eri region (Table 1) with Hipparcos and estimated parallaxes for candidate stars. Some high velocity stars are outside the figure. Open circles are stars with estimated parallaxes: (a) supposing they were on the ZAMS, showing a systematic shift relative to the central core; (b) kinematically, obtained by minimizing the velocity scattering. The stars "move" grossly in the direction away from the position (0, 0). The central concentration defines the Horologium Association. Achernar (A) is plotted for comparison.

Fig. 2.— (U, V) space velocity components for the Hipparcos stars with $\pi > 0.11$ within the ER Eri region, excluding those that are possible or probable members of the HorA (filled circles of the central core in Fig. 1). The space velocity components were calculated with an adopted radial velocity of 12 km s^{-1} for all stars. Achernar is near its position in Figure 1. Many high space velocity stars are outside the figure. There is no concentration at (U, V) = -10, -21 - that characterizes the HorA - showing this it is not a general behavior of the stars in the solar neighborhood.

Fig. 3.— Distribution in the sky of the selected X-ray sources in the ER Eri region. Right Ascension is centered at 02:15. Filled circles are the field X-ray sources and open circles are the bright star counterparts of X-ray sources not observed in this program. ER Eri is the filled circle marked with an E. Asterisks represent the stars of the HorA (probable and possible members), whereas plus signs are young field stars. The position of Achernar is marked by an A. The inner selected region is displayed with continuous lines and the broken lines define the *a posteriori* HorA region.

Fig. 4.— Evolutive diagram for the probable and possible members of the HorA. The ZAMS is indicated by a broken line. Filled circles - stars with Hipparcos parallaxes. Open circles - stars with kinematical parallaxes; Open squares - ERX 32 and ERX 53, having no parallaxes, were plotted on the 30 Myrs isochrone. The kinematical parallax of ERX 16 seems underestimated, for its position on this diagram and its greater distance. We suspect that ERX 37S may be an undetected close binary. We plot also the non-member young binary ERX 33 (filled triangles). Most of the members

have masses between 0.7 and $0.9 M_{\odot}$.

Fig. 5.— The Li equivalent widths for the selected young stars as a function of temperature. Open triangles - stars of the TW Hydra Association. Asterisks - probable and possible members of the Horologium Association. Plus signs - young field stars. The dashed lines, based on Martín (1997), define the boundary for CTT. The solid line approximately represents the upper limit for Local Association and was taken using the plots of Martín (1997) and Jeffries (1995). The HorA stars are mainly in the lower part of the post-T Tau gap, whereas the TWA stars are near the boundaries for CTT.

TABLE 1
 STELLAR COUNTERPARTS OF X-RAY SOURCES IN THE ER ERI REGION.

ERX	RX J	Identification	V	$U-B$	$B-V$	$V-R$	$V-I$	SpT	W_{Li}
1	010117.7–455632	GSC 8032-1102	11.75	...	0.86	M3Ve	0:
2	010835.3–464025	HD 6869	6.89	0.05	0.29	0.17	0.50	F0IV	–0.05
3	011031.9–492928	CD –50 320	11.82	0.58	0.91	0.53	1.05	K2Ve	0:
4*	011316.2–641142	CPD –64 120	10.29	0.46	0.86	0.52	1.01	K1Ve	–0.330
5	012221.2–564418	HD 8435; BC Phe	8.84	0.32	0.76	0.42	0.80	G8Ve	0:
6*	012320.9–572853	HD 8558	8.50	0.22	0.68	0.40	0.77	G6V	–0.205
7N	012728.7–492822	CD –50 392	11.28	0.68	0.88	0.54	1.00	G9V	0:
7S	012728.7–492822	HJ 3438B ¹	10.81	0.56	0.84	0.49	0.94	K2V	0:
8*	012808.8–523824	HD 9054; CC Phe	9.07	0.60	0.91	0.53	1.01	K2V	–0.170
9	013204.4–630254	CPD –63 121	9.62	0.04	0.57	0.35	0.69	F9V	–0.04:
10	013232.8–493138	HD 9528; AE Phe	7.78	0.11	0.64	0.36	0.71	G2V	0:
11	013721.8–491144	CD –49 451	10.41	1.22	1.42	0.93	1.86	M0Ve	0:
12	014613.2–465651	HD 10922 ¹	9.35	0.22	0.75	0.43	0.79	G9V	0:
13	014841.1–483057	GSC 8044-0859	12.51	0.92	1.43	1.01	2.14	M2Ve	0:
14*	015215.7–521939	GSC 8047-0232	10.87	0.63	0.95	0.56	1.08	K3V	–0.350
15	015557.3–513634	HD 11937; χ Eri	3.69	0.46	0.84	0.46	0.90	G8V	0:
16*	020154.6–523453	CD –53 386	11.02	0.61	0.96	0.58	1.12	K3Ve	–0.315
17	020212.0–623532	HD 12695	8.10	0.04	0.58	0.35	0.69	F9V	–0.02:
18	020355.4–452446	HD 12759	7.30	0.22	0.69	0.39	0.76	G5V	0:
19*	020436.7–545320	HD 12894	6.43	–0.04	0.36	0.21	0.43	F4V	–0.08:
20	020438.7–530722	GSC 8483-1076	11.30	0.24	0.80	0.46	0.91	K0Ve	0:
21*	020718.6–531155	HD 13183	8.63	0.12	0.65	0.39	0.76	G5V	–0.241
22N*	020729.1–594014	CD –60 416	10.68	0.94	1.16	0.73	1.43	K5Ve	–0.268
22S*	020729.1–594014	HD 13246	7.50	–0.01	0.52	0.30	0.60	F7V	–0.145
23	020938.1–483556	HD 13397	7.79	0.76	1.01	0.54	1.04	G9III	0:
24	021008.2–543036	PDS 1; ER Eri ²	9.90	0.68	1.04	0.60	1.18	K3Ve	–0.185
25	021024.3–505001	HD 13445	6.12	0.45	0.81	0.42	0.86	G9V	0:
26	021055.5–460403	GSC 8042-1050	11.24	0.31	0.91	0.59	1.16	K2Ve	–0.260
27	021742.4–501450	HD 14366	9.29	0.30	0.75	0.43	0.83	G3V	0:
28	021851.4–695306	HD 14706	8.73	...	0.53	...	0.67	G0V	–0.12
29	022553.2–525749	HD 15279	8.37	0.11	0.64	0.35	0.69	G2V	–0.05:
30	022818.9–611817	HD 15638	8.88	–0.01	0.41	0.26	0.50	F3IV/V	...
31	022939.1–471336	HD 15661	9.78	0.38	0.74	0.43	0.85	G7V	0:
32*	023651.8–520300	GSC 8056-0482	12.11	1.05	1.48	1.08	2.33	M3Ve	–0.400
33N	023845.4–525710	HD 16699 ¹	7.86	0.03	0.52	0.30	0.58	F8V+K0	–0.055
33S	023845.4–525710	SAO 232842	8.36	0.23	0.72	0.42	0.81	G7V	–0.255
34	023901.2–581112	CD –58 538	9.53	1.14	1.44	0.93	1.86	M0Ve	0:
35	024039.0–543252	HD 16920; ζ Hor	5.22	–0.02	0.40	0.24	0.47	F4V	–0.065
36	024118.8–572522	GSC 8494-0369	13.68	1.11	1.49	1.16	2.53	M3Ve	0:
37N*	024146.8–525943	GSC 8491-1194	12.21	1.21	1.49	1.09	2.40	M3Ve	–0.08:
37S*	024146.8–525943	CD –53 544	10.21	1.04	1.26	0.82	1.60	K6Ve	–0.27
38	024233.5–504756	HD 17051; ι Hor	5.40	0.08	0.56	0.32	0.62	F9V	–0.040
39*	024234.4–573932	GSC 8497-0995	10.97	0.96	1.23	0.77	1.48	K6Ve	–0.120
40	025111.7–475314	GSC 8054-0859	11.78	0.81	1.11	0.66	1.22	K2Ve	0:
41	025153.5–613704	HD 18134; VZ Hor	8.84	0.49	0.85	0.51	1.00	K0V	0:
42	025346.9–613529	GSC 8859-0633	12.20	0.84	1.50	1.05	2.32	M3Ve	0:

TABLE 1—*Continued*

ERX	RX J	Identification	<i>V</i>	<i>U</i> − <i>B</i>	<i>B</i> − <i>V</i>	<i>V</i> − <i>R</i>	<i>V</i> − <i>I</i>	SpT	<i>W</i> _{Li}
43	025432.4−510829	GSC 8057-0342	12.21	1.34	1.52	1.07	2.29	M3Ve	0:
44	025636.8−530934	CD −53 596	10.17	1.07	1.10	0.68	1.26	K4V	0:
45*	025804.6−624115	GSC 8862-0019	11.67	0.77	1.04	0.65	1.25	K4Ve	−0.395
46	025848.7−544811	CD −55 479	10.31	1.29	1.32	0.83	1.59	K5V	0:
47E	030432.9−511904	CD −51 706 ¹	8.61	0.59	0.86	0.48	0.88	K1V	−0.020
47W	030432.9−511904	HD 19330	7.57	0.12	0.56	0.31	0.60	G1V	−0.030
48	030605.0−484748	HD 19491 ¹	8.27	−0.03	0.52	0.34	0.67	F8V	−0.055
49*	030618.8−652109	CD −65 149	<i>10.19</i>	...	<i>0.83</i>	K2V(e)	−0.32
50	031113.6−570132	GSC 8495-0384	11.35	0.81	1.09	0.64	1.25	K4Ve	−0.09
51	031546.3−542501	GSC 8493-1147	11.72	0.79	1.03	0.63	1.25	K3Ve	0:
52	032220.8−612253	GSC 8860-0497	11.39	0.92	1.12	0.74	1.45	M0Ve	0:
53*	032414.3−590101	GSC 8499-0304	12.09	1.13	1.25	0.78	1.54	M0Ve	−0.245
54	032738.6−580937	CD −58 693	<i>11.17</i>	...	<i>1.08</i>	K4Ve	−0.100
55	032837.0−665504	GSC 8873-0040	<i>10.70</i>	...	<i>0.73</i>	G5V(e?)	−0.04:
Bright stars not observed in this program									
	010605.9−464309	HD 6595; β Phe	<i>3.32</i>	<i>0.57</i>	<i>0.89</i>	<i>0.47</i>	<i>0.90</i>	<i>G8III</i>	...
	010719.3−614633	HD 6793; ι Tuc	<i>5.37</i>	...	<i>0.88</i>	...	<i>0.80</i>	<i>G5III</i>	...
	011545.7−685221	HD 7788; κ Tuc	<i>4.25</i>	<i>0.01</i>	<i>0.48</i>	<i>0.28</i>	<i>0.55</i>	<i>F6IV</i>	...
	013237.6−494336	HD 9544	<i>6.27</i>	...	<i>0.46</i>	...	<i>0.53</i>	<i>F4V</i>	...
	015710.3−472307	HD 12055	<i>4.83</i>	<i>0.52</i>	<i>0.86</i>	...	<i>0.89</i>	<i>G8III</i>	...
	015846.3−613410	HD 12311; α Hyi	<i>2.87</i>	<i>0.14</i>	<i>0.28</i>	<i>0.18</i>	<i>0.34</i>	<i>F0V</i>	...
	015936.9−652656	HD 12452	<i>7.30</i>	...	<i>0.42</i>	...	<i>0.49</i>	<i>F3V</i>	...
	024208.5−463121	HD 17006	<i>6.10</i>	...	<i>0.88</i>	...	<i>0.86</i>	<i>K1III</i>	...
	031743.8−623452	HD 20766; ζ^1 Ret	<i>5.53</i>	<i>0.06</i>	<i>0.64</i>	<i>0.36</i>	<i>0.71</i>	<i>G2.5V</i>	...
	032920.0−625624	HD 22001; κ Ret	<i>4.71</i>	<i>−0.04</i>	<i>0.39</i>	<i>0.24</i>	<i>0.47</i>	<i>F5IV-V</i>	...

¹Double line SB²Double line SB (P = 5.9255 days).

NOTE.—Data in italic were taken from the literature (from SIMBAD or Hipparcos). A colon denotes less precise values. *W*_{Li} measures with three decimals come from FEROS spectra. Asterisks mark probable and possible members of the Horologium Association.

TABLE 2
STELLAR COUNTERPARTS OF X-RAY SOURCES IN THE SOUTHERN REGION

SRX	RX J	Identification	<i>V</i>	<i>U</i> − <i>B</i>	<i>B</i> − <i>V</i>	<i>V</i> − <i>R</i>	<i>V</i> − <i>I</i>	SpT	<i>W</i> _{<i>L</i>i}
1	062230.9−601301	HD 45270	<i>6.50</i>	<i>0.05</i>	<i>0.60</i>	<i>0.34</i>	<i>0.66</i>	G1V	−0.149
2	062555.6−600325	GSC 8894-0426	<i>12.7:</i>	M5Ve	0:
3	063105.2−590017	HD 46697;TZ Pic	<i>6.63</i>	<i>1.00</i>	<i>1.14</i>	<i>0.61</i>	<i>1.13</i>	K1III	0:
4	063107.3−563914	CD −56 1546	<i>9.79</i>	...	<i>0.54</i>	G1V	0:
5	063536.4−623824	RR Pic	<i>12.39</i>	...	− <i>0.03</i>	...	− <i>0.02</i>	Nova	...
6	063800.7−613156	HD 48189	<i>6.19</i>	<i>0.10</i>	<i>0.62</i>	...	<i>0.69</i>	G0V	−0.150
7	063951.7−612842	CD −61 1439	<i>9.69</i>	...	<i>1.38</i>	...	<i>1.53</i>	M1V	0:
8N	064207.0−643126	HD 49078N	<i>8.16</i>	...	<i>0.34</i>	...	<i>0.45</i>	F0V	−0.027
8S	064207.0−643126	HD 49078S	<i>10.62</i>	...	<i>0.72</i>	K0V	−0.30
9	064345.6−642436	GSC 8903-0978	<i>13:</i>	M5Ve	0:
10	064558.4−550147	HD 49595	<i>8.76</i>	...	<i>0.46</i>	F5V	−0.05
11	064752.6−571332	GSC 8544-1037	<i>11:</i>	K5V	−0.169
12	064812.0−615623	HD 50241; α Pic	<i>3.24</i>	<i>0.12</i>	<i>0.22</i>	<i>0.14</i>	<i>0.28</i>	A7V	0:
13	064854.4−600857	GSC 8895-1353	<i>12:</i>	K4Ve	0:
14	065000.0−601459	HD 50571	<i>6.11</i>	...	<i>0.46</i>	...	<i>0.53</i>	F5V	0:
15	065459.1−563914	CD −56 1687	<i>9.54</i>	...	<i>0.47</i>	F7V	−0.13
16	070000.6−612008	HD 53143	<i>6.81</i>	<i>0.43</i>	<i>0.79</i>	<i>0.39</i>	<i>0.82</i>	G9V	0:
17	070002.4−570421	GSC 8545-0020	<i>12:</i>	F7V	0:
18	070350.6−582719	GSC 8549-0141 ¹	<i>11.69</i>	...	<i>0.62</i>	G8V	−0.219
19	070414.7−625404	GSC 8912-1753	<i>11:</i>	K2V	0:
20	070512.3−573402	GSC 8545-1235	<i>9.87</i>	...	<i>0.80</i>	G9V	−0.070
21N	070922.9−572953	HD 55402	<i>9.18</i>	...	<i>0.44</i>	F7V	0:
21S	070922.9−572953	GSC 8558-1141	<i>11.08</i>	...	<i>0.48</i>	G2V	0:
22	071051.1−573637	GSC 8558-1148	<i>10.45</i>	...	<i>0.68</i>	G6V	−0.170
23	071059.9−563256	GSC 8558-1964	<i>13:</i>	M2Ve	0:
24	071212.8−611627	CD −61 1583	<i>9.60</i>	...	<i>0.48</i>	F8V	−0.08
25	071450.8−591600	HD 56785	<i>9.32</i>	...	<i>0.54</i>	F8V	0:
26	071529.4−583228	HD 56910	<i>6.85</i>	...	<i>0.23</i>	...	<i>0.30</i>	A8V	0:
27S	071827.7−572101	HD 57555	<i>7.90</i>	...	<i>0.66</i>	...	<i>0.72</i>	G5V	−0.05
27N	071827.7−572101	Rst 244B	<i>13.6:</i>	F0V	−0.03
28	072123.9−572034	GSC 8559-1016	<i>10.73</i>	...	<i>0.64</i>	K0V	−0.16
29	072700.5−561439	HD 59487	<i>9.18</i>	...	<i>0.48</i>	...	<i>0.55</i>	F7V	−0.08
30	073028.9−563525	GSC 8559-1282	<i>11.48</i>	...	<i>0.71</i>	F5V	0:
31	073113.7−623051	CPD −62 837	<i>9.77</i>	...	<i>0.68</i>	G8V	0:
32	073233.9−562630	GSC 8559-1292	<i>10.76</i>	...	<i>0.75</i>	G6V(e?)	0:
33N	073341.0−574555	CD −57 1785A	<i>10.13</i>	...	<i>0.36</i>	F6V	−0.10
33S	073341.0−574555	CD −57 1785B	<i>11:</i>	F4V	−0.08

¹Double line SB

NOTE.—Data in italic were taken from the literature (from SIMBAD or Hipparcos). A colon denotes less precise values. *W*_{*L*i} measures with three decimals come from FEROS spectra.

TABLE 3
STELLAR COUNTERPARTS OF X-RAY SOURCES IN THE NORTHERN REGION

NRX	RX J	Identification	<i>V</i>	<i>U</i> − <i>B</i>	<i>B</i> − <i>V</i>	<i>V</i> − <i>R</i>	<i>V</i> − <i>I</i>	SpT	<i>W</i> _{<i>L</i>i}
1	094414.2−115218	HD 84323	9.29	0.14	0.66	0.38	0.74	G3V	−0.223
2	094454.2−122047	G 161-71	13.80	1.26	1.67	1.49	3.37	M5Ve	0:
3	094458.6−134707	GSC 5486-0589	<i>13:</i>	G0V	0:
4	095013.6−104255	GSC 5479-0365	10.96	0.42	0.86	0.49	0.94	K0V	0:
5	095039.1−053029	BD −04 2739 ¹	9.94	−0.04	0.59	0.36	0.72	G0V	−0.03:
6	095048.5−090657	G 161-78	13.31	1.33	1.66	1.15	2.63	M4Ve	0:
7	095128.5−145041	HD 85444	<i>4.12</i>	<i>0.65</i>	<i>0.92</i>	...	<i>0.92</i>	G7III	0:
8	095351.8−072001	GSC 4902-0210	10.79	1.19	1.50	0.96	1.91	M1Ve	0:
9	095433.6−125711	HD 85883	<i>6.87</i>	...	<i>0.55</i>	...	<i>0.62</i>	F8V	−0.065
10	095509.2−081921	GSC 5475-0507	11.60	1.19	1.48	0.96	1.97	M1Ve	0:
11A	095737.7−123003	GSC 5484-0936A	12.78J	1.48	1.55	1.05	2.18	M2Ve	0:
11B	095737.7−123003	GSC 5484-0936B	12.78J	1.48	1.55	1.05	2.18	M4Ve	0:
12	095916.0−073515	GSC 5476-0149 ²	<i>12:</i>	K0V	0:
13	100035.3−085442	GSC 5476-0910	11.55	0.20	0.78	0.47	0.98	G5Ve	0:
14	100755.8−133146	HD 87916	<i>8.59</i>	...	<i>1.06</i>	G7III	0:
15	100826.4−110708	HD 87978	<i>8.15</i>	...	<i>0.69</i>	...	<i>0.74</i>	G5V	0:
16	100839.4−053502	GSC 4909-0801	<i>13:</i>	K0Ve	0:
17	101317.3−082702	HD 88654	<i>7.68</i>	<i>0.53</i>	<i>0.87</i>	<i>0.47</i>	<i>0.87</i>	G7V	0:
18	101332.2−050512	HD 88682	<i>7.45</i>	...	<i>0.42</i>	...	<i>0.48</i>	F4V	0:
19	101344.5−072301	HD 88697	<i>7.21</i>	<i>0.03</i>	<i>0.50</i>	...	<i>0.57</i>	F7V	0:
20A	101628.3−052026	GSC 4910-1132A	<i>13.58</i>	...	<i>1.47</i>	<i>0.96</i>	...	M1V(e?)	0:
20B	101628.3−052026	GSC 4910-1132B	<i>14.15</i>	...	−0.18	−0.05	...	<i>DAe</i>	...
21	101731.3−080900	BD −07 3000	<i>9.87</i>	...	<i>0.71</i>	G7V	−0.078
22	101933.4−050610	HD 89490	<i>6.37</i>	<i>0.60</i>	<i>0.90</i>	...	<i>0.85</i>	G8V	0:
23	101956.8−084150	RW Sex	10.69	−0.65	−0.01	0.07	0.26	<i>DAe</i>	...

¹Double line SB

²Simbad gives a better association for this X-ray source with a QSO

NOTE.—Data in italic were taken from the literature (from SIMBAD or Hipparcos). A colon denotes less precise values. *W*_{*L*i} measures with three decimals come from FEROS spectra.

TABLE 4
 STELLAR COUNTERPARTS OF X-RAY SOURCES IN THE TW HYA REGION

TRX	RX J	Identification	<i>V</i>	<i>U</i> − <i>B</i>	<i>B</i> − <i>V</i>	<i>V</i> − <i>R</i>	<i>V</i> − <i>I</i>	SpT	<i>W</i> _{<i>Li</i>}
1	104728.2–372159	GSC 7198-0901	12.47	1.01	1.16	0.71	1.40	K5Ve	−0.029
2	104809.7–320106	CD −31 8535	9.46	0.04	0.60	0.37	0.73	G2V:	0:
3	105057.2–284954	CD −28 8475	10.63	0.06	0.60	0.35	0.69	GOV	0:
4	105604.3–265309	GSC 6643-1097	10.54	0.24	0.69	0.38	0.73	G5V	0:
5	105632.3–343404	HD 94853;OR Hya	8.40	0.14	0.64	0.35	0.68	G1V	0:
6	105813.0–292611	GSC 6647-0463	10.94	0.26	0.77	0.48	0.96	K2Ve	0:
7	105841.3–290813	HD 95135	8.26	0.58	0.90	0.53	1.03	K1III	0:
8	110110.9–313250	CD −30 8896	10.34	0.56	0.93	0.53	1.04	K3Ve	0:
9	110152.0–344212	TW Hya	11.07	−0.49	0.97	1.01	1.70	K8Ve	−0.440
10N	110327.5–324527	GSC 7204-0998	13.84	0.92	1.52	0.97	2.18	M2Ve	−0.05:
10S	110327.5–324527	GSC 7204-1026	11.62	0.94	1.22	0.77	1.53	K7Ve	−0.055
11	110913.5–300133	CD −29 8887;PDS 45	11.07	1.13	1.48	1.06	2.24	M2Ve	−0.560
12	111028.9–373204	Hen 600;PDS 50	12.04J	0.74	1.49	1.35	2.94	M4Ve	−0.610
13	111502.0–331905	HD 97840	<i>7.02</i>	...	<i>0.37</i>	...	<i>0.44</i>	F2V	0:
14	111737.6–344420	HD 98221	<i>6.48</i>	...	<i>0.41</i>	...	<i>0.47</i>	F3V	0:
15	111743.6–363210	HD 98233	<i>6.67</i>	...	<i>0.95</i>	...	<i>0.96</i>	G8III	0:
16N	112117.1–344644	CD −34 7390N	10.93J	1.00	1.44	0.95	1.97	M2Ve	−0.650
16S	112117.1–344644	CD −34 7390S	10.93J	1.00	1.44	0.95	1.97	M1Ve	−0.590
17	112149.4–241116	HD 98764	8.19	0.18	0.66	0.37	0.71	G2V	−0.120
18	112205.4–244632	HD 98800;TV Crt	8.90	1.09	1.24	0.77	1.51	K7V	−0.430
19	112256.2–254538	GSC 6654-0317	11.35	0.92	1.14	0.66	1.21	K4V	0:
20	113155.7–343632	CD −33 7795;PDS 55 ¹	11.37	0.93	1.47	1.10	2.36	M3Ve	−0.540
21	113241.7–265155	GSC 6659-1080;TWA 8	12.23	0.85	1.46	1.08	2.41	M2Ve	−0.520
22	113259.7–315127	HD 100407;ξ Hya	<i>3.65</i>	<i>0.70</i>	<i>0.95</i>	...	<i>0.92</i>	G8III	0:
23	114247.8–354832	HD 101799; V572 Cen	9.32	0.04	0.57	0.35	0.68	F8V	0:

¹SB, discrepant radial velocities measured in PDS and now.

NOTE.—Data in italic were taken from the literature (from SIMBAD or Hipparcos). A colon denotes less precise values. *W*_{*Li*} measures with three decimals come from FEROS spectra.

TABLE 5
 PROPERTIES OF THE YOUNG STARS OF THE ER ERI REGION

ERX	μ_α	μ_δ	RV	π	U	V	W	M_V	$[F_x/F_b]$	T_{eff}	$W_{H\alpha}$	N_{Li}	$v\sin i$
Probable members of the HorA													
6	+93.0	-36.9	+9.2	20.3	-10.4	-22.8	-0.9	5.04	-3.39	5700	-3.3	3.2	15
8	+104.8	-43.4	+7.7	26.9	-8.8	-19.5	-1.0	6.22	-3.31	4900	-0.2	2.1	6
14	+47.0	-7.5	+13.8	<i>11.2</i>	-11.3	-20.7	-6.3	6.12	-3.08	4750	-0.2	3.2	19
16	+35.0	-9.0	+11.9	<i>8.5</i>	-9.6	-21.1	-3.2	5.67	-3.08	4750	+0.1	2.9	19
21	+85.2	-22.5	+9.9	19.9	-9.8	-21.0	-0.6	5.12	-3.24	5775	-3.7	3.3	23
22N	+85.9	-24.0	+10.0	22.2	-7.9	-20.0	-0.3	7.41	<i>-3.22</i>	4350	+0.7	1.8	12
37S	+101.6	-9.9	+9	<i>25.0</i>	-10.4	-18.6	1.1	7.20	-2.97	4200	+1.4	1.9	80
39	+82.2	-8.9	+12.4	<i>20.8</i>	-9.1	-20.6	-1.3	7.56	-3.52	4250	+0.4	0.9	5
45	+39.4	-0.4	+16.3	<i>10.2</i>	-8.3	-22.8	-3.6	6.71	-3.13	4600	+0.2	3.3	6
49	+44.7	+7.1	+14	<i>11.9</i>	-9.9	-20.4	-2.6	5.57	-3.02	4900	-0.3	3.1	75
Possible members of the HorA													
4	+72.0	-6.0	+6.3	<i>17.1</i>	-13.5	-15.9	-2.3	6.47	-2.91	5000	+0.3	3.4	32
19	+75.8	-24.7	+13.3	21.2	-6.9	-20.7	-4.3	3.06	-4.94	6600	-7.0	3.0:	110
22S	+93.0	-18.8	+11	22.2	-9.6	-20.9	-1.2	4.23	<i>-4.26</i>	6280	-4.4	3.3	35
32	+16	<i>24</i>	9.01	-2.90	3500	+5.0	1.3	37
37N	+97.5	-13.7	+14	<i>25.0</i>	-9.2	-21.2	-3.2	9.20	-3.22	3500	+5.8	-1.2:	8:
53	+16.6	<i>13</i>	7.66	-3.42	4000	+0.8	1.0	11
Stars rejected as members of the HorA													
24 ¹	+39.8	+24.6	+9.0	<i>3</i>	-70.8	-22.6	-4.9	2.29	-2.90	4700	+0.6	1.7	38
26*	+56.1	-22.2	+25.4	<i>12</i>	-9.6	-30.5	-13.9	6.64	-3.00	4900	+0.2	2.6	36
28	+1.6	+1.2	+4	14.8	+0.5	-2.8	-2.9	4.58	-4.13	6030	-3.5	3.0	...
33N	+72.7	+48.8	+15.9	16.3	-24.0	-15.1	-9.9	3.92	<i>-3.75</i>	6200	-5.6	2.5	22
33S	+72.7	+48.8	+15.9	16.3	-24.0	-15.1	-9.9	4.42	<i>-3.58</i>	5600	-3.5	3.1	8
35	+32.8	+5.2	+5.8	20.5	-5.4	-7.6	-2.2	<i>2.53</i>	-5.29	6600	-6.6	2.9	8
38*	+333.7	+219.2	+16.6	58.0	-31.2	-17.2	-8.5	4.22	-4.98	6125	-4.3	2.4	7
47E*	+85.4	+72.2	+20.0	18.2	-28.8	-17.2	-11.0	4.91	<i>-4.76</i>	5100	-2.5	1.0:	6
47W*	+88.2	+71.9	+20.1	18.6	-28.6	-17.6	-10.9	3.92	<i>-5.10</i>	5950	-4.5	1.6:	6
50	+2.6	+24.9	+57	<i>16</i>	-4.7	-33.2	-46.7	7.37	-3.21	4600	+1.0	1.2	35
54	+78.6	+74.8	+81	<i>18</i>	-24.8	-60.2	-56.0	7.45	-2.67	4590	+2.0	1.3	35

¹This star, ER Eri, is a RS CVn-like system and, therefore, not young.

NOTE.—Data in italic are evolutive or kinematical parallaxes, as explained in the text, or deduced values, for M_V and $[F_x/F_b]$, taking into account duplicity. A colon denotes less precise values. Asterisks mark stars in the northern strip.

TABLE 6
 PROPERTIES OF THE YOUNG STARS OF THE CONTROL REGIONS

*	μ_α	μ_δ	RV	π	U	V	W	M_V	$[F_x/F_b]$	T_{eff}	$W_{H\alpha}$	N_{Li}	$v\sin i$	
Southern Region														
1	-11.2	+64.2	+32	42.6	-7.5	-28.4	-14.6	4.65	-4.16	5950	-4.0	3.1	18	
6	-50.1	+72.7	+35.0	46.2	-7.7	-30.3	-18.1	4.51	-3.71	6030	-3.5	3.2	18	
8S	-56.0	-21.0	+30:	<i>10</i>	+6	-15	-38	5.62	-4.28	5200	-0.4	3.5	...	
11	+2.2	+18.5	+30.5	<i>20</i>	-5.6	-28.5	-10.4	7.5:	-3.72	4500	-0.6	1.4	7	
18	+8.0	+31.1	+30:	<i>1</i>	4	-33	-35	+8	4.70	-3.38	5570	-2.7	3.2	60
20	+2.4	-21.1	-45.2	<i>17</i>	+7.3	+42.4	14.9	6.02	-3.74	5410	-2.1	1.9	25	
22	-10.7	+8.4	+29.7	<i>9</i>	-6.9	-26.4	-13.8	5.22	-3.28	5700	-2.8	3.0	27	
28	-1.6	+22.1	+30:	<i>11</i>	-10	-29	-7	5.94	-3.05	5200	-2.0	2.5	...	
Northern Region														
1	-42.2	-30.7	+19.6	12.5	-12.1	-24.0	-7.2	4.77	-3.32	5830	-3.3	3.5	30	
21	-23.0	+3.2	+49.8	<i>13</i>	-20.2	-37.5	+27.1	5.44	-3.48	5640	-1.8	2.2	9	
TWHya Region														
9	-66.9	-12.4	+13.0	17.7	-12.0	-18.2	-5.0	7.31	-2.52	4000	+300	2.5	14	
10N	+31.7	-33.2	-7.6	<i>20</i>	+9.4	+6.7	-6.6	10.35	-2.52	3500	+5	-0.7:	...	
10S	+31.7	-33.2	-7.6	<i>20</i>	+9.4	+6.7	-6.6	8.12	-3.12	4060	+1.0	0.2	3	
11	-90.1	-21.1	+12.0	<i>23.2</i>	-12.4	-18.0	-4.9	7.89	-3.20	3600	+2.1	3.5	15	
12	-112	-11	+14.0	<i>28.7</i>	-12.5	-19.3	-3.5	10.08	-3.18	3400	+10	3.5	15	
16N	+11.3	<i>20</i>	8.19	-2.86	3600	+3.5	3.6	10	
16S	+11.6	<i>20</i>	8.19	-2.86	3700	+2.7	3.6	12	
17	+40.5	-33.7	+0.3	20.8	+11.7	-0.4	-2.5	4.78	-4.37	5900	-3.8	2.8	4	
18	-85.4	-33.4	+13	21.4	-11.4	-20.7	-4.8	6.30	-3.42	4060	+0	2.4	10	
20	-81.6	-29.4	+14	<i>20.9</i>	-10.1	-21.3	-5.2	7.97	-2.97	3500	+10	3.7	58	
21	+7.5	<i>20</i>	8.74	-2.69	3450	+25	3.3	5	

¹Double line SB, velocity adopted as explained in the text.

NOTE.—Data in italic: for RV, we adopted a typical value of the measured stars in the Southern Region or, for TWX10, the one of the bright component; for parallaxes, estimated or kinematical values, as explained in the text; for M_V and $[F_x/F_b]$, deduced values taking into account the duplicity. A colon denotes less precise values.

TABLE 7
NON-X RAY STARS POSSIBLY BELONGING TO THE HORA

Identification	V	$B-V$	$V-I$	SpT	π
HD 8077	8.88	0.55	0.62	F6V	11.6
HD 10144	0.45	-0.16	-0.17	B3Vpe	22.7
HD 10269	7.08	0.47	0.54	F5V	20.4
HD 10472	7.62	0.42	0.49	F2IV/V	15.0
HD 14228	3.56	-0.12	-0.11	B8IV-V	21.1
HD 20888	6.03	0.13	0.15	A3V	17.2

NOTE.—Data from Hipparcos.

TABLE 8
SUMMARY OF THE STATISTICS OF THE OBSERVED REGIONS

Region	Area	X-ray	D ₁	XY	Fraction	D ₂	XA	D ₃
I-ERX	115	19	0.17	8	0.42	0.07	7	0.06
O-ERX	388	46	0.12	11	0.24	0.03	6	0.02
ERX	503	65	0.13	19	0.29	0.04	13	0.03
NERX	174	24	0.14	2	0.08	0.01
HRX	270	38	0.14	16	0.42	0.06	13	0.05
SRX	100	33	0.33	8	0.24	0.08
NRX	100	23	0.23	2	0.09	0.02
TRX	182	23	0.13	9	0.39	0.05	7	0.05

NOTE.—The first two regions are the inner and outer ER Eri regions, as defined in section 2. NERX and HRX are the northern strip and the HorA region as defined in section 5. In column 3 the number of all selected X-ray sources with stellar counterparts is given, as explained in section 2; D₁ is their sky density in each area (number of stars per square degrees). XY means the number of the X-ray sources associated with possible young stars later than F; in column 6 we present their fraction relative to the selected X-ray sources; D₂ is their sky density in each area. XA is the number of X-ray sources associated with probable and possible young stars later than F belonging to the HorA or to the TWA in the observed regions, and D₃ is their density.

TABLE 9
PHOTOMETRIC DATA OF OTHER POSSIBLE MEMBERS OF THE TWA

name	RX J	Identification	V	$U-B$	$B-V$	$V-R$	$V-I$
TWA 6	101828.7–315002	GSC 7183-1477	11.62	1.02	1.31	0.84	1.68
TWA 7	104230.3–334014	GSC 7190-2111	11.65	0.98	1.46	1.09	2.44
TWA 9A	114824.5–372838	CD –36 7429A	11.26	0.97	1.26	0.80	1.62
TWA 9B	114824.5–372838	CD –36 7429B	14.00	1.4:	1.43	1.17	2.58
TWA 10	123504.4–413629	GSC 7766-0743	12.96	0.81	1.43	1.09	2.47
Sterzik 1 ¹	110940.0–390657	CD –38 6968	9.92	0.07	0.59	0.34	0.67
Sterzik 2	(1121.1–3845)	GSC 7739-2180	12.85	1.0:	1.53	1.08	2.36

¹Not a member

NOTE.—These objects come from Webb et al. (1999) and Sterzik et al. (1999). A colon denotes less precise values.









