

Hard X-ray Emission and the Ionizing Source in LINERs

Yuichi Terashima¹, Luis C. Ho², and Andrew F. Ptak³

ABSTRACT

We report X-ray fluxes in the 2–10 keV band from LINERs (low-ionization nuclear emission-line regions) and low-luminosity Seyfert galaxies obtained with the *ASCA* satellite. Observed X-ray luminosities are in the range between 4×10^{39} and 5×10^{41} ergs s^{-1} , which are significantly smaller than that of the “classical” low-luminosity Seyfert 1 galaxy NGC 4051. We found that X-ray luminosities in 2–10 keV of LINERs with broad $H\alpha$ emission in their optical spectra (LINER 1s) are proportional to their $H\alpha$ luminosities. This correlation strongly supports the hypothesis that the dominant ionizing source in LINER 1s is photoionization by hard photons from low-luminosity AGNs. On the other hand, the X-ray luminosities of most LINERs without broad $H\alpha$ emission (LINER 2s) in our sample are lower than LINER 1s at a given $H\alpha$ luminosity. The observed X-ray luminosities in these objects are insufficient to power their $H\alpha$ luminosities, suggesting that their primary ionizing source is other than an AGN, or that an AGN, if present, is obscured even at energies above 2 keV.

Subject headings: galaxies: active — galaxies: nuclei — galaxies: Seyfert — X-rays: galaxies

1. Introduction

LINERs (low-ionization nuclear emission-line regions; Heckman 1980) are fairly common in nearby galaxies. A recent optical spectroscopic survey of nearby galactic nuclei has shown that $\sim 30\%$ of bright galaxies have LINERs (Ho, Filippenko, & Sargent 1997a, 1997b; Ho et al. 1997c). The ionization mechanism of LINERs is still controversial (see Filippenko 1996 for a review). There are several ionization mechanisms, which can explain LINER type optical emission lines, such as (1) photoionization by low-luminosity AGNs (LLAGNs), (2) photoionization by very hot Wolf-Rayet stars or O-stars, (3) shocks, (4) cooling flows, and so on.

¹NASA Goddard Space Flight Center, Code 662, Greenbelt, MD 20771

²The Observatories of the Carnegie Institution of Washington, 813 Santa Barbara St., Pasadena, CA 91101-1292

³Department of Physics, Carnegie Mellon University, 5000 Forbes Ave., Pittsburgh, PA 15213

Recent multiwavelength observations argue that a significant fraction of LINERs are genuine AGNs (e.g., Ho 1999a). Detection of an X-ray nucleus is one of the most convincing pieces of evidence for the presence of an AGN. From X-ray observations, the presence of LLAGNs in LINERs is reported for several objects (Ho 1999a; Terashima et al. 2000b, and references therein) and their luminosities ranges from 4×10^{39} to 5×10^{41} ergs s⁻¹, which is about 1–3 orders of magnitude lower than typical Seyfert galaxies. If such LLAGNs are the dominant ionizing source of the LINER-type emission lines, the optical emission-line luminosities are expected to be proportional to the luminosities of the ionizing photons from the LLAGNs. Indeed, a strong positive correlation between optical emission-line (H α , H β , [O III]) luminosities and X-ray luminosities is observed for luminous AGNs (Kriss et al. 1982; Elvis, Soltan, & Keel 1984; Ward et al. 1988; Mulchaey et al. 1994). Koratkar et al. (1995) reported that this correlation extends to intrinsically faint objects based on *ROSAT* observations of five low-luminosity Seyfert galaxies in the soft X-ray band. A small number of LINERs have been studied with *Einstein*, and a similar correlation is suggested (Halpern & Steiner 1983; Elvis et al. 1984).

The ionization mechanisms of LINERs, however, could be heterogeneous, and it is also possible that ionization sources other than AGNs are present. If other ionization mechanisms play a role, the X-ray luminosities are expected to be small compared to given optical emission-line luminosities (Pérez-Olea & Colina 1996).

In this paper, we present *ASCA* measurements of X-ray fluxes of LINERs and low-luminosity Seyfert galaxies in the 2–10 keV band and examine the probable ionization mechanism in LINERs by comparing X-ray luminosities with H α luminosities.

2. The Data

We have systematically analyzed *ASCA* archival and proprietary data of LINERs in the bright galaxy sample of Ho et al. (1997a). We also analyzed low-luminosity Seyfert galaxies (hereafter referred as LLSeyfert) with $\log L_{\text{H}\alpha} < 41$ ergs s⁻¹, which is smaller than or comparable to that of the well-studied low-luminosity Seyfert galaxy NGC 4051, to compare their X-ray properties with those of LINERs and to discriminate the ionizing source in LINERs. Detailed results will be presented in a future paper (Terashima et al. 2000b). We also use several objects in the southern hemisphere with good *ASCA* data (Iyomoto et al. 1996, 1997). We use only galaxies for which optical emission-line luminosities are published in the literature. Note that our sample is not complete and that some of the galaxies in our sample are selected based on large H α fluxes (see Terashima et al. 2000a). Therefore, our sample is not an X-ray selected sample, and so it is not significantly biased toward X-ray bright objects.

The optical classifications, adopted distances, broad and narrow H α luminosities, and X-ray luminosities in the 2–10 keV band are summarized in Table 1. We refer to objects with and without broad H α emission as Type 1 (LINER 1s and LLSeyfert 1s) and Type 2 objects (LINER 2s and

LLSeyfert 2s), respectively, by analogy with Seyfert 1s and Seyfert 2s. Two objects (NGC 4192 and NGC 4569) are classified as a transition object between a LINER and an H II nucleus. Although NGC 1097 was a LINER 2 historically (Phillips et al. 1984), we adopt a classification of Seyfert 1.5 based on Storchi-Bergmann, Baldwin, & Wilson (1993) since the *ASCA* observation was performed after the appearance of double-peaked broad H α . H α luminosities are taken from Ho et al. (1997a, c) except for several objects (see Table 1 for references). We estimated the amount of reddening by the Balmer decrement for narrow lines and corrected for reddening using the reddening curve of Cardelli et al. (1989), where we assumed the theoretical value of H α /H β =3.1. In the case that the observed H α /H β is less than 3.1, no correction was made. The reddening corrections for broad H α lines were made by same amount as narrow lines.

X-ray emission is detected from all the objects except for NGC 404. The X-ray spectra of most of the objects are represented by a two-component model consisting of an optically-thin thermal plasma with a temperature of ~ 0.7 keV and a hard component. Spectra of a few objects do not require a soft thermal component. The hard component is well represented by a power law with a photon index of ~ 1.5 – 2.0 . Detailed results of spectral fits are given in Terashima et al. (2000b). The absorption column density for the hard component ranges from 10^{20} to 10^{24} cm $^{-2}$. We will make use of the intrinsic luminosities of the hard component corrected for the absorption in the 2–10 keV band. Among the objects in our sample, seven objects (NGC 1052, 1365, 2639, 4258, 2273, 2655, and 4941) have a heavily absorbed ($N_{\text{H}} = 10^{23-24}$ cm $^{-2}$) transmitted continuum in their spectra, and the correction of the absorption is significant. Adopted column densities of these objects are 2.0, 2.8, 4.8, 1.3, 4.6, 4.0, 10×10^{23} cm $^{-2}$, respectively, as measured in Terashima et al. (2000b) and references in Table 1. If the absorption column is greater than 10^{24} cm $^{-2}$ and a transmitted hard X-ray continuum is not seen, corrections for the absorption cannot be made. In such a case, we utilize the observed luminosities corrected for only the Galactic absorption. These luminosities are summarized in Table 1. The X-ray luminosities range from 4×10^{39} ergs s $^{-1}$ to 5×10^{41} ergs s $^{-1}$. Results of *ASCA* observations for several objects have been already published, and references for them are also shown in Table 1.

3. Results and Discussion

3.1. LINER 1s

In order to examine whether optical emission lines in LINER 1s are photoionized by high-energy photons from an AGN, we search for a correlation between X-ray luminosities in the 2–10 keV band (L_X) and H α luminosities ($L_{\text{H}\alpha}$) for objects with broad H α in their optical spectra. It is known that a significant positive correlation between these two quantities exists for luminous AGNs (e.g., Ward et al. 1988). If photoionization by an LLAGN is the dominant ionization mechanism in LINER 1s, an L_X - $L_{\text{H}\alpha}$ correlation is expected. Figure 1a shows the correlation between L_X and $L_{\text{H}\alpha}$ for LINER 1s and LLSeyfert 1s in our sample and luminous Seyfert 1s and QSOs taken

from Ward et al. (1988); the $H\alpha$ luminosities shown represent the sum of the narrow and broad components of the line. It is clear that the correlation extends to lower luminosities. The same correlation using fluxes is shown in Figure 1b. The correlation is still significant in this plot. This correlation strongly suggests that the dominant ionization source in LINER 1s is photoionization by LLAGNs with luminosities less than $\sim 5 \times 10^{41}$ ergs s^{-1} . The scatter in the plots could be due to X-ray and $H\alpha$ variability of a factor of a few or more since X-ray and $H\alpha$ luminosities were not obtained simultaneously. A drastic example is an appearance of a double-peaked broad $H\alpha$ in NGC 1097, NGC 4203, and NGC 4450 (Storchi-Bergmann et al. 1993, 1995 ; Shields et al. 2000; Ho et al. 2000). Note that the Seyfert 1.8 galaxy NGC 1365 is probably Compton thick, and the X-ray luminosity is not corrected for absorption. Therefore, the data point is less luminous in X-rays by about two orders of magnitude than the correlation.

The X-ray images in the 2–10 keV band of most LINER 1s are unresolved within the spatial resolution of *ASCA* (FWHM $\approx 0.5'$) and consistent with the LLAGN interpretation as the origin of hard X-rays. Note that hard X-ray emission from NGC 4636 is clearly extended compared to the point-spread function, and an AGN is not the dominant X-ray source (Matsushita et al. 1994; Terashima et al. 2000b). Therefore, the X-ray luminosity of this object should be regarded as an upper limit. This upper limit is still consistent with the correlation above. If we exclude NGC 1365, 4636, and 4438, the latter having a very small $L_X/L_{H\alpha}$ ratio (§3.2), the correlation for the sample of LINER 1s, LLSeyfert 1s, and luminous AGNs plotted in Figure 1a is expressed as $\log L_X = (0.97 \pm 0.07) \log L_{H\alpha} + (2.3 \pm 2.7)$ (errors are 90% confidence).

3.2. LINER 2s

The X-ray emission from LINER 1s is most likely dominated by AGNs. If LINER 2s are also LLAGNs, they might be obscured, by analogy with Seyfert 2s, and the X-ray continuum emission is expected to be absorbed by a large column density. On the other hand, with other ionization mechanism such as hot stars or shocks resulting from starburst activity, an $L_X/L_{H\alpha}$ value is expected to be small compared to those for AGN (Pérez-Olea & Colina 1996). In Figure 2, we compare the luminosity ratio $L_X/L_{H\alpha}$ for the different object classes to examine whether the ionization mechanism in LINER 2s is the same as in LINER 1s and LLSeyferts; in this comparison we make use only of the narrow component of the $H\alpha$ line. The mean of the $L_X/L_{H\alpha}$ distributions for LLSeyfert 1s, LLSeyfert 2s, LINER 1s, and LINER 2s are 52.8, 10.3, 67.3, and 7.7, respectively. Thus the mean of $L_X/L_{H\alpha}$ for LINER 2s is about one order of magnitude smaller than those for LLSeyfert 1s and LINER 1s. A few LINER 2s (NGC 4261, NGC 4594, and NGC 4736) have an $L_X/L_{H\alpha}$ ratio similar to LINER 1s and LLSeyfert 1s, while many LINER 2s have a small $L_X/L_{H\alpha}$. Some LLSeyferts with small $L_X/L_{H\alpha}$ are most probably Compton thick objects in which only scattered X-rays are observed (NGC 1365, Iyomoto et al. 1997; NGC 1386, Maiolino et al. 1998, Bassani et al. 1999; NGC 5194, Terashima et al. 1998b) since these objects have a strong Fe-K emission line with an equivalent width of > 1 keV, and the intrinsic $L_X/L_{H\alpha}$ values could be 1–2

orders of magnitude larger than observed. There is no clear X-ray evidence for the presence of heavily obscured AGN in NGC 3079 and NGC 7743, although such a possibility cannot be ruled out.

If we assume a spectral energy distribution from UV to X-ray is power-law shape with an index of -1 ($f_\nu \propto \nu^{-1}$) (Ho 1999b), $\log L_X/L_{H\alpha}$ should be greater than 1.4 to drive $H\alpha$ luminosities by photoionization under Case B recombination and a covering fraction of unity (Osterbrock 1989). Since X-ray luminosities of most LINER 2s are small ($L_X < 2 \times 10^{40}$ ergs s^{-1}), X-ray binaries in the host galaxy might also contribute to observed X-ray fluxes. Actually some objects such as NGC 3607, NGC 4111, NGC 4374, and NGC 4569 have hard band (>2 keV) images extended to several kpc (Terashima et al. 2000a, b), which implies a lower value of $L_X/L_{H\alpha}$ for the nuclear component. Therefore the objects with low $\log L_X/L_{H\alpha}$ (<1) are too X-ray weak to ionize optical emission lines, even if X-ray variability of a factor of a few is also taken into account. If an AGN is present and is the dominant ionizing source in these objects, it should be obscured even at energies above 2 keV. Although no clear evidence for the presence of heavily obscured AGN (e.g. heavily absorbed X-ray continuum and/or strong Fe-K emission line) has been obtained so far, the current data cannot rule out such a possibility. Alternatively, there might be other ionization sources. The *Hubble Space Telescope* UV spectra of NGC 404 and NGC 4569 actually show large number of hot stars concentrated in pc-scale nuclear regions (Maoz et al. 1998), and these objects could be examples of LINERs ionized by stellar sources (Terashima et al. 2000a).

3.3. Fraction of AGNs in Bright Galaxies

According to the optical spectroscopic survey by Ho et al. (1997a, b), Seyferts, LINER 1s and LINER 2s are detected in 11%, 5%, 28% of northern ($\delta > 0^\circ$) bright ($B_T \leq 12.5$ mag) galaxies. If we assume an extreme case that all Seyferts and LINER 1s are AGNs and all LINER 2s are not AGNs, the fraction of AGN is estimated to be 16% of bright galaxies. This percentage, however, should be regarded as a lower limit because extremely weak broad $H\alpha$ is difficult to detect unambiguously (Ho et al. 1997c). Furthermore, some LINER 2s clearly indicate AGN-like activity. For example, NGC 4261 has prominent radio jets and kinematic evidence for a massive black hole (Ferrarese, Ford, & Jaffe 1996). Other examples include the ‘‘Sombrero’’ galaxy (NGC 4594) and NGC 4736. The $L_X/L_{H\alpha}$ values for these object are similar to LINER 1s and LLSeyferts. This fact also indicates that the $L_X/L_{H\alpha}$ ratio is a good indicator of the presence of AGNs. Hard X-ray surveys conducted at high angular resolution, such as afforded by *Chandra*, would be crucial to refine the true AGN fraction in nearby galaxies.

The authors are grateful to all the *ASCA* team members. YT is supported by the Japan Society for the Promotion of Science Postdoctoral Fellowships for Research Abroad. The research of LCH is partially supported by NASA grant NAG 5-3556 and by NASA grants GO-06837.01-95A and AR-07527.02-96A from the Space Telescope Science Institute (operated by AURA, Inc., under

NASA contract NAS5-26555). AFP is partially supported by NASA grant NAG 5-8093.

REFERENCES

- Allen, S. W., Di Matteo, T., & Fabian, A. C. MNRAS, 2000, 311, 493
- Awaki, H., et al. 1994, PASJ, 46, L65
- Bassani, L., et al. 1999, ApJS, 121, 473
- Buote, D. A., & Fabian, A. C. 1998, MNRAS, 296, 977
- Cardelli, J. A., Clayton, G. C., Mathis, J. S. 1989, ApJ, 345, 245
- Dahlem, M., Weaver, K. A., & Heckman, T. M. 1998, ApJS, 118, 401
- Elvis, M., Soltan, A., & Keel, W. C. 1984, ApJ, 283, 479
- Ferrarese, L., Ford, H. C., & Jaffe, W. 1996, ApJ, 470, 444
- Filippenko, A. V. 1996, in *The Physics of LINERs in View of Recent Observations*, ed. M. Eracleous et al. (San Francisco: ASP), 17
- Guainazzi, M., & Antonelli, L. A. 1999, MNRAS, 304, L15
- Halpern, J. P., & Steiner, J. E. 1983, ApJ, 269, L37
- Heckman, T. M. 1980, A&A, 87, 152
- Ho, L. C. 1999a, *Advances in Space Research*, 23(5-6), 813
- Ho, L. C. 1999b, ApJ, 516, 672
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1993, ApJ, 417, 63
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997a, ApJS, 112, 315
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997b, ApJ, 487, 568
- Ho, L. C., Filippenko, A. V., Sargent, W. L. W., & Peng, C. Y. 1997c, ApJS, 112, 391
- Ho, L. C., Ptak, A., Terashima, Y., Kunieda, H., Serlemitsos, P. J., Yaqoob, T., & Koratkar, A. P. 1999, ApJ, 525, 168
- Ho, L. C., Rudnick, G., Rix, H.-W., Shields, J. C., McIntosh, D. H., Filippenko, A. V., & Sargent, W. L. W. 2000, ApJ, in press
- Ishisaki, Y., et al. 1996, PASJ, 48, 237

- Iyomoto, N., Makishima, K., Fukazawa, Y., Tashiro, M., Ishisaki, Y., Nakai, N., & Taniguchi, Y. 1996, PASJ, 48, 231
- Iyomoto, N., Makishima, K., Fukazawa, Y., Tashiro, M., & Ishisaki, Y. 1997, PASJ, 49, 425
- Iyomoto, N., Makishima, K., Matsushita, K., Fukazawa, Y., Tashiro, M., & Ohashi, T. 1998, ApJ, 503, 168
- Keel, W. C. 1983, ApJ, 269, 466
- Koratkar, A. P., Deustua, S. E., Heckman, T. M., Filippenko, A. V., Ho, L. C., & Rao, M. 1995, ApJ, 440, 132
- Kriss, G. A., & Canizares, C. R. 1982, ApJ, 261, 51
- Maiolino, R. et al. 1998, A&A, 338, 781
- Makishima, K., et al. 1994, ApJ, 46, L77
- Maoz, D., Koratkar, A. P., Shields, J. C., Ho, L. C., Filippenko, A. V., & Sternberg, A. 1998, AJ, 116, 55
- Matsushita, K., et al. 1994, ApJ, 436, L41
- Matsumoto, H., Koyama, K., Awaki, H., Tsuru, T., Loewenstein, M., & Matsushita, K. 1997, ApJ, 482, 133
- Matsumoto, Y., et al. 2000, in preparation
- Mizuno, T., Ohnishi, T., Kubota, A., Makishima, K., & Tashiro, M. 1999, PASJ, 51, 663
- Mulchaey, J. S., Koratkar, A. P., Ward, M. J., Wilson, A. S., Whittle, M., Antonucci, R. R. J., Kinney, A. L., & Hurt, T. 1994, ApJ, 436, 586
- Nicholson, K. L., Reichert, G. A., Mason, K. O., Puchnarewicz, E. W., Ho, L. C., Shields, J. C., & Filippenko, A. V. 1998, MNRAS, 300, 893
- Osterbrock, D. E. 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (Mill Valley: Univ. Science Books)
- Pérez-Olea, D. E., & Colina, L. 1996, ApJ, 468, 191
- Phillips, M. M., Pagel, B. E. J., Edmunds, M. G., & Díaz, A. 1984, MNRAS, 210, 701
- Ptak, A., Serlemitsos, P. J., Yaqoob, T., & Mushotzky, R. 1999, ApJS, 120, 179
- Ptak, A., Yaqoob, T., Serlemitsos, P. J., Kunieda, H., & Terashima, Y. 1996, ApJ, 459, 542
- Roberts, T. P., Warwick, R. S., & Ohashi, T. 1999, MNRAS, 304, 52

- Sambruna, R. M., Eracleous, M., & Mushotzky, R. F. 1999, *ApJ*, 526, 60
- Shields, J. C., Rix, H-W., McIntosh, D. H., Ho, L. C., Rudnick, G., Filippenko, A. V., Sargent, W. L. W., & Sarzi, M. 2000, *ApJ*, 534, L27
- Storchi-Bergmann, T., Baldwin, J. A., & Wilson, A. S. 1993, *ApJ*, 410, L11
- Storchi-Bergmann, T., Eracleous, M., Livio, M., Wilson, A. S., Filippenko, A. V., & Halpern, J. P. 1995, *ApJ*, 443, 617
- Storchi-Bergmann, T. & Pastriza, M.G. 1989, *ApJ*, 347, 195
- Taniguchi, Y., Ohyaama, Y., Yamada, T., Mouri, H., & Yoshida, M. 1996, *ApJ*, 467, 215
- Terashima, Y., et al. 2000b, in preparation
- Terashima, Y., Ho, L C., Ptak, A. F., Mushotzky, R. F., Serlemitsos, P. J., Yaqoob, T., & Kunieda, H. 2000a, *ApJ*, 533, 729
- Terashima, Y., Kunieda, H., Misaki, K. 1999, *PASJ*, 51, 277
- Terashima, Y., Kunieda, H., Misaki, K., Mushotzky, R. F., Ptak, A. F., & Reichert, G. A. 1998a, *ApJ*, 503, 212
- Terashima, Y., Ptak, A., Fujimoto, R., Itoh, M., Kunieda, H., Makishima, K., & Serlemitsos, P. J. 1998b, *ApJ*, 496, 210
- Véron-Cetty, M.-P., & Véron, P. 1986, *A&AS*, 66, 335
- Ward, M. J., Done, C., Fabian, A. C., Tennant, A. F., & Shafer, R. A. 1988, *ApJ*, 324, 767
- Weaver, K. A., Wilson, A. S., Henkel, C., & Braatz, J. A. 1999, *ApJ*, 520, 130
- Wilson, A. S., et al. 1998, *ApJ*, 505, 587

Table 1. X-ray and H α luminosities for observed galaxies

| Name | Distance | Class | $\log L(\text{H}\alpha)$ (Broad) | $\log L(\text{H}\alpha)$ (Narrow) | $\log L_X$ (2–10 keV) | $\log L_X/L_{\text{H}\alpha}$ | reference | X | H α |
|----------|----------|-----------|-------------------------------------|--------------------------------------|--------------------------|-------------------------------|-----------|-----|------------|
| | [Mpc] | | [ergs s $^{-1}$] | [ergs s $^{-1}$] | [ergs s $^{-1}$] | | | | |
| NGC 315 | 65.8 | L1.9 | 39.92 | 39.61 | 41.70 | 2.09 | 1 | | |
| NGC 1052 | 17.8 | L1.9 | 39.54 | 39.45 | 41.58 | 2.13 | 2,3 | 1,2 | |
| NGC 3998 | 21.6 | L1.9 | 40.59 | 40.43 | 41.67 | 1.24 | 4 | | |
| NGC 4203 | 9.7 | L1.9 | 38.59 | 38.34 | 40.40 | 2.06 | 5 | | |
| NGC 4438 | 16.8 | L1.9 | 39.54 | 40.04 | 39.96 | −0.08 | | | |
| NGC 4450 | 16.8 | L1.9 | 38.48 | 38.51 | 40.34 | 1.83 | | | 1,2 |
| NGC 4579 | 16.8 | S1.9/L1.9 | 39.49 | 39.53 | 41.18 | 1.65 | 4,6 | | |
| NGC 4636 | 17.0 | L1.9 | 38.37 | 38.27 | 40.22 | 1.95 | 7–11 | | |
| NGC 5005 | 21.3 | L1.9 | 40.05 | 39.48 | 40.59 | 1.11 | | | |
| NGC 404 | 2.4 | L2 | ... | 37.82 | < 37.66 | < −0.16 | 12 | | |
| NGC 3507 | 19.8 | L2 | ... | 39.60 | 39.92 | 0.32 | | | |
| NGC 3607 | 19.9 | L2 | ... | 39.53 | 40.16 | 0.63 | | | |
| NGC 4111 | 17.0 | L2 | ... | 39.85 | 39.94 | 0.09 | 12 | | |
| NGC 4192 | 16.8 | T2 | ... | 40.49 | 39.58 | −0.91 | 12 | | |
| NGC 4261 | 35.1 | L2 | ... | 39.82 | 41.18 | 1.36 | 1, 13 | | |
| NGC 4374 | 16.8 | L2 | ... | 39.35 | 40.33 | 0.98 | 8–10 | | |
| NGC 4457 | 17.4 | L2 | ... | 39.79 | 39.98 | 0.20 | 12 | | |
| NGC 4569 | 16.8 | T2 | ... | 40.66 | 40.08 | −0.58 | 12 | 3 | |
| NGC 4594 | 20.0 | L2 | ... | 39.82 | 41.00 | 1.18 | 4,14 | | |
| NGC 4736 | 4.3 | L2 | ... | 38.12 | 39.64 | 1.52 | 15 | 4 | |
| NGC 7217 | 16.0 | L2 | ... | 39.78 | 39.86 | 0.08 | | | |
| NGC 1097 | 14.5 | S1.5 | 40.55 | 39.53 | 40.63 | 1.10 | 16 | 5 | |
| NGC 1365 | 16.9 | S1.8 | 42.02 | 41.08 | 40.50 | −0.58 | 17 | 6 | |
| NGC 2639 | 42.6 | S1.9 | 39.79 | 40.48 | 41.66 | 1.18 | 18 | 1,2 | |
| NGC 3031 | 1.4 | S1.5 | 38.41 | 38.44 | 39.60 | 1.16 | 19 | | |
| NGC 4258 | 6.8 | S1.9 | 38.90 | 38.60 | 40.83 | 2.23 | 4,20 | | |
| NGC 4565 | 9.7 | S1.9 | 38.38 | 38.46 | 39.77 | 1.31 | 21 | | |
| NGC 4639 | 16.8 | S1.0 | 39.75 | 38.39 | 40.54 | 2.15 | 22 | | |
| NGC 5033 | 18.7 | S1.5 | 40.32 | 39.70 | 41.37 | 1.67 | 23 | | |
| NGC 1386 | 16.9 | S2 | ... | 40.94 | 40.34 | −0.60 | 17 | 7 | |
| NGC 2273 | 28.4 | S2 | ... | 41.00 | 41.93 | 0.93 | | | |

Table 1—Continued

| Name | Distance | Class | $\log L(\text{H}\alpha)$ (Broad) [ergs s ⁻¹] | $\log L(\text{H}\alpha)$ (Narrow) [ergs s ⁻¹] | $\log L_X$ (2–10 keV) [ergs s ⁻¹] | $\log L_X/L_{\text{H}\alpha}$ | reference X | H α |
|----------|----------|-------|--|---|---|-------------------------------|----------------|------------|
| NGC 2655 | 24.4 | S2 | ... | 40.04 | 41.46 | 1.42 | | |
| NGC 3079 | 20.4 | S2 | ... | 41.04 | 40.40 | -0.64 | 4,24 | |
| NGC 3147 | 40.9 | S2 | ... | 40.02 | 41.53 | 1.51 | 4,25 | |
| NGC 4501 | 16.8 | S2 | ... | 39.08 | 40.30 | 1.22 | | |
| NGC 4941 | 6.4 | S2 | ... | 39.81 | 40.69 | 0.87 | | 7 |
| NGC 5194 | 7.7 | S2 | ... | 40.00 | 39.86 | -0.14 | 4,26 | 3 |
| NGC 7743 | 24.4 | S2 | ... | 40.29 | 39.57 | -0.72 | | |

References. — X-ray luminosities: (1) Matsumoto et al. 2000; (2) Guainazzi & Antonelli 1999; (3) Weaver et al. 1999; (4) Ptak et al. 1999; (5) Iyomoto et al. 1998; (6) Terashima et al. 1998a; (7) Awaki et al. 1994; (8) Matsushita et al. 1994; (9) Matsumoto et al. 1996; (10) Boute & Fabian 1998; (11) Allen, Di Matteo, & Fabian 1999; (12) Terashima et al. 2000a; (13) Sambruna et al. 1999; (14) Nicholson et al. 1998; (15) Roberts et al. 1999; (16) Iyomoto et al. 1996; (17) Iyomoto et al. 1997; (18) Wilson et al. 1998; (19) Ishisaki et al. 1996; (20) Makishima et al. 1994; (21) Mizuno et al. 1999; (22) Ho et al. 1999; (23) Terashima et al. 1999; (24) Dahlem et al. 1998; (25) Ptak et al. 1996; (26) Terashima et al. 1998b

H α luminosities: (1) L. C. Ho 2000, private communication; (2) Ho et al. 1997b; (3) Keel 1983; (4) Taniguchi et al. 1996; (5) Storchi-Bergmann et al. 1993; (6) Véron-Cetty & Véron 1986; (7) Storchi-Bergmann & Pastriza 1989; others Ho et al. 1997a, b

Figure Captions

Fig. 1.— (a) Correlation between X-ray and $H\alpha$ luminosities for LINER 1s, low-luminosity Seyfert 1s, and luminous type 1 AGNs taken from Ward et al. (1998)., (b) Correlation between X-ray and $H\alpha$ fluxes for the same sample as (a). The triangle and the cross correspond to NGC 1365 and NGC 4438, respectively.

Fig. 2.— $\log L_X/L_{H\alpha}$ for low-luminosity Seyfert 1s, low-luminosity Seyfert 2s, LINER 1s, and LINER 2s. The symbol " $<$ " denotes an upper limit.





