

# New observations of the halo radial temperature structure in NGC7662

C. Sandin, D. Schönberner, M.M. Roth, M. Steffen,  
A. Monreal-Ibero, P. Böhm, U. Tripphahn

Astrophysikalisches Institut Potsdam, An der Sternwarte 16, D-14482 Potsdam, Germany  
email: CSandin@aip.de

**Abstract.** We report on our studies of the physical structure of the planetary nebula (PN) NGC7662. Using (3D) Integral Field Spectroscopy we have been able to measure the electron temperature more accurately and at a larger number of radial locations than before. Here we briefly present our method by which we find a strong positive temperature gradient with increasing radius. According to hydrodynamic models a hot halo, when compared to the central star, can be the product of the passage of an ionization front (e.g. Marten 1993). Such a gradient is not found in equilibrium models, and this finding – when confirmed for other objects – strongly advocates the use of hydrodynamic models when modeling PN halos.

**Keywords.** hydrodynamics, instrumentation: spectrographs, methods: data analysis, techniques: spectroscopic, (ISM:) planetary nebulae: individual (NGC7662)

---

## 1. Observations

The data presented here was obtained with the PMAS instrument placed on the 3.5m telescope at Calar Alto in September 2005. In three different pointings, forming a *mosaic*, the PMAS integral field unit (IFU) covered a distance out to 40'' west of the central star. The wavelength coverage was 3436–5093 Å with a spectral resolution of 3.6 Å. The IFU holds 16 × 16 separate fibers that each represents a so called *spaxel* – covering 1'' × 1'' on the sky. Any number of spaxels can be combined to create a final spectrum with an increased S/N ratio, compared to that of a single spaxel. We have extracted six bands of spaxels from the three pointings. The flux in the spectra of the consecutive bands at increasing radius is a sum of (the spatial width in arcseconds of each band is specified as a subscript): 2<sub>1</sub>, 4<sub>1</sub>, 16<sub>2</sub>, 36<sub>3</sub>, 63<sub>4</sub>, and 128<sub>8</sub> spaxels, respectively.

## 2. Results

To calculate the electron temperature  $T_e$  at each of the six radial positions represented by the bands we use the three [OIII] 4363, 4959, 5007 Å lines out of which [OIII] 4363 Å is the weakest. Since we do not have a suitable sky frame the flux of the Hg 4359 Å sky line is removed from that of [OIII] 4363 Å by decomposition. We note that Middlemass et al. (1991) in a previous observation of NGC7662 removed the same sky line using a sky frame in which a significant amount of flux appears to be present in the [OIII] line (see Fig. 3b *ibid.*). This most likely indicates that the halo extends to radii larger than where the sky was measured. By accounting for this “missing” flux we have found that their measured [OIII] temperature increases by a factor of about 2.5 (see below).

The radial temperature structure is presented together with the surface brightness (SB) of [OIII] 4363 Å and error estimates in Fig. 1. A strong temperature gradient is seen




Figure 1. Radial temperature structure

Middlemass, D., Clegg, R.E.S., Walsh, J.R. & Harrington, J.P. 1991 *MNRAS* 251, 284

Tylenda, R. 2003, in: Kwok, S., Dopita, M. & Sutherland, R. (eds.), *Hot Haloes of Planetary Nebulae*, Proc. IAU Symposium No. 209 (San Francisco: ASP), p. 451