

Required experimental accuracy to select between supersymmetrical models

David Grellscheid

DA C n v s t o Ca br M
Ca br MCB A K

December 24, 2018

Abstract

We present a method to determine the required experimental accuracy to select between supersymmetrical models. The method is based on the observation that the required accuracy is determined by the number of parameters to be determined. The method is applied to the Minimal Supersymmetric Standard Model (MSSM) and the Next-to-Minimal Supersymmetric Standard Model (NMSSM). The results show that the required accuracy is significantly higher for the NMSSM than for the MSSM. The method is also applicable to other supersymmetrical models.


Background

In this paper we discuss the required experimental accuracy to select between supersymmetrical models. The method is based on the observation that the required accuracy is determined by the number of parameters to be determined. The method is applied to the Minimal Supersymmetric Standard Model (MSSM) and the Next-to-Minimal Supersymmetric Standard Model (NMSSM). The results show that the required accuracy is significantly higher for the NMSSM than for the MSSM. The method is also applicable to other supersymmetrical models.

At the root of our procedure stands the observation that an n -parameter supersymmetrical model can be described by a set of n parameters. The required accuracy to determine these parameters is determined by the number of parameters. The method is applied to the Minimal Supersymmetric Standard Model (MSSM) and the Next-to-Minimal Supersymmetric Standard Model (NMSSM). The results show that the required accuracy is significantly higher for the NMSSM than for the MSSM. The method is also applicable to other supersymmetrical models.

Supersymmetrical models are characterized by a large number of parameters. The required accuracy to determine these parameters is determined by the number of parameters. The method is applied to the Minimal Supersymmetric Standard Model (MSSM) and the Next-to-Minimal Supersymmetric Standard Model (NMSSM). The results show that the required accuracy is significantly higher for the NMSSM than for the MSSM. The method is also applicable to other supersymmetrical models.

The natural outcome to be probed at or stan ar n at on a or t s
 such as MINUIT bas on ot v r r u ar natur ot otpr nts bou ar s w
 or nat s n t k us on ooup s, a pnts A or pro s n appoa w w ar
 urrit wor n on st Mus o G M A or t s w ar or robust a a nst t
 o urrit u pnts o o obsrvab s to b p ott a a nst t Max s n M
 turn out to b so w at rt, a o un t to b us to pot ass rat os rat t an
 t ass s r t to nat t M p h h oot p ots on t Nov a s oot h Y
 ass p t n s



Fur Footprints at b s ann n ran o pnts n t at on o nat on
 t or a oot t r r ons M o s p at ur o

In w av s own t at t s parat on ot t r r o s s poss b n t at on
 o nat s as s ur w on oot nput para s at a x u
 $\theta = 90^\circ$ o st n us t o s a o b n t or h a an xp m ta a ura o o
 about s r qu r t M nat on o bot $m_{\tilde{g}}/m_{\tilde{q}_{avg}}$ an $m_{\tilde{e}_L}/m_{\tilde{e}_R}$ w ou
 b a ab n a o b n n ar o t LHC ana s

t our n w auto at appoa w w ab to t t s r s u t to s o s parat on
 or $0^\circ \leq \theta \leq 90^\circ$ an ar urrit wor n on t t n t towar s t o s on a t
 s an ov a our nput para s

oss b otur r t ons or t s wor ou n u Ma stu oot t s a o s
 w o not n s sar av to b str n o s Lar o prov nts an probab b a
 n t s n oot n n at on a or t an t o o n v st at obsrvab s

References

- e.g. A LA Co laborat on *Detector and Physics Performance Technical Design Report* CE LHC C apt an r s t
- G A Bar oro Z was Eur s J C27 hep-ph/0210058
- B C A ana D Gr s an F u o JHE 0205 hep-ph/0111057

B C A ana Comput s Co un 143 hep-ph/0104145

CE ro ra L brar D <http://wwwinfo.cern.ch/asd/index.html>

e.g. DE Go b *Genetic Algorithms in Search, Optimization & Machine Learning*
A son

B C A ana Kra oro JHE 0303 hep-ph/0302102