

Dynamical screening away from equilibrium: hard photon production and collisional energy loss

R. Baier and M. Dirks*

Fakultät für Physik, Universität Bielefeld, D-33501 Bielefeld, Germany

K. Redlich

Institute for Theoretical Physics, University of Wrocław,

PL-50204 Wrocław, Poland

and

GSI, PF 110552, D-64220 Darmstadt, Germany

We investigate the production rate for hard real photons and the collisional energy loss in the quark-gluon plasma away from chemical equilibrium. Applying the Hard-Thermal-Loop resummation scheme away from equilibrium, we can show that Landau damping provides dynamical screening for both fermion and boson exchange present in the two quantities.

I. INTRODUCTION

Considerable effort is being made to experimentally realize a new deconfinement phase of hadronic matter, the quark-gluon plasma, in ultra-relativistic heavy ion collisions. In parallel important progress has also been made in the theoretical understanding of this phase which is central for the experimental detection to be successful. For an equilibrated system in particular the Hard-Thermal-Loop (HTL) resummation scheme [1–3] allows to take consistently into account collective effects on the soft scale within (resummed) perturbation theory [4]. A number of important quantities, with unscreened mass singularities preventing any prediction within a fixed order perturbative calculation, can be obtained consistently in this scheme.

It has been realized that a valid comparison of theoretical predictions with experimental data should include non-equilibrium effects beyond the simplified assumption of thermal equilibrium, because a fully equilibrated plasma might not be realized. In this work we therefore explore the role of collective soft scale phenomena away from equilibrium and the generalization of the HTL resummation scheme. This will be done on the relevant examples of hard photon production [5–11] and the collisional energy loss [12–14] which are known to be sensitive to soft scale physics. The following discussion is based on [15,16] where further details can be found.

II. APPROXIMATION SCHEME: CHEMICAL NON-EQUILIBRIUM

Our analysis assumes the scenario of chemical non-equilibrium that has been found to be a relevant, tractable approximation to the pre-equilibrium dynamics in heavy ion collisions [17–20]. It is inspired from an analysis of the equilibration process in a heavy ion collision where the elastic momentum exchange is found to be much more effective than the inelastic particle production. While local thermal equilibrium might therefore be quickly established the parton densities are predicted to stay away from equilibrium through to the hadronization of the plasma [18]. Thereby the undersaturation of quarks will be especially dramatic as has been formulated in the hot-gluon scenario [17]. Following [18] and as in [10,21] we parameterize this scenario in terms of factorized fugacities inserted into the distributions which are assumed to be equilibrated in momentum space

$$\begin{aligned}\tilde{n}(X, p) &= \begin{cases} \lambda_q(X)n_F(|p_0|), & p_0 > 0 \\ 1 - \lambda_q n_F(|p_0|), & p_0 < 0 \end{cases}, \\ n(X, p) &= \begin{cases} \lambda_g(X)n_B(|p_0|), & p_0 > 0 \\ -[1 + \lambda_g(X)n_B(|p_0|)], & p_0 < 0. \end{cases} \quad (1)\end{aligned}$$

We do not attempt to investigate the microscopic non-equilibrium evolution of the distribution functions in the quark-gluon plasma, the evolution of the fugacities $\lambda(X)$ on the large scale X of inelastic processes is taken as input e.g. from [18]. On this background we calculate the emission and absorption rates for probes external to the plasma such as electromagnetic rates and the collisional energy loss of a projectile. For the description of the dynamics we rely on the Closed-Time-Path formulation of statistical field theory [22,23]. The simplified scenario of chemical non-equilibrium allows for the standard approximation scheme of neglecting any but the second order non-equilibrium correlations and performing a gradient expansion [23–26], here in terms of the large scale $X \sim 1/g^4 T$ of the inelastic particle production. Scales fast with respect to this, in particular both the hard ($\sim 1/T$) and soft ($\sim 1/gT$), $g < 1$, scales involved in the HTL resummation scheme, can be transformed to momentum space. To lowest order in the gradients a

*talk presented at 5th International Workshop on Thermal Field Theories and their Applications, Regensburg, Germany, August 1998

perturbative expansion can be set up very similar to the equilibrium case with free in medium propagators depending on the modified distributions Eq. (1). In terms of Wigner variables (X, p) :

$$\begin{aligned} iD_{21} &= 2\pi\varepsilon(p_0)\delta(p^2)(1 + n(X, p_0)), \\ iS_{21}(X, p) &= \not{p} 2\pi\delta(p^2)\varepsilon(p_0)(1 - \tilde{n}(X, p_0)). \end{aligned} \quad (2)$$

Predictions for absorption and emission rates as calculated in this approximation scheme will incorporate the dependence on $\lambda(X)$ locally. As in the equilibrium situation, finite results for the real hard photon spectrum and the collisional energy loss can be expected only after resummation of leading HTL corrections. Away from equilibrium taking higher order corrections into account brings about additional terms in the perturbative expansion [27,28] which vanish due to detailed balance in the equilibrium situation. A careful analysis is therefore required in order to demonstrate that Landau damping provides dynamical screening of the singularities also away from equilibrium.

III. SOFT FERMION EXCHANGE IN HARD PHOTON PRODUCTION

The production of hard real photons to lowest order arises from annihilation and Compton scattering. The rate can be obtained from the absorptive part of the two-loop approximation to the photon selfenergy as shown in Fig. 1.




FIG. 1. Diagrams contributing to hard real photon production in fixed leading order for hard exchanged momentum (a) and in HTL-resummed perturbation theory for soft exchanged momentum (b). The parameter k_c , separating hard and soft scales is introduced in intermediate steps of the calculation [29]

The exchange of a massless quark in these processes induces a logarithmic mass singularity when calculating to fixed lowest order, Fig. 1(a). In equilibrium a solution to this problem is found within the HTL resummation scheme that accounts for the fact that for soft exchanged momentum the dispersion of the exchanged particle is modified by leading interaction effects. Within the resummed theory in particular Landau damping effects are included that dynamically screen the singularity on the soft scale $m_q \sim gT$ [5,6]. Away from equilibrium, in

leading order of the gradient expansion, the resummed fermion propagator can be written

$$\begin{aligned} S_{12}^*(p) &= -\tilde{n}(X, p_0) \left(\frac{1}{\not{p} - \Sigma_R + i\varepsilon p_0} - \frac{1}{\not{p} - \Sigma_R^* - i\varepsilon p_0} \right) \\ &\quad - \frac{1}{\not{p} - \Sigma_R + i\varepsilon p_0} \left[(1 - \tilde{n}(X, p_0)) \Sigma_{12} \right. \\ &\quad \left. + \tilde{n}(X, p_0) \Sigma_{21} \right] \frac{1}{\not{p} - \Sigma_R^* - i\varepsilon p_0}, \end{aligned} \quad (3)$$

with a first term having the structure of the equilibrium expression and a second term being nonzero only when detailed balance is violated, $\Sigma_{12} \neq -e^{-p_0/T} \Sigma_{21}$. Two remarks have to be added about the modified structure of the propagator, Eq. (3). First we find that the non-equilibrium modification of the dispersion as defined by the pole of the retarded propagator in the first term of Eq. (3) can be absorbed into a redefinition of the soft scale parameter used in the equilibrium situation

$$\tilde{m}_q^2 = \frac{g^2}{2\pi^2} C_F \int_0^\infty$$

Given the close analogy of this effective resummed propagator with the equilibrium expression, the remaining steps are performed in parallel with the equilibrium calculation [5,6] taking only the modified dispersion into account. We find the rate for hard photon production to depend on the modified scale parameter Eq. (4) that enters as a prefactor and screens the logarithmic singularity:

$$E_\gamma \frac{dR}{d^3q} = e_q^2 \frac{\alpha\alpha_s}{2\pi^2} \lambda_q T^2 e^{-E_\gamma/T} \times \left[\frac{2}{3} \left(\lambda_g + \frac{\lambda_q}{2} \right) \ln \left(\frac{2E_\gamma T}{\tilde{m}_q^2(\lambda_q, \lambda_g)} \right) + \frac{4}{\pi^2} C(E_\gamma, T, \lambda_q, \lambda_g) \right], \quad (9)$$

where the nonsingular contribution $C(E_\gamma, T, \lambda_q, \lambda_g)$ has been calculated explicitly in [15].

The result Eq. (9) is independent of the separating scale k_c introduced in intermediate steps of the calculation by matching the soft partial result with the result from a fixed order calculation for hard exchanged momentum. In this context it is important to point out that no additional singularities are showing up away from equilibrium although at one loop order the extra term in the fermion propagator analogous to Eq. (3) is pinch singular on the mass shell [27]. The important observation is that for hard real photon production the singular configuration lies only on the boundary of the available phase space that is restricted to spacelike momentum exchange. The effective propagator for the hard quark exchange to be used in the upper part of Fig. 1(a) follows from Eq. (3) in the one loop approximation,

$$\delta S_{12}(p)|_{p^2 \leq -k_c^2} \hat{=} -\frac{1}{(p^2)^2} \not{p} \Sigma_{12} \not{p}. \quad (10)$$

There remains no explicit dependence on the distribution $\tilde{n}(X, p_0)$. A mass singularity is induced, which, however, is dynamically screened upon matching with the contribution from the soft exchange with the propagator Eq. (8).

IV. SOFT (GAUGE) BOSON EXCHANGE – COLLISIONAL ENERGY LOSS

The collisional energy loss of a propagating heavy fermion with mass M [13,14], which is not thermalized, is obtained from the absorptive part of the fermion self-energy, i.e. from its 21-component [4]. The corresponding diagrams are shown in Fig. 2. To lowest order the loss arises from the elastic interaction with the medium constituents via the exchange of one gauge boson, Fig. 2(a); again mass singularities are known to arise for massless photon/gluon exchange.

It is important to recall that the energy loss is obtained from these diagrams only upon weighting the loop integration with one extra power of the energy exchanged:

$$-\frac{dE}{dx} = -\frac{1}{4E} \text{Tr}[(\not{q} + M) i \Sigma'_{21}(X, q)] = \frac{e^2}{4E} \int \frac{d^3\vec{p}}{(2\pi)^4} \int \frac{dp_0 p_0}{v} \times \text{Tr}[(\not{q} + M) \gamma_\mu i S_{12}^{T=0}(p-q) \gamma_\nu] i D_{21}^{\mu\nu}(X, p). \quad (11)$$

The extra power of p_0/v distinguishes dE/dx from the interaction/damping rate and makes it well defined in equilibrium within the resummed theory.




FIG. 2. Diagrams contributing to the collisional energy loss in fixed leading order for hard exchanged momentum (a) and in HTL-resummed perturbation theory for soft exchanged momentum (b).

Away from equilibrium the resummed propagator for soft gauge boson-exchange has a modified structure similar to the fermion propagator [30]; longitudinal and transverse components in Coulomb gauge read

$$D_{21}^{*L/T} = (1 + n(X, p_0))(D_R^{*L/T} - D_A^{*L/T}) + D_R^{*L/T} [n(X, p_0) \Pi_{21}^{L/T} - (1 + n(X, p_0)) \Pi_{12}^{L/T}] D_A^{*L/T}.$$

The discussion follows the lines of Sec. III. The modifications of the dispersion relation in HTL approximation can again be absorbed into a redefinition of the plasma frequency according to

$$m_\gamma^2 = \frac{e^2 T^2}{9} \quad \rightarrow \quad \tilde{m}_\gamma^2 = \frac{4e^2}{3\pi^2} \int_0^\infty dk k \tilde{n}(X, k) = \lambda_f m_\gamma^2 \quad (12)$$

in QED. For the role of the extra term however an important difference with the fermionic case arises. In the bosonic case the contribution from $\Pi^+ = \frac{1}{2}(\Pi_{12} + \Pi_{21})$ is found to dominate over $\Pi^- = \frac{1}{2}(\Pi_{12} - \Pi_{21}) = iIm\Pi_R$ for both longitudinal and transverse components [30]

$$-\Pi_{L/T}^+ = \frac{1}{p_0} R \Pi_{L/T}^-, \quad R = \frac{\int_0^\infty dk k^2 \tilde{n}(k)(1 - \tilde{n}(k))}{\int_0^\infty dk k \tilde{n}(k)} \sim T, \quad (13)$$

and therefore cannot be neglected in general. However, both contributions have a symmetry under $p_0 \rightarrow -p_0$, while Π^- is odd, Π^+ is even. Given the additional power of p_0 present in the energy loss rate the contribution of Π^+ therefore integrates to zero in this special case as

$$\int_{-vp}^{vp} dp_0 p_0 \Pi^+(X, p_0) = 0. \quad (14)$$

As a result the dominant contribution to dE/dx comes from Π^- and the corresponding terms in the resummed propagator rearrange into

$$D_{21}^{*L/T} \Big|_{dE/dx} \stackrel{\Delta}{=} \frac{1}{2}(D_R^{*L/T} - D_A^{*L/T}) \quad \text{in } dE/dx, \quad (15)$$

which is similar in structure to the equilibrium expression and allows to perform the remaining steps as in [13]. As for hard photon production no additional (pinch) singularities arise away from equilibrium for spacelike momentum exchange. For energies $E \ll M^2/T$ we find the energy loss away from equilibrium to be well defined

$$-\frac{dE}{dx} = \frac{3e^2 \tilde{m}_\gamma^2}{8\pi v} \left(1 - \frac{1-v^2}{2v} \ln \frac{1+v}{1-v} \right) \left[\ln \frac{ET}{3M\tilde{m}_\gamma} + A(v) \right], \quad (16)$$

with Landau damping dynamically screening the logarithmic mass singularity. Following the calculation for the equilibrium case it turns out that the regular contribution $A(v)$ does not depend on the fugacity factor λ_f , and that it is therefore the same as discussed in [13].

The generalization for a heavy quark propagating through a QCD-plasma is obtained [16] with the help of the expressions given in [14],

$$-\frac{dE}{dx} = \frac{g^2 \tilde{m}_g^2}{2\pi v} \left(1 - \frac{1-v^2}{2v} \ln \frac{1+v}{1-v} \right) \left[\ln \frac{ET}{3M\tilde{m}_g} - \frac{\ln 2}{(1 + \lambda_q N_f / 6\lambda_g)} + A(v) \right], \quad (17)$$

introducing the QCD coupling g and the thermal gluon mass, including the proper quark and gluon fugacities: $\tilde{m}_g^2 = g^2 T^2 (\lambda_g + \lambda_q \frac{N_f}{6}) / 3$. Because in the QCD case contributions from the scatterings off quarks and gluons in the plasma have to be added, a dependence on λ_q and λ_g is found also in the regular part of Eq. (17).

V. DISCUSSION AND CONCLUSIONS

From the results given in Eqs. (9) and (17), respectively, both the hard photon production rate and the collisional energy loss are found to be suppressed by powers of the fugacity in an undersaturated plasma that is likely to be created in a relativistic heavy ion experiment [17,18]. Taking into account the extra off-equilibrium term in the propagator there is no dependence in the final result on the distribution of the exchanged particle that would have introduced additional powers of fugacities. In the case of the photon production rate this result is especially important phenomenologically as higher powers of the small quark fugacity could seriously question its observability. It is encouraging that both rates can be obtained consistently in perturbation theory away from

equilibrium with dynamical screening occurring also for $\lambda \neq 1$. We argued that no additional pinch singularities [27] do arise in the quantities considered that involve spacelike momentum exchange only. The treatment given here however is not applicable to quantities involving on-shell momentum exchange as e.g. virtual photon production [31–33]. Further progress in these cases requires to take consistently into account the explicit time variations in the distributions [34,35] via the inclusion of gradient terms which could be neglected in the presented cases.

VI. ACKNOWLEDGEMENTS

We like to thank D. Schiff and M.H. Thoma for helpful discussions and critical comments. M.D. is supported by DFG, Contract BA 915/4-2.

-
- [1] R. D. Pisarski, *Physica* **A158**, 146 (1989).
 - [2] E. Braaten and R. D. Pisarski, *Nucl. Phys.* **B337**, 569 (1990).
 - [3] J. Frenkel and J. C. Taylor, *Nucl. Phys.* **B334**, 199 (1990).
 - [4] M. Le Bellac, *Thermal Field Theory* (Cambridge University Press, Cambridge, 1996).
 - [5] J. Kapusta, P. Lichard, and D. Seibert, *Phys. Rev.* **D44**, 2774 (1991).
 - [6] R. Baier, H. Nakkagawa, A. Niégawa, and K. Redlich, *Z. Phys.* **C53**, 433 (1992).
 - [7] E. Shuryak and L. Xiong, *Phys. Rev. Lett.* **70**, 2241 (1993).
 - [8] M. Strickland, *Phys. Lett.* **B331**, 245 (1994).
 - [9] B. Kämpfer and O. Pavlenko, *Z. Phys.* **C62**, 491 (1994).
 - [10] M. H. Thoma and C. T. Traxler, *Phys. Rev.* **C53**, 1348 (1996).
 - [11] D. K. Srivastava, M. G. Mustafa, and B. Müller, *Phys. Rev.* **C56**, 1064 (1997).
 - [12] J. D. Bjorken, Fermilab Report PUB-82/59-THY (unpublished) (1982).
 - [13] E. Braaten and M. H. Thoma, *Phys. Rev.* **D44**, 1298 (1991).
 - [14] E. Braaten and M. H. Thoma, *Phys. Rev.* **D44**, 2625 (1991).
 - [15] R. Baier, M. Dirks, K. Redlich, and D. Schiff, *Phys. Rev.* **D56**, 2548 (1997).
 - [16] R. Baier, M. Dirks, and K. Redlich, in preparation.
 - [17] E. Shuryak, *Phys. Rev. Lett.* **68**, 3270 (1992).
 - [18] T. S. Biro *et al.*, *Phys. Rev.* **C48**, 1275 (1993).
 - [19] K. Geiger and J. I. Kapusta, *Phys. Rev.* **D47**, 4905 (1993).
 - [20] S. M. H. Wong, hep-ph/9801427 (unpublished) (1997).
 - [21] B. Kämpfer, O. P. Pavlenko, A. Peshier, and G. Soff, *Phys. Rev.* **C52**, 2704 (1995).

- [22] N. P. Landsman and C. G. van Weert, Phys. Rept. **145**, 141 (1987).
- [23] K.-C. Chou, Z.-B. Su, B.-L. Hao, and L. Yu, Phys. Rept. **118**, 1 (1985).
- [24] S. Mrowczynski and U. Heinz, Ann. Phys. **229**, 1 (1994).
- [25] E. Calzetta and B. L. Hu, Phys. Rev. **D37**, 2878 (1988).
- [26] P. A. Henning, Phys. Rept. **253**, 235 (1995).
- [27] T. Altherr and D. Seibert, Phys. Lett. **B333**, 149 (1994).
- [28] T. Altherr, Phys. Lett. **B341**, 325 (1995).
- [29] E. Braaten and T. C. Yuan, Phys. Rev. Lett. **66**, 2183 (1991).
- [30] M. E. Carrington, H. Defu, and M. H. Thoma, hep-ph/9708363 (unpublished) (1997).
- [31] R. Baier, M. Dirks, and K. Redlich, Phys. Rev. **D55**, 4344 (1997).
- [32] M. Le Bellac and H. Mabilat, Z. Phys. **C75**, 137 (1997).
- [33] A. Niégawa, Eur. Phys. J. **C5**, 345 (1998).
- [34] P. F. Bedaque, Phys. Lett. **B344**, 23 (1995).
- [35] C. Greiner and S. Leupold, hep-ph/9804239 (unpublished) (1998).