

On $N = 1$ gauge models from geometric engineering in M-theory

A. Belhaj^{1,2,*}, L.B. Drissi¹, J. Rasmussen^{2,3,†}

¹ National Grouping of High Energy Physics, GNPHE, *and*
Lab/UFR High Energy Physics, Department of Physics
Faculty of Sciences, Rabat, Morocco

² Department of Mathematics and Statistics, Concordia University
Montréal, PQ, Canada H4B 1R6

³ Centre de Recherches Mathématiques, Université de Montréal
C.P. 6128, succursale centre-ville, Montréal, PQ, Canada H3C 3J7

October 29, 2018

Abstract

We study geometric engineering of four-dimensional $N = 1$ gauge models from M-theory on a seven-dimensional manifold with G_2 holonomy. The manifold is constructed as a K3 fibration over a three-dimensional base space with ADE geometry. The resulting gauge theory is discussed in the realm of (p, q) webs. We discuss how the anomaly cancellation condition translates into a condition on the associated affine ADE Lie algebras.

*belhaj@crm.umontreal.ca

†rasmusse@crm.umontreal.ca

1 Introduction

Type IIA superstring compactification on a K3 space with A_{n-1} singularities develops an enhanced $SU(n)$ gauge symmetry in six dimensions [1]. This can be further compactified on a real two-dimensional base space, resulting in a four-dimensional $N = 2$ quantum field theory (QFT₄). The gauge fields and (massless) matter fields are described by wrappings of D-branes around cycles in the Calabi-Yau threefold defined by the K3 fibrations. The gauge group and matter content of the QFT₄ thus depend on the non-trivial base geometry. This program is referred to as geometric engineering of $N = 2$ QFT₄. These models, which give exact results for the moduli space of the Coulomb branch, may be represented by quiver diagrams similar to Dynkin diagrams of ordinary or affine Lie algebras [2, 3, 4, 5, 6, 7, 8]. Let the base space consist of a collection of intersecting two-spheres, each of which will give rise to an SU gauge-group factor. To each of these is associated a quiver or Dynkin diagram node, and for each pair of groups with matter in bi-fundamental representations, the two corresponding nodes are connected by a line.

One should distinguish three cases: models with negative beta function, models with conformal invariance, requiring the vanishing of the beta function, and models with positive beta function. The first models are represented by ordinary *ADE* Dynkin diagrams with only bi-fundamental matter. A K3 fiber with A_{n_i-1} singularities generates an $SU(n_i)$ gauge-group factor, resulting in the gauge group

$$G = \bigotimes_{i=1}^m SU(n_i), \quad (1)$$

where m is the number of intersecting two-spheres. The second models are classified by two categories according to the type of intersecting geometry of the base space: (i) $N = 2$ superconformal field theory in four dimensions (SCFT₄), based on *finite ADE* Lie algebra singularities. It has gauge group similar to the first class but matter in both fundamental (n_i) and bi-fundamental ($n_i, \bar{n}_j$) representations of G . (ii) $N = 2$ SCFT₄ with gauge group

$$G = \bigotimes_{i=1}^m SU(s_i n) \quad (2)$$

and bi-fundamental matter only. Here n is an unspecified positive integer. This second category of scale invariant field models is classified by *affine ADE* Lie algebras. The positive integers s_i appearing in G are the Coxeter labels (sometimes referred to as marks or Dynkin

numbers); they form a positive-definite integer vector (s_j) satisfying

$$\sum_{j=1}^m K_{ij} s_j = 0. \quad (3)$$

K is the Cartan matrix of the affine Lie algebra, and $m = r + 1$ with r being the rank of the underlying Lie algebra. Models with positive beta function are discussed in [9].

Quite recently, four-dimensional quiver gauge models preserving only four supercharges have attracted a lot of attention. Work has been done using either type II superstrings on Calabi-Yau threefolds [10, 11, 12, 13, 14, 15], or M-theory on G_2 manifolds [16, 17]. The aim of this work is to contribute to this program. We study M-theory on seven-dimensional manifolds with G_2 holonomy group. They are realized explicitly as K3 fibrations over a three-dimensional base space. First, though, we review the geometric engineering of $N = 2$ QFT₄ in type IIA superstring theory. We then extend this to geometric engineering of $N = 1$ QFT₄ in M-theory, where particular emphasis is put on the three-dimensional ADE base geometry. Comparing our construction to the related D6-brane scenario described using the method of (p, q) webs, we determine the gauge group and matter content of our $N = 1$ QFT₄.

2 Geometric engineering of $N = 2$ gauge theory

A well-known way to get supersymmetric quantum field theory (QFT) from superstrings, M- or F-theory, is to consider the compactification on a *singular* manifold X with K3 fibration over a base space B . The gauge group G and matter content of the QFT are defined by the singularities of the fiber and the non-trivial geometry of the base space. In particular, the gauge coupling g is proportional to the inverse of the square root of the volume of the base: $V(B) = g^{-2}$. The complete set of physical parameters of the QFT is related to the moduli space of the manifold. Moreover, several exact results for the Coulomb branch of $N = 2$ QFT₄ embedded in type IIA superstring theory are obtained naturally by local mirror symmetry of Calabi-Yau threefolds CY^3 . The latter are realized as a K3 fibration with ADE singularity over a projective P^1 complex curve or a collection of intersecting P^1 curves.

Let us briefly outline the main steps in obtaining $N = 2$ QFT₄ from type IIA superstring theory on CY^3 . The idea is subsequently illustrated by a simple example. Depending on the sought gauge group, one first specifies the ADE singularity of the K3 fiber. Then one considers the limit in which the volume $V(B)$ of the base of CY^3 is very large so that gravitational effects may be ignored. This allows one to consider locally (i.e., near the singularities) the K3 fibers as

non-compact ALE spaces with the same singularities. Finally, one examines the propagation of the type IIA superstrings on CY^3 in the presence of D2-branes wrapping the vanishing two-cycles in the ALE spaces. This enables one to make conclusions about the gauge group and matter content.

As an illustration, we now consider the situation where the K3 fiber has an A_1 singularity. In the vicinity of the singularity, such a fiber may be viewed as an ALE space with A_1 singularity and described by

$$z_1^2 + z_2^2 + z_3^2 = 0. \quad (4)$$

Using a simple change of variables, this is equivalent to

$$xy = z^2, \quad (5)$$

where x , y and z are complex variables. This geometry has a nice physical realization as a two-dimensional $N = 2$ linear sigma model with $U(1)$ gauge symmetry. To see this, consider the three chiral fields ϕ_i with charges $(1, -2, 1)$. The coordinates of the ALE space can then be expressed in terms of $U(1)$ gauge invariants:

$$\begin{aligned} x &= \phi_1^2 \phi_2, \\ y &= \phi_3^2 \phi_2, \\ z &= \phi_1 \phi_2 \phi_3. \end{aligned} \quad (6)$$

This is related to the so-called D -term in the bosonic potential $U(\phi_1, \phi_2, \phi_3)$ in supersymmetric theories with four supercharges:

$$U(\phi_1, \phi_2, \phi_3) = (\phi_1 \phi_1 + \phi_3 \phi_3 - 2\phi_2 \phi_2 - R)^2. \quad (7)$$

In this equation, R is the coupling parameter of the $U(1)$ Fayet-Iliopoulos (FI) term one may introduce in the Lagrangian model. In the superfield language, the action then reads

$$S(\Phi, V) = \int d^2x d^4\theta \bar{\Phi} e^V \Phi - R \int d^2x d^4\theta V, \quad (8)$$

where Φ and V are the chiral and gauge superfields, respectively. From this, one sees that the $U(1)$ Cartan subgroup of the $SU(2)$ symmetry of the singularity of K3 carries the gauge symmetry of the $N = 2$ supersymmetric linear sigma model. The presence of the FI term resolves the singularity of the potential $U(\phi_1, \phi_2, \phi_3)$. Geometrically, this corresponds to replacing the singular point $x = y = z = 0$ by a P^1 curve parameterized by a new variable x' defined as

$x' = x/z$. In the new local coordinates (x', y, z) , the equation of the original A_1 singularity becomes

$$x'y = z \tag{9}$$

thus removing the singularity (which is now “hidden” in the singular transformation $x' = x/z$). In the field theory language, this corresponds to a positive value of the FI coupling R .

The next step is to consider the propagation of type IIA superstrings in this background. In this case, D2-branes wrapping around the blown-up P^1 curves give rise to a pair of vector particles, $W^\pm$, one for each of the two possible orientations of the wrappings. The particles have masses proportional to the volume of the blown-up (real) two-sphere described by x' . $W^\pm$ are charged under the $U(1)$ field Z obtained by decomposing the type IIA superstring three-form in terms of the harmonic form on the two-sphere and a one-form not in the K3 fiber. In the limit where the blown-up two-sphere shrinks to a point, the three vector particles become massless, and they form an $SU(2)$ adjoint representation. We thus obtain an $N = 2$ $SU(2)$ gauge symmetry in six dimensions. A further compactification on the base B_2 (here chosen as a real two-sphere), gives $N = 2$ pure $SU(2)$ Yang-Mills theory in four dimensions.

The geometric $SU(2)$ gauge theory analysis is extended straightforwardly to higher $SU(n)$ gauge groups and products thereof. Extensions to general simply-laced ADE gauge groups are obtained using more general Calabi-Yau threefolds [6, 7]. Up to some pertinent details, non-simply-laced $BCFG$ groups may also be constructed [8, 18], see also [19, 20]. The incorporation of matter was alluded to in the Introduction, and is governed by a non-trivial base space geometry of the Calabi-Yau threefold. For example, consider a two-dimensional locus with an $SU(n)$ singularity and another locus with an $SU(m)$ singularity. If they intersect or meet at a point, the (mixed) wrapped two-cycles give rise to bi-fundamental matter of the $SU(n) \otimes SU(m)$ gauge group in four dimensions. This extends readily to multiple intersecting loci. It was also indicated in the Introduction how the associated toric quiver diagrams of $N = 2$ QFT₄'s are similar to the usual Dynkin diagrams of ordinary and affine Lie algebras.

3 Geometric engineering of $N = 1$ gauge theory

Here we discuss a straightforward way of elevating the geometric engineering of $N = 2$ gauge models in string theory to a similar construction of $N = 1$ gauge models in M-theory. Alternative studies of four-dimensional $N = 1$ gauge models in the framework of F- or M-theory may be found in refs. [21, 22, 23].

3.1 Outline of method

The method we will be using here is quite similar to the geometric engineering of $N = 2$ QFT₄ discussed above. Our starting point is a local description of M-theory compactification on a G_2 manifold with K3 fibration over a real three-dimensional base space B_3 . In this way, the scenario of type IIA superstring theory in six dimensions appears in seven dimensions in M-theory. The D2-branes are replaced by M2-branes, and we obtain an $N = 2$ gauge theory in seven dimensions. If we compactify this seven-dimensional theory on T^3 to four dimensions, we obtain an $N = 4$ gauge theory, while a compactification on $S^2 \times S^1$ leads to a system with eight supercharges (corresponding to $N = 2$) [22].

We are, however, interested in models with only four supercharges in four dimensions, i.e., $N = 1$ QFT₄. In order to reach such a model we need to compactify on a manifold whose three-dimensional base preserves 1/4 of the remaining 16 supercharges. The seven-dimensional K3 fibration must also have vanishing first Betti number, $b_1 = 0$, to meet the requirement of G_2 holonomy. An example of such a geometry has the three-dimensional sphere S^3 as base, in which case we obtain only *pure* $N = 1$ Yang-Mills theory [22, 23]. We shall, instead, consider (a collection of) non-trivial S^2 fibrations over a finite line segment.

As in the $N = 2$ case, the incorporation of matter may be achieved by introducing a non-trivial geometry in the base of the K3 fibration. This leads us to consider a three-dimensional intersecting geometry to describe a product gauge group with bi-fundamental matter.

3.2 Construction of the base geometry

Our construction of the non-trivial base geometry B_3 is quite simple and motivated by recent work on Lagrangian submanifolds in Calabi-Yau manifolds [24]. We shall consider the three-dimensional base space as a two-dimensional fibration over a one-dimensional base, where the fiber and the base each preserves half of the seven-dimensional M-theory supercharges. The entire base space B_3 is embedded in a three-dimensional complex Calabi-Yau manifold. As discussed below, this ensures that the elements of the intersection matrix of a collection of such base space constituents are integers. The Calabi-Yau manifolds are realized explicitly as a family of (deformed) *ADE* singular K3 surfaces over the complex plane. This geometry extends the family of elliptic curves on the complex plane used in F-theory compactifications down to eight dimensions [25]. In this way, non-trivial three-cycles, satisfying the constraints of the G_2 base geometry, constitute an intersecting geometry of *ADE* K3 surfaces fibered over a line

segment in the complex plane. The “*ADE* part” corresponds to a two-dimensional geometry obtained by resolving the *ADE* singularities of the K3 surfaces. As already mentioned, the *ADE* geometry is then either an (i) ordinary (finite Lie algebraic) *ADE* geometry, or an (ii) affine (infinite Lie algebraic) *ADE* geometry.

We start by examining the model with an ordinary A_1 singularity in the K3 fiber. This is subsequently extended to more general *ADE* geometries. The deformed A_1 geometry is given by

$$z_1^2 + z_2^2 + z_3^2 = \mu \quad (10)$$

where μ is a complex parameter. The real part of this parameter, being the radius squared of the sphere, is related to the FI terms (7) and (8), while the imaginary part can be identified with the NS-NS B -field in superstring compactifications. The A_1 threefolds may be obtained by varying the parameter μ over the complex plane parametrized by w :

$$z_1^2 + z_2^2 + z_3^2 = \mu(w). \quad (11)$$

The base space B_3 can then be viewed as a finite line segment with an S^2 fibration, where the radius r of S^2 vanishes at the two interval end points, and at the end points only [11]. The latter requirement ensures that no unwanted singularities are introduced. One way to realize this geometry is

$$r \sim \sin x \quad (12)$$

where x is a real variable parameterizing the interval $[0, \pi]$ in the w -plane.

This construction may be extended to more general *ADE* geometries where we have intersecting spheres according to affine *ADE* Dynkin diagrams. Here we consider a general model describing a family of deformed elliptic *ADE* singularities of K3 surfaces fibered over the complex plane, and their mirror partners in type II superstring compactifications. In type IIB superstring theory, the blow-up of these singularities can be described in a two-dimensional $N = 2$ linear sigma model with $U(1)^{\otimes(r+1)}$ gauge symmetry. It has $r + 5$ chiral multiplets ϕ_ℓ with vector charges q_ℓ^i , where r is the rank of the associated affine *ADE* Lie algebra. The D -terms corresponding to such classical vacua are given by

$$U = \sum_{i=1}^{r+1} \sum_{\ell=1}^{r+5} (q_\ell^i |\phi_\ell|^2 - R_i)^2, \quad (13)$$

where the FI terms are denoted R_i . The q_ℓ^i 's, being the quantum charges of ϕ_ℓ under the $U(1)^{\otimes(r+1)}$ gauge symmetry, are identified with the intersection matrix (known as Mori vectors in toric geometry) of the non-compact and compact divisors of the elliptic *ADE* K3

surfaces. For the compact divisors¹ used in the resolution of the elliptic *ADE* singularities, the intersection numbers are given by minus the affine *ADE* Cartan matrix K :

$$q_i^j = -K_{ij}, \quad i, j = 1, \dots, r+1. \quad (14)$$

This link stems from a nice correspondence between the *ADE* roots and the two-cycles, C , involved in the deformation of the elliptic singularities of the K3 surfaces [6]. In this way, the cohomology class of the deformed elliptic curve, T^2 , is represented by

$$[T^2] = \sum_{i=1}^{r+1} s_i C_i, \quad (15)$$

where s_i are the Coxeter labels of the affine Lie algebra.

The elliptic *ADE* threefolds may now be described using the toric geometry data of these threefolds and local mirror symmetry [7, 8, 18]. Applying this symmetry in the K3 fiber, the elliptic *ADE* type IIA mirror geometries, X^* , may be described by an algebraic geometry equation of the form

$$P(X^*) = \sum_{\ell=1}^{r+5} a_\ell(w) y_\ell. \quad (16)$$

$a_\ell = a_\ell(w)$ are varying complex moduli² in the w -plane, while the y_ℓ 's are complex variables satisfying

$$\prod_{\ell=1}^{r+5} y_\ell^{q_\ell^i} = 1, \quad \forall i = 1, \dots, r+1. \quad (17)$$

This defines the mirror elliptic *ADE* K3 fiber. Eq. (17) can be solved by a quasi-homogeneous polynomial in some variables (x, y, z, v) , and eq. (16) becomes

$$p_0(x, y, z; \lambda(w)) + \sum_{i=1}^{r+1} p_i(x, y, z, v; a_i(w)) = 0. \quad (18)$$

Here (x, y, z) are the homogeneous coordinates of the weighted projective space $WP_{3,2,1}^2$, while the index on a_i refers to a (physical) compact divisor. Parameterized by the complex structure modulus λ , $p_0 = 0$ now describes a family of elliptic curves over the w -plane:

$$x^2 + y^3 + z^6 + \lambda(w)xyz = 0. \quad (19)$$

The v -dependent term depends on the structure of the affine Lie algebra. In particular, the powers of the complex variable v appearing in the *ADE* mirror geometries are given by the

¹We shall be using a convention where the first $r+1$ divisors are compact, i.e., $q_1^j, \dots, q_{r+1}^j$ correspond to the compact divisors.

²Note that only $r+1$ of these are physical.

Coxeter labels of the corresponding affine Lie algebra. The geometry of affine A_2 , for instance, is given by

$$x^2 + y^3 + z^6 + \lambda(w)xyz + v[a_1(w)z^3 + a_2(w)yz + a_3(w)x] = 0. \quad (20)$$

A similar analysis of the simple A_1 model discussed in eqs. (10)-(12) could be done using the so-called mirror map in the Calabi-Yau moduli spaces [6]. Indeed, mirror symmetry relates the complexified FI terms T_j in type IIB geometry X to the coordinates, Z_j , of the complex moduli space of type IIA geometry:

$$T_j = B_j + iV(C_j) \sim -i \ln(Z_j). \quad (21)$$

$V(C_j)$ is the volume of the two-cycle C_j , whereas Z_j is given by

$$Z_j = \prod_{\ell=1}^{r+5} a_{\ell}^{q_{\ell}^j}. \quad (22)$$

For vanishing integrated NS-NS B -fields B_j , equation (21) reduces to

$$V(C_j) \sim -\ln(Z_j). \quad (23)$$

Under mirror symmetry, the constraints on the Kähler parameters of the two-cycles in type IIB geometry translate into conditions on the complex structures in type IIA geometry.

In the geometric setting, our choice of base geometry is given as a line segment fibered by intersecting spheres according to ADE affine Dynkin diagrams.

4 Gauge group and matter content

Having specified the base geometry of the G_2 manifold, we will discuss the corresponding gauge theory of the compactified M-theory. Our analysis here will be based on a dual type IIA superstring description. It turns out that, in addition to being dual to heterotic string compactification, M-theory on G_2 manifolds is also dual to type IIA string theory on Calabi-Yau threefolds in the presence of D6-branes wrapping Lagrangian sub-manifolds and filling the four-dimensional Minkowski space [10]. In particular, a local description of M-theory near the A_{n-1} singularity of K3 surfaces is equivalent to n units of D6-branes [26]. Indeed, on the seven-dimensional world-volume of each D6-brane we have a $U(1)$ symmetry. When the n D6-branes approach each other, the gauge symmetry is enhanced from $U(1)^{\otimes n}$ to $U(n)$. An extra compactification of M-theory down to four-dimensional space-time is equivalent to

wrapping D6-branes on the same geometry. In this way, the physics content of the resulting D6-brane scenario can be computed explicitly using the method of the so-called (p, q) webs [11, 12, 13, 14, 15]. Using the results of this method, we expect that the final gauge theory in four-dimensional Minkowski space has gauge group

$$G = \bigotimes_{i=1}^m U(n_i). \quad (24)$$

m is the number of intersecting three-cycles in the base geometry. The integers n_i are specified by the anomaly cancellation condition. This means that they should form a null vector of the intersection matrix of the three-cycles:

$$\sum_{j=1}^m I_{ij} n_j = 0. \quad (25)$$

In the infra-red limit the $U(1)$ factors decouple and one is left with the gauge group

$$G = \bigotimes_{i=1}^m SU(n_i). \quad (26)$$

As already mentioned, the gauge group and matter content depend on the intersecting geometry in the three-dimensional base of the G_2 manifold. The intersection matrix of real N -dimensional spheres S^N (constructed as blow-ups of singularities in Calabi-Yau spaces) is symmetric for N even and anti-symmetric for N odd. In more general geometries, the intersection matrix may be written as a linear combination of a symmetric and an anti-symmetric term.

Here we identify the base with a collection of the three-cycles discussed in Section 3. We recall that they correspond to S^2 fibrations over a line segment. Thus, the intersection matrix of the three-cycles is equivalent to the intersection matrix formed by two-spheres only. From the obvious similarity with the $N = 2$ scenario discussed in Section 2, we see that information on our intersecting geometry is naturally encoded in the Cartan matrix, K , of an affine ADE Lie algebra:

$$I_{ij} = -K_{ij}. \quad (27)$$

It follows that the anomaly cancellation condition is translated into a condition on the affine Lie algebra, thus solving (25) by

$$n_i = s_i n. \quad (28)$$

Here s_i , $i = 1, \dots, r + 1$, are the Coxeter labels of the associated affine Lie algebra of rank r , and $r + 1$ is equal to the number of intersecting three-cycles. As in (2), n is an unspecified

positive integer. The gauge group finally reads

$$G = \bigotimes_{i=1}^{r+1} SU(s_i n). \quad (29)$$

5 Conclusion

We have studied the geometric engineering of $N = 1$ four-dimensional models from M-theory on a manifold with G_2 holonomy. This manifold is a K3 fibration over a three-dimensional base space with ADE geometry. To that end we have first constructed the base geometry as intersecting two-cycles over a line segment. This special base has been embedded in a family of K3 surfaces over the complex plane. Based on the relation between M-theory and D6-branes in type IIA superstrings, we have discussed the physics content of M-theory compactified on our G_2 manifold. In particular, the resulting gauge theory has been discussed in the framework of (p, q) webs. In this scenario, the anomaly cancellation condition was seen to translate into a condition on the associated affine Lie algebra.

Acknowledgments. AB thanks Department of Mathematics and Statistics, Concordia University, Montréal, for its kind hospitality during the preparation of this work. He is very grateful to J. McKay for the invitation, discussions and encouragement, and thanks E.H. Saidi for previous collaborations related to this work. JR thanks F. Lesage for helpful discussions.

References

- [1] S. Kachru, C. Vafa, *Exact results for $N = 2$ compactifications of heterotic strings*, Nucl. Phys. **B450** (1995) 69, hep-th/9505105.
- [2] A. Klemm, W. Lerche, P. Mayr, C. Vafa, N. Warner, *Self-dual strings and $N = 2$ supersymmetric field theory*, Nucl. Phys. **B477** (1996) 746, hep-th/9604034.
- [3] S. Katz, C. Vafa, *Matter from geometry*, Nucl. Phys. **B497** (1997) 146, hep-th/9606086.
- [4] S. Katz, A. Klemm, C. Vafa, *Geometric engineering of quantum field theories*, Nucl. Phys. **B497** (1997) 173, hep-th/9609239.
- [5] M. Bershadsky, A. Johansen, T. Pantev, V. Sadov, C. Vafa, *F-theory, geometric engineering and $N = 1$ dualities*, Nucl. Phys. **B505** (1997) 153, hep-th/9612052.

- [6] S. Katz, P. Mayr, C. Vafa, *Mirror symmetry and exact solution of 4d $N = 2$ gauge theories I*, Adv. Theor. Math. Phys. **1** (1998) 53, hep-th/9706110.
- [7] P. Mayr, *Geometric construction of $N = 2$ gauge theories*, Fortsch. Phys. **47** (1999) 39, hep-th/9807096.
- [8] A. Belhaj, A.E. Fallah, E.H. Saidi, *On the non-simply mirror geometries in type II strings*, Class. Quant. Grav. **17** (2000) 515.
- [9] M. AitBenhaddou, A. Belhaj, E.H. Saidi, to appear.
- [10] M. Cvetič, G. Shiu, A.M. Uranga, *Three-family supersymmetric standard-like models from intersecting brane worlds*, Phys. Rev. Lett. **87** (2001) 201801, hep-th/0107143; *Chiral four-dimensional $N = 1$ supersymmetric type IIA orientifolds from intersecting D6-branes*, Nucl. Phys. **B615** (2001) 3, hep-th/0107166; *Chiral type II orientifold constructions as M theory on G_2 holonomy spaces*, hep-th/0111179.
- [11] A. Hanany, A. Iqbal, *Quiver theories from D6-branes via mirror symmetry*, JHEP **0204** (2002) 009, hep-th/010813.
- [12] B. Feng, A. Hanany, Y.-H. He, A. Iqbal, *Quiver theories, soliton spectra and Picard-Lefschetz transformations*, hep-th/0206152.
- [13] S. Franco, A. Hanany, *Geometric dualities in 4d field theories and their 5d interpretation*, hep-th/0207006; *Toric duality, Seiberg duality and Picard-Lefschetz transformations*, hep-th/0212299.
- [14] A.M. Uranga, *Chiral four-dimensional string compactifications with intersecting D-branes*, hep-th/0301032.
- [15] D. Cremades, L.E. Ibanez, F. Marchesano, *Yukawa couplings in intersecting D-brane models*, hep-th/0302105.
- [16] Y.-H. He, *G_2 quivers*, JHEP **0302** (2003) 023, hep-th/0210127.
- [17] A. Belhaj, *Comments on M-theory on G_2 manifolds and (p,q) webs*, hep-th/0303198.
- [18] A. Belhaj, E.H. Saidi, *Toric geometry, enhanced non simply laced gauge symmetries in superstrings and F-theory compactifications*, hep-th/0012131.

- [19] M. Bershadsky, T. Pantev, V. Sadov, *F-theory with quantized fluxes*, Adv. Theor. Math. Phys. **3** (1999) 727, hep-th/9805056.
- [20] P. Berglund, A. Klemm, P. Mayr, S. Theisen, *On type IIB vacua with varying coupling constant*, Nucl. Phys. **B558** (1999) 178, hep-th/9805189.
- [21] C. Vafa, *On $N = 1$ Yang-Mills in four dimensions*, Adv. Theor. Math. Phys. **2** (1998) 497, hep-th/9801139.
- [22] B.S. Acharya, *M theory, Joyce orbifolds and super Yang-Mills*, Adv. Theor. Math. Phys. **3** (1999) 227, hep-th/9812205.
- [23] T. Friedmann, E. Witten, *Unification scale, proton decay, and manifolds of G_2 holonomy*, hep-th/0211269.
- [24] S. Gukov, C. Vafa, E. Witten, *CFT's from Calabi-Yau four-folds*, Nucl. Phys. **B584** (2000) 69, Erratum-ibid. **B608** (2001) 477, hep-th/9906070.
- [25] C. Vafa, *Evidence for F-theory*, Nucl. Phys. **B469** (1996) 403, hep-th/9602022.
- [26] N.C. Leung, C. Vafa, *Branes and toric geometry*, Adv. Theor. Math. Phys. **2** (1998) 91, hep-th/9711013.