

p-*sd* shell gap reduction in neutron-rich systems and cross-shell excitations in ^{20}O

M. Wiedeking, S.L. Tabor, J. Pavan, A. Volya, A.L. Aguilar, I.J. Calderin, D.B. Campbell, W.T. Cluff, E. Diffenderfer, J. Fridmann, C.R. Hoffman, K.W. Kemper, S. Lee, M.A. Riley, B.T. Roeder, C. Teal, V. Tripathi, and I. Wiedenhöver
Florida State University, Tallahassee, FL 32306, USA
(Dated: November 20, 2018)

Excited states in ^{20}O were populated in the reaction $^{10}\text{Be}(^{14}\text{C},\alpha)$ at Florida State University. Charged particles were detected with a particle telescope consisting of 4 annularly segmented Si surface barrier detectors and γ radiation was detected with the FSU γ detector array. Five new states were observed below 6 MeV from the α - γ and α - γ - γ coincidence data. Shell model calculations suggest that most of the newly observed states are core-excited 1p-1h excitations across the $N = Z = 8$ shell gap. Comparisons between experimental data and calculations for the neutron-rich O and F isotopes imply a steady reduction of the *p*-*sd* shell gap as neutrons are added.

PACS numbers: 23.20.Lv, 25.60.-t, 21.60.Cs, 27.30.+t

The mean field is a central concept in all macroscopic many-body phenomena; in nuclei it leads to shell structure and magic numbers and it provides the basis for the most powerful theoretical techniques such as the nuclear shell model. In recent years attention has been focused on light nuclei, which are intermediate between macroscopic and few-body nucleonic systems, because they can exhibit extreme modifications or even complete breakdown of the mean field approach. This behavior emphasizes the underlying question of nature of how combinations of a few simple building blocks can lead to the astonishing diversity and complexity of many-body phenomena. Interest in this subject is fueled by experimental advances which allow for the exploration of nuclear structure at the limit of stability with critical proton to neutron ratios.

The most significant manifestation of shell breaking is seen in nearly semi-magic nuclei, where the mean-field shell structure breaks down as a function of the proton to neutron ratio. “Islands of inversion” [1, 2] have been observed around the neutron shell closures at $N = 20$ [3] and $N = 28$ [4], where the mean-field shell structure is weakened due to the proton-neutron imbalance and residual interactions dominate [5]. This leads to structures where intruder configurations dominate low-lying states [6].

The oxygen isotopes, which lie at the $Z = 8$ shell boundary, provide an excellent opportunity to investigate the effects of neutron excess on the shell gap and cross shell excitations. Possible modifications of the shell structure with neutron excess raise important questions related to the surprisingly large shift in the neutron drip line between the O and F isotopes [7, 8]. Cross shell *p*-*sd* excitations are known to be essential in the structure of nuclei in this region. Examples include the 110 keV 1^- state in ^{19}F [9] and a recent experimental investigation unveiling low-lying intruder states in ^{22}F [10].

The availability of suitable reactions has limited the study of the neutron-rich O isotopes. Most of the previous work on ^{20}O has involved one of the earliest radioac-

tive beams through the $^{18}\text{O}(t,p)$ reaction [11, 12, 13]. Fragmentation reactions were recently used to determine γ decays from some previously known states [14]. Intruder states, which have an odd number of nucleons promoted across the *p*-*sd* shell gap, provide the clearest evidence for cross shell excitations. Only 2 intruder states, a 1^- at 5.35(10) MeV [15] and a tentative (3^-) at 5.614(3) MeV [16, 17], had been identified in ^{20}O prior to the present work. However, shell model calculations including *p* shell excitations predict significantly more states in the region of 4.5 to 6 MeV. Due to this discrepancy the present experiment was performed to study the shell structure of ^{20}O .

In this letter, we report on the addition of five new states in ^{20}O below an excitation energy of 6 MeV. With the additional experimental information, shell model calculations were performed for neutron rich O and F isotopes. These provide strong evidence for a reduction of the *p*-*sd* shell gap as neutrons are added.

To populate ^{20}O , a long-lived radioactive beam and target were employed through the $^{10}\text{Be}(^{14}\text{C},\alpha)$ reaction at 21.4 MeV. The ^{14}C beam was provided by the Florida State University (FSU) Superconducting Accelerator Laboratory. The $113\ \mu\text{g}/\text{cm}^2$ ^{10}Be target on a $1\ \text{mg}/\text{cm}^2$ thick Pt backing was specially made [18]. Neutron evaporation dominated the reaction, and ^{20}O could only be observed by detecting the α particles. These were detected in an E- Δ E Si telescope, consisting of a $150\ \mu\text{m}$ Δ E detector and a stack of three $1500\ \mu\text{m}$ E detectors. All four Si wafers were divided into 4 annular segments and placed at 0° relative to the beam. A $32\ \text{mg}/\text{cm}^2$ thick gold foil was placed between the target and detector telescope to stop the beam. γ radiation was detected in coincidence with the α particles using the FSU Compton suppressed γ -detector array. During the experiment 4.3×10^6 α - γ events were recorded.

The data were analyzed using GNUSCOPE [19]. The α spectra from the 5 annular segments were kinematically corrected, and a Doppler correction was applied




FIG. 1: Transitions in coincidence with α particles and the 1674 keV γ ray ($2_1^+ \rightarrow 0_1^+$). The lines in this spectrum marked by their energies correspond to decays in ^{20}O .

to the γ spectra. Although the target contained significant amounts of ^{16}O and ^{12}C , the corresponding reaction products, ^{26}Mg and ^{22}Ne , are well known and their lines were removed from consideration.

A portion of the α - γ - γ spectrum in coincidence with the lowest $2_1^+ \rightarrow 0_1^+$ transition at 1674 keV is shown in Fig. 1. The α - γ - γ coincidences with most of the newly assigned lines are shown in Fig. 2. Clear coincidences with the 1674 keV transition ($2_1^+ \rightarrow 0_1^+$) can be seen in all these spectra. Placement of the γ transitions was verified by the coincident α energies, as described in Refs. [20, 21]. Some α spectra obtained by gating on γ lines are shown in Fig. 3. An extensive search for possible contaminants was performed for all transitions before a firm assignment to ^{20}O was made. The level scheme from this and previous work is shown in Fig. 4. The uncertainties in γ -ray energies range from 1 keV up to 6 keV for the weaker new transitions.

Although the 5 new states at 3895, 4353, 4598, 5115, and 5873 keV have not been previously reported, a careful examination of previous studies of ^{20}O does reveal hints of their existence. Very small peaks are visible in the $^{18}\text{O}(t,p)$ spectrum shown in Fig. 1 of Ref. [16] at 3.9, 4.6, 5.1, and possibly 5.9 MeV. While examination of only one spectrum at one angle cannot rule out contaminants, the weakness of these peaks is consistent with their weak population in the present experiment. Interestingly, a tentative state was reported at 5.83(20) MeV in an early paper [22] and included in an earlier compilation [23]. A state was reported at 6.02 MeV in another measurement [11]. The 5.83 MeV state was dropped from the subsequent compilation [24], possibly because it was not confirmed in Ref. [16]. The state observed here at 5873 keV may correspond to one or both of these previously




FIG. 2: Spectra in coincidence with α particles and newly observed transitions in ^{20}O .

reported states.

While shell model calculations restricted to the p shell or to the sd shell have been very successful in describing the structure of nuclei in these regions, the description of excitations between the p and sd shells has been much more limited. The lack of experimental data and increased dimensionality of such calculations together with the center-of-mass issue [25] resulted in the lack of well tested interactions. To gain an understanding of the current experimental results, p - sd shell model calculations were performed for neutron rich O and F isotopes using and slightly modifying (as discussed below) cross-shell interactions from Refs. [1, 26]. Calculations were also conducted taking excitations into the fp shell into account. However, at excitation energies of interest fp configurations proved to be of no importance.

For the calculations, an m -scheme shell model code developed at FSU [27] was used which allows for a full diagonalization of the p - sd shell simultaneously. However, the interactions based on a perturbative merging of the sd and p shells imply a particle-hole hierarchy. Establishing a trend in reproducing the experimental data is a key step in validating the model calculations. The monopole corrected single-particle energies in the sd shell were adjusted to reproduce single-nucleon excitations in ^{17}O . Thus, the model is in good agreement with USD




FIG. 3: α spectra for two γ -ray pairs originating from states with similar excitation energies. The arrows indicate the cut-off location.




FIG. 4: The level scheme as deduced from current and previous work. The widths of the lines indicate their intensities normalized to the 1674 keV transition. Branching ratios are given where applicable. States and transitions marked with solid circles are new additions. γ transitions with open circles were tentatively reported in Ref. [13] and are confirmed in the present work.

shell model results. The cross shell excitations in ^{16}O are also well reproduced. The perturbative combined p - sd shell model essentially has one parameter, the shell gap. The role of this gap becomes more essential in establishing the relative positions of cross shell excitations as the neutron number N increases above the shell closure at $N = 8$. Considering negative-parity states in $^{18,19,20,22}\text{O}$ and using the original interaction, a significant trend of deviation is found. For example the 1p-1h states in ^{22}O




FIG. 5: Systematic reduction of the p - sd shell gap as neutrons are added to O and F nuclei. Shell model calculations were performed and the shell gap adjusted based on the comparison to available experimental data from the current work and Refs. [10, 28, 29]. The error bars reflect the current experimental knowledge available for the respective nuclei.

are predicted to be more than 1 MeV above the first tentatively observed negative-parity state at 5.8 MeV [28]. The agreement with experimentally observed states can be improved significantly by reducing the p - sd shell gap with increasing in neutron number. The necessity to reduce the shell gap to obtain the best agreement between experimental and theoretical states is also observed in the neutron-rich $^{19,20,22}\text{F}$ isotopes. This suggests a general p - sd shell gap decrease as more neutrons are added, in agreement with theoretical predictions [30]. The shell gap dependence on neutron number for O and F isotopes is shown in Fig. 5. The diminishing of the shell gap emphasizes the approach to a region of intruder dominated structures.

The agreement with experimentally observed states in ^{20}O can be improved significantly by reducing the p - sd shell gap by 850 keV. Fig. 6 compares the calculated and experimentally observed states in ^{18}O and ^{20}O , where the closed shell ^{16}O single-particle energies are used for ^{18}O and a reduced p - sd shell gap of 850 keV is used for ^{20}O . The theoretical states shown in blue are mainly the pure sd ones involving only valence neutrons outside the closed ^{16}O core using the USD interaction. Two states at 3743 and 5470 keV in ^{18}O and at 3486 and 4848 keV in ^{20}O are due to 2p-2h excitations. Previously known experimental positive-parity states are also shown in blue. There is good agreement for the positive-parity states in ^{18}O and ^{20}O . The tendency to overpredict the energy of the first 2^+ state is somewhat greater in ^{20}O . Negative-parity 1p-1h states (red) are predicted to start at about 4.5 MeV in both nuclei. There are good candidates in

