Comm. in Asteroseismology
Contribution to the Proceedings of the Wroclaw HELAS Workshop, 2008

Spectroscopy of the sdB pulsator HS 2201+2610

R. Kruspe[1], S. Schuh[1], R. Silvotti[2] and I. Traulsen[1]

[1]Institut für Astrophysik, Georg-August-Universität Göttingen,
Friedrich-Hund-Platz 1, 37077 Göttingen, Germany
[2]INAF (Istituto Nazionale di AstroFisica), Osservatorio Astronomico di
Capodimonte, via Moiariello 16, 80131 Napoli, Italy

Abstract

We present time resolved echelle spectra of the planet-hosting subdwarf B pulsator HS 2201+2610 and report on our efforts to extract pulsational radial velocity measurements from this data.

Session: Poster
Individual Objects: HS 2201+2610, HS 0702+6043

HS 2201+2610 – a planet-hosting pulsating subdwarf B star

Pulsating subdwarf B stars oscillate in short-period p-modes or long-period g-modes. The sdB star HS 2201+2610 (V 391 Peg) is one of the three known hybrids (Lutz et al. 2008), but it has become famous for different reasons. From its p-modes (Østensen et al. 2001, Silvotti et al. 2002), a secular period change due to the star's evolution has recently been measured over a period of 10 years. Furthermore the O–C diagram has revealed a sinusoidal component which is explained by the presence of a planetary-mass companion (Silvotti et al. 2007). To determine the mass of the companion object, the inclination of the orbit needs to be determined. Assuming alignment of the orbital, rotational, and pulsational axes, it should be possible to derive the stellar inclination from a combination of rotational splitting in the photometric frequency spectrum and the projected rotational velocity $v \sin i$ from measured rotational broadening of spectral lines. The measured overall broadening must be corrected for the broadening contribution due to pulsational radial velocities in order to derive the rotational broadening. We therefore initially concentrate on extracting pulsational radial velocities, which requires pulsation phase resolved data.

HET HRS echelle spectra

During two nights each in May and September 2007, spectra of HS 2201+2610 were taken with the HRS echelle spectrograph at the 9m-class Hobby-Eberly-Telescope. The spectra in September were obtained as two series of 95 and 84 20 s-exposures at a resolution of ≈30 000. The resulting signal-to-noise at 5 000 Å is slightly above 3 for individual spectra. When dividing the main pulsation period (349.5 s) into ten phase bins and summing (on average 15) individual spectra corresponding to one phase bin, we obtain a pulsation phase resolved series of ten spectra, each at a signal-to-noise of ≈ 9.

A similar data set has been obtained for HS 0702+6043 in January 2008.

Preliminary results and outlook

The series of spectra covers the hydrogen Balmer series from Hβ to Hδ. These observed spectral lines can be cross correlated with those of an appropriate model spectrum. Our template has been chosen at the spectroscopic parameters $T_{\rm eff}$=30 000 K and $\log{(g/\rm cm\,s^{-2})}$=5.5, close to those given by Østensen et al. 2001. From the cross correlation results for Hβ we do not reliably detect a sinusoidal variation with the phase, and instead give an upper limit of $16 \rm km\,s^{-1}$ for the pulsational radial velocity amplitude corresponding to the main photometric period. Results for the other spectral lines and for further pulsation periods are work in progress. This preliminary result however already implies that the broadening due to the pulsational radial velocities is small, and it looks like the same is true for the rotational broadening. So while the average of all spectra (including the May 2007 series) will help us to refine the stellar parameters of HS 2201+2610 in a future spectral analysis, it remains a challenge to use them to put constraints on the orbit solution for the planet.

Acknowledgments. The authors thank H. Edelmann for his help in obtaining the HET spectra, and U. Heber for kindly providing grids of model spectra. This work has further benefitted from the advice of S. Dreizler and R. Lutz. R. Kruspe thanks HELAS for financially supporting the presentation of this poster through a conference fee waiver.

References

Lutz R., Schuh S., Silvotti R., et al. 2008, A&A, submitted

Østensen R., Solheim J.-E., Heber U., et al. 2001, A&A, 368, 175

Silvotti R., Janulis R., Schuh S., et al. 2002, A&A, 389, 180

Silvotti R., Schuh S., Janulis R., et al. 2007, Nature, 449, 189