Experimental search for radiative decays of the pentaquark baryon $\Theta^+(1540)$

DIANA Collaboration

V.V. Barmin[a], A.E. Asratyan[a], V.S. Borisov[a], C. Curceanu[b], G.V. Davidenko[a], A.G. Dolgolenko[a,*], C. Guaraldo[b], M.A. Kubantsev[a], I.F. Larin[a], V.A. Matveev[a], V.A. Shebanov[a], N.N. Shishov[a], L.I. Sokolov[a], G.K. Tumanov[a], and V.S. Verebryusov[a]

[a] *Institute of Theoretical and Experimental Physics, Moscow 117259, Russia*

[b] *Laboratori Nazionali di Frascati dell' INFN, C.P. 13-I-00044 Frascati, Italy*

January 14, 2019

Abstract

The data on the reactions $K^+\mathrm{Xe} \to K^0\gamma X$ and $K^+\mathrm{Xe} \to K^+\gamma X$, obtained with the bubble chamber DIANA, have been analyzed for possible radiative decays of the $\Theta^+(1540)$ baryon: $\Theta^+ \to K^0p\gamma$ and $\Theta^+ \to K^+n\gamma$. No signals have been observed, and we derive the upper limits $\Gamma(\Theta^+ \to K^0p\gamma)/\Gamma(\Theta^+ \to K^0p) < 0.032$ and $\Gamma(\Theta^+ \to K^+n\gamma)/\Gamma(\Theta^+ \to K^+n) < 0.041$ which, using our previous measurement of $\Gamma(\Theta^+ \to KN) = (0.39\pm0.10)$ MeV, translate to $\Gamma(\Theta^+ \to K^0p\gamma) < 8$ keV and $\Gamma(\Theta^+ \to K^+n\gamma) < 11$ keV at 90% confidence level. We have also measured the cross sections of K^+-induced reactions involving emission of a neutral pion: $\sigma(K^+n \to K^0p\pi^0) = (68 \pm 18)\ \mu$b and $\sigma(K^+N \to K^+N\pi^0) = (30 \pm 8)\ \mu$b for incident K^+ momentum of 640 MeV.

---

[*] Corresponding author. E-mail address: dolgolenko@itep.ru.

Possible existence of multiquark hadrons, and pentaquark baryons in particular, has been discussed for many years [1]. Fairly definite predictions for the antidecuplet of light pentaquark baryons with spin–parity $1/2^+$ have been formulated by Diakonov, Petrov, and Polyakov in the framework of the chiral quark–soliton model [2]. In particular, they predicted $m \simeq 1530$ MeV and $\Gamma < 15$ MeV for the mass and width of the explicitly exotic baryon with $S = +1$ and $I = 0$, the $\Theta^+(uudd\bar{s})$ that should decay to the nK^+ and pK^0 final states. Narrow peaks near 1540 MeV in the nK^+ and pK^0 mass spectra were initially detected in low-energy photoproduction by LEPS [3] and in the charge-exchange reaction $K^+n \to pK^0$ by DIANA [4]. Other searches for the Θ^+ baryon in different reactions and experimental conditions yielded positive evidence as well as null results casting doubt on its existence, see the review papers [5] and [6]. However, both LEPS and DIANA were able to confirm their initial observations [7, 8, 9].

Using the unique properties of the charge-exchange reaction that forms Θ^+ baryons in the s-channel, DIANA reported a direct measurement of the Θ^+ intrinsic width: $\Gamma = 0.39 \pm 0.10$ MeV [8, 9]. The extreme smallness of the Θ^+ width, as compared with well known baryons, indicates a strong dynamical suppression of the "fall-apart" mechanism of Θ^+ decays to pK^0 and nK^+. This does not necessarily imply that radiative decays to γpK^0 and γnK^+ should be equally suppressed, so that the relative probability of radiative decay may prove to be higher for the Θ^+ baryon than for ordinary hadrons. Indeed, partial widths of the radiative decays $\Theta^+ \to K^0p\gamma$ and $\Theta^+ \to K^+n\gamma$ have been computed in [10] as $\Gamma(\Theta^+ \to K^+n\gamma) = 34$–$41$ keV and $\Gamma(\Theta^+ \to K^0p\gamma) = 0.25{\times}\Gamma(\Theta^+ \to K^+n\gamma)$, so that $\Gamma(\Theta^+ \to K^+n\gamma)/\Gamma(\Theta^+ \to K^+n) \simeq 0.15$ if $\Gamma(\Theta^+ \to K^+n) = 0.5{\times}\Gamma(\Theta^+ \to KN)$ and the measured value of $\Gamma(\Theta^+ \to KN)$ [9] is substituted. On the other hand, the Θ^+ radiative widths predicted in [11, 12] using different assumptions are smaller than in [10] by nearly two orders of magnitude[1]. Anyway, the simple arguments brought above and the prediction [10] for $\Gamma(\Theta^+ \to KN\gamma)$ provide adequate motivation for a search for radiative decays $\Theta^+ \to K^0p\gamma$ and $\Theta^+ \to K^+n\gamma$ in the DIANA experiment. The results of the search are reported in this paper.

The non-magnetic bubble chamber DIANA [13] was filled with liquid Xenon and exposed to a separated K^+ beam with incident momentum of 850 MeV at the proton synchrotron at ITEP, Moscow. The fiducial volume of the chamber was $700 \times 700 \times 1400$ mm^3, and the density and radiation length of the fill were 2.2 g/cm^3 and 3.2 cm,

---

[1] An important conclusion reached in [12] was that the parity of the Θ^+ baryon can be determined from the shape of the γ energy spectrum.

respectively. The total number of incident K^+ mesons was $\sim 10^6$, and the data were recorded on $\sim 4 * 10^5$ stereo-frames. In the fiducial volume of the bubble chamber, K^+ momentum varies from $\sim$ 730 MeV for entering kaons to zero for those that range out through ionization. Throughout this momentum interval, all collisions and decays of incident K^+ mesons are efficiently detected. The K^+ momentum at interaction point is determined from the spatial distance between the detected vertex and the center of the observed maximum due to decays of stopping K^+ mesons. Secondary protons and charged pions are identified by ionization, charged kaons — by ionization and decays, K^0_S mesons — by the ranges and emission angles of decay pions, and γ-quanta — by conversion to an e^+e^- pair. Charged particles are momentum-analyzed by range, and γ-quanta — by the summed length of the electron–positron shower. The apparatus efficiency for γ-quanta with $E_\gamma > 30$ MeV is close to 100%. Further details on the experimental procedure may be found in [14, 15, 8].

Radiative decays of the Θ^+ baryon were searched for among the collisions with one or more detected γ-quanta in the final state: $K^+\mathrm{Xe} \to K^+(m\gamma)X$ and $K^+\mathrm{Xe} \to K^0_S(m\gamma)X$, $K^0_S \to \pi^+\pi^-$, where $m \geq 1$. We scanned for and then analyzed the K^+Xe collisions that showed either a secondary K^+ or a Vee from $K^0_S \to \pi^+\pi^-$, plus at least one e^+e^- pair from γ conversion that pointed back to the primary vertex. The collisions involving $K^0_S \to \pi^+\pi^-$ were selected in the full data sample, and those with a secondary K^+ — in nearly 30% of the data. The events were selected in a double scan of the film, and a portion of the film was scanned three times for better understanding of the scanning efficiency. All selected events were then carefully measured and reconstructed in space using specially designed stereo-projectors. In the interval $350 < p(K^+) < 600$ MeV of the incident K^+ momentum, we found 3 events of the reaction $K^+\mathrm{Xe} \to K^0_S\gamma X$, $K^0_S \to \pi^+\pi^-$ and 11 events of the reaction $K^+\mathrm{Xe} \to K^+\gamma X$ with $E_\gamma > 30$ MeV. We also found 20 events of the reaction $K^+\mathrm{Xe} \to K^0_S X$ with $K^0_S \to \pi^+\pi^-\gamma$.

The dominant source of background for the radiative decays $\Theta^+ \to K^0p\gamma$ and $\Theta^+ \to K^+n\gamma$ are the radiative non-resonant transitions $K^+n \to K^0p\gamma$, $K^+n \to K^+n\gamma$, and $K^+p \to K^+p\gamma$. We estimate the probabilities of these transitions relative to corresponding binary reactions $K^+n \to K^0p$, $K^+n \to K^+n$, and $K^+p \to K^+p$ using the formalism developed in [16] (see also [12]). The cross-section ratios $\sigma(K^+n \to K^0p\gamma)/\sigma(K^+n \to K^0p)$, $\sigma(K^+n \to K^+n\gamma)/\sigma(K^+n \to K^+n)$, and $\sigma(K^+p \to K^+p\gamma)/\sigma(K^+p \to K^+p)$ increase with incident momentum $p(K^+)$ by a factor ~ 2 between the boundary values $p(K^+) = 350$ and 600 MeV. For our experimental conditions, the yields of the radiative reactions $K^+n \to K^0p\gamma$, $K^+n \to K^+n\gamma$, and $K^+p \to K^+p\gamma$ relative to $K^+n \to K^0p$, $K^+n \to K^+n$, and $K^+p \to K^+p$ are computed as 3.8×10^{-4}, 5.6×10^{-4}, and 7.6×10^{-4}, respectively.

The experimental determination of the probabilities of radiative processes $K^+\mathrm{Xe} \to K^0_S\gamma X$ and $K^+\mathrm{Xe} \to K^+\gamma X$ relative to the charge-exchange and elastic reactions $K^+\mathrm{Xe} \to K^0_S X$ and $K^+\mathrm{Xe} \to K^+X$ relies on the scanning information for the total numbers of events with detected $K^0_S \to \pi^+\pi^-$ decays or secondary K^+ mesons. In the interval $350 < p(K^+) < 600$ MeV of the incident K^+ momentum, the scan found 12750 ± 750 events of the charge-exchange reaction with $K^0_S \to \pi^+\pi^-$. With three events of the radiative reaction $K^+\mathrm{Xe} \to K^0_S\gamma X$ detected with an efficiency $\epsilon(K^0_S\gamma X) = 0.65 \pm 0.06$, we obtain the relative probability

$$W(K^+\mathrm{Xe} \to K^0_S\gamma X)/W(K^+\mathrm{Xe} \to K^0_S X) = (3.6 \pm 2.1) \times 10^{-4}$$

which agrees with the theoretical prediction. When extracting the total number of events of the elastic reaction $K^+\mathrm{Xe} \to K^+X$ from the scanning information, the events due to coherent scattering $K^+\mathrm{Xe} \to K^+\mathrm{Xe}$ are rejected by the cut $T' < 0.9T$ where T and T' are kinetic energies of the incident and secondary kaons. The number of events of the elastic reaction corrected for the cut $T' < 0.9T$ is estimated as 29500 ± 2100 events for $350 < p(K^+) < 600$ MeV. With 11 events of the corresponding radiative reaction $K^+\mathrm{Xe} \to K^+\gamma X$ detected with an efficiency $\epsilon(K^+\gamma X) = 0.75 \pm 0.05$, we obtain the relative probability

$$W(K^+\mathrm{Xe} \to K^+\gamma X)/W(K^+\mathrm{Xe} \to K^+X) = (5.0 \pm 1.7) \times 10^{-4}.$$

The latter ratio is in fact the mean value of relative probabilities for the proton and neutron, because the numbers of elastic K^+p and K^+n collisions are expected to be virtually equal (the proportion between protons and neutrons in the Xenon nucleus proves to be very close to the inverse ratio between the K^+p and K^+n elastic cross sections [18, 19].) Again, the measured value agrees with the theoretical prediction of 6.6×10^{-4} (the mean value for the proton and neutron, see above). The number of detected charge-exchange events with $K^0_S \to \pi^+\pi^-\gamma$ (20 ± 5) agrees with the results of a simulation[2] that predicts 23 events of this type. We may conclude that the detected $K^+\mathrm{Xe} \to K^0_S\gamma X$

---

[2] The simulation reproduces the experimental conditions and selections, relies on the total number of detected events with $K^0_S \to \pi^+\pi^-$ decays for normalization, and accounts for the branching fraction of the decay $K^0_S \to \pi^+\pi^-\gamma$ [20].

and $K^+\mathrm{Xe} \to K^+\gamma X$ events are consistent with being entirely due to the non-resonant radiative transitions $K^+n \to K^0p\gamma$, $K^+n \to K^+n\gamma$, and $K^+p \to K^+p\gamma$.

In estimating the probabilities of radiative decays $\Theta^+ \to K^0p\gamma$ and $\Theta^+ \to K^+n\gamma$, we rely on our previous observation of 71.7 ± 13.2 decays $\Theta^+ \to K^0p$, $K^0_S \to \pi^+\pi^-$ in the K^+ momentum range of 445–525 MeV [9]. Correcting for the scanning efficiency of 0.83 for the reaction $K^+\mathrm{Xe} \to K^0_S X$ and for unmeasurable events whose fraction reaches 0.40 ± 0.07, we obtain a "reference" signal of

$$N_0(\Theta^+ \to K^0p,\ K^0_S \to \pi^+\pi^-) = (145 \pm 27) \text{ events}$$

for $445 < p(K^+) < 525$ MeV. To derive the "reference" number of $\Theta^+ \to K^+n$ decays, we assume equal branching fractions for $\Theta^+ \to K^0p$ and $\Theta^+ \to K^+n$, correct for the probability of $K^0 \to \pi^+\pi^-$ and for the difference of detection efficiencies for events with secondary K^+ and $K^0_S \to \pi^+\pi^-$, and take into account that the final state $K^+p\gamma$ has been analyzed for only 30% of the data. Thereby, for $445 < p(K^+) < 525$ MeV we obtain

$$N_0(\Theta^+ \to K^+n) = (127 \pm 23) \text{ events.}$$

Of the three detected events of the reaction $K^+\mathrm{Xe} \to K^0_S\gamma X$, only one lies within the interval 445–525 MeV of incident K^+ momentum. The background from the the non-resonant radiative transition $K^+n \to K^0p\gamma$ is estimated as 1.0 events. The final state of the observed event is $K^0_Sp\gamma$ as expected for $\Theta^+ \to K^0p\gamma$, but the effective mass of this three-body system ($m_3 = (1515 \pm 12)$ MeV) is below $m(\Theta^+) = (1538 \pm 2)$ MeV [9] by almost 2σ. Therefore, we disregard this event and conclude that the number of detected $\Theta^+ \to K^0p$ decays is less than 2.3 at 90% confidence level. Dividing by the detection efficiency $\epsilon(K^0_S\gamma X)$ and by the reference signal $N_0(\Theta^+ \to K^0p,\ K^0_S \to \pi^+\pi^-)$ as quoted above, we obtain

$$\Gamma(\Theta^+ \to K^0p\gamma)/\Gamma(\Theta^+ \to K^0p) < 0.032 \text{ at 90\% confidence level.}$$

Of the 11 detected events of the reaction $K^+\mathrm{Xe} \to K^+\gamma X$, three fall within the incident momentum interval of $445 < p(K^+) < 525$ MeV (two events are at $350 < p(K^+) < 445$ MeV and six — at $525 < p(K^+) < 600$ MeV). The background from non-resonant transitions $K^+N \to K^+N\gamma$ is estimated as 4.7 events, so the number of detected $\Theta^+ \to K^+n$ decays is less than 3.0 at 90% confidence level [21]. Dividing by the detection efficiency $\epsilon(K^+\gamma X)$ and by the corresponding reference signal $N_0(\Theta^+ \to K^+n)$, we obtain

$$\Gamma(\Theta^+ \to K^+n\gamma)/\Gamma(\Theta^+ \to K^+n) < 0.041 \text{ at 90\% confidence level.}$$

Substituting $\Gamma(\Theta^+ \to K^0p) = \Gamma(\Theta^+ \to K^+n) = (0.19 \pm 0.05)$ MeV as measured in [9], we are able to restrict the absolute partial widths of Θ^+ radiative decays: $\Gamma(\Theta^+ \to K^0p\gamma) < 8$ keV and $\Gamma(\Theta^+ \to K^+n\gamma) < 11$ keV at 90% confidence level. These measurements are in obvious disagreement with the "optimistic" predictions $\Gamma(\Theta^+ \to K^0 p\gamma) = 8$–$10$ keV and $\Gamma(\Theta^+ \to K^+ n\gamma) = 34$–$41$ keV [10]. On the other hand, the sensitivity of our experiment is insufficient for a substantiative comparison with lower predictions [11, 12].

Apart from the radiative reactions $K^+\mathrm{Xe} \to K^0_S\gamma X$ and $K^+\mathrm{Xe} \to K^+\gamma X$, we have also searched for the inelastic collisions involving formation of neutral pions. In the interval $540 < p(K^+) < 660$ MeV of the incident K^+ momentum, 22 events of the reaction $K^+\mathrm{Xe} \to K^0_S\pi^0 X$, $K^0_S \to \pi^+\pi^-$ and 27 events of the reaction $K^+\mathrm{Xe} \to K^+\pi^0 X$ have been found. The distribution of these inelasic events in incident K^+ momentum steeply increases with $p(K^+)$ from the threshold of $p(K^+) \simeq 520$ MeV to the maximum measured value of $p(K^+) = 660$ MeV. The energy spectrum of detected π^0 mesons agrees with that for the simulated reaction $K^+N \to KN\pi^0$ on a bound nucleon [22]. Selecting the beam-momentum interval $600 < p(K^+) < 660$ MeV with $\langle p(K^+)\rangle \simeq 640$ MeV and taking into account the total numbers of events due to the charge-exchange and elastic reactions $K^+\mathrm{Xe} \to K^0_S X$ and $K^+\mathrm{Xe} \to K^+X$ (not quoted), for the reactions with π^0 emission on the Xenon nucleus we obtain $W(K^+\mathrm{Xe} \to K^0_S\pi^0 X)/W(K^+\mathrm{Xe} \to K^0_S X) = (0.90 \pm 0.23)\%$ and $W(K^+\mathrm{Xe} \to K^+\pi^0 X)/W(K^+\mathrm{Xe} \to K^+X) = (0.31 \pm 0.07)\%$ at $p(K^+) = 640$ MeV. These relative yields have been corrected for the scanning efficiency for the reactions of π^0 emission and for π^0 absorption in the Xenon nucleus [23]. Substituting the measured cross sections of the binary reactions at $p(K^+) = 640$ MeV as $\sigma(K^+n \to K^0p) = (7.5 \pm 0.5)$ mb, $\sigma(K^+n \to K^+n) = (8 \pm 1)$ mb, and $\sigma(K^+p \to K^+p) = (12.5 \pm 1.0)$ mb [17, 18, 19] and taking into account that the Xe nucleus consists of $Z = 54$ protons and $A - Z = 77$ neutrons, we obtain

$$\sigma(K^+n \to K^0p\pi^0) = (68 \pm 18)\ \mu\mathrm{b},$$
$$\sigma(K^+N \to K^+N\pi^0) = (30 \pm 8)\ \mu\mathrm{b},$$

where the latter refers to the cross section averaged over constituent nucleons of the Xe nucleus. The cross sections of these reactions for $p(K^+) \simeq 640$ MeV were previously measured in a single experiment that used a deuterium bubble chamber [24]: $\sigma(K^+n \to K^0p\pi^0) = (80 \pm 20)$ μb, $\sigma(K^+d \to K^+\pi^0pn) = (0 \pm 20)$ μb. The former value is in good agreement with our measurement.

In summary, we have analyzed the data on the reactions $K^+\mathrm{Xe} \to K^0\gamma X$ and $K^+\mathrm{Xe} \to K^+\gamma X$, obtained with the bubble chamber DIANA, for possible radiative decays of the $\Theta^+(1540)$ baryon: $\Theta^+ \to K^0p\gamma$ and $\Theta^+ \to K^+n\gamma$. No signals have been observed, and we derive the upper limits $\Gamma(\Theta^+ \to K^0p\gamma)/\Gamma(\Theta^+ \to K^0p) < 0.032$ and $\Gamma(\Theta^+ \to K^+n\gamma)/\Gamma(\Theta^+ \to K^+n) < 0.041$ which, upon using our previous measurement of $\Gamma(\Theta^+ \to KN)$, translate to $\Gamma(\Theta^+ \to K^0 p\gamma) < 8$ keV and $\Gamma(\Theta^+ \to K^+ n\gamma) < 11$ keV at 90% confidence level. We have also measured the cross sections of K^+-induced reactions involving emission of a neutral pion: $\sigma(K^+ n \to K^0 p\pi^0) = (68 \pm 18)$ μb and $\sigma(K^+ N \to K^+ N\pi^0) = (30 \pm 8)$ μb for incident K^+ momentum of 640 MeV.

It is a pleasure to thank L. N. Bogdanova, B. L. Ioffe, Yu. S. Kalashnikova, A. V. Samsonov, and I. I. Strakovsky for useful discussions and comments. This work is supported by the Russian Foundation for Basic Research (grant 07-02-00684).

References

[1] R.L. Jaffe, "Multi-quark hadrons", Phys. Rev. D15, 267 (1977).

[2] D. Diakonov, V. Petrov, and V. Polyakov, Z. Phys. A359, 305 (1997).

[3] T. Nakano *et al.* (LEPS Coll.), Phys. Rev. Lett. 91, 012002 (2003).

[4] V. V. Barmin *et al.* (DIANA Coll.), Yad. Fiz. 66, 1763 (2003) and Phys. At. Nucl. 66, 1715 (2003).

[5] Volker D. Burkert, Int. J. Mod. Phys. A21, 1764 (2006), arXiv:hep-ph/0510309.

[6] M.V. Danilov and R.V. Mizuk, Phys. Atom. Nucl. 71, 605 (2008), arXiv:0704.3531 [hep-ex].

[7] T. Nakano *et al.* (LEPS Coll.), Phys. Rev. C79, 025210 (2009), arXiv:0812.1035 [nucl-ex].

[8] V.V. Barmin *et al.* (DIANA Coll.), Yad. Fiz. 70, 39 (2007) and Phys. At. Nucl. 70, 35 (2007), arXiv:hep-ex/0603017.

[9] V.V. Barmin *et al.* (DIANA Coll.), arXiv:0909.4183 [hep-ex].

[10] Deog Ki Hong, arXiv:hep-ph/0412132.

[11] Xiao-Gang He *et al.,* Phys. Rev. D71, 014006 (2005).

[12] B.L. Ioffe and A.V. Samsonov, Phys. Lett. B633, 298 (2006).

[13] V.V. Barmin *et al.,* Prib. Tekhn. Eksp. 4, 61 (1984).

[14] V.V. Barmin *et al.,* Nucl. Phys. A556, 409 (1993).

[15] V.V. Barmin *et al.,* Phys. Atom. Nucl. 57, 1655 (1994).

[16] I.Ya. Pomeranchuk and I.M. Shmushkevich, Dokl. Akad. Nauk USSR 64, 499 (1949).

[17] W. Slater *et al.,* Phys. Rev. Lett. 15, 378 (1961).

[18] T. Bowen *et al.,* Phys. Rev. D7, 22 (1973).

[19] U. Casadei *et al.,* CERN–HERA 75-1.

[20] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B667, 1 (2008).

[21] G.J. Feldman and R.D. Cousins, Phys. Rev. D57, 3873 (1998).

[22] B. Sechi Zorn and G.T. Zorn, Phys. Rev. 120, 1898 (1960).

[23] J. Hornbostel and G.T. Zorn, Phys. Rev. 109, 165 (1958).

[24] G. Giacomelli *et al.,* Nucl. Phys. B37, 577 (1972).