A dendrite generated from $\{0,1\}^{\Lambda}$, $Card\Lambda \succ \aleph$

Akihiko Kitada[1], Tomoyuki Yamamoto[1,2,*], Shousuke Ohmori[2]

[1]*Institute of Condensed-Matter Science, Comprehensive Resaerch Organization, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan*
[2]*Faculty of Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan*

Abstract

The existence of a decomposition space with a dendritic structure of a topological space $(\{0,1\}^{\Lambda}, \tau_0^{\Lambda})$ is discussed. Here, Λ is any set with the cardinal number $\succ \aleph$, $\{0,1\}^{\Lambda} = \{\varphi : \Lambda \to \{0,1\}\}, \tau_0$ is the discrete topology for $\{0,1\}$ and the topology τ_0^{Λ} for $\{0,1\}^{\Lambda}$ is the topology with the base $\beta = \{< G_{\lambda_1}, \ldots, G_{\lambda_n} > \ ; \ G_{\lambda_1} \in \tau_0, \ldots, G_{\lambda_n} \in \tau_0, \{\lambda_1, \ldots, \lambda_n\} \subset \Lambda, n \in \mathbf{N}\}$ where the notation $< E_{\lambda_1}, \ldots, E_{\lambda_n} >$ concerning the subset $E_{\lambda_i}, i = 1, \ldots, n$ of $\{0,1\}$ denotes the set $\{\varphi : \Lambda \to \{0,1\} \ ; \ \varphi(\lambda_1) \in E_{\lambda_1}, \ldots, \varphi(\lambda_n) \in E_{\lambda_n}, \varphi(\lambda) \in \{0,1\}, \lambda \in \Lambda - \{\lambda_1, \ldots, \lambda_n\}\}$.

1 Introduction

In the present paper, the existence of a dendrite as a decomposition space of a topological space each of whose elements is a map φ from a set Λ whoes cardinal number is greater than or equal to $\aleph$ to $\{0,1\}$ will be discussed.

Some mathmatical definitions and statements which appear in the text are listed below.

A) [1] Let $(\mathcal{D}, \tau(\mathcal{D}))$ be a decomposition space of a topological space (X, τ). That is, $\mathcal{D}$ is a collection of nonempty, mutually disjoint subsets of X such that $\bigcup \mathcal{D} = \bigcup_{D \in \mathcal{D}} D = X$, and $\tau(\mathcal{D})$ is the decomposition topology given by $\{\mathcal{U} \subset \mathcal{D}; \bigcup \mathcal{U} = \bigcup_{D \in \mathcal{U}} D \in \tau\}$. A decomposition $\mathcal{D}$ of (X, τ) is called usc (upper semi continuous) provided that for any element $D \in \mathcal{D}$ and for any $U \in \tau$ containing D, there exists $u \in \tau$ such that $D \subset u$ and if $A \cap u \neq \phi$ for $A \in \mathcal{D}$, then $A \subset U$.

B) [2] Let $\mathcal{D}$ be a decomposition of a set X. A subset Y of X which is a union of subcollection of $\mathcal{D}$, that is, for some $\mathcal{U} \subset \mathcal{D}, Y = \bigcup \mathcal{U} (= \bigcup_{D \in \mathcal{U}} D)$, is said to be $\mathcal{D}$-saturated.

C) Let $f : (X, \tau) \to (Y, \tau')$ be a continuous, closed, onto map. Then, i) [3] The decomposition space $(\mathcal{D}_f, \tau(\mathcal{D}_f))$ of (X, τ) due to f is usc. Here, the decomposition $\mathcal{D}_f$ of X is given by $\mathcal{D}_f = \{f^{-1}(y) \subset X; y \in Y\}$. ii) [4] For any pair $f^{-1}(y)$ and $K \in \Im$ (K is a closed set of (X, τ)) such that $f^{-1}(y) \cap K = \phi$, there exists a $\mathcal{D}_f$-saturated $V \in \tau$ satisfying the relations $f^{-1}(y) \subset V$ and $V \cap K = \phi$.

D) [5] Let X be a zero-dimensional (abbreviated as 0-dim), perfect, compact T_1-space. Then, for any compact metric space Y, there exists a continuous map f from X onto Y.

E) [6] If $f : (X, \tau) \to (Y, \tau')$ is a quotient map, that is, an onto map f which is characterized by the relation $\tau' = \tau_f (= \{u' \subset Y; f^{-1}(u') \in \tau\})$, then $h : (Y, \tau') \to (\mathcal{D}_f, \tau(\mathcal{D}_f))$,

$y \mapsto f^{-1}(y)$ is a homeomorphism. Namely, the decomposition space $\mathcal{D}_f$ of X is topologically equivalent to Y.
F) [7] A locally connected, connected, compact metric space containing no simple closed curve as its subspace is called a dendrite.
G) [8] Let (X, τ) be compact. If any two distinct points x and y are separated in X, that is, there exists a closed and open (clopen) set G such that $x \in G$ and $y \notin G$, then (X, τ) is 0-dim.

2 A dendrite as a decomposition space of a 0-dim, perfect, compact T_1-space

Let (X, τ) be a 0-dim, perfect, compact T_1-space. Since the dendrite (Y, τ_d) is a compact metric space by its definition 1,F), according to 1,D), there exists a continuous map f from (X, τ) onto (Y, τ_d). Any continuous map from a compact space onto a T_2-space is easily verified to be closed. A continuous, closed, onto map is obviously a quotient map. We are convinced from 1,E) that the decomposition space $(\mathcal{D}_f, \tau(\mathcal{D}_f))$ of (X, τ) due to the map f is homeomorphic to the dendrite (Y, τ_d) under the homeomorphism $h : (Y, \tau_d) \to (\mathcal{D}_f, \tau(\mathcal{D}_f)), y \mapsto f^{-1}(y)$. The decomposition space $(\mathcal{D}_f, \tau(\mathcal{D}_f))$ is metrized by means of the homeomorphism h and the metric d of Y as $\rho(D, D') = d(h^{-1}(D), h^{-1}(D')), D, D' \in \mathcal{D}_f$. That is, the decomposition topology $\tau(\mathcal{D}_f)$ is identical with the metric topology τ_ρ. Since the metric space $(\mathcal{D}_f, \tau_\rho)$ is a locally connected, connected, compact metric space which contains no simple closed curve owing to the homeomorphism h, it must be a dendrite. From 1,C),i), this dendrite $(\mathcal{D}_f, \tau_\rho)$ based on the continuous, closed, onto map f is usc. Therefore, according to 1,C),ii) each element $f^{-1}(y)$ of this dendrite $\mathcal{D}_f$ is topologically characterized by the property that, for any pair $f^{-1}(y)$ and $K \in \Im$ such that $f^{-1}(y) \cap K = \phi$, there exists $\mathcal{D}_f$-saturated $V \in \tau$ such that $f^{-1}(y) \subset V$ and $V \cap K = \phi$.

3 Topological space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$

Let Λ be a set with $Card\Lambda \succ \aleph$ and let τ_0 be the discrete topology for $\{0,1\}$. τ_0^Λ is a topology for $\{0,1\}^\Lambda = \{\varphi : \Lambda \to \{0,1\}\}$ with the base $\beta = \{< G_{\lambda_1}, \ldots, G_{\lambda_n} >; G_{\lambda_1} \in \tau_0, \ldots, G_{\lambda_n} \in \tau_0, \ \{\lambda_1, \ldots, \lambda_n\} \subset \Lambda, n \in \mathbf{N}\}$. Here, the set $\{\varphi : \Lambda \to \{0,1\}; \varphi(\lambda_1) \in E_{\lambda_1}, \ldots, \varphi(\lambda_n) \in E_{\lambda_n}, \varphi(\lambda) \in \{0,1\}, \lambda \in \Lambda - \{\lambda_1, \ldots, \lambda_n\}\}$ concerning the subset $E_{\lambda_i}, i = 1, \ldots, n$ of $\{0,1\}$, is denoted by $< E_{\lambda_1}, \ldots, E_{\lambda_n} >$.
In the followings, we will show that the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ is 0-dim, perfect, compact T_1-space. First, we will check $\dim\{0,1\}^\Lambda = 0$. Since $(\{0,1\}, \tau_0)$ is a compact space, the product space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ is also compact due to Tychonoff. If φ_p and φ_q are distinct two points of $\{0,1\}^\Lambda$, there exists an index $\lambda_0 \in \Lambda$ such that $\varphi_p(\lambda_0) \neq \varphi_q(\lambda_0)$. Without loss of generality, we can put $\varphi_p(\lambda_0) = 0$ and $\varphi_q(\lambda_0) = 1$. The open set $< G_{\lambda_0} = \{0\} >= \{\varphi : \Lambda \to \{0,1\}; \varphi(\lambda_0) \in \{0\}, \varphi(\lambda) \in \{0,1\}, \lambda \in \Lambda - \{\lambda_0\}\}$ of $\{0,1\}^\Lambda$ clearly contains the point φ_p and does not contain the point φ_q, that is, $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ is a T_1-space. According to 1,G), it is sufficient for the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ to be 0-dim, that the

open set $< G_{\lambda_0} = \{0\} >$ is, at the same time, a closed set. Any point $\varphi \in < G_{\lambda_0} = \{0\} >$ is not contained in the open set $< G'_{\lambda_0} = \{1\} >$ and any point $\varphi \notin < G'_{\lambda_0} = \{1\} >$ must be contained in the set $< G_{\lambda_0} = \{0\} >$. Therefore, the set $< G_{\lambda_0} = \{0\} >$ is equal to the complement of the open set $< G'_{\lambda_0} = \{1\} >$. This means that $< G_{\lambda_0} = \{0\} >$ is a colsed set. Next, to prove that the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ is perfect, let us consider $H \in \tau_0^\Lambda - \{\phi\}$. For a point $\varphi_a \in H$, there exists a base $< G_{\lambda_1}, \ldots, G_{\lambda_n} >$ containing φ_a such that $< G_{\lambda_1}, \ldots, G_{\lambda_n} > \subset H$. Since the set Λ is not a finite set, there exists $\bar{\lambda} \in \Lambda - \{\lambda_1, \ldots, \lambda_n\}$ such that $\varphi(\bar{\lambda}) \in \{0,1\}$. Let φ'_a be a point of $\{0,1\}^\Lambda$ defined as $\varphi_a(\lambda) = \varphi'_a(\lambda)$ for $\lambda \neq \bar{\lambda}$ and $\varphi_a(\bar{\lambda}) \neq \varphi'_a(\bar{\lambda})$. It is clear that φ'_a is contained in $< G_{\lambda_1}, \ldots, G_{\lambda_n} >$ as well as φ_a. Therefore, the nonempty open set H contains at least two points. This means that the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ is perfect.

4 Conclusion

Since the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ is confirmed to be a 0-dim, perfect, compact T_1-space, the discussions in the section 2 can be applied to this space. As a consequence, there exists a decomposition space $\mathcal{D}$ of $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ which has a dendrite structure. Although the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ where $Card\Lambda \succ \aleph$ is not metrizable[9], its decomposition space $\mathcal{D}$ is metrizable as a dendite. Finally, we note that there exists a decomposition space of the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ one of whose decomposition space has the dendritic structure again (See Appendix).

Appendix

Using the following lemma, we will confirm the existence of a decomposition space $\mathcal{D}_f$ of $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ which has a decomposition space with the dendritic structure again.

[Lemma] Let (X, τ) be a 0-dim, perfect, compact T_0-space. Then, there exists a non empty 0-dim, perfect, compact T_0-proper subspace (Y, τ_Y) of (X, τ).

Proof. Since X is a perfect T_0-space, there exist at least two distinct points x and x' and there exists an open set U such that, without loss of generality, $x \in U$ and $x' \notin U$. Since X is 0-dim, there exists a set $u \in \tau \cap \Im$ which contains x and is contained in U. Taking this clopen set u as Y, we obtain a 0-dim, perfect, compact proper subset (Y, τ_Y) of (X, τ). □

Now for the above space (X, τ) and its subspace (Y, τ_Y), let us define a map $f : (X, \tau) \to (Y, \tau_Y)$ as

$$f(x) = \begin{cases} x \ , & x \in Y \\ q \ , & x \in X - Y \end{cases}$$

Here, q is an arbitrarily chosen point of Y. Since 0-dim T_0-space is necessarily a T_2-space, the singleton $\{q\}$ is a closed set. Thus, it is easy to see that the map f is a continuous, closed map from X onto Y[10], that is, the map f is a quotient map. Let us note that the decomposition space $\mathcal{D}_f$ due to this *not one to one* map f is not the trivial decomposition

$\{\{x\}; x \in X\}$. From 1,E), $(\mathcal{D}_f, \tau(\mathcal{D}_f))$ is homeomorphic to 0-dim, perfect, compact T_0-space (Y, τ_Y), and then, $(\mathcal{D}_f, \tau(\mathcal{D}_f))$ must be a 0-dim, perfect, compact T_0-space. Since any 0-dim, T_0-space is a T_2-space, all of the discussions in the text concerning the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ can be repeated for the decomposition space $(\mathcal{D}_f, \tau(\mathcal{D}_f))$ of $(\{0,1\}^\Lambda, \tau_0^\Lambda)$.

Acknowledgement

The authors are grateful to Dr. H. Ryo, Dr. Y. Yamashita, Dr. K. Yoshida, Dr. Y. Watayoh and Dr. K. Shikama for useful discussions and encouragements.

References and notes

[1] S.B.Nadler Jr., *Continuum theory*, Marcel Dekker 1992, p.36, p38.
[2] Ref.[1], p.38, p.39.
[3] proof of i) It is sufficient for $\mathcal{D}_f$ to be usc that the natural map $\pi : X \to \mathcal{D}_f, x \mapsto D$ (there exists unique $D \in \mathcal{D}$ such that $x \in D$) is closed (See, for example Ref.[1],p.39). Since f is continuous and $f(K)$ is closed for $K \in \Im$, the set $\bigcup\limits_{x \in K} \pi(x)$ must be closed from the relation $\bigcup\limits_{x \in K} \pi(x) = f^{-1}(f(K))$ which can be easily verified. This means that π is a closed map. In fact, $\{\pi(x); x \in A\}$ is closed in $(\mathcal{D}_f, \tau(\mathcal{D}_f)) \Leftrightarrow \bigcup\limits_{x \in A} \pi(x) \in \Im$.
[4] Ref.[1], p.39.
[5] Ref.[1], p.105, p.106 and p.109 in which the space X is supposed to be a metric space. But the 0-dim, perfect, compact space X is not necessarily a metric space. It is sufficient for the space X to have the property that any compact metric space is a continuous image of it that X is a T_1-space.
[6] Ref.[1], p.46.
[7] a) Ref.[1], p.165, b) A.Kitada, Y.Ogasawara, T.Yamamoto, Chaos, Solitons & Fractals **34**, 1732 (2007).
[8] proof of G) Let U be an open set containing x. The complement U^c in X of U is a compact set. For each point $y \in U^c$, there exists a clopen set $u(y)$ containing y such that $x \notin u(y)$. We can choose a finite sub cover $\{u(y_1), \ldots, u(y_n)\}$ of U^c from the open cover $\{u(y); y \in U^c\}$. The clopen set $(\bigcup\limits_{i \in \bar{n}} u(y_i))^c$ containing x is contained in U. (X, τ) is, therefore, 0-dim.
[9] R.Engelking, *General Topology*, Heldermann Verlag Berlin 1989, p.260. If $Card\Lambda \prec \aleph_0$, the space $(\{0,1\}^\Lambda, \tau_0^\Lambda)$ is well-known to be metrizable. In the present study, it is emphasized that the metrizability of the space is not necessary for its decomposition space to be a metric space "dendrite".
[10] proof) At the point $x \in Y, f(x) = x$. For any $U \in \tau_Y$ containing $f(x)$, there exists U containing x such that $f(U) = U \subset U$. Since Y is an open set, $U \in \tau$. At the point $x \in Y^c$, for any $U \in \tau_Y$ containing $f(x) = q$, there exists $Y^c \in \tau$ containing x such that $f(Y^c) = \{q\} \subset U$. Thus the map f is continuous at any point of X. Next, let K

be a closed set of X. $f(K) = f(K \cap X) = f(K \cap (Y \cup Y^c)) = f((K \cap Y) \cup (K \cap Y^c))$
$= f(K \cap Y) \cup f(K \cap Y^c) = (K \cap Y) \cup \{q\} \in \Im_Y$. Therefore, f is a closed map.