Mass of the bottom quark from Upsilon(1S) at NNNLO: an update[1]

César Ayala[1], **Gorazd Cvetič**[2], **Antonio Pineda**[3]

[1]Department of Theoretical Physics, IFIC, University of Valencia - CSIC, Valencia, Spain
[2]Physics Department, Universidad Técnica Federico Santa María, Valparaíso, Chile
[3]Physics Department and IFAE, Universitat Autònoma de Barcelona, Barcelona, Spain

Abstract. We update our perturbative determination of $\overline{\rm MS}$ mass $\overline{m}_b(\overline{m}_b)$, by including the recently obtained four-loop coefficient in the relation between the pole and $\overline{\rm MS}$ mass. First the renormalon subtracted (RS or RS') mass is determined from the known mass of the $\Upsilon(1S)$ meson, where we use the renormalon residue N_m obtained from the asymptotic behavior of the coefficient of the 3-loop static singlet potential. $\overline{\rm MS}$ mass is then obtained using the 4-loop renormalon-free relation between the RS (RS') and $\overline{\rm MS}$ mass. We argue that the effects of the charm quark mass are accounted for by effectively using $N_f = 3$ in the mass relations. The extracted value is $\overline{m}_b(\overline{m}_b) = 4222(40)$ MeV, where the uncertainty is dominated by the renormalization scale dependence.

1. Introduction

The ($\overline{\rm MS}$) mass of the bottom (b) quark, $\overline{m}_b \equiv \overline{m}_b(\overline{m}_b)$, is an important quantity in particle physics, free of renormalon ambiguities, and appears in many physical observables. Since it is relatively high, ~ 4 GeV, perturbative QCD methods are suitable for its extraction. The mass of the ground state of the $b\bar{b}$ quarkonium, $\Upsilon(1S)$, is one of the best quantities for such an extraction, $M^{\rm (th)}_{\Upsilon(1S)} = 2m_b + E_{\Upsilon(1S)} = 9.460$ GeV, where m_b is the pole mass of the bottom quark, and $E_{\Upsilon(1S)}$ is the binding energy. We use the available perturbative expansions of $2m_b/\overline{m}_b$ and of $E_{\Upsilon(1S)}/\overline{m}_b$ in powers of QCD coupling $a(\mu) \equiv \alpha_s(\mu)/\pi$ and thus extract the value of $\overline{m}_b$. In the extraction, we use the fact that the leading infrared (IR) renormalon ambiguity of $2m_b$ cancels out with that of $E_{\Upsilon(1S)}$ [1, 2, 3].

These proceedings are a brief review of our previous work [4], which we update by including in the analysis the recently calculated [5] four-loop coefficient of the relation between the pole mass and the $\overline{\rm MS}$ mass. Here we outline: (1) The correct treatment of charm quark mass effects in the perturbation expansion of $m_b/\overline{m}_b$; (2) Asymptotic expressions for the coefficients in the perturbation expansion of the ratio $m_b/\overline{m}_b$ and of the static singlet potential $V(r)$, and the extraction of the renormalon residue N_m; (3) The construction of the (modified) renormalon-subtracted mass $m_{b,{\rm RS}^{(\prime)}}$ (using N_m), and the renormalon-free relation between $m_{b,{\rm RS}^{(\prime)}}$ and $\overline{m}_b$; (4) Renormalon-free perturbation expansion for $M^{\rm (th)}_{\Upsilon(1S)}$ in terms of $m_{b,{\rm RS}^{(\prime)}}$, and extraction, from $M^{\rm (th)}_{\Upsilon(1S)} = 9.460$ GeV, of the values of $m_{b,{\rm RS}^{(\prime)}}$ ($\Rightarrow \overline{m}_b$).

---

[1] preprint USM-TH-343; presented at ACAT 2016, 18-22 Jan. 2016, UTFSM, Valparaíso, Chile

2. Charm mass effects in the bottom pole mass

The pole mass m_b and the $\overline{\rm MS}$ mass $\overline{m}_b$ are related:

$$m_b = \overline{m}_b \left(1 + S(N_f)\right) + \delta m_c^{(+)} \ , \tag{1}$$

$$\text{where} \qquad S(N_f) = \frac{4}{3} a_+(\mu) \left[1 + r_1^{(+)}(\mu) a_+(\mu) + r_2^{(+)}(\mu) a_+^2(\mu) + r_3^{(+)}(\mu) a_+^3(\mu) + \mathcal{O}(a_+^4)\right] \tag{2}$$

and the evaluation is usually performed in QCD with $N_f = N_l + 1 = 4$ active flavors: $r_j^{(+)}(\mu) \equiv r_j(\mu; N_f)$, $a_+(\mu) = a(\mu; N_f)$. The coefficients $R_0 = 4/3$ and r_j $(j = 1, 2)$ were obtained in Refs. [6], [7], [8, 9], respectively. Recently, numerical values of the 4-loop coefficient r_3 were obtained [5], and we incorporate them here in the form given in [10].

These coefficients have a specific dependence on the renormalization scale μ, dictated by μ-independence of $S(N_f)$

$$r_1(\mu; N_f) = r_1(N_f) + \beta_0 L_m(\mu) \ , \text{ etc.} \tag{3}$$

where $L_m(\mu) = \ln(\mu^2/\overline{m}_b^2)$, and we maintain, for simplicity, the notation $r_j \equiv r_j(\overline{m}_b)$. We will use the notations $\beta_0 = (1/4)(11 - 2N_f/3)$ and $\beta_1 = c_1\beta_0 = (102 - 38N_f/3)/16$ for the first two coefficients of the RGE of $a(\mu)$

$$\frac{da(Q)}{d \ln Q^2} = -\beta_0 a^2(Q) \left(1 + c_1 a(Q) + c_2 a^2(Q) + c_3 a^3(Q) + \cdots \right) \ . \tag{4}$$

Finite-mass charm quark effects are incorporated in

$$\delta m_c^{(+)} = \delta m_{(c,+)}^{(1)} a_+^2(\overline{m}_b) + \delta m_{(c,+)}^{(2)} a_+^3(\overline{m}_b) + \mathcal{O}(a_+^4) \, , \tag{5}$$

which vanishes in the $m_c \to 0$ limit. We have

$$\delta m_{(c,+)}^{(1)} = \frac{4}{3} \overline{m}_b \Delta[\overline{m}_c/\overline{m}_b] = 1.9058 \text{ MeV [7]}, \quad \delta m_{(c,+)}^{(2)} = 48.6793 \text{ MeV [11]}, \tag{6}$$

$$\Rightarrow \quad \delta m_{(c,+)}^{(1)} a_+^2(\overline{m}_b) = 9.3 \text{ MeV}, \qquad \delta m_{(c,+)}^{(2)} a_+^3(\overline{m}_b) = 18.1 \text{ MeV}, \tag{7}$$

so $\delta m_c^{(+)}$ is badly divergent. Why? At loop order n, the natural scale of the loop integral for m_b is $m_b e^{-n}$ [12], which for n large enough is: $m_b e^{-n} < m_c$. Therefore, for large n (> 2) charm quark appears as very heavy (decoupled), leading to the effective number of flavors being $N_l = 3$ and not $N_f = N_l + 1 = 4$. Therefore, it is convenient to rewrite the relation between the pole and the $\overline{\rm MS}$ mass in terms of $a_-(\mu) = a(\mu; N_l)$ and $r_j^{(-)}(\mu) \equiv r_j(\mu; N_l)$ $[N_l = 3]$

$$m_b = \overline{m}_b \left(1 + S(N_l)\right) + \delta m_c \ , \tag{8}$$

$$\text{where} \quad S(N_l) \ = \ \frac{4}{3} a_-(\mu) \left[1 + r_1^{(-)}(\mu) a_-(\mu) + r_2^{(-)}(\mu) a_-^2(\mu) + r_3^{(-)}(\mu) a_-^3(\mu) + \mathcal{O}(a_-^4)\right], \tag{9}$$

and $r_j^{(-)}(\overline{m}_b) = 7.74,\ 87.2,\ 1265.3 \pm 16.1$, for $j = 1, 2, 3$. The effects of the decoupling of S $(N_f \mapsto N_l = 3)$ are absorbed in the new δm_c

$$\delta m_c \ = \ \left[\delta m_{(c,+)}^{(1)} + \delta m_{(c,\text{dec.})}^{(1)}\right] a_-^2(\overline{m}_b) + \left[\delta m_{(c,+)}^{(2)} + \delta m_{(c,dec.)}^{(2)}\right] a_-^3(\overline{m}_b) + \mathcal{O}(a_-^4) \ , \tag{10}$$

where $\delta m^{(j)}_{(c,\text{dec.})}$ are generated by this decoupling and read

$$\delta m^{(1)}_{(c,\text{dec.})} = \frac{2}{9}\overline{m}_b \left(\ln \left(\frac{\overline{m}_b^2}{\overline{m}_c^2} \right) - \frac{71}{32} - \frac{\pi^2}{4} \right) \tag{11}$$

and $\delta m^{(2)}_{(c,\text{dec.})}$ can be found in Ref. [4].

Numerical evaluation gives for $\left[\delta m^{(1)}_{(c,+)} + \delta m^{(1)}_{(c,\text{dec.})}\right] a_-^2(\overline{m}_b) = -1.6$ MeV and $\left[\delta m^{(2)}_{(c,+)} + \delta m^{(2)}_{(c,\text{dec.})}\right] a_-^3(\overline{m}_b) = -0.3$ MeV. This means that the previous divergent series (in $\text{QCD}_{N_f=4}$) $\delta m_c^{(+)} = (9.3 + 18.1 + \ldots)$ MeV [Eq. (7)] now tranforms (in $\text{QCD}_{N_l=3}$) to

$$\delta m_c = (-1.6 - 0.3 + \ldots) \text{ MeV}. \tag{12}$$

The series for δm_c in $\text{QCD}_{N_l=3}$ formulation is convergent, strong cancellation takes place between $\delta m^{(j)}_{(c,+)}$ and $\delta m^{(j)}_{(c,\text{dec.})}$, as expected.

3. Leading renormalon of the pole mass

The asymptotic behaviour of r_N is determined by the leading IR renormalon:

$$\frac{4}{3} r_N^{\text{asym}}(\mu) \simeq \pi N_m \frac{\mu}{\overline{m}_b} (2\beta_0)^N \frac{\Gamma(\nu+N+1)}{\Gamma(\nu+1)} \left[1 + \sum_{s=1}^{3} \frac{\nu \cdots (\nu-s+1)}{(N+\nu)\cdots(N+\nu-s+1)} \widetilde{c}_s + \mathcal{O}\left(N^{-4}\right) \right]. \tag{13}$$

$$\frac{4}{3} r_N(\mu) = \pi N_m \frac{\mu}{\overline{m}_b} (2\beta_0)^N \frac{\Gamma(\nu+N+1)}{\Gamma(\nu+1)} \left[1 + \sum_{s\geq 0} \frac{\nu \cdots (\nu-s+1)}{(N+\nu)\cdots(N+\nu-s+1)} \widetilde{c}_s \right] + h_N(\mu), \tag{14}$$

where h_N is dominated by subleading renormalons, and the coefficients $\widetilde{c}_s$ $(s = 1, 2, 3)$ are given in [13, 14, 15, 4] ($\widetilde{c}_0 = 1$ by convention).

Determining the pole mass from $\Upsilon(1S)$ mass has large uncertainties due to the pole mass renormalon ambiguity $\delta m_b \sim \Lambda_{\text{QCD}}$ [13]. In order to avoid this problem, we work with the renormalon-subtracted (RS) bottom mass $m_{b,\text{RS}}$ instead [14]. Then, $\overline{m}_b$ is obtained from its stable (renormalon-free) relation with the m_{RS} mass.

The use of m_{RS} in the theoretical evaluation of the $\Upsilon(1S)$ mass is convenient because it has no leading IR renormalon ambiguity, and the renormalon cancellation in the quarkonium mass $M_{\Upsilon(1S)} = 2m_b + E_{\Upsilon(1S)}$ is implemented automatically and explicitly.

4. Determination of the renormalon residue N_m and N_V

The asymptotic behavior of the coefficients $v_N(\mu)$ of the static singlet potential,

$$V(r) = -\frac{4\pi}{3}\frac{1}{r} a_-(\mu) \left[1 + v_1(\mu) a_-(\mu) + v_2 a_-(\mu)^2 + v_3 a_-(\mu)^3 + \ldots \right], \tag{15}$$

can be determined in complete analogy with those of r_N

$$-\frac{4}{3} v_N(\mu) = N_V \mu r (2\beta_0)^N \sum_{s\geq 0} \widetilde{c}_s \frac{\Gamma(\nu+N+1-s)}{\Gamma(\nu+1-s)} + d_N(\mu) \quad \Rightarrow \tag{16}$$

$$-\frac{4}{3} v_N^{\text{asym}}(\mu) \approx N_V \mu r (2\beta_0)^N \frac{\Gamma(\nu+N+1)}{\Gamma(\nu+1)} \left[1 + \sum_{s=1}^{3} \frac{\nu \cdots (\nu-s+1)}{(N+\nu)\cdots(N+\nu-s+1)} \widetilde{c}_s \right], \tag{17}$$

where in Eq. (17) $d_N = 0$ was taken. We can determine the "strength", N_V, of the leading IR renormalon by approximating the asymptotic $v_N^{\rm asym}(\mu)$ with the exact $v_N(\mu)$ ($N = 0, 1, 2, 3$): $v_N^{\rm asym}(\mu) \approx v_N(\mu) \ \Rightarrow$

$$N_V \approx -\frac{4}{3} v_N(\mu) \Big/ \left\{ \mu r (2\beta_0)^N \frac{\Gamma(\nu+N+1)}{\Gamma(\nu+1)} \left[1 + \sum_{s=1}^{3} \frac{\nu \cdots (\nu - s + 1)}{(N+\nu)\cdots(N+\nu-s+1)} \widetilde{c}_s \right] \right\}. \tag{18}$$

The result for N_V should be the best for the highest available N ($N = 3$) and should also have reduced spurious μ-dependence. At present, the v_j are known up to N^{3}LO (v_3) [16, 17, 18, 19, 20, 21, 22, 23, 24, 25].

In the sum $2m_b + V(r)$ the leading IR renormalon gets cancelled. N_V is then related with N_m by the renormalon cancellation of the sum $2m_b + V(r)$: $2N_m + N_V = 0$. Determining N_m via N_V gives us the value that we use [4]

$$N_m = -N_V/2 = 0.56255(260) \ (N_l = 3). \tag{19}$$

5. Renormalon-subtracted (RS, RS') mass of bottom

The RS mass is defined by subtracting the leading IR renormalon singularity from the pole mass [14]:

$$m_{b,{\rm RS}}(\nu_f) = m_b - N_m \pi \nu_f \sum_{N=0}^{\infty} a_-^{N+1}(\nu_f)(2\beta_0)^N \sum_{s \geq 0} \widetilde{c}_s \frac{\Gamma(\nu+N+1-s)}{\Gamma(\nu+1-s)} \tag{20}$$

Equation (20) is still formal. In practice, one rewrites m in terms of $\overline{m}$ using Eqs. (8)-(9)

$$m_b = \overline{m}_b(1 + (4/3)a_-(\nu_f) + \ldots), \tag{21}$$

and reexpands the perturbation series in Eq. (20) around the same coupling $a_-(\mu)$, at fixed but otherwise arbitrary scale μ:

$$m_{b,{\rm RS}}(\nu_f) = \overline{m}_b \left[1 + \sum_{N=0}^{\infty} h_N(\nu_f) a_-^{N+1}(\nu_f) \right] \Rightarrow m_{b,{\rm RS}}(\nu_f) = \overline{m}_b \left[1 + \sum_{N=0}^{\infty} \widetilde{h}_N(\nu_f;\mu) a_-^{N+1}(\mu) \right], \tag{22}$$

where $h_N(\nu_f)$ is determined from Eq. (14) (with $\mu = \nu_f$ and with the sum truncated at $\widetilde{c}_3$) for $N = 0, 1, 2, 3$. For $N \geq 4$ we take $h_N(\overline{m}_b) = 0$. The coefficients $\widetilde{h}_N(\nu_f;\mu)$ in Eq. (22) are obtained by expanding $a_-(\nu_f)$ in the expansion in powers of $a_-(\mu)$. Note that $m_{b,{\rm RS}}(\nu_f)$ will only marginally depend on μ when we truncate the infinite sum in Eq. (22). On the other hand, the coefficients h_N are functions of ν_f, μ, and $\overline{m}_b$, and are much smaller than $r_N(\mu)$.

A variant of the RS mass is the modified renormalon-subtracted (RS') mass $m_{b,{\rm RS}'}$, where subtractions start at $\sim a^2$ [14]. Specifically in this case, Eqs. (20) and (22) are repeated, with the replacements $m_{b,{\rm RS}}(\nu_f) \mapsto m_{b,{\rm RS}'}(\nu_f)$ and $\sum_{N=0}^{\infty} \mapsto \sum_{N=1}^{\infty}$.

6. Bottom mass from heavy quarkonium

The perturbation expansion of $M_{\Upsilon(1S)}^{(th)}$ is presently known up to $\mathcal{O}(m_b a^5)$ [19, 20, 21, 22]:

$$\begin{aligned} M_{\Upsilon(1S)}^{(th)} &= 2m_b - \frac{4\pi^2}{9} m_b a_-^2(\mu) \Bigg\{ 1 + a_-(\mu) \left[K_{1,0} + K_{1,1} L_p(\mu) \right] + a_-^2(\mu) \sum_{j=0}^{2} K_{2,j} L_p(\mu)^j \\ &\quad + a_-^3(\mu) \Big[K_{3,0,0} + K_{3,0,1} \ln a_-(\mu) + \sum_{j=1}^{3} K_{3,j} L_p(\mu)^j \Big] + \mathcal{O}(a_-^4) \Bigg\}, \end{aligned} \tag{23}$$

μ is the renormalization scale, $L_p(\mu) = \ln(\mu/\mu_b)$ where $\mu_b = (4\pi/3)m_b a_-(\mu)$. $K_{i,j}(N_f)$ and $K_{3,0,j}$ are given, e.g., in [4]. We then rewrite m_b in terms of $m_{b,\mathrm{RS}}$ to implement the leading IR renormalon cancellation. This gives

$$\frac{M^{(th)}_{\Upsilon(1S)}}{m_{b,\mathrm{RS}}(\nu_f)} = 2 + \left[2\pi N_m b a \mathcal{K}_0 - \frac{4\pi^2}{9}a^2\right] + \left[2\pi N_m b a^2\left(\mathcal{K}_1 + z_1\mathcal{K}_0\right) - \frac{4\pi^2}{9}a^3\left(K_{1,0} + K_{1,1}L_{\mathrm{RS}}\right)\right]$$
$$+ \left[2\pi N_m b a^3\left(\mathcal{K}_2 + 2z_1\mathcal{K}_1 + z_2\mathcal{K}_0\right) - \frac{4\pi^2}{9}\left(a^4\sum_{j=0}^{2}K_{2,j}L^j_{\mathrm{RS}} + ba^3\pi N_m\mathcal{K}_0\right)\right] + \mathcal{O}(ba^4, a^5). \quad (24)$$

The terms $\mathcal{O}(ba^4, a^5)$ have a similar structure and were written in [4]. The notations are

$$a \equiv a_-(\mu) = a(\mu, N_f = 3)\ ; \quad b \equiv b(\nu_f) = \nu_f/m_{b,\mathrm{RS}}(\nu_f)\ , \quad N_m = N_m(N_l = 3)\ , \qquad (25a)$$

$$L_{\mathrm{RS}} \equiv L_{\mathrm{RS}}(\mu) = \ln\left(\frac{\mu}{(4\pi/3)m_{b,\mathrm{RS}}(\nu_f)a_-(\mu)}\right),\ \mathcal{K}_N = (2\beta_0)^N\sum_{s=0}^{3}\widetilde{c}_s\frac{\Gamma(\nu + N + 1 - s)}{\Gamma(\nu + 1 - s)}\ . \qquad (25b)$$

In the expression (24) for $M_{\Upsilon(1S)}$, the terms of the same order $(\nu_f/m_{b,\mathrm{RS}})a^n$ and a^{n+1} were combined in common brackets $[\ldots]$, in order to account for the renormalon cancellation.

If using the RS' mass in our approach instead, the above expressions are valid without changes, except that $m_{b,\mathrm{RS}} \mapsto m_{b,\mathrm{RS}'}$ and $\mathcal{K}_0 \mapsto 0$ (and: $h_0(\mu) \mapsto 4/3$).

We note that we take $N_l = 3$ active flavours, as the charm quark mass effects in the binding energy $E_{\Upsilon(1S)}$ are negligible [26].

We extract the bottom masses from the condition $M^{(th)}_{\Upsilon(1S)} = M^{(exp)}_{\Upsilon(1S)} (= 9.460\ \mathrm{GeV})$. The error estimates are made assuming $\mu = 2.5^{+1.5}_{-1.0}$ GeV [we varied μ in Eq. (24) but not in Eq. (22)], $\nu_f = 2 \pm 1$ GeV, $\alpha_s(M_z) = 0.1184(7)$ (and decoupling at $\overline{m}_b = 4.2$ GeV and at $\overline{m}_c = 1.27$ GeV), $N_m = 0.56255(260)$, and $(4/3)r_3(\overline{m}_b; N_l) = 1687.1 \pm 21.5$ [10].

In RS and RS' approaches we extract, in MeV, respectively

$$\begin{aligned}
m_{b,\mathrm{RS}}(2\mathrm{GeV}) &= 4\,437^{-11}_{+43}(\mu)^{-3}_{+5}(\nu_f)^{-2}_{+2}(\alpha_s)^{-41}_{+41}(N_m)^{-0}_{+0}(r_3); && (26a)\\
\Rightarrow \quad \overline{m}_b &= 4\,217^{-10}_{+39}(\mu)^{-3}_{+5}(\nu_f)^{-5}_{+5}(\alpha_s)^{-1}_{+1}(N_m)^{-4}_{+4}(r_3). && (26b)\\
m_{b,\mathrm{RS}'}(2\ \mathrm{GeV}) &= 4\,761^{-16}_{+41}(\mu)^{-3}_{+5}(\nu_f)^{+4}_{-3}(\alpha_s)^{-26}_{+26}(N_m)^{-0}_{+0}(r_3); && (26c)\\
\Rightarrow \quad \overline{m}_b &= 4\,223^{-14}_{+36}(\mu)^{-2}_{+4}(\nu_f)^{-4}_{+4}(\alpha_s)^{-1}_{+1}(N_m)^{-4}_{+4}(r_3). && (26d)
\end{aligned}$$

The uncertainties in $\overline{m}_b$ are dominated by the variation of the renormalization scale μ.

The renormalon cancellations are reflected numerically in Eq. (24) [we take $\mu = 2.5$ GeV]:

$$\begin{aligned}
\mathrm{RS} : M_{\Upsilon(1S)} &= (8874 + 431 + 167 + 18 - 30)\ \mathrm{MeV}\,, && (27a)\\
\mathrm{RS}' : M_{\Upsilon(1S)} &= (9521 - 150 + 112 + 8 - 31)\ \mathrm{MeV}\,, && (27b)
\end{aligned}$$

The convergence is good; except for the last (NNNLO) term $\mathcal{O}(a^5, ba^4)$, where the factorization scale dependence becomes stronger, which may signal the importance of ultrasoft effects.

The relations between RS (RS') mass and $\overline{\mathrm{MS}}$ mass are reasonably convergent:

$$\begin{aligned}
m_{b,\mathrm{RS}}(2\ \mathrm{GeV}) &= (4217 + 191 + 36 + 12 - 19)\ \mathrm{MeV}\ , && (28a)\\
m_{b,\mathrm{RS}'}(2\ \mathrm{GeV}) &= (4223 + 478 + 60 + 18 - 17)\ \mathrm{MeV}\ , && (28b)
\end{aligned}$$

where the expansion parameter is taken to be $a(2.5\ \mathrm{GeV})$. A bigger value for the renormalization scale, closer to the bottom quark mass, makes the last term smaller.

Until now we have approximated $\delta m_c = 0$ in Eq. (20). However, $\delta m_c \approx -2$ MeV, Eq.(12). Hence, we have to add 2 MeV to the values of $\bar{m}_b$ obtained in Eqs. (26b) and (26b) (in Ref. [4] it was incorrectly subtracted), leading to the final average of the RS and RS' extractions

$$\overline{m}_b = 4222(40)\ \text{MeV}\ . \tag{29}$$

where we have rounded the ± variation of each parameter to the maximum and added them in quadrature.

7. Conclusions

(i) We presented strong numerical indications that the charm quark decouples in the relation between m_b and $\overline{m}_b$ ($\Rightarrow N_l = 3$).
(ii) An improved determination of the residue of the leading renormalon for the bottom pole mass (and static potential with $N_l = 3$) was performed: $N_m = 0.56255(260)$.
(iii) Use of the 3-loop ($\sim a^5 \overline{m}_b$) corection to the $\Upsilon(1S)$ binding energy, and 4-loop relation between m_b and $\overline{m}_b$, allowed us to perform extraction of $m_{b,\mathrm{RS}^{(\prime)}}$ and $\overline{m}_b$ to NNNLO, with the resulting values Eq. (29). The uncertainties are dominated by the variation of the renormalization scale.

Acknowledgments

This work was supported in part by FONDECYT (Chile) Grant No. 1130599 (GC); Spanish Government and ERDF funds (EU Commission) FPA2014-53631-C2-1-P and by CONICYT "Becas Chile" Grant No. 74150052 (CA); and Spanish Government FPA2014-55613-P, FPA2013-43425-P (AP).

References

[1] Pineda A 1998 "Heavy quarkonium and nonrelativistic effective field theories". *PhD. thesis*
[2] Hoang A H, Smith M C, Stelzer T and Willenbrock S 1999 *Phys. Rev.* D **59** 114014 (arXiv:hep-ph/9804227)
[3] Beneke M 1998 *Phys. Lett.* B **434** 115 (arXiv:hep-ph/9804241)
[4] Ayala C, Cvetič G and and Pineda A 2014 *JHEP* **1409** 045 (arXiv:1407.2128 [hep-ph])
[5] Marquard P, Smirnov A V, Smirnov V A and Steinhauser M 2015 *Phys. Rev. Lett.* **114** 142002 (arXiv:1502.01030 [hep-ph])
[6] Tarrach R 1981 *Nucl. Phys.* B **183** 384
[7] Gray N, Broadhurst D J, Grafe W and Schilcher K 1990 *Z. Phys.* C **48** 673
[8] K. G. Chetyrkin K G and M. Steinhauser M 1999 *Phys. Rev. Lett.* **83** 4001 (arXiv:hep-ph/9907509)
[9] Melnikov K and van Ritbergen T 2000 *Phys. Lett.* B **482** 99 (arXiv:hep-ph/9912391)
[10] Kiyo Y, Mishima G and Sumino Y 2015 *JHEP* **1511** 084 (arXiv:1506.06542 [hep-ph])
[11] Bekavac S, Grozin A, Seidel D and Steinhauser M 2007 *JHEP* **0710** 006 (arXiv:0708.1729 [hep-ph])
[12] Ball P, Beneke M and Braun V M 1995 *Nucl. Phys.* B **452** 563 (arXiv:hep-ph/9502300)
[13] Beneke M 1995 *Phys. Lett.* B **344** 341 (arXiv:hep-ph/9408380)
[14] Pineda A 2001 *JHEP* **0106** 022 (arXiv:hep-ph/0105008)
[15] Contreras C, Cvetič G and Gaete P 2004 *Phys. Rev.* D **70** 034008 (arXiv:hep-ph/0311202)
[16] Fischler W 1977 *Nucl. Phys.* B **129** 157
[17] Billoire A 1980 *Phys. Lett.* B **92** 343
[18] Schröder Y 1999 *Phys. Lett.* B **447** 321 (arXiv:hep-ph/9812205)
[19] Pineda A and Yndurain F J 1998 *Phys. Rev.* D **58** 094022 (arXiv:hep-ph/9711287)
[20] Brambilla N, Pineda A, Soto J and Vairo A 1999 *Phys. Lett.* B **470** 215 (arXiv:hep-ph/9910238)
[21] Kniehl B A, Penin A A, Smirnov V A and Steinhauser M 2002 *Nucl. Phys.* B **635** 357 (arXiv:hep-ph/0203166)
[22] Penin A A and Steinhauser M 2002 *Phys. Lett.* B **538** 335 (arXiv:hep-ph/0204290)
[23] Smirnov A V, Smirnov V A and Steinhauser M 2008 *Phys. Lett.* B **668** 293 (arXiv:0809.1927 [hep-ph])
[24] Anzai C, Kiyo Y and Sumino Y 2010 *Phys. Rev. Lett.* **104** 112003 (arXiv:0911.4335 [hep-ph])
[25] Smirnov A V, Smirnov V A and Steinhauser M 2010 *Phys. Rev. Lett.* **104** 112002 (arXiv:0911.4742 [hep-ph])
[26] Brambilla N, Sumino Y and Vairo A 2002 *Phys. Rev.* D **65** 034001 (arXiv:hep-ph/0108084)