Noname manuscript No.
(will be inserted by the editor)

A convergent relaxation of the Douglas-Rachford algorithm

Nguyen Hieu Thao

Received: date / Accepted: date

Abstract This paper proposes an algorithm for solving structured optimization problems, which covers both the backward-backward and the Douglas-Rachford algorithms as special cases, and analyzes its convergence. The set of fixed points of the algorithm is characterized in several cases. Convergence criteria of the algorithm in terms of general fixed point operators are established. When applying to nonconvex feasibility including the inconsistent case, we prove local linear convergence results under mild assumptions on regularity of individual sets and of the collection of sets which need not intersect. In this special case, we refine known linear convergence criteria for the Douglas-Rachford algorithm (DR). As a consequence, for feasibility with one of the sets being affine, we establish criteria for linear and sublinear convergence of convex combinations of the alternating projection and the DR methods. These results seem to be new. We also demonstrate the seemingly improved numerical performance of this algorithm compared to the RAAR algorithm for both consistent and inconsistent sparse feasibility problems.

Keywords Almost averagedness · Picard iteration · projection method · Krasnoselski-Mann relaxation · metric subregularity · transversality

Mathematics Subject Classification (2000) Primary 49J53 · 65K10
Secondary 49K40 · 49M05 · 49M27 · 65K05 · 90C26

The research leading to these results has received funding from the German-Israeli Foundation Grant G-1253-304.6 and the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement N° 339681.

Nguyen Hieu Thao
Institut für Numerische und Angewandte Mathematik, Universität Göttingen, 37083 Göttingen, Germany
Delft Center for Systems and Control, Delft University of Technology, 2628CD Delft, The Netherlands
E-mail: h.nguyen@math.uni-goettingen.de, h.t.nguyen-3@tudelft.nl

1 Introduction

Convergence analysis has been one of the central and very active applications of variational analysis and mathematical optimization. Examples of recent contributions to the theory of the field that have initiated efficient programs of analysis are [1,2,7,39]. It is the common recipe emphasized in these and many other works that there are two key ingredients required in order to derive convergence of a numerical method 1) regularity of the individual functions or sets such as *convexity* and *averagedness*, and 2) regularity of families of functions or sets at their critical points such as *transversality*, *Kurdyka-Lojasiewicz property* and *metric subregularity.* The question of convergence for a given method can therefore be reduced to checking regularity properties of the problem data. There have been a considerable number of works studying the two ingredients of convergence analysis in order to provide sharper tools in various circumstances, especially in nonconvex cases, e.g., [5,13,14,20,27,28,32–34,39,43,46].

This paper proposes an algorithm called T_λ, which covers both the backward - backward/alternating projection and DR algorithms as special cases of choosing the parameter $\lambda \in [0,1]$, and reports its convergence results. When applied to feasibility for two sets with one of the sets being affine, T_λ is a convex combination of the two fundamental projection methods. On the other hand, T_λ can be viewed as a relaxation of DR. Motivation for relaxing the DR algorithm comes from its lack of stability for inconsistent feasibility. This phenomenon has been observed for the *phase retrieval problem* via the Fourier transform which is essentially inconsistent due to the reciprocal relationship between the spatial and frequency variables of the Fourier transform [36,37]. To address this issue, a relaxation of DR, often known as RAAR, was introduced and applied to phase retrieval problems by Luke in the aforementioned papers. In the framework of feasibility, RAAR is described as convex combinations of the basic DR operator and one projector. Preliminary numerical tests on the performance of T_λ in comparison with RAAR look promising. This has motivated the study of convergence analysis of T_λ in this current work.

After introducing basic notation and proving preliminary results in Section 2, we introduce T_λ in terms of fixed point operators, discuss its fixed point set (Proposition 1), and establish general convergence criteria for T_λ (Theorem 2) in Section 3. We discuss T_λ in the framework of feasibility problems in Section 4. The fixed point set of T_λ is characterized for convex inconsistent feasibility (Proposition 3). For consistent feasibility we prove almost averagedness of T_λ (Proposition 4) and metric subregularity of $T_\lambda - \mathrm{Id}$ (Lemma 3) under regular properties of individual sets and of collections of sets, respectively. The two ingredients are combined to obtain local linear convergence of T_λ (Theorem 4). Section 5 is devoted to demonstrate the improved numerical performance of T_λ compared to the RAAR algorithm for both consistent and inconsistent feasibility problems.

2 Notation and preliminary results

Our notation is standard, c.f. [12,41,48]. The setting throughout this paper is a finite dimensional Euclidean space $\mathbb{E}$. The norm $\|\cdot\|$ denotes the Euclidean norm. The open unit ball and the unit sphere in a Euclidean space are denoted $\mathbb{B}$ and $\mathbb{S}$, respectively. $\mathbb{B}_\delta(x)$ stands for the open ball with radius $\delta > 0$ and center x. The distance to a set $A \subset \mathbb{E}$ with respect to the bivariate function $\operatorname{dist}(\cdot,\cdot)$ is defined by

$$\operatorname{dist}(\cdot, A)\colon \mathbb{E} \to \mathbb{R}\colon\, x \mapsto \inf_{y\in A} \operatorname{dist}(x,y).$$

We use the convention that the distance to the empty set is $+\infty$. The set-valued mapping

$$P_A : \mathbb{E} \rightrightarrows \mathbb{E} : x \mapsto \{y \in A \mid \operatorname{dist}(x,y) = \operatorname{dist}(x,A)\}$$

is the *projector* on A. An element $y \in P_A(x)$ is called a *projection.* This exists for any closed set $A \subset \mathbb{E}$, as can be deduced by the continuity and the coercivity of the norm. Note that the projector is not, in general, single-valued, and indeed uniqueness of the projector defines a type of regularity of the set A: local uniqueness characterizes *prox-regularity* [47] while global uniqueness characterizes convexity [9]. Closely related to the projector is the *prox* mapping corresponding to a function f and a stepsize $\tau > 0$ [42]

$$\operatorname{prox}_{\tau,f}(x) := \operatorname{argmin}_{y\in\mathbb{E}} \left\{ f(y) + \tfrac{1}{2\tau}\|y-x\|^2 \right\}.$$

When $f = \iota_A$ is the *indicator function* of A, that is $\iota_A(x) = 0$ if $x \in A$ and $\iota_A(x) = +\infty$ otherwise, then $\operatorname{prox}_{\tau,\iota_A} = P_A$ for all $\tau > 0$. The value function corresponding to the prox mapping is known as the *Moreau* envelope, which we denote by $e_{\tau,f}(x) := \inf_{y\in\mathbb{E}} \left\{ f(y) + \frac{1}{2\tau}\|y-x\|^2 \right\}$. When $\tau = 1$ and $f = \iota_A$ the Moreau envelope is just one-half the squared distance to the set A: $e_{1,\iota_A}(x) = \frac{1}{2}\operatorname{dist}^2(x,A)$. The *inverse projector* P_A^{-1} is defined by

$$P_A^{-1}(y) := \{x \in \mathbb{E} \mid y \in P_A(x)\}.$$

The *proximal normal cone* to A at $\bar{x}$ is the set, which need not be either closed or convex,

$$N_A^{\mathrm{prox}}(\bar{x}) := \operatorname{cone}\left(P_A^{-1}\bar{x} - \bar{x}\right). \tag{1}$$

If $\bar{x} \notin A$, then $N_A^{\mathrm{prox}}(\bar{x})$ is defined to be empty. Normal cones are central to characterizations both of the regularity of individual sets and of the regularity of collections of sets. For a refined numerical analysis of projection methods, one can define the *Λ-proximal normal cone* to A at $\bar{x}$

$$N_{A|\Lambda}^{\mathrm{prox}}(\bar{x}) := \operatorname{cone}\left((P_A^{-1}(\bar{x}) \cap \Lambda) - \bar{x}\right).$$

When $\Lambda = \mathbb{E}$, it coincides with the proximal normal cone (1). We refer the reader to [5] for a thorough discussion on the restricted versions of various normal cones.

For $\varepsilon \geq 0$ and $\delta > 0$, a set A is (ε,δ)-*regular relative to* Λ at $\bar{x} \in A$ [14, Definition 2.9] if for all $x \in \mathbb{B}_\delta(\bar{x})$, $a \in A \cap \mathbb{B}_\delta(\bar{x})$ and $v \in N^{\mathrm{prox}}_{A|\Lambda}(a)$,

$$\langle x - a, v\rangle \leq \varepsilon \|x - a\| \, \|v\| .$$

When $\Lambda = \mathbb{E}$, the quantifier "relative to" is dropped.

For a set-valued operator $T : \mathbb{E} \rightrightarrows \mathbb{E}$, its *fixed point set* is defined by $\mathsf{Fix}\, T := \{x \in \mathbb{E} \mid x \in Tx\}$. We denote the λ-*reflector* of T by $R_{T,\lambda} := (1 + \lambda)T - \lambda\,\mathrm{Id}$. A frequently used example in this paper corresponds to T being a projector.

In the context of convergence analysis of Picard iterations, the following generalization of the Fejér monotonicity of sequences appears frequently, see, for example, the book [4] or the paper [40] for the terminology.

Definition 1 (linear monotonicity) The sequence (x_k) is *linearly monotone* with respect to a set $S \subset \mathbb{E}$ with rate $c \in [0, 1]$ if

$$\mathrm{dist}(x_{k+1}, S) \leq c\, \mathrm{dist}(x_k, S) \quad \forall k \in \mathbb{N}.$$

Our analysis follows the abstract analysis program proposed in [39] which requires the two key components of the convergence: *almost averagedness* and *metric subregularity.*

Definition 2 (almost nonexpansive/averaged mappings) [39] Let $T : \mathbb{E} \to \mathbb{E}$ and $U \subset \mathbb{E}$.

(i) T is *pointwise almost nonexpansive* on U at $y \in U$ with violation ε if for all $x \in U$, $x^+ \in Tx$, and $y^+ \in Ty$,

$$\left\|x^+ - y^+\right\| \leq \sqrt{1+\varepsilon}\, \|x - y\| .$$

(ii) T is *pointwise almost averaged* on U at $y \in U$ with violation ε and averaging constant $\alpha > 0$ if for all $x \in U$, $x^+ \in Tx$, and $y^+ \in Ty$,

$$\left\|x^+ - y^+\right\|^2 \leq (1+\varepsilon) \|x - y\|^2 - \frac{1-\alpha}{\alpha} \left\|(x^+ - x) - (y^+ - y)\right\|^2 . \quad (2)$$

When a property holds on U at every point $z \in U$, we simply say the property holds on U.

From Definition 2, the almost nonexpansive property is actually the almost averaged property with the same violation and averaging constant $\alpha = 1$.

It is worth noting that if the iteration $x_{k+1} \in Tx_k$ is linearly monotone with respect to $\mathsf{Fix}\, T$ and T is almost averaged, then (x_k) converges R-linearly to a fixed point [40, Proposition 3.5].

We next prove a fundamental preliminary result for our analysis regarding almost averaged mappings.

Lemma 1 *Let $T : \mathbb{E} \rightrightarrows \mathbb{E}$, $U \subset \mathbb{E}$ and $\lambda \in [0, 1]$. The following statements are equivalent.*

(i) T *is almost averaged on* U *with violation* ε *and averaging constant* $\alpha \le \frac{1}{1+\lambda}$.
(ii) *The* λ*-reflector* $R_{T,\lambda} := (1+\lambda)T - \lambda\,\mathrm{Id}$ *is almost averaged on* U *with violation* $(1+\lambda)\varepsilon$ *and averaging constant*

$$\alpha' := \frac{1}{1+(1+\lambda)\left(\frac{1-\alpha}{\alpha}-\lambda\right)}. \tag{3}$$

That is, for all $x, y \in U$, $\tilde{x} \in R_{T,\lambda}x$ *and* $\tilde{y} \in R_{T,\lambda}y$,

$$\|\tilde{x}-\tilde{y}\|^2 \le (1+(1+\lambda)\varepsilon)\|x-y\|^2 - \frac{\frac{1-\alpha}{\alpha}-\lambda}{1+\lambda}\|(\tilde{x}-x)-(\tilde{y}-y)\|^2. \tag{4}$$

Proof Let $x, y \in U$, $x^+ \in Tx$, $y^+ \in Ty$, $\tilde{x} = (1+\lambda)x^+ - \lambda x \in R_{T,\lambda}x$ and $\tilde{y} = (1+\lambda)y^+ - \lambda y \in R_{T,\lambda}y$. We have by definition of $R_{T,\lambda}$ and [4, Corollary 2.14] that

$$\begin{aligned}
&\|\tilde{x}-\tilde{y}\|^2 \\
&= \left\|(1+\lambda)(x^+-y^+) - \lambda(x-y)\right\|^2 \\
&= (1+\lambda)\left\|x^+-y^+\right\|^2 - \lambda\|x-y\|^2 + \lambda(1+\lambda)\left\|(x^+-y^+)-(x-y)\right\|^2.
\end{aligned} \tag{5}$$

Substituting (2) into (5) and noting that

$$\|(\tilde{x}-x)-(\tilde{y}-y)\| = (1+\lambda)\left\|(x^+-x)-(y^+-y)\right\|,$$

we obtain

$$\begin{aligned}
&\|\tilde{x}-\tilde{y}\|^2 \\
&\le (1+\varepsilon(1+\lambda))\|x-y\|^2 - (1+\lambda)\left(\frac{1-\alpha}{\alpha}-\lambda\right)\left\|(x^+-x)-(y^+-y)\right\|^2 \\
&= (1+\varepsilon(1+\lambda))\|x-y\|^2 - \frac{\frac{1-\alpha}{\alpha}-\lambda}{1+\lambda}\|(\tilde{x}-x)-(\tilde{y}-y)\|^2,
\end{aligned}$$

which is exactly (4).

Conversely, substituting (4) into (5), we obtain

$$\begin{aligned}
&(1+\lambda)\left\|x^+-y^+\right\|^2 - \lambda\|x-y\|^2 + \lambda(1+\lambda)\left\|(x^+-y^+)-(x-y)\right\|^2 \\
&\le (1+(1+\lambda)\varepsilon)\|x-y\|^2 - \frac{\frac{1-\alpha}{\alpha}-\lambda}{1+\lambda}\|(\tilde{x}-x)-(\tilde{y}-y)\|^2 \\
&= (1+(1+\lambda)\varepsilon)\|x-y\|^2 - (1+\lambda)\left(\frac{1-\alpha}{\alpha}-\lambda\right)\left\|(x^+-x)-(y^+-y)\right\|^2,
\end{aligned}$$

which is equivalent to (2). The proof is complete. □

Lemma 1 generalizes [14, Lemma 2.4] where the result was proved for $\alpha = 1/2$ and $\lambda = 1$.

The next lemma recalls facts regarding the almost averagedness of projectors and reflectors associated with regular sets.

Lemma 2 *Let $A \subset \mathbb{E}$ be (ε,δ)-regular at $\bar{x} \in A$ and define*

$$U := \{x \in \mathbb{E} \mid P_A x \subset \mathbb{B}_\delta(\bar{x})\}.$$

(i) The projector P_A is pointwise almost nonexpansive on U at every point $z \in A \cap \mathbb{B}_\delta(\bar{x})$ with violation $2\varepsilon + \varepsilon^2$.

(ii) The projector P_A is pointwise almost averaged on U at every point $z \in A \cap \mathbb{B}_\delta(\bar{x})$ with violation $2\varepsilon + 2\varepsilon^2$ and averaging constant $1/2$.

(iii) The λ-reflector $R_{P_A,\lambda}$ is pointwise almost averaged on U at every point $z \in A \cap \mathbb{B}_\delta(\bar{x})$ with violation $(1+\lambda)(2\varepsilon + 2\varepsilon^2)$ and averaging constant $\frac{1+\lambda}{2}$, that is, for all $x \in U$, $x^+ \in R_{P_A,\lambda}x$ and $z \in A \cap \mathbb{B}_\delta(\bar{x})$,

$$\left\|x^+ - z\right\|^2 \le \left(1 + (1+\lambda)(2\varepsilon + 2\varepsilon^2)\right)\|x - z\|^2 - \frac{1-\lambda}{1+\lambda}\left\|x^+ - x\right\|^2.$$

Proof Statements (i) and (ii) were proved in [14, Theorem 2.14]. Statement (iii) follows from (ii) and Lemma 1 applied to $T = P_A$ and $\alpha = 1/2$. ☐

The following concept of *metric subregularity with functional modulus* has played a central role, implicitly and explicitly, in the analysis of convergence of Picard iterations [1,14,39,40]. Recall that a function $\mu : [0,\infty) \to [0,\infty)$ is a *gauge function* if μ is continuous and strictly increasing with $\mu(0) = 0$ and $\lim_{t\to\infty} \mu(t) = \infty$.

Definition 3 (metric subregularity with functional modulus) A mapping $F : \mathbb{E} \rightrightarrows \mathbb{E}$ is *metrically subregular with gauge μ on $U \subset \mathbb{E}$ for y relative to $\Lambda \subset \mathbb{E}$* if

$$\mu\left(\operatorname{dist}\left(x, F^{-1}(y) \cap \Lambda\right)\right) \le \operatorname{dist}\left(y, F(x)\right) \quad \forall x \in U \cap \Lambda. \tag{6}$$

When μ is a linear function, that is $\mu(t) = \kappa t$, $\forall t \in [0,\infty)$, one says "with constant κ" instead of "with gauge $\mu = \kappa\,\mathrm{Id}$". When $\Lambda = X$, the quantifier "relative to" is dropped.

Metric subregularity has many important applications in variational analysis and mathematical optimization, see the monographs and papers [12,16–19, 21,22,26,41,45]. For the discussion of metric subregularity in connection with subtransversality of collections of sets, we refer the reader to [24,25,30,31].

The next theorem serves as the basic template for the quantitative convergence analysis of fixed point iterations. By the notation $T : \Lambda \rightrightarrows \Lambda$ where Λ is a subset of $\mathbb{E}$, we mean that $T : \mathbb{E} \rightrightarrows \mathbb{E}$ and $Tx \subset \Lambda$ for all $x \in \Lambda$. This simplification of notation should not lead to any confusion if one keeps in mind that there may exist fixed points of T that are not in Λ. For the importance of the use of Λ in isolating the desirable fixed point, we refer the reader to [1, Example 1.8]. In the following, $\operatorname{ri}\Lambda$ denotes the *relative interior* of Λ.

Theorem 1 *[39, Theorem 2.1] Let $T : \Lambda \rightrightarrows \Lambda$ for $\Lambda \subset \mathbb{E}$ and let $S \subset \operatorname{ri}\Lambda$ be closed and nonempty with $Ty \subset \mathsf{Fix}\ T \cap S$ for all $y \in S$. Let $\mathcal{O}$ be a neighborhood of S such that $\mathcal{O} \cap \Lambda \subset \operatorname{ri}\Lambda$. Suppose that*

(a) T is pointwise almost averaged at all points $y \in S$ with violation ε and averaging constant $\alpha \in (0,1)$ on $\mathcal{O} \cap \Lambda$, and
(b) there exists a neighborhood $\mathcal{V}$ of $\mathsf{Fix}\ T \cap S$ and a constant $\kappa > 0$ such that for all $y \in S$, $y^+ \in Ty$ and all $x^+ \in Tx$ the estimate

$$\kappa \operatorname{dist}(x, S) \le \left\| \left(x - x^+\right) - \left(y - y^+\right) \right\| \tag{7}$$

holds whenever $x \in (\mathcal{O} \cap \Lambda) \setminus (\mathcal{V} \cap \Lambda)$.

Then for all $x^+ \in Tx$

$$\operatorname{dist}\left(x^+, \mathsf{Fix}\ T \cap S\right) \le \sqrt{1 + \varepsilon - \frac{(1-\alpha)\kappa^2}{\alpha}}\operatorname{dist}(x, S)$$

whenever $x \in (\mathcal{O} \cap \Lambda) \setminus (\mathcal{V} \cap \Lambda)$.

In particular, if $\kappa > \sqrt{\frac{\varepsilon\alpha}{1-\alpha}}$, then for any initial point $x_0 \in \mathcal{O} \cap \Lambda$ the iteration $x_{k+1} \in Tx_k$ satisfies

$$\operatorname{dist}\left(x_{k+1}, \mathsf{Fix}\ T \cap S\right) \le c^k \operatorname{dist}(x_0, S)$$

with $c := \sqrt{1 + \varepsilon - \frac{(1-\alpha)\kappa^2}{\alpha}} < 1$ for all k such that $x_j \in (\mathcal{O} \cap \Lambda) \setminus (\mathcal{V} \cap \Lambda)$ for $j = 1, 2, \ldots, k$.

Remark 1 In the case of $S = \mathsf{Fix}\ T$ condition (7) reduces to metric subregularity of the mapping $F := T - \mathrm{Id}$ for 0 on the annular set, that is

$$\kappa \operatorname{dist}(x, F^{-1}(0)) \le \operatorname{dist}(0, F(x)) \quad \forall x \in (\mathcal{O} \cap \Lambda) \setminus (\mathcal{V} \cap \Lambda).$$

The inequality $\kappa > \sqrt{\frac{\varepsilon\alpha}{1-\alpha}}$ then states that the constant of metric subregularity κ is sufficiently large relative to the violation of the averaging property to guarantee a linear progression of the iterates through that annular region.

For a comprehensive discussion on the roles of S and Λ in the analysis program of Theorem 1, we would like to refer the reader to the paper [39].

For the sake of simplification in terms of presentation, we have chosen to reduce the number of technical constants appearing in the analysis. It would be obviously analogous to formulate more theoretically general results by using more technical constants in appropriate places.

3 T_λ as a fixed point operator

We consider the problem of finding a fixed point of the operator

$$T_\lambda := T_1\left((1+\lambda)T_2 - \lambda\,\mathrm{Id}\right) - \lambda\left(T_2 - \mathrm{Id}\right), \tag{8}$$

where $\lambda \in [0,1]$ and $T_i : \mathbb{E} \rightrightarrows \mathbb{E}$ $(i = 1,2)$ are assumed to be easily computed.

Examples of T_λ include the backward-backward and the DR algorithms [8,11,35,37,44] for solving the structured optimization problem

$$\underset{x\in\mathbb{E}}{\text{minimize}}\; f_1(x) + f_2(x)$$

under different assumptions on the functions f_i $(i = 1,2)$. Indeed, when T_i are the prox mappings of f_i with parameters $\tau_i > 0$, then T_λ with $\lambda = 0$ and 1 takes the form $T_\lambda = \mathrm{prox}_{\tau_1,f_1} \circ \mathrm{prox}_{\tau_2,f_2}$, and $T_\lambda = \mathrm{prox}_{\tau_1,f_1}\left(2\,\mathrm{prox}_{\tau_2,f_2} - \mathrm{Id}\right) - \mathrm{prox}_{\tau_2,f_2} + \mathrm{Id}$, respectively.

We first describe the fixed point set of T_λ via those of the constituent operators T_i $(i = 1,2)$.

Proposition 1 *Let $T_1, T_2 : \mathbb{E} \to \mathbb{E}$, $\lambda \in [0,1]$ and consider*

$$T_\lambda = T_1\left((1+\lambda)T_2 - \lambda\,\mathrm{Id}\right) - \lambda(T_2 - \mathrm{Id}). \tag{9}$$

The following statements hold true.

(i) $(1+\lambda)T_\lambda - \lambda\,\mathrm{Id} = \left((1+\lambda)T_1 - \lambda\,\mathrm{Id}\right) \circ \left((1+\lambda)T_2 - \lambda\,\mathrm{Id}\right)$. *As a consequence,*

$$\mathsf{Fix}\; T_\lambda = \mathsf{Fix}\;\left((1+\lambda)T_1 - \lambda\,\mathrm{Id}\right) \circ \left((1+\lambda)T_2 - \lambda\,\mathrm{Id}\right).$$

(ii) Suppose that $T_1 = P_A$ is the projector associated with an affine set A and T_2 is single-valued. Then

$$\begin{aligned}\mathsf{Fix}\; T_\lambda &= \{x \in \mathbb{E} \mid P_A x = \lambda T_2 x + (1-\lambda)x\}\\ &\subset \{x \in \mathbb{E} \mid P_A x = P_A T_2 x\}.\end{aligned}$$

Proof (i).

$$\begin{aligned}(1+\lambda)T_\lambda - \lambda\,\mathrm{Id} &= (1+\lambda)\left(T_1\left((1+\lambda)T_2 - \lambda\,\mathrm{Id}\right) - \lambda(T_2 - \mathrm{Id})\right) - \lambda\,\mathrm{Id}\\ &= (1+\lambda)T_1\left((1+\lambda)T_2 - \lambda\,\mathrm{Id}\right) - \lambda\left[(1+\lambda)T_2 - \lambda\,\mathrm{Id}\right]\\ &= \left((1+\lambda)T_1 - \lambda\,\mathrm{Id}\right) \circ \left((1+\lambda)T_2 - \lambda\,\mathrm{Id}\right).\end{aligned}$$

(ii).

$$\begin{aligned}x = T_\lambda x &= P_A\left((1+\lambda)T_2 x - \lambda x\right) - \lambda(T_2 x - x)\\ &\Leftrightarrow \lambda T_2 x + (1-\lambda)x = P_A\left((1+\lambda)T_2 x - \lambda x\right).\end{aligned} \tag{10}$$

In particular, $\lambda T_2 x + (1-\lambda)x \in A$. This together with equality (10) and P_A being an affine operator imply

$$\begin{aligned} P_A\left(\lambda T_2 x + (1-\lambda)x\right) &= P_A\left((1+\lambda)T_2 x - \lambda x\right) \\ \Leftrightarrow \lambda P_A T_2 x + (1-\lambda)P_A x &= (1+\lambda)P_A T_2 x - \lambda P_A x \\ \Leftrightarrow P_A x &= P_A T_2 x. \end{aligned} \tag{11}$$

Substituting (11) into (10) yields

$$\begin{aligned} \lambda T_2 x + (1-\lambda)x &= (1+\lambda)P_A T_2 x - \lambda P_A x \\ &= (1+\lambda)P_A x - \lambda P_A x = P_A x. \end{aligned}$$

The proof is complete. □

The next proposition shows that the almost averagedness of T_λ naturally inherits from that of T_1 and T_2 via Krasnoselski–Mann relaxations.

Proposition 2 (almost averagedness of T_λ) *Let T_i $(i=1,2)$ be almost averaged on $U_i \subset \mathbb{E}$ with violation ε and averaging constant α and define*

$$U = \{x \in U_2 \mid R_{T_2,\lambda}x \subset U_1\}.$$

Then T_λ is almost averaged on U with violation $2\varepsilon + (1+\lambda)\varepsilon^2$ and averaging constant

$$\tilde{\alpha} := \frac{1}{(1+\lambda)\left(1+\frac{1+\lambda}{2}\left(\frac{1-\alpha}{\alpha}-\lambda\right)\right)}. \tag{12}$$

Proof By Lemma 1, the operators $R_{T_i,\lambda} := (1+\lambda)T_i - \lambda\,\mathrm{Id}$ $(i=1,2)$ are almost averaged on U_i with violation $(1+\lambda)\varepsilon$ and averaging constant α' given by (3). Then thanks to [39, Proposition 2.10 (iii)], the operator $T := R_{T_1,\lambda}R_{T_2,\lambda}$ is almost averaged on U with violation $(1+\lambda)\left(2\varepsilon + (1+\lambda)\varepsilon\right)$ and averaging constant $\frac{2\alpha'}{1+\alpha'}$. Note that $T_\lambda = (1+\lambda)T - \lambda\,\mathrm{Id}$ by Proposition 1. We have, again by Lemma 1, that T_λ is almost averaged on U with violation $2\varepsilon + (1+\lambda)\varepsilon$ and averaging constant $\tilde{\alpha}$ given by (12) as claimed. □

We next discuss convergence of T_λ based on the abstract results established in [39]. Our agenda is to verify the assumptions of Theorem 1. To simplify the exposure in terms of presentation, we have chosen to state the results corresponding to $S = \mathsf{Fix}\ T_\lambda$ and $\Lambda = \mathbb{E}$ in Theorem 1.

Theorem 2 (convergence of T_λ with metric subregularity) *Let T_λ be defined at (9) with T_i $(i=1,2)$ being almost averaged with violation ε and averaging constant α. Denote $S_\rho := \mathsf{Fix}\ T_\lambda + \rho\mathbb{B}$ for a nonnegative real ρ. Suppose that there are $\delta > 0$ and $\gamma \in (0,1)$ such that, for each $n \in \mathbb{N}$, the mapping $F := T_\lambda - \mathrm{Id}$ is metrically subregular with gauge μ_n on $R_n := S_{\gamma^n\delta} \setminus S_{\gamma^{n+1}\delta}$ for 0, where μ_n satisfies*

$$\inf_{x \in R_n} \frac{\mu_n\left(\mathrm{dist}\left(x, \mathsf{Fix}\ T_\lambda\right)\right)}{\mathrm{dist}\left(x, \mathsf{Fix}\ T_\lambda\right)} \geq \kappa_n > \sqrt{\frac{\tilde{\alpha}\varepsilon'}{1-\tilde{\alpha}}}, \tag{13}$$

where $\tilde{\alpha}$ *is given by* (12) *and*

$$\varepsilon' := 2\varepsilon + (1+\lambda)\varepsilon^2. \tag{14}$$

Then, for any initial point $x_0 \in S_\delta$*, the iterates* $x_{k+1} \in T_\lambda x_k$ *satisfy*

$$\operatorname{dist}(x_k, \mathsf{Fix}\, T_\lambda) \to 0, \tag{15}$$

and

$$\operatorname{dist}(x_{k+1}, \mathsf{Fix}\, T_\lambda) \le c_n \operatorname{dist}(x_k, \mathsf{Fix}\, T_\lambda) \quad \forall\, x_k \in R_n, \tag{16}$$

where $c_n := \sqrt{1+\varepsilon' - \frac{(1-\tilde{\alpha})\kappa_n^2}{\tilde{\alpha}}} < 1.$

In particular, if (κ_n) *is bounded below by* $\underline{\kappa} > \sqrt{\frac{\tilde{\alpha}\varepsilon'}{1-\tilde{\alpha}}}$ *for all* n *large enough, then* (x_k) *is eventually R-linearly convergent to a fixed point with rate at most* $c := \sqrt{1+\varepsilon' - \frac{(1-\tilde{\alpha})\underline{\kappa}^2}{\tilde{\alpha}}} < 1.$

Proof For each $n \in \mathbb{N}$, we verify the assumptions of Theorem 1 for $\mathcal{O} = S_{\gamma^n\delta}$ and $\mathcal{V} = S_{\gamma^{n+1}\delta}$. Since T_i $(i=1,2)$ are almost averaged with violation ε and averaging constant α, Proposition 2 ensures that T_λ is almost averaged with violation ε' given by (14) and averaging constant $\tilde{\alpha}$ given by (12). In other words, condition (a) of Theorem 1 is satisfied with $\varepsilon = \varepsilon'$ and $\alpha = \tilde{\alpha}$. On the other hand, the metric subregularity of F with gauge μ_n satisfying (13) for 0 on R_n fulfills condition (b) of Theorem 1 with $\kappa = \kappa_n$ in view of Remark 1. Theorem 1 then yields the conclusion of Theorem 2 after a straightforward care of the involving technical constants. ☐

The first inequality in (13) essentially says that the gauge function μ_n can be bounded from below by a linear function on the reference interval.

4 Applications to feasibility

We consider T_λ for solving feasibility involving two closed sets $A, B \subset \mathbb{E}$,

$$\begin{aligned} x^+ \in T_\lambda x &= P_A\left((1+\lambda)P_B x - \lambda x\right) - \lambda\left(P_B x - x\right) \\ &= P_A R_{P_B,\lambda}(x) - \lambda\left(P_B x - x\right). \end{aligned} \tag{17}$$

Note that T_λ with $\lambda = 0$ and 1 corresponds to the alternating projection $P_A P_B$ and the DR methods $\frac{1}{2}(R_A \circ R_B + \mathrm{Id})$, respectively.

It is worth recalling that feasibility for $m \ge 2$ sets can be reformulated as feasibility for two constructed sets on the product space $\mathbb{E}^m$ with one of the later sets is a linear subspace, and the regularity properties in terms of both individual sets and collections of sets of the later sets are inherited from those of the former ones [3,33].

When A is an affine set, then the projector P_A is affine and T_λ is the convex combination of the alternating projections and the DR since

$$\begin{aligned} T_\lambda x &= P_A\left((1-\lambda)P_Bx + \lambda(2P_Bx - x)\right) - \lambda\left(P_Bx - x\right) \\ &= (1-\lambda)P_AP_Bx + \lambda\left(x + P_A(2P_Bx - x) - P_Bx\right) \\ &= (1-\lambda)T_0(x) + \lambda T_1(x). \end{aligned}$$

In this case, we establish convergence results for all convex combinations of the alternating projections and DR. To our best awareness, this kind of results seems to be new.

Recall that for inconsistent feasibility the DR operator has no fixed points. We next show that the fixed point set of T_λ with $\lambda \in [0,1)$ for convex inconsistent feasibility is nonempty. This result follows the lines of [37, Lemma 2.1] where the fixed point set of the RAAR algorithm was characterized.

Proposition 3 (Fixed points of T_λ for convex inconsistent feasibility) *For closed convex sets A, B, let $G = \overline{B-A}$, $g = P_G0$, $E = A \cap (B-g)$ and $F = (A+g) \cap B$. Then*

$$\mathsf{Fix}\ T_\lambda = E - \frac{\lambda}{1-\lambda}g \quad \forall t \in [0,1).$$

Proof We first show that $E - \frac{\lambda}{1-\lambda}g \subset \mathsf{Fix}\ T_\lambda$. Pick any $e \in E$ and denote $f = e + g \in F$ as definitions of E and F. We are checking that

$$x := e - \frac{\lambda}{1-\lambda}g \in \mathsf{Fix}\ T_\lambda.$$

Since $x = f - \frac{1}{1-\lambda}g$ and $-g \in N_B(f)$, we get $P_Bx = f$.

Analogously, since $g \in N_A(e)$ and

$$(1+\lambda)P_Bx - \lambda x = (1+\lambda)f - \lambda x = e + \frac{1}{1-\lambda}g,$$

we have $P_A((1+\lambda)P_Bx - \lambda x) = e$.

Hence,

$$\begin{aligned} x - T_\lambda x &= x - P_A\left((1+\lambda)P_Bx - \lambda x\right) + \lambda\left(P_Bx - x\right) \\ &= x - e + \lambda\left(f - x\right) = 0. \end{aligned}$$

That is $x \in \mathsf{Fix}\ T_\lambda$.

We next show that $\mathsf{Fix}\ T_\lambda \subset E - \frac{\lambda}{1-\lambda}g$. Pick any $x \in \mathsf{Fix}\ T_\lambda$. Let $f = P_Bx$ and $y = x - f$. Thanks to $x \in \mathsf{Fix}\ T_\lambda$ and the definition of T_λ,

$$\begin{aligned} P_A((1+\lambda)P_Bx - \lambda x) &= \lambda(P_Bx - x) + x \\ &= -\lambda y + y + f = f + (1-\lambda)y. \end{aligned} \tag{18}$$

Now, for any $a \in A$, since A is closed and convex, we have

$$\begin{aligned} 0 &\geq \langle a - P_A((1+\lambda)P_Bx - \lambda x), (1+\lambda)P_Bx - \lambda x - P_A((1+\lambda)P_Bx - \lambda x)\rangle \\ &= \langle a - (f + (1-\lambda)y), (1+\lambda)f - \lambda x - (f + (1-\lambda)y)\rangle \\ &= \langle a - f - (1-\lambda)y, -y\rangle = \langle -a + f, y\rangle + (1-\lambda)\|y\|^2. \end{aligned}$$

On the other hand, for any $b \in B$, since B is closed and convex, we have

$$\langle b - f, y\rangle = \langle b - f, x - f\rangle = \langle b - P_Bx, x - P_Bx\rangle \leq 0.$$

Combining the last two inequalities yields

$$\langle b - a, y\rangle \leq -(1-\lambda)\|y\|^2 \leq 0.$$

Take a sequence (a_n) in A and a sequence (b_n) in B such that $g_n := b_n - a_n \to g$. Then

$$\langle g_n, y\rangle \leq -(1-\lambda)\|y\|^2 \leq 0 \quad \forall n. \tag{19}$$

Taking the limit and using the Cauchy–Schwarz inequality yields

$$\|y\| \leq \frac{1}{1-\lambda}\|g\|.$$

Conversely, by (18) with noting that $f \in B$ and $P_A((1+\lambda)P_Bx - \lambda x) \in A$,

$$\|y\| = \frac{1}{1-\lambda}\|f - P_A((1+\lambda)P_Bx - \lambda x)\| \geq \frac{1}{1-\lambda}\|g\|.$$

Hence $\|y\| = \frac{1}{1-\lambda}\|g\|$, and taking the limit in (19), which yields $y = -\frac{1}{1-\lambda}g$. Since $f \in B$ and $f - g = f + (1-\lambda)y = P_A((1+\lambda)P_Bx - \lambda x) \in A$, we have $f - g \in A \cap (B - g) = E$ and, therefore,

$$x = f + y = f - \frac{1}{1-\lambda}g = f - g - \frac{\lambda}{1-\lambda}g \in E - \frac{\lambda}{1-\lambda}g.$$

□

We next discuss the two key ingredients for convergence of T_λ when applied to feasibility problems: 1) almost averagedness of T_λ, and 2) metric subregularity of $T_\lambda - \mathrm{Id}$. The two properties will be verified under natural assumptions on (ε, δ)-regularity of the sets and the transversality of the collection of sets, respectively.

The next proposition shows averagedness of T_λ for feasibility involving (ε, δ)-regular sets.

Proposition 4 *Let A and B be (ε,δ)-regular at $\bar{x} \in A \cap B$ and define the set*

$$U := \{x \in \mathbb{E} \mid P_B x \subset \mathbb{B}_\delta(\bar{x}) \text{ and } P_A R_{P_B,\lambda} x \subset \mathbb{B}_\delta(\bar{x})\}. \tag{20}$$

Then T_λ is pointwise almost averaged on U at every point $z \in S$ with violation $\tilde{\varepsilon}$ and averaging constant $\frac{2}{3+\lambda}$, where $S := A \cap B \cap \mathbb{B}_\delta(\bar{x})$ and

$$\tilde{\varepsilon} := 2(2\varepsilon + 2\varepsilon^2) + (1+\lambda)(2\varepsilon + 2\varepsilon^2)^2. \tag{21}$$

That is, for all $x \in U$, $x^+ \in T_\lambda x$ and $z \in S$,

$$\left\|x^+ - z\right\|^2 \le (1+\tilde{\varepsilon})\left\|x - z\right\|^2 - \frac{1+\lambda}{2}\left\|x - x^+\right\|^2.$$

Proof Let us define

$$U_A := \{y \in \mathbb{E} \mid P_A y \subset \mathbb{B}_\delta(\bar{x})\}, \quad U_B := \{x \in \mathbb{E} \mid P_B x \subset \mathbb{B}_\delta(\bar{x})\}$$

and note that $x \in U$ if and only if $x \in U_B$ and $R_{P_B,\lambda} x \subset U_A$. Thanks to Lemma 2 (iii), $R_{P_A,\lambda}$ and $R_{P_B,\lambda}$ are pointwise almost averaged on U at every point $z \in S$ with violation $(1+\lambda)(2\varepsilon + 2\varepsilon^2)$ and averaging constant $\frac{1+\lambda}{2}$. Then due to [39, Proposition 2.10 (iii)], the operator $T := R_{P_A,\lambda} R_{P_B,\lambda}$ is pointwise almost averaged on U at every point $z \in S$ with violation $(1+\lambda)\tilde{\varepsilon}$ and averaging constant $\frac{2(1+\lambda)}{3+\lambda}$, where $\tilde{\varepsilon}$ is given by (21). Note that $T_\lambda = (1+\lambda)T - \lambda \mathrm{Id}$ by Proposition 1. Thanks to Lemma 1, T_λ is pointwise almost averaged on U at every point $z \in S$ with violation $\tilde{\varepsilon}$ and averaging constant $\frac{2}{3+\lambda}$ as claimed. □

Remark 2 It follows from Lemma 2 (i) & (iii) that the set U defined by (20) contains at least the ball $\mathbb{B}_{\delta'}(\bar{x})$, where

$$\delta' := \frac{\delta}{2(1+\varepsilon)\sqrt{1 + (1+\lambda)(2\varepsilon + 2\varepsilon^2)}} > 0.$$

We first integrate Proposition 4 into Theorem 2 to obtain convergence of T_λ for consistent feasibility involving regular sets.

Corollary 1 (convergence of T_λ for feasibility) *Consider T_λ for feasibility (17) and suppose that $\mathsf{Fix}\, T_\lambda = A \cap B \neq \emptyset$. Denote $S_\rho := \mathsf{Fix}\, T_\lambda + \rho\mathbb{B}$ for a nonnegative real ρ. Suppose that there are $\delta > 0$, $\varepsilon \ge 0$ and $\gamma \in (0,1)$ such that A and B are (ε,δ')-regular at avery point $z \in A \cap B$, where*

$$\delta' := 2\delta(1+\varepsilon)\sqrt{1 + (1+\lambda)(2\varepsilon + 2\varepsilon^2)},$$

and for each $n \in \mathbb{N}$, the mapping $F := T_\lambda - \mathrm{Id}$ is metrically subregular with gauge μ_n for 0 on $R_n := S_{\gamma^n\delta} \setminus S_{\gamma^{n+1}\delta}$, where μ_n satisfies

$$\inf_{x \in R_n} \frac{\mu_n(\mathrm{dist}(x, A \cap B))}{\mathrm{dist}(x, A \cap B)} \ge \kappa_n > \sqrt{\frac{2\tilde{\varepsilon}}{1+\lambda}},$$

where $\tilde{\varepsilon}$ is given at (21).

Then for any initial point $x_0 \in S_\delta$*, the iterates* $x_{k+1} \in T_\lambda x_k$ *satisfy* (15) *and* (16) *with* $c_n := \sqrt{1+\tilde{\varepsilon}-\frac{(1+\lambda)\kappa_n^2}{2}} < 1$.

In particular, if (κ_n) *is bounded from below by* $\underline{\kappa} > \sqrt{\frac{2\tilde{\varepsilon}}{1+\lambda}}$ *for all* n *large enough, then* (x_k) *is eventually R-linearly convergent to a fixed point with rate at most* $c := \sqrt{1+\tilde{\varepsilon}-\frac{(1+\lambda)\underline{\kappa}^2}{2}} < 1$.

Proof Let any $x \in R_n$, for some $n \in \mathbb{N}$, $x^+ \in T_\lambda x$ and $\bar{x} \in P_{A\cap B}x$. Proposition 4 and Remark 2 imply that T_λ is pointwise almost averaged on $\mathbb{B}_\delta(\bar{x})$ at every point $z \in A \cap B \cap \mathbb{B}_\delta(\bar{x})$ with violation $\tilde{\varepsilon}$ given by (21) and averaging constant $\frac{2}{3+\lambda}$. In other words, condition (a) of Theorem 1 is satisfied. Condition (b) of Theorem 1 is also fulfilled for the same reason as that of Theorem 2. The desired conclusion now follows from Theorem 1. ☐

In practice, the metric subregularity assumption is often more challenging to verify than the averagedness one. In the concrete example of consistent alternating projections $P_A P_B$, that condition holds true if and only if the collection of sets is subtransversal. Our main goal in the rest of this study is to verify the metric subregularity of $T_\lambda - \mathrm{Id}$ under the assumption of transversality. As a result, if the sets are also sufficiently regular, then local linear convergence of the iterates $x_{k+1} \in T_\lambda x_k$ is guaranteed.

We first describe the concept of relative transversality of collections of sets. In the sequel, we set $\Lambda := \mathrm{aff}(A \cup B)$, the smallest affine set in $\mathbb{E}$ containing both A and B.

Assumption 3 *The collection* $\{A, B\}$ *is transversal at* $\bar{x} \in A \cap B$ *relative to* Λ *with constant* $\bar{\theta} < 1$*, that is, for any* $\theta > \bar{\theta}$*, there exists* $\delta > 0$ *such that*

$$\langle u, v\rangle \geq -\theta \|u\| \cdot \|v\|,$$

for all $a \in A \cap \mathbb{B}_\delta(\bar{x})$*,* $b \in B \cap \mathbb{B}_\delta(\bar{x})$*,* $u \in N^{\mathrm{prox}}_{A|\Lambda}(a)$ *and* $v \in N^{\mathrm{prox}}_{B|\Lambda}(b)$.

Thanks to [23, Theorem 1] and [29, Theorem 1], Assumption 3 also ensures subtransversality of $\{A, B\}$ at $\bar{x}$ relative to Λ with constant at least $\sqrt{\frac{1-\theta}{2}}$ on the neighborhood $\mathbb{B}_\delta(\bar{x})$, that is

$$\sqrt{\frac{1-\theta}{2}}\,\mathrm{dist}(x, A \cap B) \leq \max\{\mathrm{dist}(x, A), \mathrm{dist}(x, B)\} \quad \forall x \in \Lambda \cap \mathbb{B}_\delta(\bar{x}). \tag{22}$$

The next lemma is at the heart of our subsequent discussion.

Lemma 3 *Under Assumption 3, for any* $\theta \in (\bar{\theta}, 1)$*, there exists a number* $\delta > 0$ *such that for all* $x \in \mathbb{B}_\delta(\bar{x})$ *and* $x^+ \in T_\lambda x$*,*

$$\kappa\,\mathrm{dist}(x, A \cap B) \leq \left\|x - x^+\right\|, \tag{23}$$

where κ *is defined by*

$$\kappa := \frac{(1-\theta)\sqrt{1+\theta}}{\sqrt{2}\max\left\{1, \lambda + \sqrt{1-\theta^2}\right\}} > 0. \tag{24}$$

Proof Take any $x \in \mathbb{B}_\delta(\bar{x})$, $b \in P_B x$, $y = (1+\lambda)b - \lambda x$, $a \in P_A y$ and $x^+ = a - \lambda(b-x) \in T_\lambda x$. By choosing a smaller δ if necessary, we can assume without loss of generality that $a, b \in \mathbb{B}_\delta(\bar{x})$. For example, one can choose $\delta/6$ in place of δ.

We first note that $x - b \in N^{\text{prox}}_{B|A}(b)$ and $y - a \in N^{\text{prox}}_{A|A}(a)$. Assumption 3 then yields

$$\langle x - b, y - a\rangle \geq -\theta \|x - b\| \cdot \|y - a\|. \tag{25}$$

By the definition of T_λ, we have

$$\begin{aligned}\left\|x - x^+\right\|^2 &= \|x - b + y - a\|^2\\ &= \|x - b\|^2 + \|y - a\|^2 + 2\langle x - b, y - a\rangle\\ &\geq \|x - b\|^2 + \|y - a\|^2 - 2\theta\|x - b\| \cdot \|y - a\|\\ &\geq \left(1 - \theta^2\right)\|x - b\|^2 = \left(1 - \theta^2\right)\operatorname{dist}^2(x, B),\end{aligned} \tag{26}$$

where the first inequality follows from (25).

We next consider the two cases regarding $\operatorname{dist}(x, A)$.

Case 1. $\operatorname{dist}(x, A) \leq \left(\lambda + \sqrt{1-\theta^2}\right)\operatorname{dist}(x, B)$. Thanks to (26) we get

$$\left\|x - x^+\right\|^2 \geq \frac{1-\theta^2}{\left(\lambda + \sqrt{1-\theta^2}\right)^2}\operatorname{dist}^2(x, A). \tag{27}$$

Case 2. $\operatorname{dist}(x, A) > \left(\lambda + \sqrt{1-\theta^2}\right)\operatorname{dist}(x, B)$. By the triangle inequality and the construction of T_λ, we get

$$\begin{aligned}\left\|x - x^+\right\| &\geq \|x - a\| - \left\|a - x^+\right\| = \|x - a\| - \lambda\|x - b\|\\ &\geq \operatorname{dist}(x, A) - \lambda\operatorname{dist}(x, B) \geq \left(1 - \frac{\lambda}{\lambda + \sqrt{1-\theta^2}}\right)\operatorname{dist}(x, A).\end{aligned} \tag{28}$$

Since

$$\frac{1-\theta^2}{\left(\lambda + \sqrt{1-\theta^2}\right)^2} = \left(1 - \frac{\lambda}{\lambda + \sqrt{1-\theta^2}}\right)^2,$$

we always have from (27) and (28) that

$$\left\|x - x^+\right\|^2 \geq \frac{1-\theta^2}{\left(\lambda + \sqrt{1-\theta^2}\right)^2}\operatorname{dist}^2(x, A). \tag{29}$$

Combining (26), (29) and (22), we obtain

$$\begin{aligned}\left\|x - x^+\right\|^2 &\geq \frac{1-\theta^2}{\max\left\{1, \left(\lambda + \sqrt{1-\theta^2}\right)^2\right\}}\max\left\{\operatorname{dist}^2(x, A), \operatorname{dist}^2(x, B)\right\}\\ &\geq \frac{(1-\theta^2)(1-\theta)}{2\max\left\{1, \left(\lambda + \sqrt{1-\theta^2}\right)^2\right\}}\operatorname{dist}^2(x, A \cap B),\end{aligned}$$

which yields (23) as claimed. □

In the special case of $\lambda = 1$, Lemma 3 improve [14, Lemma 3.14] and [46, Lemma 4.2] where the result was proved for DR with an additional assumption on regularity of the sets.

The next result is the final preparation for our linear convergence result.

Lemma 4 *[46, Proposition 2.11] Let $T : \mathbb{E} \rightrightarrows \mathbb{E}$, $S \subset \mathbb{E}$ be closed and $\bar{x} \in S$. Suppose that there are $\delta > 0$ and $c \in [0, 1)$ such that for all $x \in \mathbb{B}_\delta(\bar{x})$, $x^+ \in Tx$ and $z \in P_S x$,*

$$\left\|x^+ - z\right\| \le c \left\|x - z\right\|. \tag{30}$$

Then all iterates $x_{k+1} \in Tx_k$ converge R-linearly to a point $\tilde{x} \in S \cap \mathbb{B}_\delta(\bar{x})$ provided that x_0 is sufficiently close to $\bar{x}$. In particular,

$$\|x_k - \tilde{x}\| \le \frac{\|x_0 - \bar{x}\| (1 + c)}{1 - c} c^k.$$

We are now ready to prove local linear convergence of T_λ which generalizes the corresponding results established in [14, 46] for DR.

Theorem 4 (linear convergence of T_λ) *In addition to Assumption 3, suppose that A and B are (ε, δ)-regular at $\bar{x}$ with $\tilde{\varepsilon} < \frac{(1+\lambda)\kappa^2}{2}$, where $\tilde{\varepsilon}$ and κ are given by (21) and (24), respectively. Then the iteration $x_{k+1} \in T_\lambda x_k$ converges R-linearly to a point in $A \cap B$ provided that x_0 is sufficiently close to $\bar{x}$.*

Proof Assumption 3 yields the existence of $\delta_1 > 0$ such that Lemma 3 holds true. In view of Proposition 4 and Remark 2, one can find a number $\delta_2 > 0$ such that T_λ is pointwise almost averaged on $\mathbb{B}_{\delta_2}(\bar{x})$ at every point $z \in A \cap B \cap \mathbb{B}_{\delta_2}(\bar{x})$ with violation $\tilde{\varepsilon}$ given by (21) and averaging constant $\frac{2}{3+\lambda}$. Let $\delta' := \min\{\delta_1, \delta_2\} > 0$.

Now we consider any $x \in \mathbb{B}_{\delta'}(\bar{x})$, $x^+ \in T_\lambda x$ and $z \in P_{A\cap B} x$. Again, by choosing a smaller δ if necessary, we can assume without loss of generality that $z \in \mathbb{B}_{\delta'}(\bar{x})$. Proposition 4 and Lemma 3 then respectively yield

$$\left\|x^+ - z\right\|^2 \le (1 + \tilde{\varepsilon}) \left\|x - z\right\|^2 - \frac{1+\lambda}{2} \left\|x - x^+\right\|^2, \tag{31}$$

$$\left\|x - x^+\right\|^2 \ge \kappa^2 \operatorname{dist}^2(x, A \cap B) = \kappa^2 \left\|x - z\right\|^2, \tag{32}$$

where κ is given by (24).

Substituting (32) into (31), we get

$$\left\|x^+ - z\right\|^2 \le \left(1 + \tilde{\varepsilon} - \frac{(1+\lambda)\kappa^2}{2}\right) \left\|x - z\right\|^2, \tag{33}$$

which yields condition (30) of Lemma 4 and the desired conclusion now follows from aforementioned lemma. □

5 Numerical experiments

Our goal in this section is to demonstrate a promising performance of T_λ in comparison with the RAAR algorithm for the example of sparse feasibility problem. We first recall the *sparse optimization problem* of

$$\min_{x\in\mathbb{R}^n} \|x\|_0 \qquad \text{subject to} \qquad Mx = b, \tag{34}$$

where $M \in \mathbb{R}^{m\times n}$ $(m < n)$ is a full rank matrix, b is a vector in $\mathbb{R}^m$, and $\|x\|_0$ is the number of nonzero entries of the vector x. The sparse optimization problem with complex variable is defined analogously by replacing $\mathbb{R}$ by $\mathbb{C}$ everywhere in the above model.

Many strategies for solving (34) have been studied. We refer the reader to the famous paper by Candès and Tao [10] for solving this problem by using convex relaxations. On the other hand, assuming to have a good guess on the sparsity of the solutions to (34), one can tackle this problem by solving the *sparse feasibility problem* [15] of finding

$$\bar{x} \in A_s \cap B, \tag{35}$$

where $A_s := \{x \in \mathbb{R}^n \mid \|x\|_0 \leq s\}$ and $B := \{x \in \mathbb{R}^n \mid Mx = b\}$.

It is worth mentioning that the initial guess s of the true sparsity is not numerically sensitive with respect to various projection methods, that is, for a relatively wide range of values of s above the true sparsity, projection algorithms perform very much in the same nature. Note also that the approach via sparse feasibility does not require convex relaxations of (34) and thus can avoid the likely expensive increase of dimensionality.

We run T_λ and RAAR to solve (35) and compare their numerical performances. By taking s smaller than the true sparsity, we can also compare their performances for inconsistent feasibility.

Since B is affine, there is the closed algebraic form for the projector P_B,

$$P_B x = x - M^\dagger(Mx - b) \quad \forall x \in \mathbb{R}^n,$$

where $M^\dagger := M^T(MM^T)^{-1}$ is the *Moore-Penrose inverse* of M. We have denoted M^T the transpose matrix of M and take into account of full rank of M. There is also a closed form for P_{A_s} [6]. For each $x \in \mathbb{R}^n$, let us denote $\mathcal{I}_x$ the set of all s-tubles of indices of s largest in absolute value entries of x. The set $\mathcal{I}_x$ can contains multiple s-tubles since A_s is intrinsically nonconvex. The projector P_{A_s} can be described by

$$P_{A_s} x = \left\{ z \in \mathbb{R}^n \mid \exists\, \mathcal{I} \in \mathcal{I}_x \;\text{ such that }\; z(k) = \begin{cases} x(k) & \text{if } k \in \mathcal{I}, \\ 0 & \text{else} \end{cases} \right\}.$$

For convenience, we recall the two algorithms in this case

$$\begin{aligned} RAAR_\beta &= \beta\,(P_{A_s}(2P_B - \mathrm{Id})) + (1 - 2\beta)P_B + \beta\,\mathrm{Id}, \\ T_\lambda &= P_{A_s}\,((1+\lambda)P_B - \lambda\,\mathrm{Id}) - \lambda(P_B - \mathrm{Id}). \end{aligned}$$

We now set up a toy example as in [10, 15] which involves an unknown true object $\bar{x} \in \mathbb{R}^{256^2}$ with $\|\bar{x}\|_0 = 328$ (the sparsity rate is .005). Let b be 1/8 of the measurements of $F(\bar{x})$, the *Fourier transform* of $\bar{x}$, with the *sample indices* denoted by $\mathcal{J}$. The *Poisson noise* was added when calculating the measurements of b. Note that since $\bar{x}$ is real, $F(\bar{x})$ is *conjugate symmetric*, we indeed have nearly a double number of measurements. In this setting, we have

$$B = \{x \in \mathbb{C}^{256^2} : F(x)(k) = b(k),\ \forall k \in \mathcal{J}\},$$

and the two prox operators, respectively, take the forms

$$P_{A_s}x = \left\{ z \in \mathbb{R}^n \mid \exists\, \mathcal{I} \in \mathcal{I}_x \text{ such that } z(k) = \begin{cases} \operatorname{Re}(x(k)) & \text{if } k \in \mathcal{I}, \\ 0 & \text{else} \end{cases} \right\},$$

$$P_B x = F^{-1}(\hat{x}), \quad \text{where } \hat{x}(k) = \begin{cases} b(k) & \text{if } k \in \mathcal{J}, \\ F(x)(k) & \text{else,} \end{cases}$$

where $\operatorname{Re}(x(k))$ denotes the real part of the complex number $x(k)$, and F^{-1} is the *inverse Fourier transform*.

The initial point was chosen randomly, and a warm up procedure with around ten DR iterates was made before running the two algorithms. The stopping criterion $\|x - x^+\| < 10^{-10}$ was used. We have used the *Matlab ProxToolbox* [38] to run this experiment. The parameters were chosen in such a way that the performance is seemingly the best for each algorithm. We chose $\beta = .65$ for RAAR and $\lambda = .45$ for T_λ in the consistent case corresponding to $s = 340$, and $\beta = .6$ for RAAR and $\lambda = .4$ for T_λ in the inconsistent case corresponding to $s = 310$.

The *change* of distances between two consecutive iterates is of interest. When linear convergence appears to be the case, it can yield useful information of the convergence rate. Under the assumption that the iterates will remain in the convergence area, one can obtain error bounds for the distance from the iterate to a solution. We also pay attention to the iterate *gaps* that in a sense measure the infeasibility at the iterates. If we think feasibility as the problem of minimizing the function that is the sum of the square of the distance functions to the sets, then iterate gaps are the values of that function evaluated at the iterates. For the two algorithms under consideration, the iterates are themselves not informative but their shadows, by which we mean the projections of the iterates on one of the sets. Hence, the iterate gaps are calculated for the iterate shadows instead of the iterates themselves.

Figure 1 summarizes the performances of the two algorithms for both consistent and inconsistent sparse feasibility. We first emphasize that the algorithms appear to be convergent in both cases of feasibility. For the consistent case, T_λ appears to perform better than RAAR in both the iterate changes and gaps. Also, the CPU time for T_λ is around 10% less than that for RAAR. For the inconsistent case, we have a similar observation except that the iterate gaps for RAAR are slightly better (smaller) than those for T_λ. Extensive



Fig. 1 Performances of RAAR and T_λ for sparse feasibility problem: iterate changes in consistent case (top-left), iterate gaps in consistent case (top-right), iterate changes in inconsistent case (bottom-left) and iterate gaps in consistent case (bottom-right).

numerical experiments in imaging illustrating the empirical performance of T_λ will be the tasks for future work.

Acknowledgements The author would like to thank Prof. Dr. Russell Luke and Prof. Dr. Alexander Kruger for their encouragement and valuable suggestions during the preparation of this work.

References

1. Aspelmeier, T., Charitha, C., Luke, D.R.: Local linear convergence of the ADMM/Douglas-Rachford algorithms without strong convexity and application to statistical imaging. SIAM J. Imaging Science **9**(2), 842–868 (2016)
2. Attouch, H., Bolte, J., Redont, P., Soubeyran, A.: Proximal alternating minimization and projection methods for nonconvex problems: An approach based on the Kurdyka-Łojasiewicz inequality. Math. Oper. Res. **35**(2), 438–457 (2010)

3. Bauschke, H.H., Borwein, J.M.: On projection algorithms for solving convex feasibility problems. SIAM Rev. **38**(3), 367–426 (1996)
4. Bauschke, H.H., Combettes, P.L.: Convex Analysis and Monotone Operator Theory in Hilbert Spaces. Springer, New York (2011)
5. Bauschke, H.H., Luke, D.R., Phan, H.M., Wang, X.: Restricted normal cones and the method of alternating projections: Applications. Set-Valued Var. Anal. **21**, 475–501 (2013)
6. Bauschke, H.H., Luke, D.R., Phan, H.M., Wang, X.: Restricted normal cones and sparsity optimization with affine constraints. Found. Comput. Math. **14**, 63–83 (2014)
7. Bolte, J., Sabach, S., Teboulle, M.: Proximal alternating linearized minimization for nonconvex and nonsmooth problems. Math. Programming **146**(1-2), 459–494 (2014)
8. Borwein, J.M. and Tam, M.K.: The cyclic Douglas-Rachford method for inconsistent feasibility problems. J. Nonlinear Convex Anal. **16**(4), 537–584 (2015)
9. Bunt, L. N. H.: Bitdrage tot de theorie der konvekse puntverzamelingen. PhD thesis. University of Groningen, Amsterdam (1934)
10. Candés, E., Tao, T.: Decoding by linear programming. IEEE Trans. Inf. Theory **51**(12), 4203–4215 (2005)
11. Combettes, P. L., Pesquet, J.-C.: Proximal splitting methods in signal processing. In Fixed-Point Algorithms for Inverse Problems in Science and Engineering. Springer Optim. Appl. **49**, 185–212 (2011)
12. Dontchev, A.L., Rockafellar, R.T.: Implicit Functions and Solution Mapppings. Srpinger-Verlag, New York (2014)
13. Drusvyatskiy, D., Ioffe, A.D., Lewis, A.S.: Transversality and alternating projections for nonconvex sets. Found. Comput. Math. **15**(6), 1637–1651 (2015)
14. Hesse, R., Luke, D.R.: Nonconvex notions of regularity and convergence of fundamental algorithms for feasibility problems. SIAM J. Optim. **23**(4), 2397–2419 (2013)
15. Hesse, R., Luke, D.R., Neumann, P.: Alternating projections and Douglas-Rachford for sparse affine feasibility. IEEE Trans. Signal. Process. **62**(18), 4868–4881 (2014)
16. Ioffe, A.D.: Metric regularity and subdifferential calculus. Russian Mathematical Surveys **55**(3), 501–558 (2000)
17. Ioffe, A.D.: Regularity on a fixed set. SIAM J. Optim. **21**(4), 1345–1370 (2011)
18. Ioffe, A.D.: Nonlinear regularity models. Math. Programming **139**(1-2), 223–242 (2013)
19. Ioffe, A.D.: Metric regularity - a survey. Part I. Theory. J. Aust. Math. Soc. **101**(2), 188–243 (2016)
20. Khanh, Phan Q., Kruger, A.Y., Thao, Nguyen H.: An induction theorem and nonlinear regularity models. SIAM J. Optim. **25**(4), 2561–2588 (2015)
21. Klatte, D., Kummer, B.: Nonsmooth Equations in Optimization. Kluwer Academic Publishers, Dordrecht (2002)
22. Klatte, D., Kummer, B.: Optimization methods and stability of inclusions in Banach spaces. Math. Programming **117**(1-2), 305–330 (2009)
23. Kruger, A.Y.: Stationarity and regularity of set systems. Pac. J. Optim. **1**(1), 101–126 (2005)
24. Kruger, A.Y.: About regularity of collections of sets. Set-Valued Anal. **14**, 187–206 (2006)
25. Kruger, A.Y.: About stationarity and regularity in variational analysis. Taiwanese J. Math. **13**(6A), 1737–1785 (2009)
26. Kruger, A.Y.: Error bounds and metric subregularity. Optimization **64**(1), 49–79 (2015)
27. Kruger, A.Y., Luke, D.R., Thao, Nguyen H.: Set regularities and feasibility problems. Math. Programming B (2016). doi:10.1007/s10107-016-1039-x
28. Kruger, A.Y., Luke, D.R., Thao, Nguyen H.: About subtransversality of collections of sets. Set-Valued Var. Anal. (2017). doi:10.1007/s11228-017-0436-5
29. Kruger, A.Y., Thao, Nguyen H.: About uniform regularity of collections of sets. Serdica Math. J. **39**, 287–312 (2013)
30. Kruger, A.Y., Thao, Nguyen H.: About $[q]$-regularity properties of collections of sets. J. Math. Anal. Appl. **416**(2), 471–496 (2014)
31. Kruger, A.Y., Thao, Nguyen H.: Quantitative characterizations of regularity properties of collections of sets. J. Optim. Theory and Appl. **164**, 41–67 (2015)
32. Kruger, A.Y., Thao, Nguyen H.: Regularity of collections of sets and convergence of inexact alternating projections. J. Convex Anal. **23**(3), 823–847 (2016)

33. Lewis, A.S., Luke, D.R., Malick, J.: Local linear convergence of alternating and averaged projections. Found. Comput. Math. **9**(4), 485–513 (2009)
34. Lewis, A.S., Malick, J.: Alternating projections on manifolds. Math. Oper. Res. **33**, 216–234 (2008)
35. Li, G., Pong, T.K.: Douglas-Rachford splitting for nonconvex feasibility problems. Math. Programming **159**(1), 371–401 (2016)
36. Luke, D.R.: Relaxed averaged alternating reflections for diffraction imaging. Inverse Problems **21**, 37–50 (2005)
37. Luke, D.R.: Finding best approximation pairs relative to a convex and a prox-regular set in Hilbert space. SIAM J. Optim. **19**(2), 714–739 (2008)
38. Luke, D.R.: ProxToolbox. http://num.math.uni-goettingen.de/proxtoolbox (2017). Accessed August 2017
39. Luke, D.R., Thao, Nguyen H., Tam, M.K.: Quantitative convergence analysis of iterated expansive, set-valued mappings. Math. Oper. Res., to appear
40. Luke, D.R., Thao, Nguyen H., Teboulle, M.: Necessary conditions for linear convergence of Picard iterations and application to alternating projections. https://arxiv.org/pdf/1704.08926.pdf (2017)
41. Mordukhovich, B.S.: Variational Analysis and Generalized Differentiation. I: Basic Theory. Springer, Berlin (2006)
42. Moreau, J.-J.: Fonctions convexes duales et points proximaux dans un espace Hilbertien. Comptes Rendus de l'Académie des Sciences de Paris **255**, 2897–2899 (1962)
43. Noll, D., Rondepierre, A.: On local convergence of the method of alternating projections. Found. Comput. Math. **16**(2), 425–455 (2016)
44. Patrinos, P., Stella, L., Bemporad, A.: Douglas-Rachford splitting: Complexity estimates and accelerated variants. In 53rd IEEE Conference on Decision and Control, 4234–4239 (2014)
45. Penot, J.-P.: Calculus Without Derivatives. Springer, New York (2013)
46. Phan, H.M.: Linear convergence of the Douglas-Rachford method for two closed sets. Optimization **65**, 369–385 (2016)
47. Poliquin, R. A., Rockafellar, R.T., Thibault, L.: Local differentiability of distance functions. Trans. Amer. Math. Soc. **352**(11), 5231–5249 (2000)
48. Rockafellar, R.T., Wets, R.J.-B.: Variational Analysis. Grundlehren Math. Wiss. Springer- Verlag, Berlin (1998)