ABSTRACT CRYSTALS FOR QUANTUM BORCHERDS-BOZEC ALGEBRAS

ZHAOBING FAN, SEOK-JIN KANG, YOUNG ROCK KIM*, AND BOLUN TONG

Abstract. In this paper, we develop the theory of abstract crystals for quantum Borcherds-Bozec algebras. Our construction is different from the one given by Bozec. We further prove the crystal embedding theorem and provide a characterization of $B(\infty)$ and $B(\lambda)$ as its application, where $B(\infty)$ and $B(\lambda)$ are the crystals of the negative half part of the quantum Borcherds-Bozec algebra $U_q(\mathfrak{g})$ and its irreducible highest weight module $V(\lambda)$, respectively.

Contents

Introduction 1
1. The quantum Borcherds-Bozec algebras 4
2. Abstract crystals 10
3. Tensor product of crystals 17
4. Crystal embedding theorem 22
References 29

Introduction

The aim of this paper is to develop the theory of *abstract crystals* for quantum Borcherds-Bozec algebras. To begin with, let us quickly review the term of Borcherds-Bozec algebras.

2010 *Mathematics Subject Classification.* 17B37, 17B67, 16G20.

Key words and phrases. quantum Borcherds-Bozec algebra, abstract crystals, crystal embedding theorem.

[*] Corresponding author. All authors contribute equally.

The Cartan matrix of a semisimple Lie algebra is positive definite, where all of its diagonal entries are 2. In [Kac68, Moody68], Kac and Moody introduced a family of infinite dimensional Lie algebras by removing the condition of positive definitiveness, which are now called the *Kac-Moody algebras*. In [Bor88], Borcherds generalized the notion of Kac-Moody algebras by allowing the diagonal entries of the Cartan matrices to be non-positive. The Lie algebras and Cartan matrices thus obtained are called the *Borcherds algebras* and Borcherds-Cartan matrices, respectively. Moreover, the Borcherds algebras are generated by positive and negative simple root vectors. Associated to Borcherds-Cartan matrices, the *Borcherds-Bozec algebras* is a further generalization of Kac-Moody algebras, which are generated by higher degree positive and negative simple root vectors.

The *quantum Borcherds algebras* (or *quantum generalized Kac-Moody algebras*) were introduced by Kang [K95], as a quantum deformation of the universal enveloping algebras of Borcherds algebras. The *quantum Borcherds-Bozec algebras* were introduced by Bozec in his research of perverse sheaves theory for quivers with loops [B15, B16, BSV16]. They are a further generalization of quantum Borcherds algebras. More precisely, for each imaginary index $i \in I^{\mathrm{im}}$, i.e., those $i \in I$ satisfying $a_{ii} \leq 0$ in the Cartan matrix, there are infinitely many generators e_{il}, f_{il} $(l \in \mathbb{Z}_{>0})$, whose degrees are l multiples of α_i and $-\alpha_i$. The commutation relations between them are rather complicated (cf. [FKKT20]). But, thanks to Bozec, there exists a set of primitive generators in quantum Borcherds-Bozec algebras with better properties and simpler commutation relations. This allows us to generalize the theories of quantum groups to quantum Borcherds-Bozec algebras.

The canonical basis theory was first introduced by Lusztig in the ADE case in [L90], arising from his geometric construction of the negative part of the quantum groups, and it has been generalized to general cases by Lusztig [L91, L10]. Meanwhile, Kashiwara constructed crystal base and global base for quantum groups associated with symmetrizable Kac-Moody algebras in an algebraic way [Kas90, Kas91]. In [GL93], Grojnowski and Lusztig proved that the global crystal base coincides with the canonical base introduced by Lusztig in [L90]. Since then, the crystal basis theory has become one of the most central themes in combinatorial and geometric representation theory of quantum groups. In [JKK05], Jeong, Kang and Kashiwara developed the crystal basis theory for quantum

Borcherds algebras. Later on, together with Shin, they further introduced the notion of abstract crystals for quantum Borcherds algebras and investigated their fundamental properties [JKKS07].

In [B16], Bozec studied the crystal basis theory for quantum Borcherds-Bozec algebra $U_q(\mathfrak{g})$. He defined the crystal basis $(L(\infty), B(\infty))$ for $U_q^-(\mathfrak{g})$ and $(L(\lambda), B(\lambda))$ for $V(\lambda)$, respectively, where $V(\lambda)$ is the irreducible highest weight $U_q(\mathfrak{g})$-module with a dominant integral highest weight λ. He also constructed *generalized crystals* and their geometric realization for $U_q^-(\mathfrak{g})$ and $V(\lambda)$ based on the theory of perverse sheaves on Lusztig's and Nakajima's quiver varieties associated with quivers with loops (cf. [KS97, S02, KKS09, KKS12]).

In this paper, we develop the theory of abstract crystals for quantum Borcherds-Bozec algebras. Our construction is different from the one given in [B16], where Bozec followed the framework of [JKK05]. Instead, we take the approach given in [JKKS07], which is simpler and more natural. We provide the abstract crystal structure on $B(\infty)$ and $B(\lambda)$ following our definition. Moreover, we define the tensor products of abstract crystals which are shown to be abstract crystals. We also show that the associativity law holds for the tensor products of abstract crystals.

Furthermore, we prove the *crystal embedding theorem*, which yields a procedure to determine the structure of the crystal $B(\infty)$ in term of elementary crystals. We provide a characterization of $B(\infty)$ and $B(\lambda)$ as an application of the crystal embedding theorem. The characterization of $B(\lambda)$ in the current paper is different from the one given by Bozec in [B16], where the method of Joseph in [J95] was employed.

Our theory of abstract crystals can be applied to various general settings. For example, Bozec's geometric construction of the crystals $B(\infty)$ and $B(\lambda)$ can be reproduced through our approach. Our construction of $B(\infty)$ and $B(\lambda)$ can also be used directly to the theory of (dual) perfect bases for quantum Borcherds-Bozec algebras. Moreover, our theory will play a crucial role in the categorification of quantum Borcherds-Bozec algebras and their highest weight modules via Khovanov-Lauda-Rouquier algebras and their cyclotomic quotients. Currently, we are working on these problems as well as other applications. We believe that our work on abstract crystals will provide a strong algebraic foundation for a wide variety of interesting developments in combinatorial and geometric representation theory of quantum Borcherds-Bozec algebras.

This paper is organized as follows. In Section 1, we review some of the basic facts on quantum Borcherds-Bozec algebras. Under the assumption (1.1), the presentation of quantum Borcherds-Bozec algebras becomes simpler. In Section 2, we define the notion of abstract crystals and investigate their fundamental properties. In Section 3, We define the tensor products of abstract crystals having all the desired properties. In Section 4, we prove the crystal embedding theorem. As an application, we provide a characterization of $B(\infty)$ and $B(\lambda)$.

Acknowledgements.

Z. Fan was partially supported by the NSF of China grant 11671108, the NSF of Heilongjiang Province grant JQ2020A001, and the Fundamental Research Funds for the central universities. S.-J. Kang was supported by Hankuk University of Foreign Studies Research Fund. Y. R. Kim was supported by the Basic Science Research Program of the NRF (Korea) under grant No. 2015R1D1A1A01059643. S.-J. Kang would like to express his sincere gratitude to Harbin Engineering University for their hospitality during his visit in July and November, 2019.

1. The quantum Borcherds-Bozec algebras

Let I be a finite or countably infinite index set. An integer-valued matrix $A = (a_{ij})_{i,j \in I}$ is called an *even symmetrizable Borcherds-Cartan matrix* if it satisfies the following conditions:

(i) $a_{ii} = 2, 0, -2, -4, \ldots,$
(ii) $a_{ij} \in \mathbb{Z}_{\le 0}$ for $i \neq j$,
(iii) there is a diagonal matrix $D = \operatorname{diag}(r_i \in \mathbb{Z}_{>0} \mid i \in I)$ such that DA is symmetric.

Let $I^{\mathrm{re}} := \{i \in I \mid a_{ii} = 2\}$, $I^{\mathrm{im}} := \{i \in I \mid a_{ii} \leq 0\}$, and $I^{\mathrm{iso}} := \{i \in I \mid a_{ii} = 0\}$. The elements of I^{re} (resp. I^{im}, I^{iso}) are called *real indices* (resp. *imaginary indices*, *isotropic indices*).

A *Borcherds-Cartan datum* consists of

(a) an even symmtrizable Borcherds-Cartan matrix,
(b) a free abelian group P, the *weight lattice*,
(c) $\Pi = \{\alpha_i \in P \mid i \in I\}$, the set of *simple roots*,
(d) $P^\vee := Hom_{\mathbb{Z}}(P, \mathbb{Z})$, the *dual weight lattice*,
(e) $\Pi^\vee = \{h_i \mid i \in I\}$, the set of *simple coroots*

satisfying the following conditions:

(i) $\langle h_i, \alpha_j \rangle = a_{ij}$ for all $i, j \in I$,
(ii) Π is linearly independent over $\mathbb{Z}$,
(iii) for each $i \in I$, there is an element $\Lambda_i \in P$, called the *fundamental weights*, such that

$$\langle h_j, \Lambda_i \rangle = \delta_{ij} \quad \text{for all } j \in I.$$

We denote by

$$P^+ := \{\lambda \in P \mid \langle h_i, \lambda \rangle \geq 0 \ \text{ for all } i \in I\},$$

the set of *dominant integral weights*. The free abelian group $Q := \bigoplus_{i \in I} \mathbb{Z}\alpha_i$ is called the *root lattice*. Set $Q^+ = \sum_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i$ and $Q^- = -Q^+$. For $\beta = \sum k_i \alpha_i \in Q^+$, we define its *height* to be $\mathrm{ht}(\beta) := \sum k_i$.

Set $\mathfrak{h} = \mathbb{Q} \otimes_{\mathbb{Z}} P^\vee$, the *Cartan subalgebra*. Since A is symmetrizable and Π is linearly independent, there is a non-degenerate symmetric bilinear form $(\ , \ )$ on $\mathfrak{h}^*$ satisfying

$$(\alpha_i, \lambda) = r_i \langle h_i, \lambda \rangle \ \text{ for all } i \in I, \ \lambda \in \mathfrak{h}^*.$$

For $i \in I^{\mathrm{re}}$, we define the *simple reflection* $\omega_i \in GL(\mathfrak{h}^*)$ by

$$\omega_i(\lambda) = \lambda - \langle h_i, \lambda \rangle \alpha_i \ \text{ for } \lambda \in \mathfrak{h}^*.$$

The subgroup W of $GL(\mathfrak{h}^*)$ generated by ω_i ($i \in I^{\mathrm{re}}$) is called the *Weyl group*. One can easily verify that the symmetric bilinear form $(\ , \ )$ is W-invariant.

Let $I^\infty := (I^{\text{re}} \times \{1\}) \cup (I^{\text{im}} \times \mathbb{Z}_{>0})$. For simplicity, we shall often write i instead of $(i,1)$ for $i \in I^{\text{re}}$. Let q be an indeterminate and set

$$q_i = q^{r_i} \text{ and } q_{(i)} = q^{\frac{(\alpha_i,\alpha_i)}{2}}.$$

Note that $q_i = q_{(i)}$ if $i \in I^{\text{re}}$. For each $i \in I^{\text{re}}$ and $n \in \mathbb{Z}_{\geq 0}$, we define

$$[n]_i = \frac{q_i^n - q_i^{-n}}{q_i - q_i^{-1}}, \quad [n]_i! = \prod_{k=1}^{n} [k]_i \quad \text{and} \quad \begin{bmatrix} n \\ k \end{bmatrix}_i = \frac{[n]_i!}{[k]_i![n-k]_i!}.$$

Let $\mathscr{F}$ be the free associative algebra over $\mathbb{Q}(q)$ generated by the symbols f_{il} for $(i,l) \in I^\infty$. By setting $\deg f_{il} = -l\alpha_i$, $\mathscr{F}$ becomes a Q^--graded algebra. For any homogeneous element u in $\mathscr{F}$, we denote by $|u|$ the degree of u, and for any $A \subseteq Q^-$, we set $\mathscr{F}_A = \{x \in \mathscr{F} \mid |x| \in A\}$.

We define a *twisted* multiplication on $\mathscr{F} \otimes \mathscr{F}$ by

$$(x_1 \otimes x_2)(y_1 \otimes y_2) = q^{-(|x_2|,|y_1|)} x_1 y_1 \otimes x_2 y_2,$$

and a comultiplication $\delta \colon \mathscr{F} \to \mathscr{F} \otimes \mathscr{F}$ by

$$\delta(f_{il}) = \sum_{m+n=l} q_{(i)}^{-mn} f_{im} \otimes f_{in} \quad \text{for } (i,l) \in I^\infty.$$

Here $f_{i0} = 1$ and $f_{il} = 0$ for $l < 0$ by convention.

Proposition 1.1. [B15, B16] For any family $\nu = (\nu_{il})_{(i,l)\in I^\infty}$ of non-zero elements in $\mathbb{Q}(q)$, there exists a symmetric bilinear form $(\ ,\ )_L \colon \mathscr{F} \times \mathscr{F} \to \mathbb{Q}(q)$ such that

(a) $(x,y)_L = 0$ if $|x| \neq |y|$.
(b) $(1,1)_L = 1$.
(c) $(f_{il}, f_{il})_L = \nu_{il}$ for all $(i,l) \in I^\infty$.
(d) $(x, yz)_L = (\delta(x), y \otimes z)_L$ for all $x,y,z \in \mathscr{F}$,

where $(x_1 \otimes x_2, y_1 \otimes y_2)_L = (x_1,y_1)_L(x_2,y_2)_L$ for any $x_1,x_2,y_1,y_2 \in \mathscr{F}$.

From now on, we will assume that

$$\nu_{il} \in 1 + q\mathbb{Z}_{\geq 0}[[q]] \quad \text{for all } (i,l) \in I^\infty. \tag{1.1}$$

Under the assumption (1.1), the radical of the bilinear form $(\ ,\ )_L$ is generated by

$$\begin{aligned} &\sum_{k=0}^{1-la_{ij}} (-1)^k \begin{bmatrix} 1-la_{ij} \\ k \end{bmatrix}_i f_i^{1-la_{ij}-k} f_{jl} f_i^k = 0 \ \text{ for } i \in I^{\mathrm{re}},\, i \neq (j,l), \\ &f_{ik}f_{jl} - f_{jl}f_{ik} = 0 \ \text{ for } a_{ij} = 0. \end{aligned} \tag{1.2}$$

Let $U^{\leq 0}$ be the associative algebra over $\mathbb{Q}(q)$ with $\mathbf{1}$ generated by f_{il} $((i,l) \in I^{\infty})$ and q^h $(h \in P^{\vee})$ with the defining relations (1.2) and

$$\begin{aligned} &q^0 = \mathbf{1}, \quad q^h q^{h'} = q^{h+h'} \ \text{ for } h, h' \in P^{\vee}, \\ &q^h f_{jl} q^{-h} = q^{-l\langle h, \alpha_j \rangle} f_{jl} \ \text{ for } h \in P^{\vee}, (j,l) \in I^{\infty}. \end{aligned} \tag{1.3}$$

The comultiplication δ induces a well-defined comultiplication $\Delta\colon U^{\leq 0} \to U^{\leq 0} \otimes U^{\leq 0}$ given by

$$\begin{aligned} &\Delta(q^h) = q^h \otimes q^h, \\ &\Delta(f_{il}) = \sum_{m+n=l} q_{(i)}^{-mn} f_{im} K_i^n \otimes f_{in}. \end{aligned} \tag{1.4}$$

By the Drinfeld double process, we have the following definition for quantum Borcherds-Bozec algebra.

Definition 1.2. The *quantum Borcherds-Bozec algebra* $U_q(\mathfrak{g})$ associated with a Borcherds-Cartan datum $(A, P, \Pi, P^{\vee}, \Pi^{\vee})$ is the associative algebra over $\mathbb{Q}(q)$ with $\mathbf{1}$ generated by the elements q^h $(h \in P^\vee)$ and e_{il}, f_{il} $((i,l) \in I^\infty)$ with the following defining relations:

$$
\begin{aligned}
& q^0 = \mathbf{1}, \quad q^h q^{h'} = q^{h+h'} \quad \text{for } h, h' \in P^\vee, \\
& q^h e_{jl} q^{-h} = q^{l\langle h,\alpha_j\rangle} e_{jl}, \quad q^h f_{jl} q^{-h} = q^{-l\langle h,\alpha_j\rangle} f_{jl} \quad \text{for } h \in P^\vee, (j,l) \in I^\infty, \\
& e_{ik} f_{jl} - f_{jl} e_{ik} = 0 \quad \text{for } i \neq j, \\
& \sum_{\substack{m+n=k \\ n+s=l}} q_{(i)}^{n(m-s)} \nu_{in} e_{is} f_{im} K_i^{-n} = \sum_{\substack{m+n=k \\ n+s=l}} q_{(i)}^{-n(m-s)} \nu_{in} f_{im} e_{is} K_i^{n} \quad \text{for } (i,l), (i,k) \in I^\infty, \\
& \sum_{k=0}^{1-la_{ij}} (-1)^k \begin{bmatrix} 1-la_{ij} \\ k \end{bmatrix}_i e_i^{1-la_{ij}-k} e_{jl} e_i^k = 0 \quad \text{for } i \in I^{\mathrm{re}},\ i \neq (j,l), \\
& \sum_{k=0}^{1-la_{ij}} (-1)^k \begin{bmatrix} 1-la_{ij} \\ k \end{bmatrix}_i f_i^{1-la_{ij}-k} f_{jl} f_i^k = 0 \quad \text{for } i \in I^{\mathrm{re}},\ i \neq (j,l), \\
& e_{ik} e_{jl} - e_{jl} e_{ik} = f_{ik} f_{jl} - f_{jl} f_{ik} = 0 \quad \text{for } a_{ij} = 0.
\end{aligned}
\tag{1.5}
$$

Here $K_i = q_i^{h_i}$ $(i \in I)$. We extend the grading by setting $|q^h| = 0$ and $|e_{il}| = l\alpha_i$.

Remark 1.3. Our presentation of $U_q(\mathfrak{g})$ is simpler than the ones in [B15, B16, BSV16] thanks to the explicit commutation relations among the generators given in [FKKT20, Appendix A].

The comultiplication on $U^{\leq 0}$ can be extended to the comultiplication $\Delta \colon U_q(\mathfrak{g}) \to U_q(\mathfrak{g}) \otimes U_q(\mathfrak{g})$ given by

$$
\begin{aligned}
\Delta(q^h) &= q^h \otimes q^h, \\
\Delta(e_{il}) &= \sum_{m+n=l} q_{(i)}^{mn} e_{im} \otimes K_i^{-m} e_{in}, \\
\Delta(f_{il}) &= \sum_{m+n=l} q_{(i)}^{-mn} f_{im} K_i^{n} \otimes f_{in}.
\end{aligned}
\tag{1.6}
$$

Let U^+ (resp. U^-) be the subalgebra of $U_q(\mathfrak{g})$ generated by e_{il} (resp. f_{il}) for $(i,l) \in I^\infty$, and U^0 the subalgebra of $U_q(\mathfrak{g})$ generated by q^h for $h \in P^\vee$. Then the quantum Borcherds-Bozec algebra $U_q(\mathfrak{g})$ has the following *triangular decomposition*

$$U_q(\mathfrak{g}) \cong U^- \otimes U^0 \otimes U^+.$$

Let $\omega\colon U_q(\mathfrak{g}) \to U_q(\mathfrak{g})$ be the involution defined by

$$\omega(q^h) = q^{-h},\ \omega(e_{il}) = f_{il},\ \omega(f_{il}) = e_{il} \ \text{ for } h \in P^\vee,\ (i,l) \in I^\infty,$$

and define a $\mathbb{Q}$-algebra involution on $U_q(\mathfrak{g})$, called the *bar involution*, by

$$\overline{e}_{il} = e_{il},\ \overline{f}_{il} = f_{il},\ \overline{q}^h = q^{-h},\ \overline{q} = q^{-1} \ \text{ for } h \in P^\vee,\ (i,l) \in I^\infty.$$

The following proposition provides a set of *primitive generators* for $U_q(\mathfrak{g})$.

Proposition 1.4. [B15, B16] For any $i \in I^{\text{im}}$ and $l \geq 1$, there exist unique elements $t_{il} \in U^-_{-l\alpha_i}$ and $s_{il} = \omega(t_{il})$ such that

(1) $\mathbb{Q}(q)\,\langle f_{il} \mid l \geq 1\rangle = \mathbb{Q}(q)\,\langle t_{il} \mid l \geq 1\rangle$ and $\mathbb{Q}(q)\,\langle e_{il} \mid l \geq 1\rangle = \mathbb{Q}(q)\,\langle s_{il} \mid l \geq 1\rangle$,

(2) $(t_{il}, z)_L = 0$ for all $z \in \mathbb{Q}(q)\,\langle f_{i1}, \cdots, f_{i,l-1}\rangle$,
$(s_{il}, z)_L = 0$ for all $z \in \mathbb{Q}(q)\,\langle e_{i1}, \cdots, e_{i,l-1}\rangle$,

(3) $t_{il} - f_{il} \in \mathbb{Q}(q)\,\langle f_{ik} \mid k < l\rangle$ and $s_{il} - e_{il} \in \mathbb{Q}(q)\,\langle e_{ik} \mid k < l\rangle$,

(4) $\overline{t}_{il} = t_{il}$, $\overline{s}_{il} = s_{il}$,

(5) $\delta(t_{il}) = t_{il} \otimes 1 + 1 \otimes t_{il}$, $\delta(s_{il}) = s_{il} \otimes 1 + 1 \otimes s_{il}$,

(6) $\Delta(t_{il}) = t_{il} \otimes 1 + K_i^l \otimes t_{il}$, $\Delta(s_{il}) = s_{il} \otimes K_i^{-l} + 1 \otimes s_{il}$,

Set $\tau_{il} = (t_{il}, t_{il})_L = (s_{il}, s_{il})_L$. We have the following commutation relations in $U_q(\mathfrak{g})$

$$s_{il}\, t_{jk} - t_{jk}\, s_{il} = \delta_{ij}\delta_{lk}\tau_{il}(K_i^l - K_i^{-l}). \tag{1.7}$$

For $i \in I^{\text{im}} \backslash I^{\text{iso}}$ (resp. $i \in I^{\text{iso}}$), denote by $\mathcal{C}_{i,l}$ the set of compositions (resp. partitions) of l, and set $\mathcal{C}_i = \bigsqcup_{l \geq 0} \mathcal{C}_{i,l}$. For $i \in I^{\text{re}}$, we put $\mathcal{C}_{i,l} = \{(l)\}$.

If $i \in I^{\text{im}}$ and $\mathbf{c} = (c_1, \cdots, c_r) \in \mathcal{C}_{i,l}$, we write $|\mathbf{c}| = l$ and define

$$t_{i,\mathbf{c}} = t_{ic_1} \cdots t_{ic_r}, \quad s_{i,\mathbf{c}} = s_{ic_1} \cdots s_{ic_r} \ \text{ and } \ \tau_{i,\mathbf{c}} = \tau_{ic_1} \cdots \tau_{ic_r}.$$

Note that $\{t_{i,\mathbf{c}} \mid \mathbf{c} \in \mathcal{C}_{i,l}\}$ forms a basis of $U^-_{-l\alpha_i}$.

For each $(i,l) \in I^\infty$, we define the linear map $e'_{il}\colon U^- \to U^-$ by

$$e'_{il}(1) = 0,\ e'_{il}(t_{jk}) = \delta_{ij}\delta_{lk} \text{ and } e'_{il}(xy) = e'_{il}(x)y + q^{l(|x|,\alpha_i)}xe'_{il}(y). \tag{1.8}$$

In [B16], Bozec used the operators e'_{il} to define the *Kashiwara operators* on U^-.

2. ABSTRACT CRYSTALS

In this section, we develop the theory of abstract crystals for quantum Borcherds-Bozec algebras. We would like to point out that we follow the outline given in [JKKS07], not in [JKK05].

Definition 2.1. Let $U_q(\mathfrak{g})$ be the quantum Borcherds-Bozec algebra associated with a given Borcherds-Cartan datum $(A, P, P^{\vee}, \Pi, \Pi^{\vee})$. An *abstract* $U_q(\mathfrak{g})$*-crystal* or simply a *crystal* is a set B together with the maps $\mathrm{wt}\colon B \to P$, $\widetilde{e}_{il}, \widetilde{f}_{il}\colon B \to B \sqcup \{0\}$ $((i,l) \in I^{\infty})$ and $\epsilon_i, \phi_i\colon B \to \mathbb{Z} \sqcup \{-\infty\}$ $(i \in I)$ satisfying the following conditions:

(i) $\mathrm{wt}(\widetilde{e}_{i,l}b) = \mathrm{wt}(b) + l\alpha_i$ if $\widetilde{e}_{il}b \neq 0$.

(ii) $\mathrm{wt}(\widetilde{f}_{i,l}b) = \mathrm{wt}(b) - l\alpha_i$ if $\widetilde{f}_{il}b \neq 0$.

(iii) For any $i \in I$ and $b \in B$, $\phi_i(b) = \epsilon_i(b) + \langle h_i, \mathrm{wt}(b) \rangle$.

(iv) For any $(i,l) \in I^{\infty}$ and $b, b' \in B$, $\widetilde{f}_{il}b = b'$ if and only if $b = \widetilde{e}_{il}b'$.

(v) For any $i \in I^{\mathrm{re}}$ and $b \in B$, we have
  (a) $\epsilon_i(\widetilde{e}_i b) = \epsilon_i(b) - 1$, $\phi_i(\widetilde{e}_i b) = \phi_i(b) + 1$ if $\widetilde{e}_i b \neq 0$,
  (b) $\epsilon_i(\widetilde{f}_i b) = \epsilon_i(b) + 1$, $\phi_i(\widetilde{f}_i b) = \phi_i(b) - 1$ if $\widetilde{f}_i b \neq 0$,
  where $\widetilde{e}_i = \widetilde{e}_{i1}$ and $\widetilde{f}_i = \widetilde{f}_{i1}$.

(vi) For any $i \in I^{\mathrm{im}}$, $l > 0$ and $b \in B$, we have
  (a) $\epsilon_i(\widetilde{e}_{il}b) = \epsilon_i(b)$, $\phi_i(\widetilde{e}_{il}b) = \phi_i(b) + l a_{ii}$ if $\widetilde{e}_{il}b \neq 0$,
  (b) $\epsilon_i(\widetilde{f}_{il}b) = \epsilon_i(b)$, $\phi_i(\widetilde{f}_{il}b) = \phi_i(b) - l a_{ii}$ if $\widetilde{f}_{il}b \neq 0$.

(vii) For any $(i,l) \in I^{\infty}$ and $b \in B$ such that $\phi_i(b) = -\infty$, we have $\widetilde{e}_{il}b = \widetilde{f}_{il}b = 0$.

Let B be an abstract crystal. For $b, b' \in B$ and $(i,l) \in I^{\infty}$, by the condition (iv), we have $\widetilde{f}_{il}b = b'$ if and only if $\widetilde{e}_{il}b' = b$. In this case,we draw a labelled arrow $b \xrightarrow{(i,l)} b'$. The directed graph thus obtained is called the *crystal graph* of B.

Definition 2.2. Let B_1 and B_2 be abstract crystals. A map $\psi\colon B_1 \to B_2$ is called a *morphism of crystals* or a *crystal morphism* if it satisfies the following conditions:

(i) for $b \in B_1$ and $i \in I$, we have
  $\mathrm{wt}(\psi(b)) = \mathrm{wt}(b)$, $\epsilon_i(\psi(b)) = \epsilon_i(b)$, $\phi_i(\psi(b)) = \phi_i(b)$,

(ii) for all $b \in B_1$, $(i,l) \in I^{\infty}$, if $\widetilde{f}_{il}b \in B_1$, then $\psi(\widetilde{f}_{il}b) = \widetilde{f}_{il}\psi(b)$.

Remark 2.3.

(a) If $b \in B_1$ and $\widetilde{e}_{il} b \in B_1$, one can deduce $\psi(\widetilde{e}_{il} b) = \widetilde{e}_{il}\psi(b)$.

(b) If a crystal morphism ψ is a bijection, then ψ^{-1} is also a crystal morphism. In particular, both ψ and ψ^{-1} commute with $\widetilde{e}_{il}$'s.

Definition 2.4. Let $\psi \colon B_1 \to B_2$ be a morphism of crystals.

(a) ψ is called a *strict morphism* if

$$\psi(\widetilde{e}_{il} b) = \widetilde{e}_{il}(\psi(b)), \quad \psi(\widetilde{f}_{il} b) = \widetilde{f}_{il}(\psi(b))$$

for all $b \in B_1$, $(i,l) \in I^\infty$. Here, we understand $\psi(0) = 0$.

(b) ψ is called a *crystal embedding* if the underlying map $\psi \colon B_1 \to B_2$ is injective. In this case, B_1 is called a *subcrystal* of B_2. If ψ is a strict embedding, B_1 is called a *full subcrystal* of B_2.

(c) ψ is called an *isomorphism* if it is a bijection.

In the following two examples, we recall the crystals $B(\infty)$ and $B(\lambda)$ ($\lambda \in P^+$) constructed in [B16].

Example 2.5.

Let $u \in U^-$ and $(i,l) \in I^\infty$.

If $i \in I^{\text{re}}$, in [Kas91], Kashiwara proved that u can be uniquely written as

$$u = \sum_{k \geq 0} f_i^{(k)} u_k,$$

where $f_i^{(k)} = f_i^k/[k]_i!$ and $e_i' u_k = 0$ for all $k \geq 0$. In this case, the *Kashiwara operators* are defined by

$$\widetilde{e}_i\, u = \sum_{k \geq 1} f_i^{(k-1)}\, u_k, \quad \widetilde{f}_i\, u = \sum_{k \geq 0} f_i^{(k+1)}\, u_k.$$

If $i \in I^{\text{im}} \setminus I^{\text{iso}}$, in [B16], Bozec showed that u can be written uniquely as

$$u = \sum_{\mathbf{c} \in \mathcal{C}_i} t_{i,\mathbf{c}} u_{\mathbf{c}},$$

where $\mathbf{c} = (c_1, \ldots, c_r)$ is a composition in $\mathcal{C}_i$ and $e'_{il}\, u_{\mathbf{c}} = 0$ for all $l \geq 1$. In this case, the Kashiwara operators are defined by

$$\widetilde{e}_{il}\, u = \sum_{\mathbf{c}:\, c_1 = l} t_{i,\mathbf{c}\setminus c_1}\, u_{\mathbf{c}}, \quad \widetilde{f}_{il}\, u = \sum_{\mathbf{c}\in\mathcal{C}_i} t_{i,(l,\mathbf{c})}\, u_{\mathbf{c}}.$$

If $i \in I^{\mathrm{iso}}$, u can be written uniquely as

$$u = \sum_{\mathbf{c}\in\mathcal{C}_i} t_{i,\mathbf{c}} u_{\mathbf{c}},$$

where $\mathbf{c} = (c_1, \ldots, c_r)$ is a partition in $\mathcal{C}_i$ and $e'_{il}\, u_{\mathbf{c}} = 0$ for all $l \geq 1$. In this case, the Kashiwara operators are defined by

$$\widetilde{e}_{il}\, u = \sum_{\mathbf{c}\in\mathcal{C}_i} \sqrt{\frac{m_l(\mathbf{c})}{l}}\, t_{i,\mathbf{c}\setminus l}\, u_{\mathbf{c}}, \quad \widetilde{f}_{il}\, u = \sum_{\mathbf{c}\in\mathcal{C}_i} \sqrt{\frac{l}{m_l(\mathbf{c})+1}}\, t_{i,\mathbf{c}\cup l}\, u_{\mathbf{c}},$$

where $m_l(\mathbf{c}) = \#\{k \mid c_k = l\}$, and $\mathbf{c} \cup l$ stands for the partition $(l, c_1, \cdots, c_r)$.

Remark 2.6. Note that the square roots appear in the above definition. So we need to consider an extension $\mathbb{F}$ of $\mathbb{Q}$ that contains all the necessary square roots in Example 2.5 and in Example 2.7. (See [B16, Remark 3.12].)

Let $\mathbb{A}_0 = \{f \in \mathbb{F}(q) \mid f \text{ is regular at } q = 0\}$ and let $L(\infty)$ be the $\mathbb{A}_0$-submodule of U^- generated by the elements of the form $\widetilde{f}_{i_1,l_1} \cdots \widetilde{f}_{i_r,l_r}\mathbf{1}$ for $r \geq 0$ and $(i_k, l_k) \in I^\infty$. Set

$$B(\infty) = \{\widetilde{f}_{i_1,l_1} \cdots \widetilde{f}_{i_r,l_r}\mathbf{1} \mod qL(\infty) \mid r \geq 0, (i_k, l_k) \in I^\infty\} \subseteq L(\infty)/qL(\infty).$$

For $b = \widetilde{f}_{i_1,l_1} \cdots \widetilde{f}_{i_r,l_r}\mathbf{1}$, we define

$$\begin{aligned} &\mathrm{wt}(b) = -(l_1\alpha_{i_1} + \cdots + l_r\alpha_{i_r}), \\ &\epsilon_i(b) = \begin{cases} \max\{k \geq 0 \mid \widetilde{e}_i^k b \neq 0\} & \text{for } i \in I^{\mathrm{re}}, \\ 0 & \text{for } i \in I^{\mathrm{im}}, \end{cases} \\ &\phi_i(b) = \epsilon_i(b) + \langle h_i, \mathrm{wt}(b)\rangle \quad \text{for any} \quad i \in I. \end{aligned} \tag{2.1}$$

Then $B(\infty)$ becomes an abstract crystal.

Note that $\mathbf{1}$ is the only element in $B(\infty)$ of weight 0 and is annihilated by all $\widetilde{e}_{il}$ for $(i,l) \in I^{\infty}$. Moreover, $B(\infty)$ is connected (as a directed graph).

Example 2.7. For a dominant integral weight $\lambda \in P^{+}$, let $V(\lambda) = U_q(\mathfrak{g})v_\lambda$ be the irreducible highest weight module with highest weight λ and highest weight vector v_λ.

Let $v \in V(\lambda)$. If $i \in I^{\mathrm{re}}$, v can be uniquely written as

$$v = \sum_{k \geq 0} f_i^{(k)} v_k,$$

where $f_i^{(k)} = f_i^k/[k]_i!$ and $e_i v_k = 0$ for all $k \geq 0$, and if $i \in I^{\mathrm{im}}$, v can be written uniquely as

$$v = \sum_{\mathbf{c} \in \mathcal{C}_i} t_{i,\mathbf{c}} v_{\mathbf{c}},$$

where $\mathbf{c} = (c_1, \dots, c_r)$ is a composition or a partition in $\mathcal{C}_i$ and $e_{il}\, v_{\mathbf{c}} = 0$ for all $l \geq 1$. The Kashiwara operators are defined in a similar manner as in Example 2.5.

Let $L(\lambda)$ be the $\mathbb{A}_0$-submodule of $V(\lambda)$ generated by the elements of the form $\widetilde{f}_{i_1,l_1} \cdots \widetilde{f}_{i_r,l_r} v_\lambda$ for $r \geq 0$ and $(i_k, l_k) \in I^{\infty}$. Set

$$B(\lambda) = \{\widetilde{f}_{i_1,l_1} \cdots \widetilde{f}_{i_r,l_r} v_\lambda \mod qL(\lambda) \mid r \geq 0, (i_k, l_k) \in I^{\infty}\} \setminus \{0\} \subseteq L(\lambda)/qL(\lambda).$$

For $b = \widetilde{f}_{i_1,l_1} \cdots \widetilde{f}_{i_r,l_r} v_\lambda$, we define

$$\begin{aligned} &\mathrm{wt}(b) = \lambda - (l_1 \alpha_{i_1} + \cdots + l_r \alpha_{i_r}), \\ &\epsilon_i(b) = \begin{cases} \max\{k \geq 0 \mid \widetilde{e}_i^k b \neq 0\} & \text{for } i \in I^{\mathrm{re}}, \\ 0 & \text{for } i \in I^{\mathrm{im}}, \end{cases} \\ &\phi_i(b) = \epsilon_i(b) + \langle h_i, \mathrm{wt}(b) \rangle \quad \text{for any} \quad i \in I. \end{aligned} \tag{2.2}$$

Then $B(\lambda)$ becomes an abstract crystal. Note that v_λ is the only element in $B(\lambda)$ of weight λ and is annihilated by all $\widetilde{e}_{il}$ for $(i,l) \in I^{\infty}$. Moreover, $B(\lambda)$ is connected.

Example 2.8.

(a) For $\lambda \in P$, set $T_\lambda = \{t_\lambda\}$ and define

$$\mathrm{wt}(t_\lambda) = \lambda, \quad \epsilon_i(t_\lambda) = \phi_i(t_\lambda) = -\infty, \quad \widetilde{e}_{il}(t_\lambda) = \widetilde{f}_{il}(t_\lambda) = 0$$

for all $i \in I$ and $l \geq 1$. Then T_λ is an abstract crystal.

(b) Let $C = \{c\}$ and define

$$\mathrm{wt}(c) = 0, \quad \epsilon_i(c) = \phi_i(c) = 0, \quad \widetilde{e}_{il}(c) = \widetilde{f}_{il}(c) = 0$$

for all $i \in I$ and $l \geq 1$. Then C is an abstract crystal which is isomorphic to $B(0)$.

As was proved in [JKK05], we have the following proposition which describes the relation between the crystals $B(\infty)$ and $B(\lambda)$.

Proposition 2.9. For every $\lambda \in P^+$, there exists an injective map $\pi_\lambda \colon B(\lambda) \to B(\infty)$ such that

(i) $\pi_\lambda(v_\lambda) = 1$,
(ii) for all $(i,l) \in I^\infty$ and $b \in B(\lambda)$ with $\widetilde{f}_{il}(b) \neq 0$, $\pi_\lambda \widetilde{f}_{il}(b) = \widetilde{f}_{il}\, \pi_\lambda(b)$,
(iii) for all $(i,l) \in I^\infty$ and $b \in B(\lambda)$, $\pi_\lambda \widetilde{e}_{il}(b) = \widetilde{e}_{il}\, \pi_\lambda(b)$,
(iv) for all $i \in I$ and $b \in B(\lambda)$, $\mathrm{wt}(\pi_\lambda(b)) = \mathrm{wt}(b) - \lambda$, $\epsilon_i(\pi_\lambda(b)) = \epsilon_i(b)$.

Remark 2.10. The map π_λ is *not* a crystal embedding because it does not preserve the functions wt and ϕ_i $(i \in I)$.

The following example provides a very important class of abstract crystals B_i $(i \in I)$ called the *elementary crystals*.

Example 2.11.

(a) If $i \in I^{\mathrm{re}}$, let $B_i = \{(l) \mid l \geq 0\}$ and define

$$\begin{aligned}
&\mathrm{wt}((l)) = -l\alpha_i,\\
&\epsilon_i((l)) = l, \quad \phi_i((l)) = -l, \quad \epsilon_j((l)) = \phi_j((l)) = -\infty \ \text{ for } j \neq i,\\
&\widetilde{e}_i((l)) = (l-1), \quad \widetilde{f}_i((l)) = (l+1),\\
&\widetilde{e}_{jk}((l)) = \widetilde{f}_{jk}((l)) = 0 \ \text{ for } j \neq i.
\end{aligned}$$

Then B_i is an abstract crystal. The crystal graph of B_i is given in Figure 1.

Figure 1

$$(0) \xrightarrow{i} (1) \xrightarrow{i} (2) \xrightarrow{i} \cdots$$

We understand $(l) = 0$ for $l < 0$.

(b) For $i \in I^{\mathrm{im}} \setminus I^{\mathrm{iso}}$, let $B_i = \{\mathbf{c} = (c_1, \dots, c_r) \mid r \geq 0,\ \mathbf{c} \text{ is a composition}\}$ and define

$$\begin{aligned}
&\mathrm{wt}(\mathbf{c}) = -|\mathbf{c}|\alpha_i, \quad \epsilon_i(\mathbf{c}) = 0, \quad \phi_i(\mathbf{c}) = -|\mathbf{c}|a_{ii}, \\
&\epsilon_j(\mathbf{c}) = \phi_j(\mathbf{c}) = -\infty \quad \text{for } j \neq i, \\
&\widetilde{e}_{il}(\mathbf{c}) = \begin{cases} \mathbf{c} \setminus l = (c_2, \dots, c_r) & \text{if } c_1 = l. \\ 0 & \text{otherwise,} \end{cases} \\
&\widetilde{f}_{il}(\mathbf{c}) = (l, \mathbf{c}) = (l, c_1, \dots, c_r), \\
&\widetilde{e}_{jk}(\mathbf{c}) = \widetilde{f}_{jk}(\mathbf{c}) = 0 \quad \text{for } j \neq i.
\end{aligned}$$

Then B_i is an abstract crystal. The crystal graph of B_i is given in Figure 2.

Figure 2



(c) For $i \in I^{\mathrm{iso}}$, let $B_i = \{\mathbf{c} = (c_1, \dots, c_r) \mid r \geq 0,\ \mathbf{c} \text{ is a partition}\}$ and define

$$\begin{aligned}
&\mathrm{wt}(\mathbf{c}) = -|\mathbf{c}|\alpha_i, \quad \epsilon_i(\mathbf{c}) = \phi_i(\mathbf{c}) = 0,\\
&\epsilon_j(\mathbf{c}) = \phi_j(\mathbf{c}) = -\infty \quad \text{for } j \neq i,\\
&\widetilde{e}_{il}(\mathbf{c}) = \begin{cases} \mathbf{c} \setminus \{l\} & \text{if } l \text{ is a part of } \mathbf{c},\\ 0 & \text{otherwise,} \end{cases}\\
&\widetilde{f}_{il}(\mathbf{c}) = \mathbf{c} \cup l = (l, c_1, \dots, c_r),\\
&\widetilde{e}_{jk}(\mathbf{c}) = \widetilde{f}_{jk}(\mathbf{c}) = 0 \quad \text{for } j \neq i.
\end{aligned}$$

Then B_i is an abstract crystal. A part of the crystal graph of B_i is given in Figure 3.

Figure 3



We will write $\mathbf{c}_i$ for $\mathbf{c} \in B_i$ when we would like to emphasize the index i.

Definition 2.12. An abstract crystal B is *normal* if

$$\epsilon_i(b) = 0, \quad \phi_i(b) \geq 0 \quad \text{for all } i \in I^{\mathrm{im}}, b \in B.$$

Remark 2.13. The crystals $B(\infty)$, $B(\lambda)$ $(\lambda \in P^+)$ and C are normal, while T_λ $(\lambda \in P)$ and B_i $(i \in I)$ are not normal.

3. Tensor product of crystals

Let B_1, B_2 be abstract crystals and let $B_1 \otimes B_2 = \{b_1 \otimes b_2 \mid b_1 \in B_1, b_2 \in B_2\}$ (as a set). Define the maps wt, ϵ_i, ϕ_i $(i \in I)$, $\widetilde{e}_{il}$, $\widetilde{f}_{il}$ $((i,l) \in I^{\infty})$ as follows.

$$\begin{aligned} &\mathrm{wt}(b_1 \otimes b_2) = \mathrm{wt}(b_1) + \mathrm{wt}(b_2), \\ &\epsilon_i(b_1 \otimes b_2) = \max(\epsilon_i(b_1), \epsilon_i(b_2) - \langle h_i, \mathrm{wt}(b_1)\rangle), \\ &\phi_i(b_1 \otimes b_2) = \max(\phi_i(b_1) + \langle h_i, \mathrm{wt}(b_2)\rangle, \phi_i(b_2)), \end{aligned} \tag{3.1}$$

If $i \in I^{\mathrm{re}}$,

$$\begin{aligned} \widetilde{e}_i(b_1 \otimes b_2) &= \begin{cases} \widetilde{e}_i b_1 \otimes b_2 & \text{if } \phi_i(b_1) \geq \epsilon_i(b_2), \\ b_1 \otimes \widetilde{e}_i b_2 & \text{if } \phi_i(b_1) < \epsilon_i(b_2), \end{cases} \\ \widetilde{f}_i(b_1 \otimes b_2) &= \begin{cases} \widetilde{f}_i b_1 \otimes b_2 & \text{if } \phi_i(b_1) > \epsilon_i(b_2), \\ b_1 \otimes \widetilde{f}_i b_2 & \text{if } \phi_i(b_1) \leq \epsilon_i(b_2). \end{cases} \end{aligned} \tag{3.2}$$

If $i \in I^{\mathrm{im}}$,

$$\begin{aligned} \widetilde{e}_{il}(b_1 \otimes b_2) &= \begin{cases} \widetilde{e}_{il} b_1 \otimes b_2 & \text{if } \phi_i(b_1) > \epsilon_i(b_2) - l a_{ii}, \\ 0 & \text{if } \epsilon_i(b_2) < \phi_i(b_1) \leq \epsilon_i(b_2) - l a_{ii}, \\ b_1 \otimes \widetilde{e}_{il} b_2 & \text{if } \phi_i(b_1) \leq \epsilon_i(b_2), \end{cases} \\ \widetilde{f}_{il}(b_1 \otimes b_2) &= \begin{cases} \widetilde{f}_{il} b_1 \otimes b_2 & \text{if } \phi_i(b_1) > \epsilon_i(b_2), \\ b_1 \otimes \widetilde{f}_{il} b_2 & \text{if } \phi_i(b_1) \leq \epsilon_i(b_2). \end{cases} \end{aligned} \tag{3.3}$$

Note that if $a_{ii} = 0$, we have $\widetilde{e}_{il}(b_1 \otimes b_2) = \widetilde{e}_{il} b_1 \otimes b_2$ when $\phi_i(b_1) > \epsilon_i(b_2)$, and $\widetilde{e}_{il}(b_1 \otimes b_2) = b_1 \otimes \widetilde{e}_{il} b_2$ when $\phi_i(b_1) \leq \epsilon_i(b_2)$.

Proposition 3.1. The set $B_1 \otimes B_2$ together with the maps wt, ϵ_i, ϕ_i $(i \in I)$, $\widetilde{e}_{il}$, $\widetilde{f}_{il}$ $((i,l) \in I^{\infty})$ is an abstract crystal.

Proof. It is clear that the conditions (i), (ii) and (iii) in Definition 2.1 hold. If $i \in I^{\mathrm{re}}$, it was shown in [Kas93] that all the conditions are satisfied. For the condition (vii), if

$\phi_i(b_1 \otimes b_2) = -\infty$, since

$$\phi_i(b_1 \otimes b_2) = \max(\phi_i(b_1) + \langle h_i, \mathrm{wt}(b_2)\rangle, \phi_i(b_2)),$$

we have $\phi_i(b_1) = \phi_i(b_2) = -\infty$. Thus

$$\widetilde{e}_{il}(b_1 \otimes b_2) = \widetilde{f}_{il}(b_1 \otimes b_2) = 0.$$

From now on, we will assume that $i \in I^{\mathrm{im}}$, $l \geq 1$. In this case, we don't have to check the condition (v). Hence we have only to verify the conditions (iv) and (vi).

Suppose $\widetilde{f}_{il}(b_1 \otimes b_2) = b_1' \otimes b_2'$.

If $\phi_i(b_1) > \epsilon_i(b_2)$, then $\widetilde{f}_{il}(b_1 \otimes b_2) = \widetilde{f}_{il}b_1 \otimes b_2 = b_1' \otimes b_2'$. Thus

$$\phi_i(b_1') = \phi_i(\widetilde{f}_{il}b_1) = \phi_i(b_1) - la_{ii} > \epsilon_i(b_2) - la_{ii} = \epsilon_i(b_2') - la_{ii},$$

which implies $\widetilde{e}_{il}(b_1' \otimes b_2') = \widetilde{e}_{il}b_1' \otimes b_2' = b_1 \otimes b_2$.

If, $\phi_i(b_1) \leq \epsilon_i(b_2)$, then $\widetilde{f}_{il}(b_1 \otimes b_2) = b_1 \otimes \widetilde{f}_{il}b_2 = b_1' \otimes b_2'$. Note that

$$\phi_i(b_1') = \phi_i(b_1) \leq \epsilon_i(b_2) = \epsilon_i(\widetilde{f}_{il}b_2) = \epsilon_i(b_2').$$

Thus $\widetilde{e}_{il}(b_1' \otimes b_2') = b_1' \otimes \widetilde{e}_{il}b_2' = b_1 \otimes b_2$.

Conversely, suppose $\widetilde{e}_{il}(b_1' \otimes b_2') = b_1 \otimes b_2$. Since $\widetilde{e}_{il}(b_1' \otimes b_2') \neq 0$, we don't have to consider the case $\epsilon_i(b_2') < \phi_i(b_1') \leq \epsilon_i(b_2') - la_{ii}$.

If $\phi_i(b_1') > \epsilon_i(b_2') - la_{ii}$, we have $\widetilde{e}_{il}(b_1' \otimes b_2') = \widetilde{e}_{il}b_1' \otimes b_2'$. Note that

$$\phi_i(b_1) = \phi_i(\widetilde{e}_{il}b_1') = \phi_i(b_1') + la_{ii} > \epsilon_i(b_2') = \epsilon_i(b_2).$$

Hence $\phi_i(b_1) > \epsilon_i(b_2)$ and we obtain $\widetilde{f}_{il}(b_1 \otimes b_2) = \widetilde{f}_{il}b_1 \otimes b_2 = b_1' \otimes b_2'$.

If $\phi_i(b_1') \leq \epsilon_i(b_2')$, then $\widetilde{e}_{il}(b_1' \otimes b_2') = b_1' \otimes \widetilde{e}_{il}b_2'$ and we have

$$\phi_i(b_1) = \phi_i(b_1') \leq \epsilon_i(b_2') = \epsilon_i(\widetilde{e}_{il}b_2') = \epsilon_i(b_2),$$

which gives $\widetilde{f}_{il}(b_1 \otimes b_2) = b_1 \otimes \widetilde{f}_{il}b_2 = b_1' \otimes b_2'$. Hence the condition (iv) is verified.

To verify the condition (vi), let $b_1 \otimes b_2 \in B_1 \otimes B_2$ such that $\widetilde{e}_{il}(b_1 \otimes b_2) \neq 0$ for all $l \geq 1$.

If $\phi_i(b_1) > \epsilon_i(b_2) - la_{ii}$, then $\widetilde{e}_{il}(b_1 \otimes b_2) = \widetilde{e}_{il}b_1 \otimes b_2$ and hence we get

$$\begin{aligned}
\epsilon_i(\widetilde{e}_{il}(b_1 \otimes b_2)) &= \epsilon_i(\widetilde{e}_{il}b_1 \otimes b_2) \\
&= \max(\epsilon_i(\widetilde{e}_{il}b_1), \epsilon_i(b_2) - \langle h_i, \mathrm{wt}(\widetilde{e}_{il}b_1)\rangle) \\
&= \max(\epsilon_i(b_1), \epsilon_i(b_2) - \langle h_i, \mathrm{wt}(b_1) + l\alpha_i\rangle) \\
&= \max(\epsilon_i(b_1), \epsilon(b_2) - \langle h_i, \mathrm{wt}(b_1)\rangle - la_{ii}) \\
&= \epsilon_i(b_1) = \epsilon_i(b_1 \otimes b_2),
\end{aligned}$$

$$\begin{aligned}
\phi_i(\widetilde{e}_{il}(b_1 \otimes b_2)) &= \phi_i(\widetilde{e}_{il}b_1 \otimes b_2) \\
&= \max(\phi_i(\widetilde{e}_{il}b_1) + \langle h_i, \mathrm{wt}(b_2)\rangle, \phi_i(b_2)) \\
&= \max(\phi_i(b_1) + la_{ii} + \phi_i(b_2) - \epsilon_i(b_2), \phi_i(b_2)) \\
&= \phi_i(b_1) + la_{ii} + \phi_i(b_2) - \epsilon_i(b_2) = \phi_i(b_1 \otimes b_2) + la_{ii}.
\end{aligned}$$

If $\phi_i(b_1) \le \epsilon_i(b_2)$, then $\widetilde{e}_{il}(b_1 \otimes b_2) = b_1 \otimes \widetilde{e}_{il}b_2$ and hence we get

$$\begin{aligned}
\epsilon_i(\widetilde{e}_{il}(b_1 \otimes b_2)) &= \epsilon_i(b_1 \otimes \widetilde{e}_{il}b_2) \\
&= \max(\epsilon_i(b_1), \epsilon_i(\widetilde{e}_{il}b_2) - \langle h_i, \mathrm{wt}(b_1)\rangle) \\
&= \max(\epsilon_i(b_1), \epsilon_i(b_2) - \langle h_i, \mathrm{wt}(b_1)\rangle) = \epsilon_i(b_1 \otimes b_2), \\
\phi_i(\widetilde{e}_{il}(b_1 \otimes b_2)) &= \phi_i(b_1 \otimes \widetilde{e}_{il}b_2) \\
&= \max(\phi_i(b_1) + \langle h_i, \mathrm{wt}(\widetilde{e}_{il}b_2)\rangle, \phi_i(\widetilde{e}_{il}b_2)) \\
&= \max(\phi_i(b_1) + \langle h_i, \mathrm{wt}(b_2)\rangle + la_{ii}, \phi_i(b_2) + la_{ii}) \\
&= \max(\phi_i(b_1) + \langle h_i, \mathrm{wt}(b_2)\rangle, \phi_i(b_2)) + la_{ii} \\
&= \phi_i(b_1 \otimes b_2) + la_{ii}.
\end{aligned}$$

By a similar argument, we can verify the condition (vi) for the Kashiwara operators $\widetilde{f}_{il}$'s. □

Corollary 3.2. If B_1 and B_2 are normal crystals, then $B_1 \otimes B_2$ is also a normal crystal.

Proof. For $i \in I^{\mathrm{im}}$, it is easy to see that

$$\begin{aligned}\epsilon_i(b_1 \otimes b_2) &= \max(\epsilon_i(b_1), \epsilon_i(b_2) - \langle h_i, \mathrm{wt}(b_1)\rangle) \\ &= \max(0, -\phi_i(b_1)) = 0, \\ \phi_i(b_1 \otimes b_2) &= \max(\phi_i(b_1) + \langle h_i, \mathrm{wt}(b_2)\rangle, \phi_i(b_2)) \\ &= \max(\phi_i(b_1) + \phi_i(b_2), \phi_i(b_2)) \geq 0,\end{aligned}$$

as desired. □

Corollary 3.3. For $\lambda, \mu \in P^+$, there exists a unique strict crystal embedding

$$\Phi_{\lambda,\mu} \colon B(\lambda+\mu) \to B(\lambda) \otimes B(\mu) \ \text{ given by } v_{\lambda+\mu} \mapsto \ v_\lambda \otimes v_\mu. \tag{3.4}$$

Proof. By the tensor product rule, $v_\lambda \otimes v_\mu$ is the only element of weight $\lambda+\mu$ in $B(\lambda) \otimes B(\mu)$. Moreover, it is annihilated by all $\widetilde{e}_{il}$'s. Hence the connected component of $B(\lambda) \otimes B(\mu)$ containing $v_\lambda \otimes v_\mu$ is the full subcrystal of $B(\lambda) \otimes B(\mu)$ which is isomorphic to $B(\lambda+\mu)$. □

In the following proposition, we will show that the tensor product of crystals satisfies the *associativity law.*

Proposition 3.4. Let B_i $(i = 1, 2, 3)$ be abstract crystals. Then there exists a unique crystal isomorphism

$$\Psi \colon (B_1 \otimes B_2) \otimes B_3 \xrightarrow{\sim} B_1 \otimes (B_2 \otimes B_3) \tag{3.5}$$

given by $(b_1 \otimes b_2) \otimes b_3 \mapsto b_1 \otimes (b_2 \otimes b_3)$, where $b_i \in B_i$ $(i = 1, 2, 3)$.

Proof. It is clear that Ψ is a bijection and preserves the function wt. We shall show that Ψ preserves ϵ_i, ϕ_i $(i \in I)$ and commutes with $\widetilde{f}_{il}$ $((i,l) \in I^\infty)$. As we have seen in (2.3), since Ψ is a bijection, we don't have to check the commutativity with $\widetilde{e}_{il}$'s.

Let $b = (b_1 \otimes b_2) \otimes b_3$ and $b' = \Psi(b) = b_1 \otimes (b_2 \otimes b_3)$. Our proof will be divided into the following cases.

Case 1: $\phi_i(b_1 \otimes b_2) > \epsilon_i(b_3)$.

(1) First, assume that $\phi_i(b_1) > \epsilon_i(b_2)$. In this case, we have

$$\phi_i(b_1 \otimes b_2) = \phi_i(b_1) + \phi_i(b_2) - \epsilon_i(b_2) > \epsilon(b_3), \quad \epsilon_i(b_1 \otimes b_2) = \epsilon_i(b_1),$$

which yields

$$\phi_i(b_1) + \phi_i(b_2) > \epsilon_i(b_2) + \epsilon_i(b_3). \tag{3.6}$$

Thus we obtain

$$\begin{aligned} \phi_i(b) &= \phi_i((b_1 \otimes b_2) \otimes b_3) = \phi_i(b_1 \otimes b_2) + \phi_i(b_3) - \epsilon_i(b_3) \\ &= \phi_i(b_1) + \phi_i(b_2) + \phi_i(b_3) - \epsilon_i(b_2) - \epsilon_i(b_3), \\ \epsilon_i(b) &= \epsilon_i((b_1 \otimes b_2) \otimes b_3) = \epsilon_i(b_1 \otimes b_2) = \epsilon_i(b_1), \\ \widetilde{f}_{il}(b) &= \widetilde{f}_{il}((b_1 \otimes b_2) \otimes b_3) = \widetilde{f}_{il}(b_1 \otimes b_2) \otimes b_3 = (\widetilde{f}_{il} b_1 \otimes b_2) \otimes b_3. \end{aligned}$$

On the other hand, to deal with b', we compare $\phi_i(b_1)$ and $\epsilon_i(b_2 \otimes b_3)$ and obtain

$$\begin{aligned} \phi_i(b') &= \phi_i(b_1 \otimes (b_2 \otimes b_3)) \\ &= \phi_i(b_1) + \phi_i(b_2) + \phi_i(b_3) - \epsilon_i(b_2) - \epsilon_i(b_3), \\ \epsilon_i(b') &= \epsilon_i(b_1 \otimes (b_2 \otimes b_3)) = \epsilon_i(b_1), \\ \widetilde{f}_{il}(b') &= \widetilde{f}_{il}(b_1 \otimes (b_2 \otimes b_3)) = \widetilde{f}_{il} b_1 \otimes (b_2 \otimes b_3). \end{aligned}$$

(2) Next, if $\phi_i(b_1) \le \epsilon_i(b_2)$, by a similar calculation, we obtain

$$\begin{aligned} \phi_i(b) &= \phi_i((b_1 \otimes b_2) \otimes b_3) = \phi_i(b_2) + \phi_i(b_3) - \epsilon_i(b_3), \\ \epsilon_i(b) &= \epsilon_i((b_1 \otimes b_2) \otimes b_3) = \epsilon_i(b_1) + \epsilon_i(b_2) - \phi_i(b_1), \\ \widetilde{f}_{il}(b) &= \widetilde{f}_{il}((b_1 \otimes b_2) \otimes b_3) = (b_1 \otimes \widetilde{f}_{il} b_2) \otimes b_3, \end{aligned}$$

and

$$\begin{aligned} \phi_i(b') &= \phi_i(b_1 \otimes (b_2 \otimes b_3)) = \phi_i(b_2) + \phi_i(b_3) - \epsilon_i(b_3), \\ \epsilon_i(b') &= \epsilon_i(b_1 \otimes (b_2 \otimes b_3)) = \epsilon_i(b_1) + \epsilon_i(b_2) - \phi_i(b_1), \\ \widetilde{f}_{il}(b') &= \widetilde{f}_{il}(b_1 \otimes (b_2 \otimes b_3)) = b_1 \otimes (\widetilde{f}_{il} b_2 \otimes b_3). \end{aligned}$$

Case 2: $\phi_i(b_1 \otimes b_2) \le \epsilon_i(b_3)$.

In this case, by comparing $\phi_i(b_1)$ and $\epsilon_i(b_2 \otimes b_3)$, we obtain

$$\begin{aligned}\phi_i(b) &= \phi_i((b_1 \otimes b_2) \otimes b_3) = \phi_i(b_3),\\ \epsilon_i(b) &= \epsilon_i((b_1 \otimes b_2) \otimes b_3) = \epsilon_i(b_1) + \epsilon_i(b_2) + \epsilon_i(b_3) - \phi_i(b_1) - \phi_i(b_2),\\ \widetilde{f}_{il}(b) &= \widetilde{f}_{il}((b_1 \otimes b_2) \otimes b_3) = (b_1 \otimes b_2) \otimes \widetilde{f}_{il} b_3,\end{aligned}$$

and

$$\begin{aligned}\phi_i(b') &= \phi_i(b_1 \otimes (b_2 \otimes b_3)) = \phi_i(b_3),\\ \epsilon_i(b') &= \epsilon_i(b_1 \otimes (b_2 \otimes b_3)) = \epsilon_i(b_1) + \epsilon_i(b_2) + \epsilon_i(b_3) - \phi_i(b_1) - \phi_i(b_2),\\ \widetilde{f}_{il}(b') &= \widetilde{f}_{il}(b_1 \otimes (b_2 \otimes b_3)) = b_1 \otimes (b_2 \otimes \widetilde{f}_{il} b_3).\end{aligned}$$

Thus we have proved all of our assertions. □

4. CRYSTAL EMBEDDING THEOREM

In this section, we prove one of the main results in this paper, the *crystal embedding theorem* for quantum Borcherds-Bozec algebras.

Theorem 4.1. For $i \in I$, there is a unique strict crystal embedding

$$\Psi_i : B(\infty) \hookrightarrow B(\infty) \otimes B_i \ \text{ given by } \ \mathbf{1} \mapsto \mathbf{1} \otimes (0)_i.$$

Proof. Since there is only one vector of weight 0 in $B(\infty) \otimes B_i$, which is $\mathbf{1} \otimes (0)_i$, Ψ_i should send $\mathbf{1}$ to $\mathbf{1} \otimes (0)_i$ if Ψ_i exists, because it is a crystal morphism.

Let $b = \widetilde{f}_{i_1,l_1} \cdots \widetilde{f}_{i_r,l_r} \mathbf{1} \in B(\infty)$. Choose $\lambda \in P^+$ such that

(i) $\langle h_j, \lambda \rangle \gg 0$ for all $j \in I$,

(ii) $b \in \mathrm{Im}\pi_\lambda$,

where $\pi_\lambda : B(\lambda) \to B(\infty)$ is the injective map given in Proposition 2.9.

Set $b_\lambda = \widetilde{f}_{i_1,l_1} \cdots \widetilde{f}_{i_r,l_r} v_\lambda \in B(\lambda)$ so that $\pi_\lambda(b_\lambda) = b \in B(\infty)$.

Let $l = \langle h_i, \lambda \rangle$ and $\mu = \lambda - l\Lambda_i \in P^+$. Then $\langle h_i, \mu \rangle = 0$ and by Corollary 3.3 there exists a unique strict crystal embedding

$$\Phi_{\mu,l\Lambda_i} : B(\lambda) \to B(\mu) \otimes B(l\Lambda_i) \ \text{ given by } \ v_\lambda \mapsto v_\mu \otimes v_{l\Lambda_i}.$$

We will show that

(1) $\Phi_{\mu,l\Lambda_i}(b_\lambda) = b' \otimes \widetilde{f}_{i,\mathbf{c}} v_{l\Lambda_i}$ for some $b' \in B(\mu)$, $\mathbf{c} \in B_i$,
(2) $\pi_\mu(b') \otimes \mathbf{c} \in B(\infty) \otimes B_i$ does not depend on the choice of $\lambda \gg 0$.

Here, if $i \in I^{\mathrm{re}}$, then $\mathbf{c} \in B_i$ is a non-negative integer c and we understand $\widetilde{f}_{i,\mathbf{c}} = \widetilde{f}_i^c$.

Once our claims are proved, we will get a well-defined map

$$\Psi_i : B(\infty) \to B(\infty) \otimes B_i \ \text{ given by } \ b \mapsto \pi_\mu(b') \otimes \mathbf{c}.$$

We will prove our assertions by induction on $r \geq 0$. When $r = 0$, our assertion is obvious. Assume that $r > 0$ and our assertion is true for $r - 1$.

Set $b_1 = \widetilde{f}_{i_2,l_2} \cdots \widetilde{f}_{i_r,l_r} v_\lambda$. By our induction hypothesis, we have

(1a) $\Phi_{\mu,l\Lambda_i}(b_1) = b_1' \otimes \widetilde{f}_{i,\mathbf{c}'} v_{l\Lambda_i}$ for some $b_1' \in B(\mu)$, $\mathbf{c}' \in B_i$,
(2a) $\pi_\mu(b_1') \otimes \mathbf{c}' \in B(\infty) \otimes B_i$ does not depend on the choice of λ.

Therefore it suffices to show that

(1b) $\widetilde{f}_{i_1,l_1}(b_1' \otimes \widetilde{f}_{i,\mathbf{c}'} v_{l\Lambda_i}) = b' \otimes \widetilde{f}_{i,\mathbf{c}} v_{l\Lambda_i}$ for some $b' \in B(\mu)$, $\mathbf{c} \in B_i$.
(2b) $\widetilde{f}_{i_1,l_1}(\pi_\mu(b_1') \otimes \mathbf{c}') = \pi_\mu(b') \otimes \mathbf{c}$.

If $i_1 = i$, then $\langle h_i, \mu \rangle = 0$ and hence

$$\phi_i(b_1') = \phi_i(\pi_\mu(b_1')),$$
$$\epsilon_i(\widetilde{f}_{i,\mathbf{c}'} v_{l\Lambda_i}) = \epsilon_i(\mathbf{c}') = \begin{cases} c' & \text{if } i \in I^{\mathrm{re}}, \\ 0 & \text{if } i \in I^{\mathrm{im}}. \end{cases}$$

Therefore, $\widetilde{f}_{i_1,l_1}$ acts on the 1st component (resp. 2nd component) of (1b) if and only if it acts on the 1st component (resp. 2nd component) of (2b), which proves our claim.

If $i_1 \neq i$, then $\epsilon_{i_1}(\mathbf{c}') = -\infty$ and

$$\phi_{i_1}(b_1') = \phi_{i_1}(\pi_\mu(b_1')) + \mu(h_{i_1}) \gg 0 = \epsilon_{i_1}(\widetilde{f}_{i,\mathbf{c}'} v_{l\Lambda_i}).$$

Hence $\widetilde{f}_{i_1,l_1}$ acts on the 1st component of (1b) and (2b), which yields our claim.

It is straightforward to verify that Ψ_i is a strict crystal morphism. $\square$

Our next goal is to provide a characterization of the crystals $B(\infty)$ and $B(\lambda)$ ($\lambda \in P^+$). To this end, we introduce an important family of crystals arising from tensor products of elementary crystals.

Let $\mathbf{i} = (i_1, i_2, \ldots)$ be an infinite sequence of indices in I such that every $i \in I$ appears infinitely many times. Set

$$B_{\mathbf{i}} = \{b = \cdots \otimes \mathbf{c}_k \otimes \cdots \otimes \mathbf{c}_1 \mid \mathbf{c}_k \in B_{i_k},\ \mathbf{c}_k = (0)_{i_k} \text{ for } k \gg 0\}. \tag{4.1}$$

Using the tensor product rule, there is a crystal structure on $B_{\mathbf{i}}$ defined as follows. (See [JKKS07] for a more rigorous and detailed treatment.)

Let $b = \cdots \otimes \mathbf{c}_k \otimes \cdots \otimes \mathbf{c}_1$, where $\mathbf{c}_k \in B_{i_k}$. Then we define

$$\begin{aligned}
\mathrm{wt}(b) &= -\sum_{k \geq 1} |\mathbf{c}_k|\, \alpha_{i_k}, \\
\epsilon_i(b) &= \begin{cases} \max\{c_k + \sum_{p>k} c_p\, a_{i,i_p} \mid k \geq 1,\ i_k = i\} & \text{if } i \in I^{\mathrm{re}}, \\ 0 & \text{if } i \in I^{\mathrm{im}}, \end{cases} \\
\phi_i(b) &= \begin{cases} \max\{-c_k - \sum_{1 \leq p<k} c_p\, a_{i,i_p} \mid k \geq 1, i_k = i\} & \text{if } i \in I^{\mathrm{re}}, \\ -\sum_{k \geq 1} |\mathbf{c}_k| a_{i,i_k} & \text{if } i \in I^{\mathrm{im}}. \end{cases}
\end{aligned}$$

To define the Kashiwara operators, we first assume $i \in I^{\mathrm{re}}$ and let s (resp. t) be the largest (resp. smallest) integer $k \geq 1$ such that

(i) $i_k = i$,
(ii) $c_k + \sum_{p>k} c_p\, a_{i,i_p} = \epsilon_i(b)$.

Then we define

$$\begin{aligned}
\widetilde{e}_i(b) &= \begin{cases} \cdots \otimes \mathbf{c}_{s+1} \otimes (c_s - 1) \otimes \mathbf{c}_{s-1} \otimes \cdots \otimes \mathbf{c}_1 & \text{if } \epsilon_i(b) > 0, \\ 0 & \text{otherwise}, \end{cases} \\
\widetilde{f}_i(b) &= \cdots \otimes \mathbf{c}_{t+1} \otimes (c_t + 1) \otimes \mathbf{c}_{t-1} \otimes \cdots \otimes \mathbf{c}_1.
\end{aligned}$$

Suppose $i \in I^{\mathrm{im}}$ and let r be the smallest integer $k \geq 1$ such that

(i) $i_k = i$,

(ii) $\sum_{p>k} |\mathbf{c}_p|\, a_{i,i_p} = 0$.

Then we define

$$\widetilde{f}_{i,l}(b) = \cdots \otimes \mathbf{c}_{r+1} \otimes (l, \mathbf{c}_r) \otimes \mathbf{c}_{r-1} \otimes \cdots \otimes \mathbf{c}_1.$$

Assume further that

(i) the 1st component of $\mathbf{c}_r$ is l or $\mathbf{c}_r$ is a partition having l as a part,

(ii) $\sum_{s<p\leq r} |\mathbf{c}_p|\, a_{i,i_p} < l a_{ii}$ for any s with $i_s = i$ and $1 \leq s < r$.

In this case, we define

$$\widetilde{e}_{i,l}(b) = \cdots \otimes \mathbf{c}_{r+1} \otimes (\mathbf{c}_r \setminus l) \otimes \mathbf{c}_{r-1} \otimes \cdots \otimes \mathbf{c}_1.$$

Otherwise, we define $\widetilde{e}_{il}(b) = 0$.

For each $N \geq 1$, along the sequence $\mathbf{i} = (i_1, i_2, \ldots)$, we apply the crystal embedding theorem repeatedly to get a strict crystal embedding

$$\Psi^{(N)} \colon B(\infty) \hookrightarrow B(\infty) \otimes B_{i_1} \hookrightarrow B(\infty) \otimes B_{i_2} \otimes B_{i_1} \hookrightarrow \cdots \hookrightarrow B(\infty) \otimes B_{i_N} \otimes \cdots \otimes B_{i_1}.$$

For each $b \in B(\infty)$, it is easy to see that there exists some $N \geq 1$ satisfying

$$\Psi^{(N)}(b) = \mathbf{1} \otimes \mathbf{c}_N \otimes \cdots \otimes \mathbf{c}_1 \in B(\infty) \otimes B_{i_N} \otimes \cdots \otimes B_{i_1}.$$

Hence to each $b \in B(\infty)$, one can associate a unique element

$$b_{\mathbf{i}} = \cdots \otimes (0)_{i_{N+1}} \otimes \mathbf{c}_N \otimes \cdots \otimes \mathbf{c}_1 \in B_{\mathbf{i}},$$

which yields a strict crystal embedding

$$\Psi_{\mathbf{i}} \colon B(\infty) \hookrightarrow B_{\mathbf{i}}.$$

Therefore $B(\infty)$ is isomorphic to the connected component of $B_{\mathbf{i}}$ containing the element $(0)_{\mathbf{i}} = \cdots \otimes (0)_{i_N} \otimes \cdots \otimes (0)_{i_1}$. In particular, $\mathbf{1}$ is mapped onto $(0)_{\mathbf{i}}$.

We will now give a characterization of $B(\infty)$ as an application of the crystal embedding theorem.

Theorem 4.2. Let B be a crystal satisfying the following conditions.

(i) $\mathrm{wt}(B) \subset Q^-$,

(ii) there exists an element $b_0 \in B$ such that $\mathrm{wt}(b_0) = 0$,

(iii) for any $b \neq b_0$, there is some $(i,l) \in I^\infty$ such that $\widetilde{e}_{il}\, b \neq 0$,
(iv) for each $i \in I$, there exists a strict crystal embedding $\Psi_i \colon B \hookrightarrow B \otimes B_i$.

Then there exists a crystal isomorphism

$$B \xrightarrow{\sim} B(\infty) \;\text{ given by }\; b_0 \mapsto \mathbf{1}.$$

Proof. Note that for any element $b \in B$ with $\mathrm{wt}(b) = 0$, we have $\widetilde{e}_{il}(b) = 0$ for all $(i,l) \in I^\infty$, for otherwise, $\mathrm{wt}(\widetilde{e}_{il}\, b) = l\alpha_i \notin Q^-$. Hence by the condition (iii), if $\mathrm{wt}(b) = 0$, then $b = b_0$, which implies b_0 is the only element in B such that $\mathrm{wt}(b) = 0$. It follows that the crystal embedding Ψ_i maps b_0 to $b_0 \otimes (0)_i$.

Take an infinite sequence $\mathbf{i} = (i_1, i_2, \ldots)$ such that every $i \in I$ appears infinitely many times. Then for each $N \geq 1$, we obtain a strict embedding

$$\Psi^{(N)} \colon B \hookrightarrow B \otimes B_{i_1} \hookrightarrow B \otimes B_{i_2} \otimes B_{i_1} \hookrightarrow \cdots \hookrightarrow B \otimes B_{i_N} \otimes \cdots \otimes B_{i_1}$$

sending b_0 to $b_0 \otimes (0)_{i_N} \otimes \cdots \otimes (0)_{i_1}$.

As we have seen in the discussion above the theorem, for each $b \in B$, there exists $N \geq 1$ such that

$$\Psi^{(N)}(b) = b_0 \otimes \mathbf{c}_N \otimes \cdots \otimes \mathbf{c}_1,$$

which yields a strict crystal embedding

$$B \hookrightarrow B(\mathbf{i}) \;\text{ given by }\; b \mapsto \cdots \otimes (0)_{i_{N+1}} \otimes \mathbf{c}_N \otimes \cdots \otimes \mathbf{c}_1 \,(\mathbf{c}_k \in B_{i_k}).$$

Thus B is isomorphic to the connected component of $B_{\mathbf{i}}$ containing the element $(0)_{\mathbf{i}} = \cdots \otimes (0)_{i_N} \otimes \cdots \otimes (0)_{i_1}$, which implies B is isomorphic to $B(\infty)$ and b_0 is mapped to $\mathbf{1}$. $\square$

Let us turn to the crystal $B(\lambda)$ ($\lambda \in P^+$). The properties of the crystals $T_\lambda = \{t_\lambda\}$ and $C = \{c\}$ introduced in Example 2.8 will play an important role in our characterization of $B(\lambda)$.

For $\lambda \in P^+$, consider the injective map

$$\psi_\lambda \colon B(\lambda) \to B(\infty) \otimes T_\lambda \;\text{ given by }\; b \mapsto \pi_\lambda(b) \otimes t_\lambda,$$

where $\pi_\lambda\colon B(\lambda) \to B(\infty)$ is the injective map given in Proposition 2.9. One can immediately see that ψ_λ preserves the function wt. Since $\epsilon_i(t_\lambda) = \phi_i(t_\lambda) = -\infty$, for $b \in B(\lambda)$, we have

$$\begin{aligned} \epsilon_i(b) &= \epsilon_i(\pi_\lambda(b)) = \epsilon_i(\pi_\lambda(b) \otimes t_\lambda), \\ \phi_i(b) &= \epsilon_i(b) + \langle h_i, \mathrm{wt}(b)\rangle \\ &= \epsilon_i(\pi_\lambda(b) \otimes t_\lambda) + \langle h_i, \mathrm{wt}(\pi_\lambda(b) \otimes t_\lambda)\rangle \\ &= \phi_i(\pi_\lambda(b) \otimes t_\lambda). \end{aligned}$$

Hence ψ_λ preserves ϵ_i, ϕ_i $(i \in I)$.

If $b \in B(\lambda)$ and $\widetilde{f}_{il}\, b \in B(\lambda)$, Proposition 2.9 yields

$$\widetilde{f}_{il}(\psi_\lambda(b)) = \widetilde{f}_{il}(\pi_\lambda(b) \otimes t_\lambda) = \widetilde{f}_{il}(\pi_\lambda(b)) \otimes t_\lambda = \pi_\lambda(\widetilde{f}_{il}\, b) \otimes t_\lambda = \psi_\lambda(\widetilde{f}_{il}\, b).$$

Moreover,

$$\widetilde{e}_{il}(\psi_\lambda(b)) = \widetilde{e}_{il}(\pi_\lambda(b) \otimes t_\lambda) = \widetilde{e}_{il}(\pi_\lambda(b)) \otimes t_\lambda = \pi_\lambda(\widetilde{e}_{il}\, b) \otimes t_\lambda = \psi_\lambda(\widetilde{e}_{il}\, b).$$

Thus ψ_λ is a crystal morphism commuting with all $\widetilde{e}_{il}$'s.

Consider the injective map

$$\iota_\lambda\colon B(\lambda) \to B(\infty) \otimes T_\lambda \otimes C \quad \text{given by } b \mapsto \psi_\lambda(b) \otimes c.$$

For $b \in B(\lambda)$, we have

$$\begin{aligned} \mathrm{wt}(\iota_\lambda(b)) &= \mathrm{wt}(\psi_\lambda(b)) + \mathrm{wt}(c) = \mathrm{wt}(b), \\ \phi_i(\iota_\lambda(b)) &= \max(\phi_i(\psi_\lambda(b)) + \langle h_i, \mathrm{wt}(c)\rangle, \phi_i(c)) \\ &= \max(\phi_i(b), 0) = \phi_i(b), \\ \epsilon_i(\iota_\lambda(b)) &= \phi_i(\iota_\lambda(b)) - \langle h_i, \mathrm{wt}(\iota_\lambda(b))\rangle \\ &= \phi_i(b) - \langle h_i, \mathrm{wt}(b)\rangle = \epsilon_i(b). \end{aligned}$$

Furthermore, since $b \in B(\lambda)$, we have $\phi_i(\psi_\lambda(b)) = \phi_i(b) \geq 0 = \epsilon_i(c)$. Thus if $i \in I^{\mathrm{re}}$, then

$$\begin{aligned}
\widetilde{e}_i(\iota_\lambda(b)) &= \widetilde{e}_i(\psi_\lambda(b) \otimes c) = \widetilde{e}_i(\psi_\lambda(b)) \otimes c \\
&= \psi_\lambda(\widetilde{e}_i\, b) \otimes c = \iota_\lambda(\widetilde{e}_i\, b), \\
\widetilde{f}_i(\iota_\lambda(b)) &= \widetilde{f}_i(\psi_\lambda(b) \otimes c) \\
&= \begin{cases} \widetilde{f}_i \psi_\lambda(b) \otimes c = \psi_\lambda(\widetilde{f}_i\, b) \otimes c & \text{if } \phi_i(b) > 0, \\ 0 & \text{if } \phi_i(b) = 0, \end{cases} \\
&= \iota_\lambda(\widetilde{f}_i\, b).
\end{aligned}$$

If $i \in I^{\mathrm{im}}$ and $\phi_i(b) = \phi_i(\psi_\lambda(b)) > -l a_{ii}$, then we have

$$\widetilde{e}_{il}(\iota_\lambda(b)) = \widetilde{e}_{il}(\psi_\lambda(b) \otimes c) = \widetilde{e}_{il}(\psi_\lambda(b)) \otimes c = \psi_\lambda(\widetilde{e}_{il}\, b) \otimes c = \iota_\lambda(\widetilde{e}_{il}\, b).$$

If $i \in I^{\mathrm{im}}$ and $0 < \phi_i(b) = \phi_i(\psi_\lambda(b)) \leq -l a_{ii}$, then

$$\widetilde{e}_{il}(\iota_\lambda(b)) = \widetilde{e}_{il}(\psi_\lambda(b) \otimes c) = 0.$$

On the other hand, since $\epsilon_i(b) = 0$, our condition implies

$$0 < \phi_i(b) = \langle h_i, \mathrm{wt}(b) \rangle \leq -l a_{ii}.$$

By the definition of category $\mathcal{O}_{\mathrm{int}}$ ([KK19b, Definition 5.1]), we have $\widetilde{e}_{il}(b) = 0$, which implies

$$\widetilde{e}_{il}(\iota_\lambda(b)) = \iota_\lambda(\widetilde{e}_{il}\, b) = 0$$

as desired.

If $i \in I^{\mathrm{im}}$ and $\phi_i(b) = \phi_i(\psi_\lambda(b)) = 0$, then $\langle h_i, \mathrm{wt}(b) \rangle = 0$, which implies $\widetilde{e}_{il}(b) = 0$ and

$$\widetilde{e}_{il}(\iota_\lambda(b)) = \widetilde{e}_{il}(\psi_\lambda(b) \otimes c) = \psi_\lambda(b) \otimes \widetilde{e}_{il}(c) = 0 = \psi_\lambda(\widetilde{e}_{il}(b)) \otimes c = \iota_\lambda(\widetilde{e}_{il}(b)).$$

For the operators $\widetilde{f}_{il}$, if $i \in I^{\mathrm{im}}$ and $\phi_i(b) = \phi_i(\psi_\lambda(b)) > 0$, then

$$\widetilde{f}_{il}(\iota_\lambda(b)) = \widetilde{f}_{il}(\psi_\lambda(b) \otimes c) = \widetilde{f}_{il}(\psi_\lambda(b)) \otimes c = \psi_\lambda(\widetilde{f}_{il}(b)) \otimes c = \iota_\lambda(\widetilde{f}_{il}\, b).$$

If $i \in I^{\text{im}}$ and $\phi_i(b) = 0$, then $\widetilde{f}_{il}(b) = 0$ and hence

$$\widetilde{f}_{il}(\iota_\lambda(b)) = \widetilde{f}_{il}(\psi_\lambda(b) \otimes c) = \psi_\lambda(b) \otimes \widetilde{f}_{il}\, c = 0 = \iota_\lambda(\widetilde{f}_{il}\, b)$$

as desired.

Therefore, $\iota_\lambda \colon B(\lambda) \to B(\infty) \otimes T_\lambda \otimes C$ is a strict crystal embedding and obtain the following characterization of $B(\lambda)$.

Theorem 4.3. Let $\lambda \in P^+$ be a dominant integral weight. Then the crystal $B(\lambda)$ is isomorphic to the connected component of $B(\infty) \otimes T_\lambda \otimes C$ containing $\mathbf{1} \otimes t_\lambda \otimes c$.

Remark 4.4. (a) As was mentioned at the end of [B16], our characterization can be applied to reproduce Bozec's geometric realization of $B(\infty)$ and $B(\lambda)$ (cf. [KKS09, KKS12]).

(b) Our constuction of $B(\infty)$ and $B(\lambda)$ can also be used directly to the theory of (dual) perfect bases for quantum Borcherds-Bozec algebras.

(c) Our theory will play a crucial role in the categorification of quantum Borcherds-Bozec algebras and their highest weight modules via Khovanov-Lauda-Rouquier algebras and their cyclotomic quotients.

References

[Bor88] R. Borcherds, *Generalized Kac-Moody algebras*, J. Algebra **115** (1988), 501–512.

[B15] T. Bozec, *Quivers with loops and perverse sheaves*, Math. Ann. **362** (2015), 773-797.

[B16] T. Bozec, *Quivers with loops and generalized crystals*, Compositio Math. **152** (2016), 1999-2040.

[BSV16] T. Bozec, O. Schiffmann and E. Vasserot, *On the number of points of nilpotent quiver varieties over finite fields*, arXiv:1701.01797.

[FKKT20] Z. Fan, S.-J. Kang, Y. R. Kim and B. Tong, *Classical limit of quantum Borcherds-Bozec algebras*, J. Pure Appl. Algebra, **225** (2021), 106502.

[GL93] I. Grojnowski and G. Lusztig, *A comparison of bases of quantized enveloping algebras*, Contemporary Math., **153** (1993), 11-19.

[HK02] J. Hong and S.-J. Kang, *Introduction to Quantum Groups and Crystal Bases*, Graduate Studies in Mathematics **42**, Amer. Math. Soc., 2002.

[JKK05] K. Jeong, S.-J. Kang and M. Kashiwara, *Crystal Bases for quantum generalized Kac-Moody algebras*, Proc. Lond. Math. Soc. **(3) 90** (2005), 395-438.

[JKKS07] K. Jeong, S.-J. Kang, M. Kashiwara and D.-U. Shin, *Abstract crystals for quantum generalized Kac-Moody algebras*, Int. Math. Res. Not. IMRN (2007), no.1. 1-19.

[J95] A. Joseph, *Quantum groups and their primitive ideals*, Springer-Verlag, Berlin, 1995

[Kac68] V. G. Kac, *Simple irreducible graded Lie algebras of finite growth*, Math. USSR. Izv. **2** (1968), 1271–1311.

[K19] S.-J. Kang, *Borcherds-Bozec algebras, root multiplicities and the Schofield construction*, Communications in Contemporary Mathematics **21** (2019), no.3.

[K95] S.-J. Kang, *Quantum deformations of generalized Kac-Moody algebras and their modules*, J. Algebra **(3) 175** (1995), 1041-1066

[KKS09] S.-J. Kang, M. Kashiwara and O. Schiffmann, *Geometric construction of crystal bases for quantum generalized Kac-Moody algebras*, Adv. Math.**222** (2009), 996-1015.

[KKS12] S.-J. Kang, M. Kashiwara and O. Schiffmann, *Geometric construction of highest weight crystals for quantum generalized Kac-Moody algebras*, Math. Ann. **354** (2012), 193–208.

[KK19b] S.-J. Kang and Y. R. Kim, *Quantum Borcherds-Bozec algebras and their integrable representations*, J. Pure Appl. Algebra **224** (2020), 106388.

[Kas90] M. Kashiwara, *Crystallizing the q-analogue of universal enveloping algebras*, Commun. Math. Phys. **133** (1990), 249–260.

[Kas91] M. Kashiwara, *On crystal bases of the q-analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), 465-516.

[Kas93] M. Kashiwara, *The crystal base and Littlemann's refined Demazure character formula*, Duke Math. J. **71** (1993), 839-858.

[KS97] M. Kashiwara and Y. Saito, *Geometric construction of crystal bases*, Duke Math. J. **89** (1997), 9-36.

[L90] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), 447-498.

[L91] G. Lusztig, *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), no. 2, 365-421.

[L10] G. Lusztig, *Introduction to quantum groups*, Modern Birkhäuser Classics, Birkhäuser/Springer, New York, 2010

[Moody68] R. V. Moody, *A new class of Lie algebras*, J. Algebra **10** (1968), 211–230.

[NZ97] T. Nakashima and A. Zelevinsky, *Polyhedral realizations of crystal bases for quantized Kac-Moody algebra*, Adv. Math. **131** (1997), 253-278.

[S02] Y. Saito, *Crystal bases and quiver varieties*, Math. Ann. **324** (2002), 675–688.

HARBIN ENGINEERING UNIVERSITY, HARBIN, CHINA
Email address: `fanz@ksu.edu`

KOREA RESEARCH INSTITUTE OF ARTS AND MATHEMATICS, ASAN-SI, CHUNGCHEONGNAM-DO, 31551, KOREA
Email address: `soccerkang@hotmail.com`

GRADUATE SCHOOL OF EDUCATION, HANKUK UNIVERSITY OF FOREIGN STUDIES, SEOUL, 02450, KOREA
Email address: `rocky777@hufs.ac.kr`

HARBIN ENGINEERING UNIVERSITY, HARBIN, CHINA
Email address: `tbl_2019@hrbeu.edu.cn`